

## About a family of Naturally Graded no $p$ -filiform Lie algebras <sup>†</sup>

L.M. CAMACHO, J.R. GÓMEZ, A.J. GONZÁLEZ

*Departamento de Matemática Aplicada I, Universidad de Sevilla  
Avda. Reina Mercedes s/n, 41012 Sevilla, Spain  
e-mail: lcamacho@us.es, jrgomez@us.es, agonzalez@us.es*

(Presented by Santos González)

AMS Subject Class. (2000): 17B45, 17B70

Received December 20, 2004

### 1. INTRODUCTION

The knowledge of naturally graded Lie algebras of a particular Lie algebras class gives a valuable information about the structure of the rest of algebras of that class. In 1970, Vergne [9] obtained the classification in finite arbitrary dimension,  $n$ , for the case of filiform (nilindex  $n - 1$ ). In [8, 7] Goze and Khakimdjanov gave the geometric description of the characteristically nilpotent filiform Lie algebras using the naturally graded filiform Lie algebras. In [6] Gómez and Jiménez-Merchán, obtained the classification in finite arbitrary dimension for the case 2-filiform (nilindex  $n - 2$ ). There are two subcases for the nilindex  $n - 3$ : 3-filiform Lie algebras and the Lie algebras with characteristic sequence  $(n - 3, 2, 1)$ . In [4, 5], Cabezas, Gómez and Pastor gave the classification of naturally graded  $p$ -filiform Lie algebras.

Consistently, for nilindex  $n - 3$ , only rest to study the case of characteristic sequence  $(n - 3, 2, 1)$ . In this work we offer the classification in arbitrary finite dimension of the family of naturally graded Lie algebras  $\mathfrak{g}$  with the above characteristic sequence such that the dimension of the derived ideal is minimum, that is, with  $\dim[\mathfrak{g}, \mathfrak{g}] = n - 3$ .

The two first acceptable dimensions are 5 and 6, but the general situation occurs only for  $n \geq 7$ .

---

<sup>†</sup> This paper has been partially supported by the PAICYT, of Junta de Andalucía (Spain), and by the Ministerio de Ciencia y Tecnología (Spain), ref. BFM 2000-1047

## 2. PRELIMINARIES

The *descending central sequence* of a Lie algebra  $\mathfrak{g}$  is defined by  $(\mathcal{C}^i(\mathfrak{g}))$ ,  $i \in \mathbb{N} \cup \{0\}$ , where  $\mathcal{C}^0(\mathfrak{g}) = \mathfrak{g}$  and  $\mathcal{C}^i(\mathfrak{g}) = [\mathfrak{g}, \mathcal{C}^{i-1}(\mathfrak{g})]$ .

A Lie algebra  $\mathfrak{g}$  is called *nilpotent* if there exists  $k \in \mathbb{N}$  such that  $\mathcal{C}^k(\mathfrak{g}) = \{0\}$ . The smallest integer verifying this equation is called the *nilindex* of  $\mathfrak{g}$ .

A Lie algebra  $\mathfrak{g}$ , with  $\dim(\mathfrak{g}) = n$ , is called *filiform* (or *1-filiform*) if it verifies  $\dim(\mathcal{C}^i(\mathfrak{g})) = n - i - 1$  for  $1 \leq i \leq n - 1$ . These algebras have maximal nilindex  $n - 1$ . The Lie algebras with a nilindex  $n - 2$  are called *quasifiliform* (or *2-filiform*) and those whose nilindex is 1 are called *abelian*.

Let  $\mathfrak{g}$  be a nilpotent Lie algebra of dimension  $n$ .

For all  $X \in \mathfrak{g} - [\mathfrak{g}, \mathfrak{g}]$ ,  $c(X) = (c_1(X), c_2(X), \dots, 1)$  is the sequence, in decreasing order, of the dimensions of the characteristic subspaces of the *nilpotent operator*  $\text{ad}(X)$ , where the adjoint operator of an element  $X \in \mathfrak{g}$ ,  $\text{ad}(X)$ , is defined by

$$\begin{aligned} \text{ad}(X) : \mathfrak{g} &\rightarrow \mathfrak{g} \\ Y &\mapsto [X, Y]. \end{aligned}$$

The finite sequence  $c(\mathfrak{g}) = \sup\{c(X) : X \in \mathfrak{g} - [\mathfrak{g}, \mathfrak{g}]\}$  is called the *characteristic sequence* or *Goze invariant* of the nilpotent Lie algebra  $\mathfrak{g}$ . The filiform, quasifiliform and abelian Lie algebras of dimension  $n$  have as their Goze invariant  $(n - 1, 1)$ ,  $(n - 2, 1, 1)$  and  $(1, 1, \dots, 1)$ , respectively. The Lie algebras with characteristic sequence  $(n - p, 1, \dots, 1)$  are known as *p-filiform* Lie algebras [3]. We know the classification of *p-filiform* for the integer values of  $p$  between  $n - 5$  and  $n - 2$  ([2, 1]). Remark that, for nilindex  $n - 3$ , there are two families with Goze invariant  $(n - 3, 1, 1, 1)$  and  $(n - 3, 2, 1)$  respectively.

Note that a complex Lie algebra  $\mathfrak{g}$  is naturally filtered by the descending central sequence. This result leads to associate any Lie algebra  $\mathfrak{g}$  with a graded Lie algebra,  $\text{gr } \mathfrak{g}$  with equal nilindex:

$$\text{gr } \mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}_i, \quad \mathfrak{g}_i = \mathcal{C}^{i-1}(\mathfrak{g}) / \mathcal{C}^i(\mathfrak{g}).$$

By nilpotency, the above graduation is finite, that is  $\text{gr } \mathfrak{g} = \mathfrak{g}_1 \oplus \mathfrak{g}_2 \oplus \dots \oplus \mathfrak{g}_k$  with  $[\mathfrak{g}_i, \mathfrak{g}_j] \subset \mathfrak{g}_{i+j}$ , for  $i+j \leq k$ . A Lie algebra  $\mathfrak{g}$  is said to be naturally graded if  $\text{gr } \mathfrak{g}$  is isomorphic to  $\mathfrak{g}$ , what will be denoted henceforth by  $\text{gr } \mathfrak{g} = \mathfrak{g}$ .

Let  $\{X_0, X_1, \dots, X_{n-3}, Y_1, Y_2\}$  be an adapted basis of  $\mathfrak{g}$ . We study the case where the dimension of the derived ideal is minimum, consistently  $\dim[\mathfrak{g}, \mathfrak{g}] = n - 3$ . Thus,  $Y_1$  is not in  $[\mathfrak{g}, \mathfrak{g}]$  and, consequently,  $Y_1 \in \mathfrak{g}_1$ . In general, if we denote as  $r$  to the position of the vector  $Y_1$  into the subspaces of the natural

graduation, we observe that the value of  $r$  is  $r = 1$ . We remark that the position of  $Y_2$  is previously determined because we have that  $[X_0, Y_1] = Y_2$  and that implies  $Y_2 \in \mathfrak{g}_{r+1}$  with  $1 \leq r \leq n - 4$ . Then, in this case  $Y_2 \in \mathfrak{g}_2$ .

From now, Jacobi identity for the vectors  $X, Y, Z$  will be denoted as  $\text{Jac}(X, Y, Z)$  and the laws of the algebras,  $\mathfrak{g}$ , of dimension  $n$  such that  $\dim[\mathfrak{g}, \mathfrak{g}]$  is minimum will be denoted as  $\mu_n$ .

### 3. STRUCTURE THEOREM

In this section, we will obtain a first approximation to the structure of naturally graded Lie algebras with Goze invariant  $(n - 3, 2, 1)$ .

Let  $\mathfrak{g}$  be a naturally graded Lie algebra of Goze's invariant  $(n - 3, 2, 1)$  and let  $\{X_0, X_1, \dots, X_{n-3}, Y_1, Y_2\}$  be an adapted basis of  $\mathfrak{g}$ , that is:

$$\begin{aligned} [X_0, X_i] &= X_{i+1} \quad (1 \leq i \leq n - 4), \\ [X_0, X_{n-3}] &= 0, \\ [X_0, Y_1] &= Y_2, \\ [X_0, Y_2] &= 0, \end{aligned}$$

where  $X_0 \in \mathfrak{g} - [\mathfrak{g}, \mathfrak{g}]$ . That implies

$$\begin{aligned} \mathcal{C}^1(\mathfrak{g}) &\supset \langle X_2, X_3, \dots, X_{n-3}, Y_2 \rangle, \\ \mathcal{C}^i(\mathfrak{g}) &\supset \langle X_{i+1}, X_{i+2}, \dots, X_{n-3} \rangle \quad (2 \leq i \leq n - 4). \end{aligned}$$

LEMMA 3.1. *Let  $\mathfrak{g}$  be a Lie algebra of dimension  $n$  and Goze's invariant  $(n - 3, 2, 1)$  and let  $\{X_0, X_1, \dots, X_{n-3}, Y_1, Y_2\}$  be an adapted basis of  $\mathfrak{g}$ . Then,*

$$X_1 \notin \mathcal{C}^1(\mathfrak{g}), \quad X_{n-3} \in \mathcal{Z}(\mathfrak{g}), \quad Y_1 \notin \mathcal{C}^{n-4}(\mathfrak{g}), \quad Y_2 \notin \mathcal{C}^{n-3}(\mathfrak{g}).$$

*Proof.* Obviously,  $X_{n-3} \in \mathcal{Z}(\mathfrak{g})$ ,  $Y_1 \notin \mathcal{C}^{n-4}(\mathfrak{g})$  and  $Y_2 \notin \mathcal{C}^{n-3}(\mathfrak{g})$  because, otherwise,  $\mathfrak{g}$  could not be of characteristic sequence  $(n - 3, 2, 1)$ . It is easy to prove that  $X_1 \notin [\mathfrak{g}, \mathfrak{g}]$  supposing that  $X_1 \in [Y_1, Y_2]$ , or  $X_1 \in [X_i, Y_j]$ ,  $1 \leq i \leq n - 4$ ,  $1 \leq j \leq 2$ , or  $X_1 \in [X_i, X_j]$ ,  $1 \leq i < j \leq n - 3 - i$ , and obtaining contradiction. ■

*Remark 3.2.* We identify each vector with its class, and we call  $\mu(n, r)$  the family of laws of Lie algebras with Goze invariant  $(n - 3, 2, 1)$  where  $n$  is the dimension and  $r$  is the position of  $Y_1$  in the subsets of the natural gradation. We remark that the position of  $Y_2$  is previously determined because we have that  $[X_0, Y_1] = Y_2$  and that implies  $Y_2 \in \mathfrak{g}_{r+1}$  with  $1 \leq r \leq n - 4$ .

*Remark 3.3.* It is easy to see that  $\mathfrak{g}_1 \supset \langle X_0, X_1 \rangle$  and  $\mathfrak{g}_i \supset \langle X_i \rangle$ ,  $2 \leq i \leq n-3$ .

Now, we obtain the general structure of laws of naturally graded Lie algebras of characteristic sequence  $(n-3, 2, 1)$  in arbitrary dimension. At first, we prove that if  $Y_1 \in \mathfrak{g}_r$ , then  $r$  is odd.

LEMMA 3.4. *If  $r$  is even, the case  $\mu(n, r)$  is not admissible in any dimension.*

*Proof.* Let  $\mathfrak{g}$  be a naturally graded Lie algebra of Goze invariant  $(n-3, 2, 1)$ , let  $\{X_0, X_1, \dots, X_{n-3}, Y_1, Y_2\}$  be an adapted basis of  $\mathfrak{g}$ , and let  $Y_1 \in \mathfrak{g}_r$  be with  $r$  even. It is easy to prove that  $Y_1 \notin [\mathfrak{g}, \mathfrak{g}]$  so  $Y_1 \in \mathfrak{g}_1$  and this is impossible because  $r$  is even. ■

THEOREM 3.5. (STRUCTURE THEOREM) *Any complex naturally graded Lie algebra  $\mathfrak{g}$  of dimension  $n \geq 5$ , with Goze invariant  $(n-3, 2, 1)$  is isomorphic to one whose law can be expressed in an adapted basis  $\{X_0, X_1, \dots, X_{n-3}, Y_1, Y_2\}$  by:*

- If  $r = 1$

$$\begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2, \\ [X_i, X_j] = a_{ij}X_{i+j} & (1 \leq i < j \leq n-3-i). \end{cases}$$

- If  $3 \leq r \leq \frac{n-5}{2}$ ,  $r$  odd

$$\begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2, \\ [X_i, X_j] = a_{ij}X_{i+j} & (i+j \notin \{r, r+1\}, 1 \leq i < j \leq n-3-i), \\ [X_i, X_{r-i}] = a_{i,r-i}X_r + (-1)^{i-1}Y_1 & (1 \leq i \leq \frac{r-1}{2}), \\ [X_i, X_{r+1-i}] = a_{i,r+1-i}X_{r+1} + (-1)^{i-1} \frac{(r+1-2i)}{2} Y_2 & (1 \leq i \leq \frac{r-1}{2}), \\ [X_i, Y_1] = \varepsilon X_{r+i} & (1 \leq i \leq n-3-r), \end{cases}$$

with  $\varepsilon \in \{0, 1\}$ .

- If  $\frac{n-4}{2} \leq r \leq n-4$ ,  $r$  odd

$$\left\{ \begin{array}{l} [X_0, X_i] = X_{i+1} \quad (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2, \\ [X_i, X_j] = a_{ij}X_{i+j} \quad (i+j \notin \{r, r+1\}, 1 \leq i < j \leq n-3-i), \\ [X_i, X_{r-i}] = a_{i,r-i}X_r + (-1)^{i-1}Y_1 \quad (1 \leq i \leq \frac{r-1}{2}), \\ [X_i, X_{r+1-i}] = a_{i,r+1-i}X_{r+1} \\ \quad + (-1)^{i-1} \frac{(r+1-2i)}{2} Y_2 \quad (1 \leq i \leq \frac{r-1}{2}), \\ [X_i, Y_1] = (c_1 - (i-1)c_2)X_{r+i} \quad (1 \leq i \leq n-3-r \leq \frac{n-2}{2}), \\ [X_i, Y_2] = c_2X_{r+1+i} \quad (1 \leq i \leq n-4-r \leq \frac{n-4}{2}), \\ [Y_1, Y_2] = hX_{n-3} \quad (h = 0 \text{ if } r \neq \frac{n-4}{2}), \end{array} \right.$$

with  $c_1, c_2 \in \mathbb{C}$ .

*Proof.* If  $\mathfrak{g}$  is in the condition of theorem, then a first general expression of  $\mathfrak{g}$  is given by:

$$\left\{ \begin{array}{l} [X_0, X_i] = X_{i+1} \quad (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2, \\ [X_i, X_j] = a_{ij}X_{i+j} \quad (i+j \notin \{r, r+1\}, 1 \leq i < j \leq n-3-i), \\ [X_i, X_{r-i}] = a_{i,r-i}X_r + b_{i1}Y_1 \quad (1 \leq i \leq \frac{r-1}{2}), \\ [X_i, X_{r+1-i}] = a_{i,r+1-i}X_{r+1} + b_{i2}Y_2 \quad (1 \leq i \leq \frac{r-1}{2}), \\ [X_1, Y_1] = c_{11}X_{r+1} + dY_2, \\ [X_i, Y_1] = c_{i1}X_{r+i} \quad (2 \leq i \leq n-3-r), \\ [X_i, Y_2] = c_{i2}X_{r+1+i} \quad (1 \leq i \leq n-4-r), \\ [Y_1, Y_2] = hX_{2r+1} \quad (\text{si } r \leq \frac{n-4}{2}). \end{array} \right.$$

Some elementary changes of basis jointly with Jacobi identity implies that:

- If  $1 \leq r \leq \frac{n-5}{2}$  the coefficients can be expressed by

$$c_{i,1} = c_1 \quad (1 \leq i \leq n-3-r) \quad \text{and} \quad c_{i,2} = 0 \quad (1 \leq i \leq n-4-r).$$

- If  $\frac{n-4}{2} \leq r \leq n-4$  the coefficients can be expressed by

$$c_{i,1} = c_1 - (i-1)c_2 \quad (1 \leq i \leq n-3-r) \quad \text{and} \quad c_{i,2} = c_2 \quad (1 \leq i \leq n-4-r).$$

By using Jacobi identity it is posible to obtain that

$$b_{i,2} = (-1)^{(i-1)} \frac{r+1-2i}{2} b_1, \quad 1 \leq i \leq \frac{r-1}{2}.$$

Furthermore,  $b_1 \neq 0$  (in other case  $Y_1 \notin \mathcal{C}^1(\mathfrak{g})$  and then  $Y_1 \notin \mathfrak{g}_r = \langle X_r, Y_1 \rangle$  with  $r \geq 3$ ). Next, an easy change of basis allows to suppose  $b_1 = 1$ . Then,

- If  $3 \leq r \leq \frac{n-5}{2}$ . As  $b_1 \neq 0$ , if  $c_1 \neq 0$  an easy change of basis allows to suppose  $c_1 = 1$ , and consistently  $c_1 \in \{0, 1\}$ .
- If  $r = 1$ , the case must be studied separately. ■

#### 4. DIMENSIONS $n = 5$ AND $n = 6$ .

Even if our main aim is to study the case of dimension  $n$  finite arbitrary, the low dimensional cases are special and we will study them previously. The lowest cases are for dimensions  $n = 5$  and  $n = 6$  and they have a special treatment.

**THEOREM 4.1.** *Any complex naturally graded Lie algebra of dimension 5 with Goze invariant  $(2, 2, 1)$  is isomorphic to one whose law can be expressed in an adapted basis  $\{X_0, X_1, X_2, Y_1, Y_2\}$  by:*

$$\mu_5 : \begin{cases} [X_0, X_1] = X_2, \\ [X_0, Y_1] = Y_2. \end{cases}$$

*Proof.* The proof is trivial. ■

**THEOREM 4.2.** *Any complex naturally graded Lie algebra of dimension 6 with Goze invariant  $(3, 2, 1)$  is isomorphic to one whose law can be expressed in an adapted basis  $\{X_0, X_1, X_2, X_3, Y_1, Y_2\}$  by:*

$$\mu_6^1 : \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq 2), \\ [X_0, Y_1] = Y_2, \end{cases} \quad \mu_6^2 : \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq 2), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_2] = X_3. \end{cases}$$

*Proof.* In dimension six the graduation is

$$\langle X_0, X_1, Y_1 \rangle \oplus \langle X_2, Y_2 \rangle \oplus \langle X_3 \rangle,$$

and by Theorem 3.5 the laws of these algebras are the following:

$$\begin{cases} [X_0, X_1] = X_2, \\ [X_0, X_2] = X_3, \\ [X_0, Y_1] = Y_2, \\ [X_1, X_2] = a_{12}X_3. \end{cases}$$

By using a generic change of basis we prove that nullity of coefficient  $a_{12}$  is an invariant.

- If  $a_{12} \neq 0$ , it is easy to obtain the algebra of law  $\mu_6^2$ .
- If  $a_{12} = 0$ , we obtain the algebra of law  $\mu_6^1$ . ■

### 5. DIMENSION $n \geq 7$ .

Now, we present the classification of the naturally graded Lie algebras with Goze invariant  $(n - 3, 2, 1)$ , dimension  $n \geq 7$  and  $\dim[\mathfrak{g}, \mathfrak{g}]$  minimum, that is, equal to  $n - 3$ . The first expression of this family is given by the following lemma:

LEMMA 5.1. *Let  $\mathfrak{g}$  be a naturally graded Lie algebra with Goze invariant  $(n - 3, 2, 1)$ ,  $\dim(\mathfrak{g}) = n \geq 7$  and  $\dim[\mathfrak{g}, \mathfrak{g}] = n - 3$ . Then, there exists a characteristic vector  $X_0$  and an adapted basis  $\{X_0, X_1, \dots, X_{n-3}, Y_1, Y_2\}$ , which lead us to express the laws of  $\mathfrak{g}$  by:*

$$\mu_n^a : \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n - 4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = aX_{i+1} & (2 \leq i \leq n - 4), \end{cases}$$

if  $n$  is odd, or

$$\mu_n^{a,b} : \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n - 4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = aX_{i+1} & (2 \leq i \leq n - 5), \\ [X_1, X_{n-4}] = (a + b)X_{n-3}, \\ [X_i, X_{n-3-i}] = (-1)^{i+1}bX_{n-3} & (2 \leq i \leq \frac{n-4}{2}), \end{cases}$$

if  $n$  is even.

*Proof.* By using Teorema 3.5 it follows that, in this case ( $r = 1$ ), there exists a characteristic vector  $X_0$  and an adapted basis,  $\{X_0, X_1, \dots, X_{n-3}, Y_1, Y_2\}$ , such that the laws of the algebra are given by

$$\mu_n : \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n - 4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = a_{ij}X_{i+j} & (2 \leq i < j \leq n - 3 - i). \end{cases}$$

Now, we use an inductive procedure on  $n$ .

DIMENSION  $n = 7$ : In dimension seven the graduation is

$$\langle X_0, X_1, Y_1 \rangle \oplus \langle X_2, Y_2 \rangle \oplus \langle X_3 \rangle \oplus \langle X_4 \rangle,$$

and by using the Jacobi identity in the family  $\mu_7$  we obtain  $\mu_7^a$ .

DIMENSION  $n = 8$ : Analogously, by using the Jacobi identity it is easy to obtain that  $\mu_8$  is  $\mu_8^{a,b}$ .

The inductive procedure is realized in function of the parity of the dimension. That is the reason why we study the cases of dimension  $n$  even and  $n$  odd separately.

DIMENSION  $n > 7$ ,  $n$  ODD: If we suppose that the result is true for  $n = k$  even, we will prove it for  $n = k + 1$  odd. If  $k$  is even, we suppose that it is possible to express  $\mu_k$  by

$$\mu_k^{a,b} : \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq k-4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = aX_{i+1} & (2 \leq i \leq k-5), \\ [X_1, X_{n-4}] = (a+b)X_{n-3}, \\ [X_i, X_{n-3-i}] = (-1)^{i+1}bX_{n-3} & (2 \leq i \leq \frac{k-4}{2}). \end{cases}$$

Now, for  $n = k + 1$ , we add the brackets

$$\begin{aligned} [X_0, X_{k-3}] &= \alpha_0 X_{k-2}, \\ [X_i, X_{k-2-i}] &= \alpha_i X_{k-2} \quad (1 \leq i \leq \frac{k-4}{2}), \\ [X_{k-3}, Y_1] &= \beta_1 X_{k-2}, \\ [X_{k-4}, Y_2] &= \beta_2 X_{k-2}. \end{aligned}$$

By using Jacobi identity we prove the result.

DIMENSION  $n > 8$ ,  $n$  EVEN: We suppose that the result is true for  $n = k$  odd and we will prove it for  $n = k + 1$  even. If  $k$  is odd, we suppose that it is possible to express  $\mu_k$  by

$$\mu_k^a : \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq k-4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = aX_{i+1} & (2 \leq i \leq k-4). \end{cases}$$

For  $n = k + 1$  it is necessary to add the same brackets as in the odd case and analogously we obtain the result. ■



## 6. CLASSIFICATION THEOREM

Finally, we give the theorem of classification for naturally graded Lie algebras with Goze invariant  $(n-3, 2, 1)$ ,  $r = 1$  and  $n \geq 7$ .

**THEOREM 6.1.** *Any complex naturally graded Lie algebra of dimension  $n$ ,  $n \geq 7$ , with Goze invariant  $(n-3, 2, 1)$  and laws  $\mu(n)$  is isomorphic to one whose law can be expressed in suitable adapted basis by*

$$\begin{aligned} \mu_{(n-3,2,1)}^1 : & \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2; \end{cases} \\ \mu_{(n-3,2,1)}^2 : & \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2, \\ [X_i, X_{n-3-i}] = (-1)^{i+1} X_{n-3} & (1 \leq i \leq \frac{n-4}{2}); \end{cases} \\ \mu_{(n-3,2,1)}^3 : & \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = X_{i+1} & (2 \leq i \leq n-4); \end{cases} \\ \mu_{(n-3,2,1)}^4 : & \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = X_{i+1} & (2 \leq i \leq n-5), \\ [X_i, X_{n-3-i}] = (-1)^i X_{n-3} & (2 \leq i \leq \frac{n-4}{2}); \end{cases} \\ \mu_{(n-3,2,1)}^5 : & \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = X_{i+1} & (2 \leq i \leq n-5), \\ [X_1, X_{n-4}] = 2X_{n-3}, \\ [X_i, X_{n-3-i}] = (-1)^{i+1} X_{n-3} & (2 \leq i \leq \frac{n-4}{2}). \end{cases} \end{aligned}$$

*Proof.* By using the above lemma we will obtain the result. In function of the dimension of the algebra it is necessary to consider two different cases.

Let  $\mathfrak{g}$  be a naturally graded Lie algebra of dimension  $n$  odd,  $n \geq 7$ , with Goze invariant  $(n-3, 2, 1)$  and laws  $\mu_n$ . Then, the natural graduation is given by

$$\langle X_0, X_1, Y_1 \rangle \oplus \langle X_2, Y_2 \rangle \oplus \langle X_3 \rangle \oplus \cdots \oplus \langle X_{n-3} \rangle.$$

• Case 1:  $n$  even,  $n \geq 8$ . If  $n$  is even the laws of the algebra can be expressed by

$$\mu_n^{a,b} : \begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n-4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = aX_{i+1} & (2 \leq i \leq n-5), \\ [X_1, X_{n-4}] = (a+b)X_{n-3}, \\ [X_i, X_{n-3-i}] = (-1)^{i+1}bX_{n-3}, & (2 \leq i \leq \frac{n-4}{2}). \end{cases}$$

The general change of basis implies three generators,  $X_0$ ,  $X_1$  and  $Y_1$ :

$$\begin{aligned} X'_0 &= \sum_{i=0}^{n-3} P_i X_i + P_{n-2} Y_1 + P_{n-1} Y_2, \\ X'_1 &= \sum_{i=0}^{n-3} Q_i X_i + Q_{n-2} Y_1 + Q_{n-1} Y_2, \\ Y'_1 &= \sum_{i=0}^{n-3} R_i X_i + R_{n-2} Y_1 + R_{n-1} Y_2. \end{aligned}$$

By using the condition of the family we obtain that

$$\begin{cases} Q_0 = 0, \\ R_i = 0 & (0 \leq i \leq n-5). \end{cases}$$

Finally, the admissible changes of basis are

$$\begin{aligned} X'_0 &= P_0 X_0 + P_1 X_1 + P_2 X_2 + \cdots + P_{n-4} X_{n-4} + P_{n-3} X_{n-3} \\ &\quad + P_{n-2} Y_1 + P_{n-1} Y_2, \\ X'_1 &= Q_1 X_1 + Q_2 X_2 + \cdots + Q_{n-4} X_{n-4} + Q_{n-3} X_{n-3} + Q_{n-2} Y_1 + Q_{n-1} Y_2, \\ X'_2 &= P_0 Q_1 X_2 + (P_0 Q_2 + a(P_1 Q_2 - P_2 Q_1)) X_3 + \cdots + (P_0 Q_{n-5} \\ &\quad + a(P_1 Q_{n-5} - P_{n-5} Q_1)) X_{n-4} + (P_0 Q_{n-4} + a(P_1 Q_{n-4} - P_{n-4} Q_1)) \\ &\quad + \sum_{i=1}^{\frac{n-4}{2}} (-1)^{i+1} (P_i Q_{n-3-i} - P_{n-3-i} Q_i) b X_{n-3} + (P_0 Q_{n-2} - P_{n-2} Q_0) Y_2, \\ X'_3 &= P_0 (P_0 + a P_1) Q_1 X_3 + (P_0 + a P_1) (P_0 Q_2 + a(P_1 Q_2 - P_2 Q_1)) X_4 + \cdots \\ &\quad + (P_0 + a P_1) ((P_0 Q_{n-6} + a(P_1 Q_{n-6} - P_{n-6} Q_1)) X_{n-4} \\ &\quad + (P_0 + a P_1) (\dots) X_{n-3}, \\ &\quad \vdots \end{aligned}$$

$$\begin{aligned}
 X'_{n-4} &= P_0(P_0 + aP_1)^{n-6}Q_1X_{n-4} \\
 &\quad + ((P_0 + aP_1)^{n-7}(P_0 + (a + b)P_1)((P_0Q_2 + a(P_1Q_2 - P_2Q_1))X_{n-3}, \\
 X'_{n-3} &= P_0(P_0 + aP_1)^{n-6}Q_1(P_0 + (a + b)P_1)X_{n-3}, \\
 Y'_1 &= R_{n-4}X_{n-4} + R_{n-3}X_{n-3} + R_{n-2}Y_1 + R_{n-1}Y_2, \\
 Y'_2 &= (P_0 + (a + b)P_1)R_{n-4}X_{n-3} + P_0R_{n-2}Y_2,
 \end{aligned}$$

with the following restrictions

$$P_0 \neq 0, \quad Q_1 \neq 0, \quad R_{n-2} \neq 0, \quad P_0 + aP_1 \neq 0, \quad P_0 + (a + b)P_1 \neq 0.$$

The nullity of  $a$  and  $b$  are invariant, because

$$a' = \frac{Q_1a}{P_0 + aP_1} \quad \text{and} \quad b' = \frac{P_0Q_1b}{(P_0 + aP_1)(P_0 + (a + b)P_1)}.$$

Furthermore, we obtain that the nullity of  $a + b$  is invariant, because

$$a' + b' = \frac{Q_1(a + b)}{P_0 + (a + b)P_1}.$$

We consider the following cases:

- Case 2.1:  $a = b = 0$ . Trivially, we obtain  $\mu_{(n-3,2,1)}^1$ .
- Case 2.2:  $a \neq 0$  and  $b = 0$ . By choosing  $P_0$ ,  $Q_1$  and  $P_1$ , we obtain  $\mu_{(n-3,2,1)}^2$ .
- Case 2.3:  $a = 0$  and  $b \neq 0$ . As in the above case, we obtain  $\mu_{(n-3,2,1)}^3$ .
- Case 2.4:  $a \neq 0$ ,  $b \neq 0$  and  $a + b = 0$ . By choosing  $P_0$ ,  $Q_1$  and  $P_1$ , we obtain  $\mu_{(n-3,2,1)}^4$ .
- Case 2.5:  $a \neq 0$ ,  $b \neq 0$  and  $a + b \neq 0$ . It is possible to choose  $P_0$ ,  $Q_1$  and  $P_1$  for to obtain the algebra  $\mu_{(n-3,2,1)}^5$ .

Furthermore, the above results prove that the algebras  $\mu_{(n-3,2,1)}^1, \mu_{(n-3,2,1)}^2, \mu_{(n-3,2,1)}^3, \mu_{(n-3,2,1)}^4$  y  $\mu_{(n-3,2,1)}^5$  are pairwise no isomorphic for  $n$  even.

• Case 2:  $n$  odd,  $n \geq 7$ . As follows from the above lemma we obtain that an algebra of this kind is isomorphic to one whose law can be expressed by

$$\begin{cases} [X_0, X_i] = X_{i+1} & (1 \leq i \leq n - 4), \\ [X_0, Y_1] = Y_2, \\ [X_1, X_i] = aX_{i+1} & (2 \leq i < j \leq n - 3 - i). \end{cases}$$

Since, the odd case is equal to even case considering  $b = 0$ . An analogous treatment of Case 1 proves that the nullity of  $a$  is an invariant and from here, the result is obtained. ■

## REFERENCES

- [1] CABEZAS, J.M., CAMACHO, L.M., GÓMEZ, J.R., NAVARRO, R.M., A class of nilpotent Lie algebras, *Comm. Algebra* **28** (9) (2000), 4489–4499.
- [2] CABEZAS, J.M., GÓMEZ, J.R.,  $(n - 4)$ -filiform Lie algebras, *Comm. Algebra* **27** (10) (1999), 4803–4819.
- [3] CABEZAS, J.M., GÓMEZ, J.R., JIMÉNEZ-MERCHÁN, A., Family of  $p$ -filiform Lie algebras, in “Algebra and Operator Theory”, Khakimjanov, Y., Goze, M., Ayupov, Sh., eds., Kluwer Academic Publishers, Tashkent, 1998, pp. 93–102.
- [4] CABEZAS, J.M., GÓMEZ, J.R., PASTOR, E., Structure theorem for naturally graded 3-filiform Lie algebras, in “I Colloquium on Lie Theory and Applications”, University of Vigo, Vigo, 2000.
- [5] CABEZAS, J.M., PASTOR, E., Naturally graded  $p$ -filiform Lie algebras in arbitrary finite dimension, *J. Lie Theory* **15** (2005), 379–391.
- [6] GÓMEZ, J.R., JIMÉNEZ-MERCHÁN, A., Naturally graded quasifiliform Lie algebras, *J. Algebra* **256** (2002), 211–228.
- [7] GOZE, M., HAKIMJANOV (KHAKIMDJANOV), Y., Sur les algèbres de Lie nilpotentes admettant un tore de dérivations, *Manuscripta Math.* **84** (1994), 115–224.
- [8] GOZE, M., KHAKIMDJANOV, Y., “Nilpotent Lie Algebras”, Kluwer Academic Publishers, 1996.
- [9] VERGNE, M., Cohomologie des algèbres de Lie nilpotentes. Application à l’étude de la variété des algèbres de Lie nilpotentes, *Bull. Soc. Math. France*, **98** (1970), 81–116.