

Pablo J. López-González

A new gorgonian genus from deep-sea Antarctic waters (Octocorallia, Alcyonacea, Plexauridae)

Received: 11 August 2004 / Revised: 19 June 2005 / Accepted: 4 July 2005 / Published online: 25 August 2005
© Springer-Verlag and AWI 2005

Abstract *Mesogligorgia scotiae* gen. nov., sp. nov. is described and illustrated from a colony collected in the Scotia Sea, 2,201–2,213 m in depth, on the ANDEEP-I cruise. The new taxon is placed in the family Plexauridae because of: 1) the presence of a horny axis with a cross-chambered central core and numerous loculi, 2) retractile polyps in calyces with distinct spicular components, and 3) armed polyps with large sclerites with a poorly-developed collaret and eight well-developed points. The irregularly distributed sclerites running along the axis, into a thick mesogloea coenenchyme, and the elongated spindles with irregular ends are the most distinctive characters of the newly proposed genus.

Keywords Cnidaria · Octocorallia · Alcyonacea · Plexauridae · Antarctica · *Mesogligorgia*

Introduction

Among the gorgonian families, Plexauridae is one of the richest in genera and species (Bayer 1956, 1981). This family is especially well represented in temperate and tropical waters (e.g. Bayer 1961; Grasshoff 1992, 1999, 2000; Fabricius and Alderslade 2001), and also in deep waters (e.g. Wright and Studer 1889; Bayer 1956; Grasshoff 1977, 1986). Although plexaurids are present in moderately high latitudes (58°–68°) of the Northern Hemisphere of the Atlantic and Pacific (e.g. Kramp 1939; Madsen 1944; Grasshoff 1977; Sánchez and Cairns 2004; Sanmartín and P.J. López-González in prep), they are scarce in the high latitudes of the Southern Hemi-

sphere, which are dominated by the families Primnoidae and Isididae (e.g. Bayer 1996a, b, 1998; Alderslade 1998; López-González and Gili 2002).

On some recent Antarctic cruises, an extensive collection of octocorals has been made in the Weddell Sea, off the Antarctic Peninsula, and Scotia Sea. Among them, two ANDEEP cruises (Antarctic benthic deep-sea biodiversity; ANDEEP-I and ANDEEP-II, both in the austral summer 2002) were carried out on board the R/V *Polarstern* in the Scotia Sea to investigate the colonization history and recent community patterns of Antarctic benthic fauna. During the ANDEEP-I cruise an undescribed gorgonian, here placed in the family Plexauridae, was collected. The aim of this paper is the description of a new genus and species based on that material.

Methods

The material was collected in the Scotia Sea (Fig. 1) during the ANDEEP-I cruise. The octocorals were sorted, labelled and fixed on board. The holotype colony studied was fixed in buffered formaldehyde (4% in sea water) and later preserved in 70% ethanol. Fragments from different parts of the colony were prepared for SEM study following the usual methodology (e.g. Bayer and Stefani 1988; Alderslade 1998), and permanent mounts were made for light microscopy. Colony and sclerite terminology follows Bayer et al. (1983). The specimen studied has been deposited in the Zoologisches Institut und Zoologisches Museum in Hamburg (ZI-ZMH).

Results

Subclass Octocorallia
Order Alcyonacea
Suborder Holaxonia
Family Plexauridae Gray, 1859

Communicated by R.M. Kristensen

P. J. López-González
Biodiversidad y Ecología de Invertebrados Marinos,
Departamento de Fisiología y Zoología,
Facultad de Biología, Universidad de Sevilla,
Reina Mercedes, 6, 41012 Sevilla, Spain
E-mail: pjlopez@us.es




Fig. 1 Type locality of *Mesogligorgia scotiae* gen. nov., sp. nov. (asterisk)

Mesogligorgia gen. nov.

Diagnosis

Plexauridae with unbranched (possibly scarcely branched) colonies; holdfast well developed, cone shaped, not calcified. Coenenchyme remarkably thick with abundant mesogloea, in which sclerites are embedded forming distinct groups. Polyps scattered on all sides of the stem; polyps completely retractile in calyces embedded in the coenenchymal mesoglea, only distinct as low dome-shaped verrucae with eight equally developed mesogloea lobes. Sclerites elongated spindles with irregular edges; calicular sclerites indistinctly converging in eight points; anthocodial armature symmetrical, forming a poorly distinct collaret and eight well-developed points; flattened sclerites are present on the main axis of tentacles, and minute bodies on pinnules.

Etymology

The generic name combines *Mesogli-* in reference to the distinct abundance of mesogloea in the thick coenenchyme, and *-gorgia*, a common suffix in gorgonian generic names. Gender: feminine.

Fig. 2 *Mesogligorgia scotiae* gen. nov., sp. nov. Holotype ZIZMH (C 11689): **a**, whole colony, showing the conical holdfast, the irregular placement of polyps and the irregular diameter of the coenenchyme bearing polyps; **b** distal end; **c** a portion of the colony, showing the low calyces, and, due to transparency of the mesogloea, the eight points of the anthocodia and coenenchymal sclerites (*cos*); **d** limit between the basal coenenchyme without polyps and the upper part of the colony with polyps, showing the strong difference in diameter and the axis which is naked basally (*ax*). Scale bars: **a** 12 mm; **b–d**, 4 mm



Type species

Mesogligorgia scotiae sp. nov. here designated.

Mesogligorgia scotiae sp. nov. (Figs. 2–5)

Type material

Holotype: ZIZMH (C 11689), one whole colony, AN-DEEP-I cruise, *Polarstern* ANT XIX/3, stn. 41–4, Scotia Sea, 59°23.48'S 60°3.54'W, 2,201–2,213 m, Agassiz trawl, 27 Jan 2002.

Description

The holotype is a complete colony with holdfast, about 40-mm high, and unbranched (Fig. 2a). The diameter is irregular along the main stem, about 3.7–8.0 mm, with a claviform distal end. The coenenchyme of the polyp-bearing distal part, which has abundant translucent mesogloea, can be remarkably thick (0.3–2.5 mm) with respect to the basal part (Fig. 2d). Horny axis, 1.65 mm in diameter at basal level, has a cross-chambered central core and numerous loculi (Fig. 4a, b). The holdfast is cone-shaped, not calcified (Fig. 2a).

The polyps are irregularly placed around the stem (Fig. 2b, c). The anthocodiae are completely retractile into the calyces. Calyces are low, embedded in the coenenchymal mesogloea, only distinct as low dome-shaped

verrucae with eight equal mesogloecal lobes, 1.8–2.5 mm in diameter and 0.26–0.43-mm tall (Figs. 2, 3).


All sclerites in the colony are elongated spindles with irregular edges, sometimes with bi- or trifurcated ends (Fig. 5). The coenenchymal sclerites (Fig. 5a) are sparsely and irregularly distributed (Figs. 2b–d, 3, 4a, b), 0.47–1.10 mm in length, embedded in the abundant mesogloea. The coenenchymal sclerites can be observed in situ, sometimes running more or less longitudinally along the main axis. The calicular sclerites (Fig. 5b), 0.55–0.79 mm in length, are indistinctly converging in eight points, forming a deep palisade embedded into the coenenchymal mesogloea, and can be observed due to the transparency of the coenenchyme (Figs. 3, 4a, b). The anthocodial armature is symmetrical, forming an indistinct collaret and eight well-developed points with numerous lines of sclerites (Fig. 4c), 0.39–0.55 mm in length (Fig. 5c). The tentacular sclerites are flattened and slightly curved (Fig. 5d), obliquely placed on both sides of the main axis (Fig. 4d), 0.26–0.39 mm in length. Minute calcareous bodies (< 0.012 mm) are present on pinnules (Fig. 4e).

The holotype colony was male, with developing spermatid cysts up to 0.22 mm.

Colour

The living holotype was violet, whitish to dirty grey in the preserved state. The sclerites are colourless.

Fig. 3 *Mesogligorgia scotiae* gen. nov., sp. nov. Holotype ZIZMH (C 11689): **a** and **b**, retracted polyps in vertical view, showing the palisade of calicular sclerites (cs) in indistinctly converging groups and calicular aperture with mesogloecal lobes, and —due to transparency of the mesogloea—cos; **c–e**, retracted polyps in lateral view, showing the indistinct converging groups of calicular sclerites in **e**, and the eight points sclerites (ps) and calicular border (cb) in **d** and **e**. Scale bars: **a–e**, 2 mm



Etymology

The specific name *scotiae* refers to the geographic area where the new species was found.

Geographical and bathymetric distribution

At present, *Mesogligorgia scotiae* is known only from the Scotia Sea, depth 2,201–2,213 m.

Discussion

The presence of a horny axis with a cross-chambered central core and numerous loculi, retractile polyps armed with large sclerites in collaret and points, and calyces with distinct spicular components clearly support the inclusion of *Mesogligorgia* in the family Plexauridae.

As stated above, Plexauridae is one of the richest families in genera and species, although, the new taxon *Mesogligorgia* shows a set of characters (outlined above) not previously found in the family. For the following comparisons, additional information of the different plexaurid genera can be found in Germanos (1896), Wright and Studer (1889), Kükenthal (1919, 1924), Bayer (1956, 1961, 1981), Carpine and Grasshoff (1975), Grasshoff (1977, 1986, 1992, 1999, 2000), Williams (1992), Fabricius and Alderslade (2001) and Sánchez and Cairns (2004).

Mesogligorgia clearly differs from genera such as *Pseudoplexaura* Wright and Studer, 1889; *Eunicea* (s.l.)

Lamouroux, 1816; *Plexaura* Lamouroux, 1812; *Muriceopsis* Aurivillius, 1931 and *Psammogorgia* Verrill, 1868 by the absence of club-shaped sclerites; and from *Bebryce* Philippi, 1841 and *Plexaurella* Valenciennes, 1855 by the absence of sclerites as stellate plates or stellate forms with four or more rays.

A number of plexaurid genera show characteristic calicular sclerites as thorn-scales or thorn-spindles, such as *Villogorgia* (s.l.) Duchassaing and Michelotti, 1860; *Acanthaxis* Deichmann, 1936; *Astromuricea* Germanos, 1895; *Dentomuricea* Grasshoff, 1977; *Paramuricea* Kölliker, 1865; *Placogorgia* Kükenthal, 1919; *Lepidomuricea* Kükenthal, 1919; *Muriceides* Studer, 1887; *Echinogorgia* Kölliker, 1865; *Echinomuricea* Verrill, 1869; *Trimuricea* Gordon, 1926 or *Spinimuricea* Grasshoff, 1992; however, these sclerites are absent in *Mesogligorgia*.

The presence of capstans clearly distinguish *Swiftia* Duchassaing and Michelotti, 1864; *Heterogorgia* Verrill, 1868; *Anthoplexaura* Kükenthal, 1908 and *Alaskagorgia* Sánchez and Cairns, 2004 from *Mesogligorgia*. *Lytrea* Bayer, 1981 is characterized by a superficial layer of small coenenchymal tuberculate spheres and double-headed sclerites, and *Menacella* Gary, 1870 by spindles with crenulated or foliate tubercles, which are absent in *Mesogligorgia*.

Menella Gray, 1870 and *Paraplexaura* Kükenthal, 1909 show surface sclerites with vertically placed leaves, or sculptured with blunt humps, respectively, completely different from the deep layer of elongated spindles present in the new taxon here described.

Fig. 4 *Mesogligorgia scotiae* gen. nov., sp. nov. Holotype ZIZMH (C 11689): **a** longitudinal section of the distal end, showing the retracted polyps, the cross-chambered *ax*, some spermatic cysts in one of the sectioned polyps, and the thick coenenchyme; **b** detail of **a** showing a sectioned polyp, *ax*, deep layer of coenenchymal sclerites (*coe*), thick layer of mesogloea (*me*); **c** detail of the poorly distinct crown and well-developed eight point of a polyp after dissolution of the soft tissue by the use of a solution of sodium hypochlorite; **d** lateral view of a tentacle showing the oblique placement of tentacular sclerites, the soft tissue on the right are the pinnules; **e** a tentacular sclerite (out of focus) surrounded by several minute bodies (< 0.012 mm, two of them arrowed) which are in the pinnules. Scale bars: **a** 4 mm; **b** 2 mm; **c** 1 mm; **d** 0.5 mm, **e** 0.1 mm





Fig. 5 *Mesogligorgia scotiae* gen. nov., sp. nov. Holotype ZIZMH (C 11689): **a** sclerites from the coenemchyme; **b** sclerites from the calyces; **c** sclerites from the crown and points; **d** sclerites from the axis of the tentacles, the two first sclerites are in lateral view. Scale bars: 0.3 mm



Hypnogorgia Duchassaing and Michelotti, 1864 and *Mesogligorgia* differ in the asymmetrical anthocodial armature of the former genus. The cylindrical or tubular calyces of *Anthomuricea* Studer, 1887 and *Muricea* (s.l.) Lamouroux, 1812 are quite different from the low mesogloecal calyces of *Mesogligorgia*. The surface sclerites in other genera are discoidal with lobate edges like in *Discogorgia* Kükenthal, 1919; or tuberculate to small prickled spindles like in *Astro-*

gorgia Verrill, 1868 or *Elasmogorgia* Wright and Studer, 1889; which are quite different from those present in *Mesogligorgia*.

The typical two-lobed calyces of *Calicogorgia* Thomson and Henderson, 1906 and *Caliacis* Deichmann, 1936 can be used as a distinguishing character (among others) from *Mesogligorgia* in which the calicular apertures have eight equally developed mesogloecal lobes. Although the calyces of *Euplexaura* Verrill, 1869

are low and inconspicuous, the sclerites are different from the sclerites of *Mesogligorgia*.

Three plexaurid genera have flattened spindles, *Thesea* Duchassaing and Michelotti, 1864; *Scleracis* Kükenthal, 1919 and *Paracis* Kükenthal 1919. However, in these genera coenenchymal sclerites are thick large plates or flattened spindles tightly fitted as in a mosaic, and there are clear differences in size and shape between coenenchymal and calicular sclerites. In *Mesogligorgia* all sclerites (except for the minute bodies found in the pinnules) are similar in shape, only varying in size range in the different parts of the colony. Coenenchymal sclerites are not arranged in a mosaic, running more or less longitudinally in different planes, embedded in the thick coenenchyme with abundant mesogloea.

Acknowledgements The author acknowledges the valuable assistance of the officers and crew of the R/V *Polarstern*, and many colleagues on board during the ANDEEP cruises. I take this opportunity to extend my thanks to the cruise leader and steering committee of both cruises (ANDEEP-I and ANDEEP-II), especially to Dieter Fütterer and Angelika Brandt, who kindly facilitated the work on board, and for the opportunity to collaborate in this Antarctic programme. The author thanks Mercedes Conradi for her valuable assistance during ANDEEP-I cruise, and Asunción Fernández (Scanning microscope services of the University of Seville) for her helpful assistance. Support for this work was provided by the MCT projects REN2001-4269-E/ANT and REN2001-4929-E/ANT. The author is also thankful for comments and suggestions given by two anonymous referees. This is ANDEEP publication n° 48

References

- Alderslade P (1998) Revisionary systematics in the gorgonian family Isididae, with description of numerous new taxa (Coelenterata, Octocorallia). *Rec W Aust Mus Suppl* 55:1–359
- Bayer FM (1956) Octocorallia. In: Moore RC (ed) *Treatise on Invertebrate Paleontology, Part F Coelenterata*. University of Kansas Press, Lawrence, KS, pp F166–F230
- Bayer FM (1961) The shallow-water Octocorallia of the West Indian region. A manual for marine biologists. *Studies on the Fauna of Curaçao and other Caribbean Islands* 12:1–373, pls. 1–27
- Bayer FM (1981) Key to the genera of Octocorallia exclusive of Pennatulacea (Coelenterata: Anthozoa), with diagnoses of new taxa. *Proc Biol Soc Wash* 94(3):901–947
- Bayer FM (1996a) The Antarctic genus *Callozostrom* and its relationship to *Primnoella* (Octocorallia: Gorgonacea: Primnoidae). *Proc Biol Soc Wash* 109(1):150–203
- Bayer FM (1996b) New primnoid gorgonians (Coelenterata: Octocorallia) from Antarctic waters. *Bull Mar Sci* 58(2):511–530
- Bayer FM (1998) A review of the circumaustral gorgonacean genus *Fannyella* Gray, 1870 with description of five new species (Coelenterata: Octocorallia: Primnoidae). *Senckenberg biol* 77(2):161–204
- Bayer FM, Stefani J (1988) Primnoidae (Gorgonacea) de Nouvelle-Calédonie. *Bull Mus natn Hist nat Paris* (4) 10(A) 3:449–476
- Bayer FM, Grasshoff M, Verseveldt J (1983) Illustrated trilingual glossary of morphological and anatomical terms applied to Octocorallia. E.J. Brill/Dr. W. Backhuys, Leiden, pp 1–75
- Carpine C, Grasshoff M (1975) Les gorgonaires de la Méditerranée. *Bull Inst océanogr. Monaco* 71(1430):1–140
- Fabricius K, Alderslade P (2001) Soft corals and sea fans. A comprehensive guide to the tropical shallow water genera of the central-west Pacific, the Indian Ocean and the Red Sea. Australian Institute of Marine Science, Townsville, Queensland, pp i–viii + 1–264
- Germanos NK (1896) Gorgonaceen von Ternate. In: Kükenthal W (ed) *Ergebnisse einer zoologischen Forschungsreise in den Molukken und Borneo, im Auftrage der Senckenbergischen naturforschenden Gesellschaft ausgeführt von Dr. Willy Kükenthal, Teil 2, Band 1. Abhandl Senckenb naturf Gesellsch* 23 (1):145–187, 9–12
- Grasshoff M (1977) Die Gorgonarien des ostlichen Nordatlantik und des Mittelmeeres. III. Die Familie Paramuriceidae (Cnidaria, Anthozoa). “Meteor” *Forsch-Ergebnisse* D27:5–76
- Grasshoff M (1986) Die Gorgonaria der Expeditionen von “Travailleur” 1880–1882 und “Talisman” 1883 (Cnidaria: Anthozoa). *Bull Mus natn Hist nat Paris* (4)8(A)1:9–38
- Grasshoff M (1992) Die Flachwasser-Gorgonarien von Europa und Westafrika (Cnidaria, Anthozoa). *Cour Forsch-Inst Senckenberg* 149:1–135
- Grasshoff M (1999) The shallow water gorgonians of New Caledonia and adjacent islands (Coelenterata, Octocorallia). *Senckenb Biol* 78(1/2):1–121
- Grasshoff M (2000) The gorgonians of the Sinai coast and the Strait of Gubal, Red Sea (Coelenterata, Octocorallia). *Cour Forsch-Inst Senckenberg* 224:1–125
- Kramp PL (1939) Octocorallia. *The Zoology of Iceland* 2(7):1–13
- Kükenthal W (1919) Gorgonaria. *Wissensch Ergebn deutschen Tiefsee-Exped “Valdivia”* 13 (2):1–946, 30–89
- Kükenthal W (1924) Gorgonaria. *Das Tierreich*, pp 1–478
- López-González PJ, Gili J-M (2002) A new primnoid genus (Anthozoa: Octocorallia) from the Southern Ocean. *Sci Mar* 66(4):383–397
- Madsen FJ (1944) Octocorallia. *Dan Ingolf-Exped* 5(13):1–65
- Sánchez JA, Cairns SD (2004) An unusual new gorgonian coral (Anthozoa: Octocorallia) from the Aleutian Islands, Alaska. *Zool Med Leiden* 78(15):265–274
- Williams GC (1992) The Alcyonacea of Southern Africa. Gorgonian Octocorals (Coelenterata, Anthozoa). *Ann S Afr Mus* 101(8):181–296
- Wright EP, Studer T (1889) Report on the Alcyonaria collected by H.M.S. Challenger during the years 1873–1876. *Rept Sci Res Challenger, Zool.* 31:i-lxxvii + 1–314