

Hexagonal Tilings: Tutte Uniqueness

D. Garijo *, A. Márquez *, M.P. Revuelta *

Abstract

We develop the necessary machinery in order to prove that hexagonal tilings are uniquely determined by their Tutte polynomial, showing as an example how to apply this technique to the toroidal hexagonal tiling.

1 Introduction

The main result in this paper has to do with graphs determined by their Tutte polynomial. This is a two variable polynomial $T(G; x, y)$ associated with any graph G , which contains interesting information about G . For instance, $T(G; 2, 0)$ is the number of acyclic orientations of G and $T(G; x, 0)$ with $x < 0$ is the chromatic polynomial associated with G . As a natural extension of the concept of chromatically unique graph, the notion of Tutte unique graph was studied in [9]. A graph G is said to be *Tutte unique* if $T(G; x, y) = T(H; x, y)$ implies $G \cong H$ for any other graph H . A common topic in the study of this invariant is the search of large families of Tutte unique graphs. In 2003, Garijo, Márquez, Mier, Noy, Revuelta [7],[8] present locally grid graphs as the first large family of graphs uniquely determined by their Tutte polynomial. In this paper we prove that the toroidal hexagonal tilings also satisfy this uniqueness property. The technique developed here can also be applied to the rest of hexagonal tilings and their dual graphs, the locally C_6 graphs. The main motivation for this study is the search of a generalization to locally plane graphs of the well known relationship existing between the Tutte polynomials of plane graphs and their duals [13].

A hexagonal tiling, H , is defined as a connected cubic graph of girth 6, having a collection of 6-cycles, C , such that every 2-path is contained in precisely one cycle of C (2-path condition). A hexagonal tiling is simple (that is, without loops and multiple edges) and every vertex belongs to exactly three hexagons (cycles of length 6)(Figure 1). Every hexagon of the tiling is called a *cell*.



Figure 1: Hexagonal structure around x

Let $G = (V, E)$ be a graph with vertex set V and edge set E . The *rank* of a subset $A \subseteq E$ is $r(A) = |A| - k(A)$, where $k(A)$ is the number of connected components of the spanning subgraph (V, A) . The Tutte polynomial is defined as follows:

$$T(G; x, y) = \sum_{A \subseteq E} (x - 1)^{r(E) - r(A)} (y - 1)^{|A| - r(A)}$$

*Dep. Matemática Aplicada I. Universidad de Sevilla. Avda. Reina Mercedes s/n. 41012 Sevilla (Spain). {dgarijo,almar,pastora}@us.es

$T(G; x, y)$ contains exactly the same information about G as the rank-size generating polynomial which is defined as:

$$R(G; x, y) = \sum_{A \subseteq E} x^{r(A)} y^{|A|}$$

The coefficient of $x^i y^j$ in $R(G; x, y)$ is the number of spanning subgraphs in G with rank i and j edges, hence the Tutte polynomial tell us for every i and j the number of edges-sets in G with rank i and size j .

The main strategy in the paper is the following. We first recall some definitions and results from [5, 12], such as, the classification theorem of hexagonal tilings. In Section 3, we show that these graphs are locally orientable, using the relationship existing between hexagonal tilings and locally grid graphs [5]. The local orientation of these graphs allows us to count edge-sets and it is one of the basic tools used to prove the Tutte uniqueness results. Thus, as an example, in Section 5 we apply the technique developed to show that for every hexagonal tiling H not isomorphic to $H_{k,m,0}$ with $2k(m+1)$ vertices there is at least one coefficient of the rank-size generating polynomial in which both graphs differ.

2 Preliminary Results

In this section we state the classification theorem of hexagonal tilings proved in [5]. There exists an extensive literature on this topic. See for instance the works done by Altshuler [1, 2], Fisk [3, 4] and Negami [10, 11]. In [5], following up the line of research given by Thomassen [12], we add two new families to the classification theorem given in [12] proving that with these families we exhaust all the cases.

Theorem 2.1. [5] *If G is a hexagonal tiling with N vertices, then one and only one of the following holds:*

- A)** $G \simeq H_{k,m,r}$ with $N = 2k(m+1)$, $0 \leq r \leq \lfloor k/2 \rfloor$, $m \geq 2$, $k \geq 3$. If $m = 1$ then $k > 3$ and $\lfloor k/2 \rfloor \geq r \geq 2$. If $m = 0$ then $k > 3$ and $3 \leq r \leq k$
- B)** $G \simeq H_{k,m,a}$ with $N = 2k(m+1)$, $m \geq 2$, $k \geq 3$.
- C)** $G \simeq H_{k,m,b}$ with $N = 2k(m+1)$, k even, m odd, $m \geq 3$, $k \geq 4$.
- D)** $G \simeq H_{k,m,c}$ with $N = 2k(m+1)$, $m \geq 1$, k even, $k \geq 6$.
- E)** $G \simeq H_{k,m,f}$ with $N = 2k(m+2)$, k odd, $m \geq 0$, $k \geq 7$.
- F)** $G \simeq H_{k,m,g}$ with $N = 2(m+1)(k+2)$, $k \geq m+1$, $m \geq 3$.
- G)** $G \simeq H_{k,m,h}$ with $N = 2(m+1)(k+1)$, $k < m-1$, $k \geq 2$.

Some examples of hexagonal tilings are given in Figures 2, 3, 4 and 5. The parameters k and m fix, respectively, the height and the breath of the structures, called *hexagonal wall of length k and breath m* (Figures 2, 3, 4) and *hexagonal ladder of length k and breath m* (Figure 5), in which it is added edges to construct the different families of hexagonal tilings (see [5]).

Given two cycles C and C' in a hexagonal tiling G , we say that C is *locally homotopic* to C' if there exists a cell, H , with $C \cap H$ connected and C' is obtained from C by replacing $C \cap H$ with $H - (C \cap H)$. A *homotopy* is a sequence of local homotopies. A cycle in G is called *essential* if it is not homotopic to a cell. Otherwise it is called *contractible*. This definition is equivalent to the one given for a graph embedded in a surface [6]. Let l_G be the minimum length of the essential cycles of G , l_G is invariant under isomorphism. An edge-set contained in a hexagonal tiling is called a *normal edge-set* if it does not contain any essential cycle.




Figure 2: a) $H_{5,5,r}$ b) $H_{5,4,a}$ c) $H_{6,5,b}$




Figure 3: a) $H_{6,4,c}$ b) Embedding of $H_{6,4,c}$ in the Klein bottle

There are two different ways of pasting together j ladders each one containing i hexagons, from which we obtain two structures, called the *ladder* $i \times j$ and the *displaced ladder* $i \times j$, shown in Figure 6. Note that the number of shortest paths between x and y or between x and z in a ladder $i \times j$ or in a displaced ladder $i \times j$ is $\binom{i+j}{j}$ and the length of these paths is $2(i+j) - 1$.

Every hexagonal tiling is obtained by adding edges to a hexagonal wall or to a hexagonal ladder (except for $H_{k,m,h}$, in which we have also added two vertices) [5]. These edges are called *exterior edges* and every essential cycle must contains at least one of these edges (the edges $\{(0, 2k + m), (0, 2k + m - 1)\}$ and $\{(m - k - 1, 3k + 2), (m - k - 1, 3k + 1)\}$ in $H_{k,m,h}$ are not considered exterior edges).




Figure 4: a) $H_{7,4,f}$ b) Embedding of $H_{7,4,f}$ in the Klein bottle




Figure 5: a) $H_{7,4,g}$ b) $H_{6,4,h}$

3 Local Orientability of Hexagonal Tilings

In this section we prove that hexagonal tilings are locally orientable. This fact allow us to count edge-sets and it is the tool to distinguish at least one coefficient of the Tutte polynomials associated to two non isomorphic hexagonal tilings. We first recall a minor relationship existing between hexagonal tilings and locally grid graphs (see [5]).

We say that a 4-regular, connected graph G is a *locally grid graph* if for every vertex x there exists an ordering x_1, x_2, x_3, x_4 of $N(x)$ and four different vertices y_1, y_2, y_3, y_4 , such that, taking the indices modulo 4,

$$\begin{aligned} N(x_i) \cap N(x_{i+1}) &= \{x, y_i\} \\ N(x_i) \cap N(x_{i+2}) &= \{x\} \end{aligned}$$

and there are no more adjacencies among $\{x, x_1, \dots, x_4, y_1, \dots, y_4\}$ than those required by this condition (Figure 7).

Every locally grid graph is a minor of a hexagonal tiling and in [5] it was proved that there exists a bijective minor relationship preserved by duality between hexagonal tilings with the same




Figure 6: a) Displaced ladder 4×2 b) Ladder 4×2



Figure 7: Locally Grid Structure

chromatic number and locally grid graphs. A perfect matching P is selected in each one of the families given in the classification theorem of hexagonal tilings, except in one case, $H_{k,m,f}$ in which the set of selected edges is not a matching. P contains two edges of each hexagon. In $H_{k,m,f}$ there are $k - 1$ hexagons in which we select four edges pairwise incident. A locally grid graph is obtained by contracting the edges of this matching, and deleting parallel edges if necessary. There are just two cases in which parallel edges must be deleted, $H_{k,m,f}$ and $H_{k,m,h}$. These perfect matchings and the set of edges of $H_{k,m,f}$ verify that if we have two hexagonal tilings with the same chromatic number, the result of the contraction of their perfect matchings (or the selected edges in $H_{k,m,f}$) are two locally grid graphs belonging to different families. This condition is going to be essential in the search of a generalization, for these families, of the relationship existing between the Tutte polynomial of a plane graph and its dual [13].

In [8] it was proved that locally grid graphs are locally orientable. Given a vertex v of a locally grid graph G , two edges incident with v are said to be *adjacent* if there is a square containing them both; otherwise they are called *opposite*. An orientation at a vertex v consists of labeling the four edges incident with v bijectively with the labels N, S, E, W in such a way that the edges labelled N and S are opposite, and so are the ones labelled as E and W .

Lemma 3.1. *Hexagonal tilings are locally orientable.*

Proof. Let H be a hexagonal tiling. By Theorem 2.1, H is isomorphic to one of the following graphs: $H_{k,m,r}, H_{k,m,a}, H_{k,m,b}, H_{k,m,c}, H_{k,m,h}, H_{k,m,g}, H_{k,m,f}$.

Suppose first that H is not isomorphic to $H_{k,m,f}$. For every vertex $v \in V(H)$ there is exactly one edge of the perfect matching P incident with v , call it $\{v, v'\}$. Consider the union of the local structures of v and v' , labeling the other two edges incident with v as e_1 and e_2 and the other two edges incident with v' as e_3 and e_4 . Contract all the edges of P belonging to this union, deleting parallel edges if necessary. As it is shown in Figure 8a we obtain a grid 3×3 , hence we can label the edges e_1, e_2, e_3 and e_4 bijectively with the labels N, S, E, W following the criterion established for the locally grid graphs. The edge $\{v, v'\}$ is considered opposite to the unique edge incident with v and belonging to the perfect matching, P' , formed by the edges $\{(i, l)(i, l + 1)\}, 0 \leq i \leq m$ not belonging to P .

If H is isomorphic to $H_{k,m,f}$, then there are either one or two edges belonging to P and incident with v . If v does not belong to a hexagon that has more than two edges of P , we follow the same process as in the previous case. Suppose, now, that v is a vertex belonging to one of the hexagons that contains four edges belonging to P . We consider two cases:




Figure 8: a) Local orientation at $v \in V(H)$ not isomorphic to $H_{k,m,f}$ b) Local orientation at $v \in V(H_{k,m,f})$

(1) There are two edges of P incident with v , call them $\{v, v'\}$ and $\{v, v''\}$. The union of the hexagonal structures of v, v' and v'' is composed by five hexagons. Label the unique edge incident with v and not belonging to P as e_1 . The edge incident with v'' not belonging to P and sharing an hexagon with e_1 is labelled e_2 . Finally, the edges incident with v' no belonging to P are labelled e_3 and e_4 . Contracting all the edges of P belonging to this union and deleting parallel edges, we also obtain a grid 3×3 and we can label the edges incident with v, v' and v'' bijectively with the labels N, S, E, W (Figure 8b). $\{v, v'\}$ and $\{v, v''\}$ are considered opposite to the unique edge incident with v' and v'' respectively, belonging to P' .

(2) There is one edge of P incident with v , call it $\{v, v'\}$. There has to be one edge, different from $\{v, v'\}$, incident with v' and belonging to P , call it $\{v', v''\}$. The union of the hexagonal structures of v and v'' is composed by five hexagons. The two edges incident with v not belonging to P are labelled e_1 and e_2 . The unique edge incident with v' not belonging to P is labelled e_3 , and the edge incident with v'' not belonging to P or to the hexagon that contains four edges of P , is labelled e_4 . Contracting all the edges of P belonging to this union and deleting parallel edges, we also obtain a grid 3×3 and we can label the edges incident with v, v' and v'' bijectively with the labels N, S, E, W . $\{v, v'\}$ and $\{v', v''\}$ are considered opposite to the edges incident with v and v'' respectively, belonging to the hexagon that contains four edges of P .

□

The previous proof give us a definition of *adjacent and opposite edges* for hexagonal tilings. Two edges incident with a vertex v and no belonging to P are called *adjacent* or *opposite* if they are so in the locally grid structure resulting from applying the operations explained above. Two edges incident with v and belonging to P are always opposite.

By the orientation (v, v', e, f) we mean that v is the origin vertex, $\{v, v'\} \in P$, e is the unique edge incident with v and opposite to the other two edges incident with v . This edge never belongs

to P and it is labelled E . Finally, f is an edge incident with v and labelled N . The orientation (v, v', v'', e, f) means that v is the origin vertex, $\{v, v'\}, \{v, v''\} \in P$, e is the unique edge incident with v and opposite to the other two edges incident with v , and f is an edge incident with v labelled N . We will denote both cases as (v, v', e, f) .

If $\alpha \in \{N, S, E, W\}$, we denote α^{-1} the label opposite to α . If we fix an orientation at $v \in V(H)$ (v is the origin vertex) then every vertex u adjacent to v is unambiguously oriented, because the orientation at v induces an orientation at u : if $\{u, v\}$ is labelled α from v , it is labelled α^{-1} from u . If $tzxyvw$ is a hexagon and $\{x, v\}, \{w, v\}$ are labelled β and α from v respectively, then $\{x, y\}$ is β from x , $\{z, y\}$ is α from y , $\{z, t\}$ is β^{-1} from z and $\{t, w\}$ is β^{-1} from t . We also can translate the orientation at v to all the vertices of a path beginning at v , but two different paths joining v and u can induce different orientation at u . This does not happen if the union of these two paths is a contractible cycle. In fact, we can transform one path into the other one using three elementary transitions and their inverse: if e, f, g, h, i, j are the edges of a hexagon ordered cyclically, we can change e, f, g, h, i by j ; e, f, g, h by i, j and e, f, g by h, i, j and these operations do not change the orientation at the endpoint.

Let A be a connected normal edge set in a hexagonal tiling H . Fixed an orientation at a vertex $v \in V(A)$, then all the vertices in $V(A)$ are unambiguously oriented. Every path in A can be described as a sequence of the labels $\{N, S, E, W\}$ (Figure 9a). This allow us to assign coordinates to every vertex in $V(A)$: v has coordinates $(0, 0)$ and $u \in V(A)$ has coordinates (i, j) if in one path in A joining v to u (taking into account that the orientation at the endpoint induced by different paths joining v to u does not change), i is the number of labels E minus the number of labels W and j is the number of labels N minus the number of labels S .

Lemma 3.2. *Let A be a connected normal edge set in a hexagonal tiling H with $|A| \leq l_H + 3$, then different vertices in $V(A)$ have different coordinates.*

Proof. First, we are going to prove that with a fixed orientation (v, v', e, f) no vertex except the origin can have coordinates $(0, 0)$. Suppose that there exists $x \in V(A)$ such that x has coordinates $(0, 0)$, then there exists a path R joining v to x with as many labels N as labels S and as many labels E as labels W . We are going to show that $x = v$ by induction on the length of this path.

If $|R| = 6$ and v is the origin vertex, the first labels of R can be: NN, NW, E, SS , or SW and using the elementary transitions described previously, we can change these labels by $ENNW, SWNN, NNESS, ESSW$ and $NWSS$ respectively. To obtain as many labels N as labels S and as many labels E as labels W it must be that $v = x$. We assume that every path R (not a cycle) with $|R| < n$ does not have as many labels N as labels S and as many labels E as labels W and we prove the result for $|R| = n$. Suppose that there exists a path R joining v and x , not a cycle, with $|R| = n$ and having as many labels N as those labelled S and as many E as W , then there exists a subsequence labelled $\beta(\alpha^{-1})^l\beta\alpha^s\beta^{-1}(\alpha^{-1})^m\beta^{-1}$ or $\beta\alpha^l\beta\alpha^s\beta^{-1}\alpha^m\beta^{-1}$ (or other analogous subsequence). In the first case we can change this subsequence by α^{s-l-m} keeping the orientation at the endpoint. Analogously, in the second case the subsequence can be changed by α^{s+l+m} . In any case, the change gives rise to a new path R' joining v and x with $|R'| < n$, in a new connected edge set A' . This set is also a normal edge set because $|A'| \leq |A| - 4 \leq l_H - 1$. By hypothesis, R' does not have as many labels N as S and E as W , hence neither does R and we obtain a contradiction.

Given two different vertices, u and w in $V(A)$, we consider the paths R_u and R_w joining u to v and w to v respectively. Since A is connected, there is a path $R_w \cup R_u$ joining v to u and keeping the orientation at u . Hence, it is easy to prove that u and w can not have the same coordinates. \square

From now on, and unless otherwise stated, we consider that all normal edge-sets have at most $l_H + 3$ elements.

4 Counting Edge-Sets

The aim of this section is to find a system which allow us to count edge-sets in order to prove that there is at least one coefficient of the rank-size generating polynomial in which two non isomorphic hexagonal tilings differ. Our first step is to codify the edge-sets of a hexagonal tiling.

Let L^∞ be the infinite plane square lattice, that is, the infinite graph having as vertices $\mathbb{Z} \times \mathbb{Z}$ and in which (i, j) is joined to $(i-1, j), (i+1, j), (i, j-1), (i, j+1)$. We delete the edges $\{(i, 2j), (i-1, 2j)\}$ if i even and $\{(i, 2j+1), (i+1, 2j+1)\}$ if i odd, obtaining a *infinite plane hexagonal tiling* H^∞ . Let Ω be the group of graph automorphisms of H^∞ . This group is generated by translations, symmetries and rotations of the plane that map vertices to vertices. Let $\Sigma(H^\infty)$ be the set of all finite connected edge-sets of H^∞ . An equivalence relation is defined in $\Sigma(H^\infty)$ as follows:

$$B_1 \sim B_2 \Leftrightarrow \exists \sigma \in \Omega, \sigma(B_1) = B_2$$

If Λ is a set of representatives of $\Sigma(H^\infty)/\sim$ such that every $B \in \Lambda$ contains the vertex $(0, 0)$, then Λ covers all the possible shapes that a normal edge-set could have.

We are going to define a set Γ of words over the alphabet $\{N, S, E, W\}$ that represents all edge-sets of $\Sigma(H^\infty)/\sim$. Label $\{(0, 0), (0, 1)\}$ as N , $\{(0, 0), (1, 0)\}$ as E , $\{(0, 1), (-1, 1)\}$ from $(0, 1)$ as W and $\{(0, 0), (-1, 0)\}$ as S . For every $B \in \Lambda$ there is a sequence γ_B over the the alphabet $\{N, S, E, W\}$ such that beginning at $(0, 0)$ and following the instructions given by γ_B the unique edges covered are those of B . Note that one edge can be covered more than once and that the sequence γ_B is not unique (Figure 9a). Since every sequence over the alphabet $\{N, S, E, W\}$ can not represent a set of Λ (for instance, a sequence with two consecutive labels E) we are going to establish some restrictions on this alphabet:

- (1) No sequence begins with the label W .
- (2) There are no two consecutive labels E (analogously W).
- (3) If a sequence begins with a label N (analogously S), there must be an odd number of labels N and S before having a label W ; and an even number if the label is E . Hence, there are an odd number of labels N and S between two labels W and an even number between two labels E .
- (4) If a sequence begins with a label E , there are an odd number of labels N and S between two labels E and an even number between two labels W .

Call Γ the set of words over the alphabet $\{N, S, E, W\}$ verifying the three previous conditions. These restrictions do not modify the fact that every $B \in \Lambda$ has a word γ_B associated. Conversely, given $\gamma \in \Gamma$ there exists $B \in \Lambda$, such that $\gamma = \gamma_B$. It will be denoted $B(\gamma)$.

The next step is to assign one word from Γ to each normal connected edge-set of a hexagonal tiling, H . Fix an orientation (v, v', e, f) at H , being $v \in V(H)$ the origin vertex. Following the code given by γ from v with orientation (v, v', e, f) , an edge set $A^H(\gamma, v, v', e, f)$ is obtained, called the *instance of γ in H* and we call (v, v', e, f) an *orientation of the instance*. An instance of γ can have different orientations associated to it, the number of them depends only on the symmetries of $B(\gamma)$ and it is called *sym*(γ). The *length of γ* is the number of labels N, S, E, W that give rise to γ .

Lemma 4.1. *Let H be a hexagonal tiling and $\gamma \in \Gamma$. If $r(B(\gamma)) = l_H - 1$ and $|B(\gamma)| \leq l_H$, then $A^H(\gamma, v, v', e, f)$ is a normal edge-set and it has the same rank and size as $B(\gamma)$.*

Proof. We are going to prove this result by induction on the length of γ . If γ has length equal to one, the result is trivial. Let γ' be the word resulting from removing the last label from γ . By hypothesis $A^H(\gamma', v, v', e, f)$ is normal and it has the same rank and size as $B(\gamma')$. When we add the last label of γ to γ' , there are three possibilities on $B(\gamma')$:

- (1) We are adding one edge between two vertices belonging to $B(\gamma')$
- (2) We are adding an isthmus.
- (3) We do not add any edge.

of A . Since A is normal, this assignment is unambiguous. We define $\varphi : V(A) \rightarrow V(H^\infty)$ by $\varphi(x) = (i, j)$ if in every path in A joining v to x , i is the number of labels E minus the number of labels W and j is the number of labels N minus the number of labels S . Furthermore, we define $\bar{\varphi} : A \rightarrow E(H^\infty)$ by $\bar{\varphi}(\{x, y\}) = \{\varphi(x), \varphi(y)\}$. Taking $B = \bar{\varphi}(A)$, it is easy to prove that φ is a graph isomorphism and that $\bar{\varphi}$ is the isomorphism induced on edges by φ . The statements (1) and (2) of the lemma are consequence of the assignment of coordinates following an orientation (v, v', e, f) in a hexagonal tiling H , and the assignment of labels in the edges incident with the origin vertex v . We prove the uniqueness of B by induction on $m = \max\{d(v, x), x \in V(A)\}$. Suppose that there exists a pair (φ', B') verifying the statement of the lemma. If $m = 1$, $d(v, x) = 1 \forall x \in V(A)$ with $x \neq v$ hence $|A| \leq 3$. By (1) and (2), it is immediate to show that $B = B'$. Assume the result for $m \leq l$ and we prove it for $m = l + 1$. There exists $x_1, \dots, x_n \in V(A)$ such that $d(v, x_i) = l + 1$ and $d(v, x) < l + 1$ for all $x \neq x_i$. By induction on n we show that B is unique for $m = l + 1$. If $n = 1$, let x_1 be the unique vertex such that $d(v, x_1) = l + 1$ and let $A(x_1)$ be the set of edges incident with x_1 and belonging to A . Call A' the connected normal edge-set resulting from deleting the vertex x_1 . By the induction hypothesis, there exists a unique pair (ψ, \bar{B}) associated to A' , then $B = \bar{B} \cup \bar{\varphi}(A(x_1))$. Suppose that B is unique for $m = l + 1$ and having $n - 1$ vertices such that $d(v, x_i) = l + 1$. Using the same argument as in the previous case, it is easy to prove the result for n vertices. \square

Using the same arguments as those given in [8] for locally grid graphs, we prove the following result.

Lemma 4.3. *Given a connected normal edge-set $A \subseteq E(H)$, there exists a unique word $\gamma \in \Gamma$ and an orientation (v, v', e, f) , not necessarily unique, such that $A = A^H(\gamma, v, v', e, f)$.*

Lemma 4.4. *For fixed $n > 0$, the number of connected normal edge-sets with rank $n - 1$ and size n is the same for all hexagonal tilings H with $2pq$ vertices and such that $n \leq l_H$.*

Proof. By the previous lemma, every connected normal edge-set A is the instance of a unique word $\gamma(A)$. By Lemma 4.2 A is isomorphic to $B(\gamma(A))$ then they have the same rank and size. By Lemma 4.1 the instance of a word $\gamma(A)$ is a normal connected edge-set. Hence, the number of connected normal edge-sets with rank $n - 1$ and size n is equal to the number of distinct instances of words corresponding to edge-sets of H^∞ with rank $n - 1$ and size n . For every γ fixed we can choose $4pq$ different orientations. Let $\Gamma^{r,s}$ be the set of words of Γ such that $r(B(\gamma)) = r$ and $|B(\gamma)| = s$. Then the number of connected normal edge-sets with rank $n - 1$ and size n is:

$$\sum_{\gamma \in \Gamma^{n-1, n}} \frac{4pq}{\text{sym}(\gamma)}$$

Since $\text{sym}(\gamma)$ depends only on the word, this quantity does not depend on the graph. \square

A non-connected version of this lemma is proved for hexagonal tilings as in [8] for locally grid graphs.

Lemma 4.5. *For fixed $n > 0$ and $t > 1$, the number of normal edge-sets with rank $n - 1$, size n and t connected components is the same for all hexagonal tilings H with $2pq$ vertices and such that $n \leq l_H$.*

As a consequence of Lemmas 4.4 and 4.5, we have the following result.

Lemma 4.6. *Fix $n > 0$. The number of edge-sets with rank $n - 1$ and size n that do not contain essential cycles is the same for all hexagonal tilings H with $2pq$ vertices and such that $n \leq l_H$.*

Corollary 4.7. *Let H, H' be a pair of hexagonal tilings with $2pq$ vertices.*

- $l_H \neq l_{H'}$ implies $T(H; x, y) \neq T(H'; x, y)$.
- If $l_H = l_{H'}$ but H and H' do not have the same number of shortest essential cycles then $T(H; x, y) \neq T(H'; x, y)$.

Proof. If $l_H < l_{H'}$, there are essential cycles of length l_H in H but not in H' . Since by the preceding lemma the number of normal edge-sets with rank $l_H - 1$ and size l_H is the same in H and H' , the coefficient of $x^{l_H-1}y^{l_H}$ in the rank-size generating polynomial is greater in H than in H' . In an analogous way the second statement is proved. \square

The process we have followed to count normal edge-sets of size smaller than $l_H + 1$, is to show that we can assign a unique set, $A^H(\gamma(A), v, v', e, f)$ to every normal edge-set, A . Our next aim is to prove that for some special cases we can count normal edge-sets with size greater than l_H . This fact is going to be essential in the next section. Denote by $N(\gamma, H, n, r)$ the number of normal edge-sets in H with rank r , size n and having $B(\gamma)$ as a subgraph. An edge-set $A \subseteq E(H)$ is a *forbidden edge-set* for γ if it contains an essential cycle and a subset $B \subseteq A$ isomorphic to $B(\gamma)$.

Corollary 4.8. *Given $\gamma \in \Gamma$ such that $B(\gamma)$ contains at least one cycle, then we have that $N(\gamma, H, n, r)$ is the same for all hexagonal tilings H with $2pq$ vertices, with no forbidden edge-set for γ of size n and such that $n \leq l_H + 3$.*

Proof. Fix $\gamma \in \Gamma$ such that $B(\gamma)$ contains at least one cycle. We just have to prove that given $\gamma' \in \Gamma$ such that $B(\gamma) \subseteq B(\gamma')$, $r(B(\gamma')) = l_H + 2$ and $|B(\gamma')| \leq l_H + 3$, then $A^H(\gamma', v, v', e, f)$ is a normal edge-set and it has the same rank and size as $B(\gamma')$. This result is analogous to Lemma 4.1 but taking size smaller than $l_H + 4$.

If $B(\gamma) \subseteq B(\gamma')$ then $B(\gamma')$ contains at least one cycle. If $n \leq l_H$, the proof is equal to the one for Lemma 4.1. If $n = l_H + 1$, we show that $A^H(\gamma', v, v', e, f)$ has to be an essential cycle (then the rest of the proof would be analogous to the one followed in Lemma 4.1). Suppose that $A^H(\gamma', v, v', e, f)$ contains an essential cycle and one more edge. Since:

$$A^H(\gamma, v, v', e, f) \simeq B(\gamma) \subseteq B(\gamma') \simeq A^H(\gamma', v, v', e, f)$$

then $A^H(\gamma', v, v', e, f)$ is a forbidden edge-set for γ of size n and we obtain a contradiction. The cases in which $n = l_H + 2$ and $n = l_H + 3$ are analogously proved. \square

5 Tutte Uniqueness

In this section we apply the results obtained in order to prove the Tutte uniqueness of $H_{k,m,0}$. First we are going to show that the local hexagonal tiling structure is preserved by the Tutte polynomial, using the following lemmas.

Lemma 5.1. [5] *The edge-connectivity of every hexagonal tiling is equal to three.*

Lemma 5.2. [9] *If G is a 2-connected simple graph and H is Tutte equivalent to G , then H is also simple and 2-connected. If G is a 2-connected simple graph, then the following parameters are determined by its Tutte polynomial:*

- (1) *The number of vertices and edges.*
- (2) *The edge-connectivity.*
- (3) *The number of cycles of length three, four and five.*

Next, we are going to prove that there is at least one coefficient, in which, the Tutte polynomial associated to $H_{k,m,0}$ and the Tutte polynomial associated to any other hexagonal tiling not isomorphic to $H_{k,m,0}$, differ. As we have explained in the first section, the Tutte polynomial associated to a graph H tell us for every i and j the number of edge-sets in H with rank i and size j . There are two different types of these edge-sets: normal edge-sets (do not contain essential cycles) and edge-sets containing at least one essential cycle. We recall two results from [5]

Lemma 5.3. [5] *Let G be a hexagonal tiling, then the length l_G of their shortest essential cycles and the number of these cycles is:*

G	l_G	
$H_{k,m,r}$	$2k$	<i>if</i> $k < m + 1$
	$2(m + 1)$	<i>if</i> $r < \lfloor (m + 1)/2 \rfloor < \lfloor k/2 \rfloor$
	$2(m + 1 + r - \lfloor (m + 1)/2 \rfloor)$	<i>if</i> $\lfloor (m + 1)/2 \rfloor \leq r \leq \lfloor k/2 \rfloor$
	$2k$	<i>if</i> $k = m + 1$
$H_{k,m,a}$	$\min(2k, 2m + 2)$	
$H_{k,m,b}$	$\min(2k, 2m + 2)$	
$H_{k,m,c}$	$\min(k + 1, 4m + 4)$	
$H_{k,m,f}$	$\min(k, 4m + 8)$	
$H_{k,m,h}$	$2k + 2$	
$H_{k,m,g}$	$2(k - m) - 2\lfloor (m + 1)/2 \rfloor + 3$	<i>if</i> $k > 2m + 1$
	$k + 2$	<i>if</i> $k \leq 2m + 1$ and k odd
	$k + 3$	<i>if</i> $k < 2m + 1$ and k even

G	number of essential cycles of length l_G	
$H_{k,m,r}$	$m + 1$	<i>if</i> $k < m + 1$
	$k \binom{m + 1}{\lfloor (m + 1)/2 \rfloor - r}$	<i>if</i> $r < \lfloor (m + 1)/2 \rfloor < \lfloor k/2 \rfloor$
	$k \binom{r + \lfloor (m + 1)/2 \rfloor}{m}$	<i>if</i> $\lfloor (m + 1)/2 \rfloor \leq r \leq \lfloor k/2 \rfloor$
	$m + 1 + k \binom{m + 1}{\lfloor (m + 1)/2 \rfloor - r}$	<i>if</i> $k = m + 1$
$H_{k,m,a}$	$m + 1$	<i>if</i> $k < m + 1$
	2^{m+1}	<i>if</i> $k > m + 1$
	$2^{m+1} + m + 1$	<i>if</i> $k = m + 1$
$H_{k,m,b}$	$m + 1$	<i>if</i> $k < m + 1$
	$2 \binom{m + 1}{(m + 1)/2} + 4 \sum_{j=1}^{(m-1)/4} \binom{m + 1}{(m + 1)/2 - 2j}$	<i>if</i> $k > m + 1$
$H_{k,m,c}$	$m + 1 + 2 \binom{m + 1}{(m + 1)/2} + 4 \sum_{j=1}^{(m-1)/4} \binom{m + 1}{(m + 1)/2 - 2j}$	<i>if</i> $k = m + 1$
	$(k/2) \binom{2m + 2}{m + 1}$	<i>if</i> $4m + 4 < k + 1$ <i>if</i> $4m + 4 > k + 1$
$H_{k,m,f}$	$2k$	
	$(k - 1)/2 \binom{2m + 4}{m + 2}$	<i>if</i> $4m + 8 < k$ <i>if</i> $4m + 8 > k$
$H_{k,m,h}$	2	
$H_{k,m,g}$	2^{k+1}	
	2	<i>if</i> $k \leq 2m + 1$ and k odd
	$2(k + 2)$	<i>if</i> $k < 2m + 1$ and k even

Lemma 5.4. [5] If G is one of the hexagonal tilings, then the chromatic number of G is given in the following table:

G	$H_{k,m,r}$	$H_{k,m,a}$	$H_{k,m,b}$	$H_{k,m,c}$	$H_{k,m,f}$	$H_{k,m,g}$	$H_{k,m,h}$
$\chi(G)$	2	2	2	3	3	3	2

Theorem 5.5. Let G be a graph Tutte equivalent to a hexagonal tiling H , then G is an hexagonal tiling.

Proof. By Theorem 2.1 and Lemma 5.1, we know that H has $2uv$ vertices for some u, v , $3uv$ edges and edge-connectivity equal to three. By Lemma 5.2, G has the same parameters, so G is cubic, has $2uv$ vertices and $3uv$ edges. Since H is a hexagonal tiling, H has girth six hence from the previous lemma, G has girth at least six.

In order to prove that G is a hexagonal tiling, we must show one more thing: that both graphs have the same number of cycles of length six (hexagons). Since G and H are Tutte equivalent, they have the same number of edge-sets with rank five and size six. We are going to prove that these sets are cycles of length six.

Let $C \subseteq E(G)$ be an edge-set with rank five and size six. If y is the number of connected components of C , then $5 = rg(C) = |V(C)| - y$. Suppose that C is not a cycle, since the girths of G and H are at least six we have $|V(C_i)| = |A(C_i)| + 1$ where C_i is a connected component of C . Hence $|V(C)| = y + 6$ and we obtain a contradiction. \square

Theorem 5.6. The graph $H_{k,m,0}$ is Tutte unique for $m \geq 2$ and $k \geq 3$.

Proof. Let G be a graph with $T(G; x, y) = T(H_{k,m,0}; x, y)$. By Theorem 2.1 and Lemma 5.4, G has to be isomorphic to one and only one of the following graphs: $H_{k',m',r}$, $H_{k',m',a}$, $H_{k',m',b}$, $H_{k',m',h}$. We prove that G is isomorphic to $H_{k',m',0}$ with $k = k'$ and $m = m'$ by assuming that G is isomorphic to each one of the previous graphs and by obtaining a contradiction in all the cases except in the aforementioned case. By Lemma 4.7, we obtain $l_{H_{k,m,0}} = l_G$ and the number of shortest essential cycles has to be the same in both graphs. The process we follow is to compare $H_{k,m,0}$ with all the graphs given in the previous list. In the cases in which the minimum length of essential cycles or the number of cycles of this minimum length are different, we would have that the pair of graphs compared are not Tutte equivalent. For the sake of brevity, we are only going to show those cases in which these quantities can coincide.

Case 1. Suppose $G \cong H_{k',m',r}$ with $r > 0$, $k < m + 1$ and $k' < m' + 1$.

As a result of Lemma 4.7, $k = k'$ and $m = m'$. Our aim is to prove that the number of edge-sets with rank $2(m + 1) - 1$ and size $2(m + 1)$ is different for each graph. This would lead to a contradiction since this number is the coefficient of $x^{2(m+1)-1}y^{2(m+1)}$ of the rank-size generating polynomial. If $H_{k,m,0}$ has $k \binom{m+1}{\lfloor (m+1)/2 \rfloor} + x$ essential cycles of length $2(m + 1)$, then $H_{k,m,r}$ has $x + \binom{m+1}{\lfloor (m+1)/2 - r \rfloor}$ such cycles, therefore if we show the existence of a bijection between the corresponding edge-sets with rank $2(m + 1) - 1$ and size $2(m + 1)$ that are not essential cycles, we would have proved the desired result.

For every α with $0 \leq \alpha \leq m - 1$, the set of edges that join a vertex (α, x) to $(\alpha + 1, x)$ is denoted E_α . Let A be an edge-set that is not an essential cycle with rank $2(m + 1) - 1$ and size $2(m + 1)$ in $H_{k,m,0}$. Define $s(A)$ as $\min\{\alpha \in [0, m - 1]; A \cap E_\alpha = \emptyset\}$. The minimum always exists for all edge-set contained in $E(H_{k,m,0})$ or $E(H_{k,m,r})$ that is not an essential cycle. For every α with $0 \leq \alpha \leq m - 1$ we define the bijection, φ_r between $\{A \subseteq E(H_{k,m,0}) | r(A) = 2(m + 1) - 1, |A| = 2(m + 1), s(A) = \alpha\}$ and $\{A \subseteq E(H_{k,m,r}) | r(A) = 2(m + 1) - 1, |A| = 2(m + 1), s(A) = \alpha\}$ as follows:

If $A \subset E(H_{k,m,0})$, $\varphi_r(A) = \cup\{\psi(\{(i,j), (i',j')\}); \{(i,j), (i',j')\} \in A\}$ where

$$\psi(\{(i,j), (i',j')\}) = \begin{cases} \{(i,j), (i',j')\} & \text{if } r+1 \leq i, i' \leq m \\ \{(i, j-2r), (i', j')\} & \text{if } i=0 \text{ and } i'=m \\ \{(i, j-2r), (i', j'-2r)\} & \text{if } i, i' \in [0, r] \end{cases}$$

Case 2. Suppose $G \cong H_{k',m',a}$ with $k < m+1$ and $k' < m'+1$.

The arguments to prove that both graphs do not have the same Tutte polynomial are analogous to those developed in the preceding case. We only want to specify two things. The first one is the number of essential cycles of length $2(m+1)$ in each graph, and the second one, is the bijection between the edge-sets with rank $2(m+1) - 1$ and size $2(m+1)$ that are not essential cycles contained in each graph.

In $H_{k,m,a}$ we have k exterior edges giving rise to displaced ladders $i \times j$ with $i = m+1-j$ and $0 \leq j \leq k < m+1$. On the other hand, in $H_{k,m,0}$ each of the k exterior edges determine a displaced ladder $(m+1)/2 \times (m+1)/2$ for odd m or $(m+2)/2 \times m/2$ if m is even. Hence, making use of the following property:

$$\binom{p}{n} < \binom{p}{\lfloor p/2 \rfloor} > \binom{p}{l} \text{ if } n < \lfloor p/2 \rfloor < l$$

the number of essential cycles of length $2(m+1)$ in $H_{k,m,a}$ is smaller than in $H_{k,m,0}$.

For every α we define φ_r between $\{A \subseteq E(H_{k,m,0}) | r(A) = 2m+1, |A| = 2m+2, s(A) = \alpha\}$ and $\{A \subseteq E(H_{k,m,a}) | r(A) = 2m+1, |A| = 2m+2, s(A) = \alpha\}$ as follows:

If $A \subset E(H_{k,m,0})$, $\varphi_r(A) = \cup\{\psi(\{(i,j), (i',j')\}); \{(i,j), (i',j')\} \in A\}$ where

$$\psi(\{(i,j), (i',j')\}) = \begin{cases} \{(i,j), (i',j')\} & \text{if } r+1 \leq i, i' \leq m \\ \{(i, 2k-1-j+3), (i', j')\} & \text{if } i=0 \text{ and } i'=m \\ \{(i, 2k-1-j+3), (i', 2k-1-j'+3)\} & \text{if } i, i' \in [0, r] \end{cases}$$

Case 3. Suppose $G \cong H_{k',m',b}$ with $k < m+1$ and $k' < m'+1$.

As in the previous case, we prove that the number of essential cycles of length $2(m+1)$ in $H_{k,m,b}$ is smaller than that in $H_{k,m,0}$.

For every α , the bijection φ_r between $\{A \subseteq E(H_{k,m,0}) | r(A) = 2m+1, |A| = 2m+2, s(A) = \alpha\}$ and $\{A \subseteq E(H_{k,m,b}) | r(A) = 2m+1, |A| = 2m+2, s(A) = \alpha\}$ is defined as follows:

If $A \subset E(H_{k,m,0})$, $\varphi_r(A) = \cup\{\psi(\{(i,j), (i',j')\}); \{(i,j), (i',j')\} \in A\}$ where

$$\psi(\{(i,j), (i',j')\}) = \begin{cases} \{(i,j), (i',j')\} & \text{if } r+1 \leq i, i' \leq m \\ \{(i, 2k-j), (i', j')\} & \text{if } i=0 \text{ and } i'=m \\ \{(i, 2k-j), (i', 2k-j')\} & \text{if } i, i' \in [0, r] \end{cases}$$

Case 4. Suppose $G \cong H_{k',m',a}$ with $k < m+1$ and $k' > m'+1$.

By Lemma 4.7, $k = m'+1$ and $k' = m+1$. We are going to prove that there are more edge-sets with rank $2k+2$ and size $2k+3$ in $H_{k,m,0}$ than in $H_{k',m',a}$. These sets can be classified into three groups:

- 1.- Essential cycles of length $2k+3$.
 - 2.- Normal edge-sets (they are edge-sets that do not contain any essential cycle).
 - 3.- Sets containing an essential cycle of length $2k$ and three more edges (Figure 10).
- (1) Since $H_{k,m,0}$ and $H_{k',m',a}$ are bipartite, they do not have any essential cycle of odd length.




Figure 10: a) Essential cycle of length $2k$ plus three edges in $H_{k,m,0}$ b) Essential cycle of length $2k = 2(m' + 1)$ plus three edges in $H_{k',m',a}$

(2) By Corollary 4.8 we know that $H_{k,m,0}$ and $H_{k',m',a}$ have the same number of normal edge-sets with rank $2k + 2$ and size $2k + 3$ that do not contain a cycle of length six. We are going to prove that the number of normal edge-sets with rank $2k + 2$ and size $2k + 3$ containing a cycle of length six is greater in $H_{k,m,0}$ than in $H_{k',m',a}$.

Again by Corollary 4.8, the number of edge-sets with rank $2k + 1$ and size $2k + 2$ containing a hexagon is the same in both graphs, call it s_{2k+1} . Add one edge to each of these sets in order to obtain a set with rank $2k + 2$ and size $2k + 3$. This set can be one of the following types depending on which edge we are adding:

- (A) A normal edge set with rank $2k + 2$.
- (B) A normal edge set containing two non essential cycles and having rank $2k + 1$.
- (C) A edge set containing an essential cycle of length $2k$ and a cycle of length six.

Let $A(H)$, $B(H)$ and $C(H)$ (where H is $H_{k,m,0}$ or $H_{k',m',a}$) be the number of edge-sets in H that belong to the groups A, B and C respectively. $A(H) + B(H) + C(H)$ is the number of possibilities of adding one edge to the edge-sets with size $2k + 2$ containing a hexagon and such that the added edge does not belong to a fixed hexagon.

$$A(H) + B(H) + C(H) = \frac{s_{2k+1}(3k(m+1) - 2k - 2)}{2k + 3 - 6}$$

Now, we are going to show that $(2k - 3)B(H) = \sum_{B \in B(H)} (2k + 3 - \delta(B))$, where $\delta(B)$ is the number of edges of B which do not belong to all cycles of length six in B . In order to obtain an edge-set $B \in B(H)$ we add one edge to an edge-set with size $2k + 2$ that contains one hexagon, forming a new contractible cycle. Since the cycle is contractible, we can consider that the added edge has to belong to a new hexagon. Hence:

$$B(H) = \frac{\sum_{B \in B(H)} (2k + 3 - \delta(B))}{2k + 3 - 6}$$

If we apply Corollary 4.8 to every $B \in B(H)$, the number of normal edge-set of size $2k + 3 - \delta(B)$ is equal in $H_{k,m,0}$ and $H_{k,m,a}$ then:

$$\sum_{B \in B(H_{k,m,0})} (2k + 3 - \delta(B)) = \sum_{B \in B(H_{k',m',a})} (2k + 3 - \delta(B))$$

Altogether and by taking in account that $C(H_{k,m,0}) = 0$ and $C(H_{k',m',a}) \neq 0$ (Figure 10) we have proved that $A(H_{k,m,0}) > A(H_{k',m',a})$.

(3) In $H_{k,m,0}$, every essential cycle of length $2k$ plus three edges has rank $2k+2$ (Figure 10a), but in $H_{k',m',a}$ there are essential cycles in which if we add three edges we obtain sets with rank $2k+1$ (Figure 10b). By hypothesis, both graphs have the same number of shortest essential cycles therefore the number of edge-sets in this case is greater in $H_{k,m,0}$ than in $H_{k',m',a}$.

Cases 5 and 6 If $G \cong H_{k',m',b}$ with $k < m+1$ and $k' > m'+1$ or $G \cong H_{k',m',h}$ with $k < m+1$ we use the same arguments as those in the previous case to prove that $H_{k,m,0}$ has more edge-sets with rank $2k+2$ and size $2k+3$ than $H_{k',m',b}$ and $H_{k',m',h}$. \square


6 Concluding Remarks

We have proved that the toroidal hexagonal tiling, $H_{k,m,0}$ is Tutte unique. Since the results given in Sections 3 and 4 are proved for all hexagonal tilings, it seems that all hexagonal tilings are Tutte unique. To verify this, we would have to do all cross-checkings between any two of these graphs as was done in Section 5. This study has been done for locally grid graphs in [7].


The technique developed in this paper can also be applied to locally C_6 graphs. This would follow from the local orientation existing in these graphs, based on the minor relationship described in [5] between locally grid graphs and locally C_6 graphs.

References


- [1] A. Altshuler, *Hamiltonian Circuits in some maps on the torus*, Discret Math. **1**(1972), 299-314.
- [2] A. Altshuler, *Construction and enumeration of regular maps on the torus*, Discret Math. **4**(1973), 201-217.
- [3] S. Fisk, *Geometric coloring theory*, Advances in Math. **24**(1977), 298-340.
- [4] S. Fisk, *Variations on coloring, surfaces and higher dimensional manifolds*, Advances in Math. **25**(1977), 226-266.
- [5] D. Garijo, I. Gitler, A. Márquez and M.P. Revuelta, *Hexagonal tilings and Locally C_6 graphs*, Preprint, 2004.
- [6] D. Garijo, *Polinomio de Tutte de Teselaciones Regulares*, Thesis (2004).
- [7] D. Garijo, A. Márquez and M.P. Revuelta, *Tutte Uniqueness of Locally Grid Graphs*, Ars Combinatoria (to appear) (2005).
- [8] A. Márquez, A. de Mier, M. Noy, M.P. Revuelta, *Locally grid graphs:classification and Tutte uniqueness*, Discr. Math. **266** (2003), 327-352.
- [9] A. de Mier, M. Noy, *On graphs determined by their Tutte polynomials*, Graphs Combin. **20** (2004), no.1, 105-119.
- [10] S. Negami, *Uniqueness and faithfulness of embedding of toroidal graphs*, Discrete Math. **44**(1983), no.2, 161-180.
- [11] S. Negami, *Classification of 6-regular Klein bottle graphs*, Res. Rep. Inf. Sci. T.I.T.A-96 (1984).
- [12] C. Thomassen, *Tilings of the Torus and the Klein Bottle and vertex-transitive graphs on a fixed surface*, Trans. Amer. Math. Soc. **323**(1991), no.2, 605-635.
- [13] W.T. Tutte *Graph Theory*, Addison Wesley, California (1984).




a)



b)



a)



b)