

Effect of Coenzyme Q₁₀ on Psychopathological Symptoms in Fibromyalgia Patients

Elísabet Alcocer-Gómez,^{1,2} Ognjen Culic,³ José M. Navarro-Pando,⁴ José A. Sánchez-Alcázar² & Pedro Bullón¹

1 Research Laboratory, Oral Medicine Department, University of Sevilla, Sevilla, Spain

2 Centro Andaluz de Biología del Desarrollo (CABD), Universidad Pablo de Olavide-CSIC-Junta de Andalucía and Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), ISCIII, Sevilla, Spain

3 Faculty of Pharmacy and Biochemistry, University of Zagreb, Zagreb, Croatia

4 Unidad de Reproducción Humana y Cirugía Endoscópica, Instituto para el Estudio de la Biología de la Reproducción Humana (INEBIR), Seville, Spain

Correspondence

E. Alcocer-Gómez, Research Laboratory, Oral Medicine Department, Universidad de Sevilla, C/Avicena s/n 41009 –Sevilla, Spain.

Tel.: +34-954-481120;

Fax: +34-954-486784;

E-mail: agelisabet@gmail.com

Received 20 September 2016; revision 24

November 2016; accepted 25 November 2016

doi: 10.1111/cns.12668

Fibromyalgia (FM) is a common chronic pain syndrome accompanied by a myriad of variable physical and psychopathological symptoms such as fatigue, muscle stiffness, sleep disorders, morning tiredness, cognitive complaints, as well as depression and anxiety [1]. Despite the fact that it affects up to 5% of the general population worldwide, its pathogenic mechanism remains elusive. Recently, the hypothesis that oxidative stress and mitochondrial dysfunction are important events in the pathogenesis of FM [2] was proposed. Furthermore, coenzyme Q₁₀ (CoQ₁₀) deficiency has been described in patients with FM [3]. CoQ₁₀ plays a critical role in the mitochondrial ATP production and cellular metabolism. CoQ₁₀ also regulates mitochondrial uncoupling proteins, mitochondrial permeability transition pore, and ROS production [2]. Recently, preliminary data showed an interesting improvement in clinical symptoms in patients with FM after oral supplementation with CoQ₁₀ [4,5], including the control of the depressive symptom with the regulation of the serotonergic system [6].

In addition to biological and clinical parameters, other psychopathological events have been involved in the pathophysiology of FM. Psychological factors, a high prevalence of depression and anxiety symptoms have been widely reported [7]. Following our earlier work about the therapeutic effects of CoQ₁₀ on FM in a clinical trial (ISRCTN 21164124) [5], here we show the effect of 40 days of CoQ₁₀ versus placebo supplementation in psychopathological profiles from patients with FM using a multidimensional psychological screening instrument, namely the Symptom Checklist-90-R' (SCL-90-R). The study protocol was reviewed and approved by the Ethical Committee of the University of Sevilla. All the participants to the study gave their written

informed consent before initiating the study. This study was carried out in compliance with the Declaration of Helsinki, and all the International Conferences on Harmonisation and Good Clinical Practice Guidelines. Twenty patients diagnosed with FM were distributed in a clinical trial as described in reference 5. Data in tables are given as means \pm SD. Data between different groups were analyzed statistically using ANOVA on Ranks with SigmaPlot and SigmaStat statistical software (SPSS for Windows, 19, 2010, SPSS, Inc). A value of $P < 0.05$ was considered significant. To compare the trial results from patients treated with CoQ₁₀ or placebo, a two-way variance (ANOVA) analysis was used. The patients were diagnosed with FM by exclusion of other diseases and syndromes, and in accordance with the American College of Rheumatology criteria. Subjects were randomized in a double-blind fashion, according to a 1:1 ratio, to CoQ₁₀ or placebo. Ten subjects (age: 44.3 ± 9.7 years) received CoQ₁₀ (Pharma Nord, Vejle, Denmark) in soft gel capsules for 40 days (300 mg/day CoQ₁₀ divided into three daily doses), while another group of ten subjects (age: 55 ± 5 years) with similar characteristics received a matching placebo (Table 1). After 40 days, no changes were observed about BMI in the patients (baseline: 27.6 ± 4 kg/m² vs. placebo: 27.3 ± 4.8 kg/m² or CoQ₁₀ group: 26 ± 4.8 kg/m²). However, important molecular changes were induced by CoQ₁₀ after treatment such as increment in mitochondrial biogenesis and antioxidants gene expression and reduction in inflammation accompanied by an improvement in the clinical symptoms determined by Fibromyalgia Impact Questionnaire (FIQ), Pittsburgh Sleep Quality Index (PSQI) and tender points [5] and corroborated in other studies [4,6]. After evaluating the effect of CoQ₁₀ on the

Table 1 Characteristics of the patients with FM pre- and posttreatments

Parameter	Baseline N = 20	Placebo N = 10	CoQ ₁₀ N = 10
Age (years)	49 ± 9	55 ± 5	44 ± 9.7
Duration of Diseases (years)	6.9 ± 3.3	6.6 ± 3	7.1 ± 3.8
Systole BP, mm Hg	134 ± 19.4	136 ± 23.5	126 ± 21.7
Diastole BP, mm Hg	76.9 ± 9.6	78 ± 13.7	80.5 ± 12.5
Pulse rate, bpm	74.2 ± 10.3	69 ± 6.8	78.8 ± 14.5
Weight (Kg)	70.5 ± 11.2	73.2 ± 13.8	67.8 ± 13.7
BMI (kg/m ²)	27.6 ± 4	27.3 ± 4.8	26 ± 4.8

BMI, body mass index. Values are means ± SD.

psychopathological symptoms, a clinically significant improvement was observed in all subscores from SCL-90-R, being statistically significant for the reduction in interpersonal sensitivity ($P < 0.001$), depression ($P < 0.001$), anxiety ($P < 0.001$), hostility ($P < 0.001$), and psychoticism ($P < 0.001$) items, and a moderate statistical significance in somatization ($P < 0.05$) and obsessive-compulsive ($P < 0.05$; Table 2).

Previously, SCL-90-R has been used to determinate the emotional distress and show psychological distress and maladaptive emotional responses in these patients [8]. Interestingly, we show a very important improvement in psychological events highly implicated in the pathophysiology of FM. Indeed, CoQ₁₀ has been implicated in one of the most important psychological symptom in FM, depression, and it has been recommended the oral use in these patients [9]. This has been argued because of the high levels of oxidative stress and mitochondrial damage [9]. Mitochondrial dysfunction and the resultant reduced bioenergetic production have been involved in somatization and depression [9,10], two of the symptoms highly shown by patients with FM. So, our preliminary data show a possible biological implication of the psychological symptoms in FM. On the other hand, it has been known since early on that patients with CoQ₁₀ benefit from oral CoQ₁₀ supplementation [9,10]. According to this, CoQ₁₀ has been shown to be a potential drug candidate in the treatment of FM for different reasons. First, it is a mitochondrial cofactor with the potential to improve mitochondrial functions and respiration. Second, CoQ₁₀ is a

Table 2 Symptom checklist revised changes in patients with FM pre- and posttreatment with CoQ₁₀

Parameter	Baseline N = 20	Placebo N = 10	CoQ ₁₀ N = 10
Somatization	2.06 ± 0.8	1.95 ± 0.6	1.39 ± 0.8*
Obsessive– compulsive	1.69 ± 0.9	1.58 ± 0.7	1 ± 0.2*
Interpersonal sensitivity	1.59 ± 0.6	1.51 ± 0.3	0.4 ± 0.2***
Depression	1.79 ± 0.3	1.73 ± 0.4	0.8 ± 0.2***
Anxiety	1.9 ± 0.6	1.84 ± 0.5	0.7 ± 0.2***
Hostility	1.75 ± 0.3	1.9 ± 0.5	0.4 ± 0.2***
Phobic anxiety	0.68 ± 0.5	0.78 ± 0.4	0.27 ± 0.2*
Paranoid ideation	0.78 ± 0.1	0.68 ± 0.4	0.2 ± 0.07**
Psychoticism	0.69 ± 0.3	0.55 ± 0.5	0.2 ± 0.05***

Values are means ± SD. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ between pre- and posttreatment.

powerful free radical scavenger that can mitigate oxidative stress [5]. Now, our study has shown a new perspective about the treatment in patients with FM using CoQ₁₀. Because psychopathological symptoms and psychological distress are highly involved in the pathophysiology of FM and modulate the perception of the disease from the patients, the effect of CoQ₁₀ in this clinical dimension proposes a biological and psychological approach from the same treatment. Furthermore, this study has appointed a new path toward understanding of FM.

Further analysis involving more patients in double-blind placebo-controlled clinical trials is required to confirm these observations. Indeed, our research group is currently working in this direction, on the basis of the conclusions of the exploratory work discussed in this article.

Acknowledgments

This work has been supported by Grupo de Investigacion Junta de Andalucía CTS113, Consejería de Salud of the Junta de Andalucía (PI-0036-2014).

Conflict of Interest

The authors declare no conflict of interest.

References

- English B. Neural and psychosocial mechanisms of pain sensitivity in fibromyalgia. *Pain Manag Nurs* 2014;**15**:530–538.
- Cordero MD, Díaz-Parrado E, Carrión AM, et al. Is inflammation a mitochondrial dysfunction-dependent event in Fibromyalgia? *Antioxid Redox Signal* 2013;**18**:800–807.
- Cordero MD, Moreno-Fernández AM, deMiguel M, et al. Coenzyme Q10 distribution in blood is altered in patients with fibromyalgia. *Clin Biochem* 2009;**42**:732–735.
- Miyamae T, Seki M, Naga T, et al. Increased oxidative stress and coenzyme Q10 deficiency in juvenile fibromyalgia: Amelioration of hypercholesterolemia and fatigue by ubiquinol-10 supplementation. *Redox Rep* 2013;**18**:12–19.
- Cordero MD, Alcocer-Gómez E, de Miguel M, et al. Can coenzyme Q10 improve clinical and molecular parameters in fibromyalgia? *Antioxid Redox Signal* 2013;**19**:1356–1361.
- Alcocer-Gómez E, Sánchez-Alcázar JA, Cordero MD. Coenzyme q10 regulates serotonin levels and depressive symptoms in fibromyalgia patients: Results of a small clinical trial. *J Clin Psychopharmacol* 2014;**34**:277–278.
- Di Tella M, Castelli L, Colonna F, et al. Theory of mind and emotional functioning in fibromyalgia syndrome: An investigation of the relationship between social cognition and executive function. *PLoS ONE* 2015;**10**:e0116542.
- Glazer Y, Cohen H, Buskila D, et al. Are psychological distress symptoms different in fibromyalgia patients compared to relatives with and without fibromyalgia? *Clin Exp Rheumatol* 2009;**27**:S11–S15.
- Moylan S, Berk M, Dean OM, et al. Oxidative & nitrosative stress in depression: Why so much stress? *Neurosci Biobehav Rev* 2014;**45**:46–62.
- Gardner A, Boles RG. Beyond the serotonin hypothesis: Mitochondria, inflammation and neurodegeneration in major depression and affective spectrum disorders. *Prog Neuropsychopharmacol Biol Psychiatry* 2011;**35**:730–743.