

Research Article

Effects of the Fourth Ventricle Compression in the Regulation of the Autonomic Nervous System: A Randomized Control Trial

Ana Paula Cardoso-de-Mello-e-Mello-Ribeiro,¹ Cleofás Rodríguez-Blanco,²
Inmaculada Riquelme-Agulló,^{3,4} Alberto Marcos Heredia-Rizo,²
François Ricard,⁵ and Ángel Oliva-Pascual-Vaca²

¹Madrid Osteopathic School, Avenida Dr. Heitor Penteado 815, Taquaral, 13075-185 Campinas, SP, Brazil

²Department of Physical Therapy, Faculty of Nursing, Physiotherapy and Podiatry, C/Avicena s/n, University of Sevilla, 41009 Sevilla, Spain

³Department of Nursing and Physiotherapy, University of the Balearic Islands, Carrera de Valldemossa, km 7'5, 07122 Palma, Spain

⁴University Institute of Health Sciences Research (IUNICS-IdISPa), University of the Balearic Islands, Carrera de Valldemossa, km 7'5, 07122 Palma, Spain

⁵Madrid Osteopathic School, C/San Félix de Alcalá 4, Alcalá de Henares, 28807 Madrid, Spain

Correspondence should be addressed to Ángel Oliva-Pascual-Vaca; angeloliva@us.es

Received 26 December 2014; Revised 1 February 2015; Accepted 1 February 2015

Academic Editor: Anwar-Ul-Hassan Gilani

Copyright © 2015 Ana Paula Cardoso-de-Mello-e-Mello-Ribeiro et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction. Dysfunction of the autonomic nervous system is an important factor in the development of chronic pain. Fourth ventricle compression (CV-4) has been shown to influence autonomic activity. Nevertheless, the physiological mechanisms behind these effects remain unclear. **Objectives.** This study is aimed at evaluating the effects of fourth ventricle compression on the autonomic nervous system. **Methods.** Forty healthy adults were randomly assigned to an intervention group, on whom CV-4 was performed, or to a control group, who received a placebo intervention (nontherapeutic touch on the occipital bone). In both groups, plasmatic catecholamine levels, blood pressure, and heart rate were measured before and immediately after the intervention. **Results.** No effects related to the intervention were found. Although a reduction of norepinephrine, systolic blood pressure, and heart rate was found after the intervention, it was not exclusive to the intervention group. In fact, only the control group showed an increment of dopamine levels after intervention. **Conclusion.** Fourth ventricle compression seems not to have any effect in plasmatic catecholamine levels, blood pressure, or heart rate. Further studies are needed to clarify the CV-4 physiologic mechanisms and clinical efficacy in autonomic regulation and pain treatment.

1. Introduction

Dysfunction of the autonomic nervous system is considered to be an important factor in the development of chronic pain [1]. Studies using muscle blood flow, heart rate, and blood pressure as outcome measures have suggested an excessive sympathetic activation in patients with different pain conditions such as fibromyalgia [2, 3], musculoskeletal and myofascial pain [1, 4], and chronic pelvic pain [5] and patients who had undergone abdominal surgery [6]. The relationship between sympathovagal balance and psychological distress has also been proven in children with abdominal pain,

irritable bowel syndrome [7, 8], and complex regional pain syndrome [9, 10].

Catecholamines are important hormones which regulate ANS activity in both central and peripheral sympathetic nerve endings [11]. Studies with animals have suggested that chronic adrenergic stimulation or impaired epinephrine homeostasis may contribute to the pathophysiological mechanisms of pain syndromes [12, 13]. On the other hand, dopamine has proved to have antinociceptive effects both in animals and in humans [14–16] and dopaminergic neurotransmission has been shown to be affected in chronic pain syndromes [17, 18].



FIGURE 1: Physiotherapist's hand position during the procedure. (a) CV-4 technique, (b) placebo technique.

Fourth ventricle compression (CV-4) is a cranial manipulation technique aiming at influencing on brain and cranial nerve function [19]. Some authors have suggested that CV-4 may produce changes in the regulation of the autonomic nervous system (ANS). For instance, Cutler et al. [20] reported muscle sympathetic nerve activity after applying the CV-4 technique. The use of CV-4 technique has also been purported to modify heart rate, blood pressure, blood flow velocity [21, 22], and cerebral tissue oxygenation [23]. Moreover, CV-4 has been observed to produce an increment of electroencephalography alpha absolute power [19], reduce sleep latency [20], and decrease anxiety [21] in healthy subjects. In addition, CV-4 has been shown to be effective in pain relief in tension type headaches [24]. Due to the established interactions between the autonomic and nociceptive systems, the potential effects of CV-4 on the autonomic regulation could represent a novel approach to control the development or maintenance of pain. Nevertheless, the physiological mechanisms undergoing CV-4 autonomic regulation remain unclear. The aim of the present study was to assess the immediate effects of the CV-4 technique compared to sham placebo intervention in plasma catecholamine levels, heart rate, and blood pressure on healthy adults.

2. Methods

2.1. Participants. Forty healthy adults (19 males and 21 females; age range = 18–33 years) were recruited in the Faculty of Physiotherapy of UNILUS (Santos, SP, Brazil). A member of the research group did a personal interview with every volunteer informing them of the objectives of the study and the procedure. The established exclusion criteria were (1) any local or systemic pathologic condition (recent fracture in the cranial base, osteitis, tumor, encephalopathy, stroke, traumatic brain injury, depression, or asthma crisis), (2) arterial hypotension, low heart rate, or parasympathetic state, (3) pregnant women or women in menstrual state on the evaluation day, (4) tobacco or drug consumption, and (5) having received a manual therapy procedure in the last month. Subjects participating in the study were instructed not to consume coffee, tea, chocolate, vanilla, fruit or fruit derivatives, soft

drinks, or alcohol for 4 hours before data collection. Subjects who reported having consumed any of these foods were excluded from the study ($n = 1$).

The study was approved in accordance with the principles of the Declaration of Helsinki by the Ethics Committee of Hospital Guilherme Álvaro de Santos, SP, Brazil.

2.2. Procedure. Intervention and assessments were performed in a university-based physiotherapy research clinic at Lusíada University (UNILUS, Santos, SP, Brazil). The assessments and intervention were performed between 7 and 9.30 a.m., in a room with a temperature of 21–23°C.

Participants were assigned randomly to one of the treatment groups: intervention group and control group. The intervention group received the CV-4 technique for 10 minutes. The CV-4 procedure was administered by an experienced physiotherapist and member of the research group according to the standards established previously in the literature [19]. The participant lay down in a supine position. The physiotherapist, situated behind the participant's head, made a slight approximation of the occipital squama lateral angles towards the posterior occipital convexity, while taking the cranium into extension. The traction was maintained until a motionless state was perceived in the cranial pulse and released when a perception of movement was noted. The procedure was repeated until the 10-minute session time expired (Figure 1(a)).

The control group underwent a bilateral nontherapeutic contact on the occipital bone for 10 minutes. The participant lay down in a supine horizontal position. The physiotherapist, situated behind the participant's head, overlapped their hands touching the occipital bone and provided random slight touch of random duration until the end of the 10-minute session time. Figure 1(b) displays physiotherapist's hands' position in the control procedure.

2.3. Assessments. Both groups (intervention and control group) underwent assessments of blood pressure, heart rate, and blood test to determine plasmatic catecholamine levels before and 5 minutes after the procedure. Blood pressure and heart rate assessments were performed by the same evaluator

TABLE 1: Mean and SD of catecholamine plasmatic levels, blood pressure, and heart rate for both groups (intervention and control) in the preintervention and postintervention.

| | Intervention group | | Control group | |
|--------------------------|--------------------|------------------|-----------------|------------------|
| | Preintervention | Postintervention | Preintervention | Postintervention |
| Dopamine | 53.73 (20.15) | 53.63 (15.62) | 36.85 (14.47) | 52.35 (19.13) |
| Norepinephrine | 119.68 (39.55) | 114.00 (44.17) | 113.85 (39.64) | 87.65 (37.63) |
| Epinephrine | 43.58 (20.16) | 49.26 (16.27) | 53.55 (14.94) | 52.15 (22.13) |
| Systolic blood pressure | 119.74 (9.50) | 116.53 (8.08) | 121.55 (13.12) | 117.00 (15.52) |
| Diastolic blood pressure | 74.00 (9.18) | 74.26 (7.12) | 81.48 (10.25) | 80.50 (12.48) |
| Heart rate | 72.39 (6.95) | 70.40 (6.93) | 80.95 (13.39) | 77.35 (7.99) |

and the blood test was performed by a second evaluator. Both evaluators entered the room only during the assessment time and were not aware which intervention was done on the participant. Before the intervention, the subject was instructed to sit on a chair in a comfortable position for 10 minutes; after this time, evaluators entered the room and started the assessment protocol. Evaluations were made of blood pressure, heart rate, and blood test, following this order. After the intervention, the participant was instructed to relax in a seated position for 5 minutes before postintervention evaluations were performed.

Plasmatic catecholamines levels were obtained by an analysis of every participant's blood sample. Epinephrine, norepinephrine, and adrenaline were measured according to the methods described by Sealey [25].

Blood pressure was determined by the use of a mercury sphygmomanometer (UNITEC). The assessment was performed 3 times in each arm, with 1-minute interval between them. The average of all measurements was made to obtain systolic and diastolic blood pressure scores.

Heart rate was assessed by stethoscope (Littmann 3M) that was placed on the participant's thorax, over the left heart ventricle. Heart rate was assessed three times with an interval of 1 minute between the measurements. An average of the three measurements was computed to obtain the heart rate score for study purposes.

2.4. Statistical Analysis. Analyses of variance (ANOVA) were performed to assess changes in heart rate, blood pressure, and plasmatic catecholamine levels, with GROUP (intervention versus control) as between-subjects factor and TIME (preintervention versus postintervention) as within-subjects factor. Post hoc analyses were performed using post hoc Bonferroni corrected test for multiple comparisons. Analysis was performed with STATA version 7.0. (StataCorp 2001, Stata Statistical Software: Release 7.0, College Station, TX: Stata Corporation, Texas, USA). Significance level was set at $P < .05$.

3. Results

Nineteen healthy adults were included in the intervention group (7 females, mean age = 23.58 ± 4.19 years) and twenty healthy adults were in the control group (14 females, mean age = 20.05 ± 1.96 years). Table 1 displays the values of

plasmatic catecholamines, blood pressure, and heart rate for both groups before and after the procedure.

Dopamine plasmatic levels displayed a significant interaction GROUP \times TIME ($F(1,37) = 7.16, P < .05$). The control group had lower dopamine levels than the intervention group in the preintervention assessment (post hoc pairwise comparison $P < .01$). Only the control group showed a statistically significant increase in dopamine levels when comparing postintervention scores to preintervention values (post hoc pairwise comparison $P < .01$), whereas no significant changes were seen in the intervention group (post hoc pairwise comparison $P > .82$). Moreover, a main effect TIME ($F(1,37) = 6.97, P < .05$) was found.

Norepinephrine levels witnessed a significant main TIME effect ($F(1,37) = 4.57, P < .05$), showing a decrease of plasmatic norepinephrine levels when comparing the mean score changes from postintervention to preintervention. No significant statistical effects were found for the main effect GROUP ($F(1,37) = 2.34, P = .14$) or the interaction GROUP \times TIME ($F(1,37) = 1.89, P = .177$). Nevertheless, pairwise comparisons found that the control group showed lower norepinephrine plasmatic levels in the postintervention assessment, compared to the preintervention values ($P < .05$), whereas no changes were found in the intervention group ($P = .60$). Even though both groups had similar norepinephrine baseline levels ($P = .65$), pairwise comparisons after intervention showed a nonsignificant tendency for the control group to have lower norepinephrine levels than the intervention group ($P = .052$).

Likewise, no significant main effects were found for epinephrine (TIME ($F(1,37) = .32, P = .36$); GROUP ($F(1,37) = 2.51, P = .12$)). Although the interaction GROUP \times TIME was not statistically significant ($F(1,37) = .87, P = .58$), post hoc pairwise comparisons showed lower baseline epinephrine levels for the intervention group than for the control group ($P < .05$), whereas no significant differences were found in the postintervention evaluation ($P = .65$). Pairwise comparisons did not show significant changes between the preintervention and postintervention levels for any of the study groups ($P > .30$ in all cases).

Systolic blood pressure had a significant main TIME effect ($F(1,37) = 10.02, P < .01$), showing a reduction of heart rate after the procedure. No statistically significant effects were found for the main factor GROUP ($F(1,37) = .10, P = .76$) or the interaction GROUP \times TIME ($F(1,37) = .30, P = .59$). Although pairwise comparisons did not show significant

differences between the control and the intervention group neither before intervention or after intervention ($P > .63$ in all cases), the control group showed a statistically significant decrease of the systolic blood pressure after the intervention ($P < .05$), whereas the intervention group showed only a nonsignificant tendency to decrease ($P = .075$).

With regard to diastolic blood pressure, the findings achieved a significant main effect GROUP ($F(1,37) = 5.30, P < .05$), revealing lower diastolic blood pressure in the intervention group than in the control group. No statistically significant effects were found for the main factor TIME ($F(1,37) = .09, P = .76$) or the interaction GROUP \times TIME ($F(1,37) = .28, P = .60$). Although pairwise comparisons did not show significant changes of the diastolic blood pressure after the intervention in any of the study groups ($P > .56$ in all cases), the control group showed higher diastolic blood pressure levels than the intervention group in the preintervention assessment ($P < .05$), whereas only a nonsignificant tendency was found for the postintervention comparison ($P = .065$).

There were significant main effects GROUP ($F(1,37) = 7.81, P < .01$) and TIME ($F(1,37) = 7.09, P < .05$) for heart rate. A lower heart rate was observed in the intervention group compared to the control group and the score changes also showed lower heart rate values in the postintervention evaluation compared to baseline scores. Although the interaction GROUP \times TIME did not show any statistical significance ($F(1,37) = .58, P = .45$), pairwise comparisons showed that only the control group decreased the heart rate after the intervention ($P < .05$). On the contrary, no changes were found in the intervention group ($P = .19$). Nevertheless, the control group showed higher heart rate than the intervention group both before and after the intervention ($P < .05$ in all cases).

4. Discussion

The present study aimed to evaluate the effects of the CV-4 technique in the regulation of the autonomic nervous system (ANS). More specifically, the objective of the study was to evaluate changes in plasma catecholamine levels, blood pressure, and heart rate in healthy adults that were randomly assigned to either an intervention group, who received the CV-4, or a control group, who underwent a placebo intervention. The present findings confirmed a reduction of norepinephrine, systolic blood pressure, and heart rate after the intervention, in both study groups. Likewise, only subjects of the control group showed an increase of dopamine blood levels after intervention.

Our results are in contrast with other studies reporting an influence of the CV-4 technique in autonomic-related parameters, such as heart rate, blood pressure, blood flow velocity, the electroencephalography alpha power, and the muscle sympathetic nerve activity [19–22]. The previously mentioned previous studies have used several tools such as microneurographic recording, skin conductance, skin temperature, or heart rate variability assessments to test the effects of the CV-4 technique in the ANS. However, to the best of authors' knowledge, this is the first study to assess the immediate effects of CV-4 in the plasmatic catecholamine levels. The present study has addressed this issue in a different way and,

therefore, the observed findings may be different. Previous research has reported that heart rate variability can be unstable in some situations and must be cautiously interpreted to characterize sympathovagal interaction [26, 27]. Other studies have not found relationships between heart rate variability and plasmatic norepinephrine levels [28]. In this sense, our design could have included catecholamines' analysis to better indicate ANS activity. It could also be argued that the lying-down position and the relaxation provided by the procedure in both groups could have had more influence in the autonomic regulation than the CV-4 technique per se. This would be in accordance with studies reporting that increments in the head-up tilt angle have been shown to increase plasma epinephrine and heart rate [29]. Plasma catecholamine activity has also been concluded to be influenced by stressors [30], and we must also take into account the influence of the procedural setting on the sample's ANS. On the other hand, previous research has also suggested that the duration of manual intervention may have a significant impact on the autonomic response [31]. According to this, higher differences between the groups may have emerged if longer procedure times had been used. This is an important area for further research, especially when no clinical present evidence has established a proper duration for the CV-4 technique. The high standard deviations found in the catecholamine levels in the present study subjects may also explain why more significant differences have not been found. Finally, our study was performed with a healthy population; the application of CV-4 in individuals with chronic pain or autonomic dysfunction or in an experimental condition using pain stimuli could have displayed different results.

In contrast with previous research, in our study, the application of the CV-4 technique was not directly related to variation in heart rate or blood pressure [21]. It has been stated that the heart rate does not quantitatively reflect the degree of autonomic activation and Ng et al. [32] have reported changes in sympathetic nerve activity without changes in heart rate or blood pressure. Mannelli et al. [33] have related changes in catecholamine plasmatic levels to the modification of the low-frequency/high-frequency ratio, which would disregard variations in changes in heart rate. In our study, both groups experienced a reduction of norepinephrine levels after intervention. Although the present findings may indicate a tendency to a higher reduction of diastolic blood pressure and heart rate in the intervention group, the absence of significant interactions does not allow us to maintain the fact that these effects are related to the technique.

Our study has some limitations that must be taken into account for the adequate interpretation of the results. Firstly, previous research has reported significant changes in the ANS after manual techniques and placebo conditions when compared to a control group with no intervention [31]. In our study the absence of a control group with no intervention does not allow us to conclude if the mere hand contact in a lying position is sufficient to elicit a different autonomic response than in a control condition. Hence, we must question whether the proposed placebo intervention is not really a powerful intervention itself and it should not be considered as a sham intervention. Secondly, catecholamine plasmatic

concentration can be influenced by factors such as age, sex, nutritional condition, emotional factors, or heat exposure [34, 35]. Although some factors have been controlled in the present research, trying to homogenize the groups and procedure conditions, some other parameters, such as the nutritional or emotional conditions, have not been controlled. The statistical analyses to observe the influence of sex in the outcome measures displayed only main statistical effects not related to the technique application. Scientific literature concludes contradictory results on the topic, going from studies reporting gender variations in catecholamine levels during rest, exercise, or stressors [35, 36] to studies not reporting gender differences [37, 38]. Our results seem to be in accordance with the latter authors, although further studies with specific designs and bigger samples are needed to control all the possible influencing variables. Moreover, heart rate measurement would have been more accurate with the use of ECG.

5. Conclusions

The CV-4 technique does not seem to have an immediate influence on catecholamine plasmatic levels. This finding is relevant as manual techniques may have concurrent effects on pain and on the ANS activity. In this sense, manual techniques likely to regulate sympathetic activity may be of paramount importance for providing analgesia and reducing pain sensitivity. Further research on the topic is warranted to better understand the physiological mechanisms underlying the CV-4 technique and its potential effects in the autonomic and nociceptive systems.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] D. M. Hallman and E. Lyskov, "Autonomic regulation, physical activity and perceived stress in subjects with musculoskeletal pain: 24-hour ambulatory monitoring," *International Journal of Psychophysiology*, vol. 86, no. 3, pp. 276–282, 2012.
- [2] S. M. Mostoufi, N. Afari, S. M. Ahumada, V. Reis, and J. L. Wetherell, "Health and distress predictors of heart rate variability in fibromyalgia and other forms of chronic pain," *Journal of Psychosomatic Research*, vol. 72, no. 1, pp. 39–44, 2012.
- [3] L. R. Oliveira, V. U. de Melo, F. N. Macedo et al., "Induction of chronic non-inflammatory widespread pain increases cardiac sympathetic modulation in rats," *Autonomic Neuroscience: Basic and Clinical*, vol. 167, no. 1-2, pp. 45–49, 2012.
- [4] C. M. Eze-Nliam, P. J. Quartana, A. M. Quain, and M. T. Smith, "Nocturnal heart rate variability is lower in temporomandibular disorder patients than in healthy, pain-free individuals," *Journal of Orofacial Pain*, vol. 25, no. 3, pp. 232–239, 2011.
- [5] D. S. Cho, J. B. Choi, Y. S. Kim et al., "Heart rate variability in assessment of autonomic dysfunction in patients with chronic prostatitis/chronic pelvic pain syndrome," *Urology*, vol. 78, no. 6, pp. 1369–1372, 2011.
- [6] L. H. Chang, T. C. Ma, S. L. Tsay, and G. P. Jong, "Relationships between pain intensity and heart rate variability in patients after abdominal surgery: a Pilot Study," *Chinese medical journal*, vol. 125, no. 11, pp. 1964–1969, 2012.
- [7] M. Jarrett, M. Heitkemper, D. Czyzewski, L. Zeltzer, and R. J. Shulman, "Autonomic nervous system function in young children with functional abdominal pain or irritable bowel syndrome," *The Journal of Pain*, vol. 13, no. 5, pp. 477–484, 2012.
- [8] E. Sowder, R. Gevirtz, W. Shapiro, and C. Ebert, "Restoration of vagal tone: a possible mechanism for functional abdominal pain," *Applied Psychophysiology Biofeedback*, vol. 35, no. 3, pp. 199–206, 2010.
- [9] P. M. Meier, M. E. Alexander, N. F. Sethna, C. C. E. de Jong-devos van Steenwijk, D. Zurakowski, and C. B. Berde, "Complex regional pain syndromes in children and adolescents: regional and systemic signs and symptoms and hemodynamic response to tilt table testing," *Clinical Journal of Pain*, vol. 22, no. 4, pp. 399–406, 2006.
- [10] M. Matsui, M. Ito, A. Tomoda, and T. Miike, "Complex regional pain syndrome in childhood: report of three cases," *Brain and Development*, vol. 22, no. 7, pp. 445–448, 2000.
- [11] M. Masuda, M. Tsunoda, and K. Imai, "Low catechol-O-methyltransferase activity in the brain and blood pressure regulation," *Biological and Pharmaceutical Bulletin*, vol. 29, no. 2, pp. 202–205, 2006.
- [12] A. Charrua, R. Pinto, A. Taylor et al., "Can the adrenergic system be implicated in the pathophysiology of bladder pain syndrome/interstitial cystitis? A clinical and experimental study," *Neurourology and Urodynamics*, 2013.
- [13] J. Kinoshita, Y. Takahashi, A. M. Watabe, K. Utsunomiya, and F. Kato, "Impaired noradrenaline homeostasis in rats with painful diabetic neuropathy as a target of duloxetine analgesia," *Molecular Pain*, vol. 9, no. 1, article 59, 2013.
- [14] S. Becker, M. Ceko, M. Louis-Foster et al., "Dopamine and pain sensitivity: neither sulpiride nor acute phenylalanine and tyrosine depletion have effects on thermal pain sensations in healthy volunteers," *PLoS ONE*, vol. 8, no. 11, Article ID e80766, 2013.
- [15] N. Altier and J. Stewart, "The role of dopamine in the nucleus accumbens in analgesia," *Life Sciences*, vol. 65, no. 22, pp. 2269–2287, 1999.
- [16] M. Ertas, A. Sagduyu, N. Arac, B. Uludag, and C. Ertekin, "Use of levodopa to relieve pain from painful symmetrical diabetic polyneuropathy," *Pain*, vol. 75, no. 2-3, pp. 257–259, 1998.
- [17] P. B. Wood, P. Schweinhardt, E. Jaeger et al., "Fibromyalgia patients show an abnormal dopamine response to pain," *The European Journal of Neuroscience*, vol. 25, no. 12, pp. 3576–3582, 2007.
- [18] N. Hagelberg, H. Forssell, S. Aalto et al., "Altered dopamine D2 receptor binding in atypical facial pain," *Pain*, vol. 106, no. 1-2, pp. 43–48, 2003.
- [19] L. Miana, V. Hugo do Vale Bastos, S. Machado et al., "Changes in alpha band activity associated with application of the compression of fourth ventricular (CV-4) osteopathic procedure: a qEEG pilot study," *Journal of Bodywork and Movement Therapies*, vol. 17, no. 3, pp. 291–296, 2013.
- [20] M. J. Cutler, B. S. Holland, B. A. Stupski, R. G. Gamber, and M. L. Smith, "Cranial manipulation can alter sleep latency and sympathetic nerve activity in humans: a pilot study," *The Journal of Alternative and Complementary Medicine*, vol. 11, no. 1, pp. 103–108, 2005.

- [21] A. M. Fernández-Pérez, M. I. Peralta-Ramírez, A. Pilat, and C. Villaverde, "Effects of myofascial induction techniques on physiologic and psychologic parameters: a randomized controlled trial," *Journal of Alternative and Complementary Medicine*, vol. 14, no. 7, pp. 807–811, 2008.
- [22] K. E. Nelson, N. Sergueef, and T. Glonek, "The effect of an alternative medical procedure upon low-frequency oscillations in cutaneous blood flow velocity," *Journal of Manipulative and Physiological Therapeutics*, vol. 29, no. 8, pp. 626–636, 2006.
- [23] X. Shi, S. Rehner, P. Prajapati, S. T. Stoll, R. G. Gamber, and H. F. Downey, "Effect of cranial osteopathic manipulative medicine on cerebral tissue oxygenation," *Journal of the American Osteopathic Association*, vol. 111, no. 12, pp. 660–666, 2011.
- [24] W. P. Hanten, S. L. Olson, J. L. Hodson, V. L. Imler, V. M. Knab, and J. L. Magee, "The effectiveness of CV-4 and resting position techniques on subjects with tension-type headaches," *Journal of Manual and Manipulative Therapy*, vol. 7, no. 2, pp. 64–70, 1999.
- [25] J. E. Sealey, "Plasma renin activity and plasma prorenin assays," *Clinical Chemistry*, vol. 37, no. 10, pp. 1811–1819, 1991.
- [26] M. P. Tulppo, T. H. Mäkikallio, T. Seppänen, J. K. E. Airaksinen, and H. V. Huikuri, "Heart rate dynamics during accentuated sympathovagal interaction," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 274, no. 3, pp. H810–H816, 1998.
- [27] J. J. Goldberger, Y.-H. Kim, M. W. Ahmed, and A. H. Kadish, "Effect of graded increases in parasympathetic tone on heart rate variability," *Journal of Cardiovascular Electrophysiology*, vol. 7, no. 7, pp. 594–602, 1996.
- [28] G. Grassi, G. Seravalle, G. Bolla et al., "Heart rate as a sympathetic marker during acute adrenergic challenge," *Journal of Hypertension*, vol. 26, no. 1, pp. 70–75, 2008.
- [29] A. Kamiya, T. Kawada, S. Shimizu, S. Iwase, M. Sugimachi, and T. Mano, "Slow head-up tilt causes lower activation of muscle sympathetic nerve activity: Loading speed dependence of orthostatic sympathetic activation in humans," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 297, no. 1, pp. H53–H58, 2009.
- [30] R. Von Känel, P. J. Mills, M. G. Ziegler, and J. E. Dimsdale, "Effect of β_2 -adrenergic receptor functioning and increased norepinephrine on the hypercoagulable state with mental stress," *American Heart Journal*, vol. 144, no. 1, pp. 68–72, 2002.
- [31] A. Moulson and T. Watson, "A preliminary investigation into the relationship between cervical snags and sympathetic nervous system activity in the upper limbs of an asymptomatic population," *Manual Therapy*, vol. 11, no. 3, pp. 214–224, 2006.
- [32] A. V. Ng, R. Callister, D. G. Johnson, and D. R. Seals, "Age and gender influence muscle sympathetic nerve activity at rest in healthy humans," *Hypertension*, vol. 21, no. 4, pp. 498–503, 1993.
- [33] M. Mannelli, C. Lazzeri, L. Ianni et al., "Dopamine and sympathoadrenal activity in man," *Clinical and Experimental Hypertension*, vol. 19, no. 1-2, pp. 163–179, 1997.
- [34] H. Zouhal, C. Jacob, P. Delamarche, and A. Gratas-Delamarche, "Catecholamines and the effects of exercise, training and gender," *Sports Medicine*, vol. 38, no. 5, pp. 401–423, 2008.
- [35] M. Botcazou, C. Jacob, A. Gratas-Delamarche et al., "Sex effect on catecholamine responses to sprint exercise in adolescents and adults," *Pediatric Exercise Science*, vol. 19, no. 2, pp. 132–144, 2007.
- [36] C. M. Wheatley, E. M. Snyder, B. D. Johnson, and T. P. Olson, "Sex differences in cardiovascular function during submaximal exercise in humans," *Springerpulus*, vol. 3, article 445, 2014.
- [37] S. E. Pettit, I. Marchand, and T. Graham, "Gender differences in cardiovascular and catecholamine responses to cold-air exposure at rest," *Canadian Journal of Applied Physiology*, vol. 24, no. 2, pp. 131–147, 1999.
- [38] P. P. Jones, M. Spraul, K. S. Matt, D. R. Seals, J. S. Skinner, and E. Ravussin, "Gender does not influence sympathetic neural reactivity to stress in healthy humans," *The American Journal of Physiology—Heart and Circulatory Physiology*, vol. 270, no. 1, pp. H350–H357, 1996.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

