

Effects of inulin and di-D-fructose dianhydride-enriched caramels on intestinal microbiota composition and performance of broiler chickens

M. J. Peinado¹, A. Echávarri¹, R. Ruiz¹, E. Suárez-Pereira², C. Ortiz Mellet²,
J. M. García Fernández³ and L. A. Rubio^{1†}

¹Depto. de Fisiología y Bioquímica de la Nutrición Animal (INAN, EEZ, CSIC), Profesor Albareda, 1, 18008 Granada, Spain; ²Depto. de Química Orgánica, Facultad de Química, Univ. de Sevilla, Apdo. 1203, E-41071 Sevilla, Spain; ³Instituto de Investigaciones Químicas (IIQ) CSIC and Univ. de Sevilla, Américo Vespucio 49, Isla de la Cartuja, E-41092 Sevilla, Spain

(Received 18 October 2012; Accepted 9 July 2013; First published online 10 September 2013)

In vitro and in vivo experiments were designed to evaluate the effectiveness of laboratory-made di-D-fructose dianhydride (DFA)-enriched caramels. The DFA-enriched caramels were obtained from D-fructose (FC), D-fructose and sucrose (FSC), or D-fructose and β-cyclodextrin (FCDC). In the in vitro experiment, raftilose and all caramels increased (P < 0.05) L-lactate concentration and decreased (P < 0.05) pH. Total short-chain fatty acid concentration was higher (P < 0.05) than controls in tubes containing raftilose, FSC, FCDC and commercial sucrose caramel (CSC). Raftilose, and all caramels tested except FSC and FC (1%), increased (P < 0.01) lactobacilli log₁₀ number of copies compared with the non-additive control. FSC, FCDC and CSC increased (P < 0.01) the bifidobacteria number of copies as compared with controls. All additives, except FCDC, decreased (P < 0.01) Clostridium coccoides/Eubacterium rectale log number of copies. Compared with controls, raftilose, FC and CSC led to lower (P < 0.01) Escherichia–Shigella and enterobacteria. For the in vivo experiment, a total of 144 male 1-day-old broiler chickens of the Cobb strain were randomly assigned to one of the three dietary treatments for 21 days. Dietary treatments were control (commercial diet with no additive), inulin (20 g inulin/kg diet) and FC (20 g FC/kg diet). Final BW of birds fed FC diet was higher (P < 0.01) than controls or inulin-fed birds, although feed: gain values were not different. Feed intake of chickens fed FC was higher (P < 0.01) than that of inulin-fed birds but not statistically different from controls. Crop pH values were lower (P < 0.01) in birds fed FC diet as compared with control diet, with inulin-fed chickens showing values not different from control- or FC-fed birds. Lower (P < 0.05) lactobacilli number of copies was determined in the crop, ileum and caeca of birds fed the inulin diet compared with the control diet. Inulin supplementation also resulted in lower (P < 0.05) C. coccoides/E. rectale, bacteroides and total bacteria in caecal contents. Addition of FC to broiler diets gave place to lower (P < 0.05) enterobacteria and Escherichia–Shigella in crop and caecal contents compared with controls. The bacteroides number of copies increased (P < 0.05) as compared with controls in the ileum, but decreased (P < 0.05) in the caeca of chickens fed the FC diet. Energy, ADF, NDF and non-starch polysaccharides faecal digestibilities were greater (P < 0.05) than controls in chickens fed diets containing inulin or FC. Fat digestibility was higher (P < 0.05) in FC-fed birds compared with controls or inulin-fed chickens. In conclusion, DFA-enriched caramels tested here, particularly FC, may represent a type of new additives useful in poultry production.

Keywords: broiler, di-D-fructose dianhydrides, inulin, microbiota, prebiotics

Implications

Di-D-fructose dianhydride (DFA)-enriched caramels that were added to a commercial broilers' diet resisted to some extent small intestinal digestion, gave place to lower counts of potentially pathogenic bacteria in the intestine *in vivo*, and increased BW, faecal energy, fat, fibre and non-starch

polysaccharide faecal apparent digestibilities as compared with controls. DFA-enriched caramels may therefore represent a type of new additives useful to improve health, productivity and well-being for poultry production.

Introduction

Worldwide concern about the development of antimicrobial resistance and about transference of antibiotic resistance

† E-mail: lrubio@eez.csic.es

genes from animal to human microbiota led to banning the use of antibiotics as growth promoters in the European Union since January 2006 (EC Regulation, 1831/2003). One of the main consequences of this ban has been a substantial increase in the use of therapeutic antibiotics (Gaggia *et al.*, 2010). As a consequence, there is a need to look for viable alternatives that could enhance the natural defence mechanisms of animals and reduce the massive use of antibiotics. One way is to use specific feed additives and/or dietary raw materials to favourably affect animal performance and welfare, particularly through the modulation of the gut microbiota, which plays a critical role in maintaining host health. In this context, probiotics, prebiotics and synbiotics, that is, combinations of prebiotics and probiotics, have been proposed as possible solutions (Huyghebaert *et al.*, 2011). The main putative effects of these feed additives are the improved resistance to pathogenic bacteria colonization and enhanced host mucosa immunity, thus resulting in a reduced pathogen load, an improved health status of the animals and a reduced risk of food-borne pathogens in foods (Williams *et al.*, 2001). However, although prebiotics seem to selectively enhance lactobacilli and bifidobacteria populations and reduce colonization by pathogenic bacteria, results on performance are often contradictory and mostly affected by the microorganisms or compound chosen, the dietary supplementation level and duration of use. In many cases, the environmental and the stress status of the animals are not reported or considered, as the experimental settings are often too far from farm conditions (Gaggia *et al.*, 2010).

The concept of prebiotics has been recently formalized by the establishment of three scientific criteria that a food ingredient must satisfy to be considered as such: (i) resistance to gastric acidity, to hydrolysis by mammalian enzymes and to gastrointestinal absorption; (ii) be a fermentable substrate by intestinal microorganisms belonging to the human (mammalian) microbiota; and (iii) selective stimulation of the growth and/or activity of intestinal bacteria associated with health and well-being (Roberfroid, 2007). Several complex oligosaccharides, such as inulin, galactooligosaccharides and lactulose, fulfil the three criteria and can be effectively considered as prebiotics (Candela *et al.*, 2010). Di- D -fructose dianhydrides (DFAs) and their glycosylated derivatives (glycosyl-DFAs) represent recent candidates to this list (Arribas *et al.*, 2010; Ortiz Mellet and García Fernández, 2010). However, the number of studies on the effects of DFAs on bacterial growth is quite limited so far, and very little effort has been made specifically in poultry nutrition.

DFAs were found to be present in the non-volatile fraction of industrial soft caramel, although in a relatively modest (15–18%) proportion. Recently, caramelization technologies, based on the use of heterogeneous acid catalysts, have been developed that allow producing DFA-enriched products (up to 70–80% DFAs and glycosyl-DFAs) from fructose or fructose-containing mixtures of food-grade carbohydrates (Suárez-Pereira *et al.*, 2010). Initial results in rats fed with a fructose-derived DFA-enriched caramel were consistent with

a prebiotic behaviour associated with the preservation of a healthy microbiota equilibrium (Arribas *et al.* 2010). Investigating the nutritional effects of these new caramels with high DFA content in farm animals was then very appealing. Accordingly, the work here described was designed to evaluate *in vitro* and *in vivo* the effects of the use of DFA-enriched caramels, in comparison with an industrial sucrose caramel or inulin, on microbial biochemical parameters, and the eventual changes in the intestinal microbiota composition of broilers fed on diets supplemented with these products.

Material and methods

Additives

Three different DFA-enriched caramels obtained from D -fructose (FC) or 1 : 1 mixtures (w/w) of D -fructose and sucrose (FSC), and D -fructose and β -cyclodextrin (FCDC), and a commercial sucrose caramel (CSC) were tested in the *in vitro* trial. FC was also tested *in vivo*. These products were obtained by using a newly developed technology to produce caramels with high DFA and glycosyl-DFA content (>60%) on the basis of the activation of D -fructose or mixtures of different carbohydrates containing D -fructose by strongly acidic ion-exchange resins (Suárez-Pereira *et al.*, 2010). The sucrose caramel was a commercial aromatic caramel produced by Nigay (Feurs, France, ref. Nigay 1395 SMA6) conforming to the AFNOR NF V 00-100 norm (Association Française de Normalisation, Paris, 1988) and having the following technical characteristics: dry matter 792 g/kg; pH (50% in demineralized water) 2.80; colour (absorbance at 520 nm) 6.04. Powdered raftilose[®] P95 (Orafti, Tienen, Belgium) used for the *in vitro* trial was a powder produced through enzymatic hydrolysis of chicory inulin and contains oligofructose (>932 g/kg) with degree of polymerization between 2 and 8, with an average of four residues. Inulin (92.8%) was obtained from Farmusal (Granada, Spain).

In vitro fermentation procedure

Fermentations were conducted in triplicate in 50 ml sterile polypropylene tubes. The composition of the semi-defined medium used for fermentations were as in Ruiz *et al.* (2010). Caecal contents from 10 chickens receiving a cereal-based diet (Table 1) free of any antimicrobial agent were collected in aseptic tubes, sealed, immediately frozen in liquid N_2 and kept at -80°C until the inoculum was prepared. Rose *et al.* (2010) showed that freezing at -80°C had minimal influence on the intestinal microbiota composition or metabolism. To prepare the inoculum, caecal contents were thawed at room temperature, immediately diluted 1 : 10 in 150 mM NaHCO_3 buffer adjusted to pH 7.4 (1 g of caecal content in 9 ml of buffer), and mixed in a stomacher for 2 min. Bags were previously treated with a N_2 stream. Blended, diluted caecal contents were filtered through miracloth (Calbiochem, Merck Millipore, Barcelona, Spain) and sealed in serum bottles under an N_2 stream. Raftilose or caramels were added to the semi-defined medium to reach a final concentration of 10 mg/ml and autoclaved. No additive (negative control),

Table 1 Ingredient and nutrient analysis (g/kg) of the experimental diet

Maize	462
Soy flour	310
Wheat	150
Vit + min mix ^a	30
Animal fat	20
Calcium carbonate	16
Calcium phosphate	3.1
Sodium chloride	4.5
Chromium oxide	2
Methionine	2.2
Lysine	0.2
Calculated analysis	
Metabolizable energy (cal/g)	2912
CP	193.0
Crude fibre	33.7
Fat	44.6
Calcium	7.1
Phosphorous	6.2
Methionine cysteine	8.6
Lysine	1.2

^aThe mineral–vitamin mix contained (per 30 kg): vitamin A, 7 500 000 IU; vitamin D₃, 1 500 000 IU; vitamin E, 25 g; vitamin B₂, 2 g; thiamin B12, 10 mg; vitamin B₆, 67 mg; calcium pantothenate, 7.5 g; nicotinic acid, 10 g; folic acid, 25 mg; vitamin K₃, 1 g; coline chloride, 250 g; Fe, 4 g; Cu, 750 mg; Co, 50 mg; Zn, 38 g; Mn, 42 g; I, 680 mg; Se, 45 mg; coccidiostate (Nistatin + Nicarbacin), 0.50 kg; BHT, 250 mg.

raffilose (positive control) and sample tubes ($n=3$) were then aseptically inoculated with 4 ml of inoculum that were added to 26 ml of medium (see above) in each tube under N₂ stream. Therefore, seven treatments (no additive, raffilose, FSC, FCDC, FC-1%, FC-2% and CSC) with three replicates per treatment were used. Sealed tubes were incubated at 37°C with continuous mixing for 24 h. After this time, tubes were placed into a –80°C freezer and freeze-dried.

Birds, diets and housing

A total of 144 male 1-day-old broiler chickens of the Cobb strain were randomly assigned to one of the three dietary treatments. Birds were weighed on arrival and raised in wire-floored batteries. Each treatment had eight replicates (cages) of six birds. Cages were provided with the convenient heating and the birds received a lighting regimen of 23 h light : 1 h darkness. Balanced commercial diets (Table 1) free of any feed antibiotics and formulated to match the requirements for growing birds of this age and genotype were used. Diets were fed *ad libitum* for 21 days. Dietary treatments were control (commercial diet with no additive), inulin (commercial diet supplemented with 20 g inulin/kg diet) and FC (commercial diet supplemented with 20 g FC/kg diet). FC was chosen for the *in vivo* trial because it was at the moment better characterized chemically than the others. The experimental protocol was reviewed and approved by the Institutional Animal Care and Use Committee of the Spanish Council for Scientific Research (CSIC, Spain), and the animals were cared for in accordance with the Spanish Ministry of Agriculture guidelines (RD 1201/2005).

Sacrifice and sample collection

Data on live BW and feed consumption were recorded at the beginning and at the end of the experiment and used to calculate feed intake (FI) and feed: gain ratio (F/G). At 21 days of age, birds (three per replicate, i.e. 18 per treatment) were randomly selected and killed by intra-thoracic injection of the euthanasic T61 (0.2 ml/bird) (Laboratorios Intervet SA, Salamanca, Spain). The pH was immediately measured in the crop content of each bird by using a Crison pH meter (Crison Instruments SA, Alella, Spain). Immediately after killing, contents from the crop, ileum (considered as the section between the Meckel's diverticulum and the ileo-cecal junction) and caeca of each bird were collected into plastic tubes, stored at –20°C and freeze-dried (Ruiz and Rubio, 2009). Samples of about 1 cm taken at the mid-point of the ileum of three randomly selected birds from each treatment were removed for histological analysis. The samples were flushed twice with PBS to remove luminal digesta and immersed in formalin (10% neutral buffered formaldehyde) for fixation. After 24 h in 10% neutral buffered formaldehyde, the tissue samples were carefully cleaned of any remaining digesta with deionized water, and then transferred to a fresh solution of 10% neutral buffered formaldehyde (Sigma, Alcobendas, Spain).

q-PCR analysis for the *in vivo* and *in vitro* trials

Total DNA was isolated from freeze-dried intestinal or fermentation samples (40 mg) using the QIAamp DNA stool kit (Qiagen, West Sussex, UK) by following the manufacturer's instructions. To increase its effectiveness, the lysis temperature was increased to 95°C and an additional step with lysozyme (10 mg/ml, 37°C, 30 min) incubation was added. Eluted DNA was treated with RNase and the DNA concentration assessed spectrophotometrically by using a NanoDrop ND-100 Spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). Purified DNA samples were stored at –20°C until use (Ruiz and Rubio, 2009). Bacterial log₁₀ number of copies was determined in faecal samples by using q-PCR. The 16S rRNA gene-targeted primers and PCR conditions used in this study are given as Supplementary Material S1.

Chemical and biochemical analysis and calculations

Amounts of DFAs in caramel samples were determined by GC analysis as in Suárez-Pereira *et al.* (2010). Their total content in monosaccharides, DFAs and higher oligosaccharides (including glycosyl DFAs) were: FC, 185 g/kg D-fructose, 316 g/kg DFAs, 480 g/kg glycosyl-DFAs; FSC, 211 g/kg D-fructose, 77 g/kg D-glucose, 272 g/kg DFAs, 410 g/kg glycosyl-DFAs and glucooligosaccharides; FCDC, 100 g/kg D-fructose, 8 g/kg D-glucose, 217 g/kg DFAs, 640 g/kg glycosyl-DFAs and glucooligosaccharides; CSC, 167 g/kg D-fructose; 348 g/kg D-glucose; 136 g/kg DAFs; 311 g/kg glycosyl-DFAs and glucooligosaccharides. For determination of DFAs in samples from the *in vitro* trial and from faecal samples, aliquots (25–40 mg) were diluted in distilled water (60 ml) and dialysed using a Jumbosep™ Centrifugal Devices system equipped

with 3 kDa membranes. The system was centrifuged at 960 g for 25 min using a Hettich Rotina 35 centrifuge (Andreas Hettich GmbH, Tuttlingen, Germany). The filtrate containing the DAFs was collected and re-dialysed five times, freeze-dried and analysed as in Suárez-Pereira *et al.* (2010).

The pH of the *in vitro* fermentation solutions was measured immediately after the 24 h fermentation period. L-lactate and raffinose concentrations were determined by using Megazyme kits (K-LATE 10/04 and K-FRUC 12/04, respectively) (Megazyme International, Wicklow, Ireland).

Short-chain fatty acids (SCFA) (acetate, propionate, butyrate, isobutyrate, valerate and isovalerate) concentrations were determined by GLC (Playne, 1985). Briefly, 2.0 ml of the fermentation product was centrifuged (4°C, 15 000 × g, 15 min) and the supernatant (0.5 ml) mixed with 0.5 ml of a mix of crotonic acid (4 g/l) + metaphosphoric acid (20 g/l) + HCl (0.5 N), and left overnight at -20°C. After centrifugation, 1 µl of the supernatant was injected in a Perkin Elmer model AutoSystem (PerkinElmer, Madrid, Spain) gas chromatograph fitted with a Supelco SP 2380 (Sigma-Aldrich Química SL, Madrid, Spain) capillary column (30 m × 0.25 mm × 0.2 µm). Samples were analysed in triplicate. Appropriate SCFA standards were produced with crotonic acid (2 g/l) as internal standard.

The N contents in feed, ileal contents and faeces were determined according to the Dumas procedure using a LECO Truspec CN analyser (LECO Corporation, St. Joseph, MI, USA). NDF and ADF were carried out according to van Soest *et al.* (1991) by using an Ankom²²⁰ fiber analyzer unit (Ankom Technology Corp., Macedon, NY, USA). NSP in diets and faecal samples was determined as in Englyst *et al.* (1982). Chromium oxide in diets and faecal samples was determined following a colorimetric micromethod (Fenton and Fenton, 1979). AMEn was calculated as in Hill and Anderson (1958), and gross energy in feed and faeces was determined in a bomb calorimeter (Parr 1356 bomb calorimeter; Parr Instruments Co., IL, USA).

Apparent ileal or faecal digestibility (AFD) for each substance (S) (N, ADF, NDF, NSP, fat, FC) was obtained from the expression $AFD = 100 \times [1 - (S_f/Cr_2O_3f)/(S_d/Cr_2O_3d)]$, where d and f indicate concentrations in diet and faeces, respectively.

Histological analysis

Fixed samples were dehydrated and embedded in paraffin wax. Three slides were prepared from each sample, and each slide contained a minimum of two sections cut at 4 µm, at least 50 µm apart. The slides were stained with haematoxylin and eosin. All measurements were made with a light microscope with the help of an image analysis system (CellA Imagen Software, Olympus, Hamburg, Germany) equipped with a monitor. Five well-oriented villi and crypts were selected on each slide to determine villus height, width and crypt depth. The villus height was determined as the distance from the tip to the bottom of the villi, and crypt depth was determined as the distance between its mouth and its base. Villus surface area was calculated as $(3.1416 \times \text{villus width}) \times \text{villus height}$. Mucosal thickness was determined as the distance

between the mucosal epithelium and the muscular layer, and the muscularis as the inner circular and outer longitudinal layers of smooth-muscle cells (Peinado *et al.*, 2012).

Statistical analysis

Data were analysed as a one-way ANOVA using the GLM procedure of SAS (SAS Institute, 2003, Cary, NC, USA), with the pen serving as the experimental unit for performance parameters, and the individual chicken or the fermentation tube as the experimental unit for biochemical or microbiological parameters. All microbiological counts were subject to base-10 logarithm transformation before analysis. Treatment means were separated using Bonferroni's multiple comparison tests. Statistical significance was declared at a probability of $P < 0.05$.

Results

Effects of caramels on biochemical parameters and microbial populations *in vitro*

The pH of the solutions (Table 2) decreased ($P < 0.05$) after *in vitro* fermentation with all additives compared with the non-additive control, the lowest values being reached with raftilose, FC (2%) and CSC. L-lactate concentration *in vitro* (Table 2) was higher ($P < 0.05$) for raftilose than for the other additives, of which the highest values were found for FC (2%) and CSC. Regarding SCFA concentrations (Table 2), acetate concentration was highest ($P < 0.05$) for FSC and FCDC followed by raftilose. Propionate values were increased ($P < 0.05$) compared with controls for all additives tested, with the highest values for FSC and raftilose. FCDC was the only additive that increased ($P < 0.05$) butyrate and valerate concentrations. No effect was detected with respect to control values on isobutyrate or isovalerate production. The total SCFA concentration ($P < 0.05$) was higher than the concentration of control in tubes containing raftilose, FSC, FCDC and CSC, with FC (1%) showing a tendency for increased values as compared with control diets.

Incubation for 24 h within the defined media, regardless of the additive, resulted in significant ($P < 0.01$) differences in all groups of bacteria examined in this study (Table 3). Raftilose and all caramels tested, except FSC and FC (1%), increased ($P < 0.01$) lactobacilli log₁₀ number of copies compared with the non-additive control. No differences in lactobacilli number of copies were found between raftilose, FC (2%) and CSC. FSC, FCDC and CSC induced increased ($P < 0.01$) bifidobacteria log₁₀ number of copies as compared with control tubes. Only FCDC induced an increase ($P < 0.01$) in the bacteroides log₁₀ number of copies respect to raftilose. All additives, except FCDC, decreased ($P < 0.01$) *Clostridium coccoides/Eubacterium rectale* number of copies. Raftilose decreased ($P < 0.01$) and FCDC increased ($P < 0.01$) *C. leptum* log number of copies. Compared with controls, raftilose, FC and CSC gave place to lower ($P < 0.01$) *Escherichia-Shigella* and enterobacteria log number of copies.

Effects of inulin and caramels on performance and crop pH

Final BW of birds fed the FC diet was higher ($P < 0.01$) than controls or inulin-fed birds (Figure 1), although feed:

Table 2 Effect of the type of additive (10 mg/ml) on pH and concentrations of L-lactate (g/l), acetate, propionate, butyrate, isobutyrate, valerate and isovalerate ($\mu\text{mol/ml}$) in vitro after 24 h fermentation

	Additive							Pooled SD	Significance level
	No additive	Raftilose	FSC	FCDC	FC-1%	FC-2%	CSC		
pH	6.9 ^a	4.2 ^d	4.9 ^c	5.2 ^b	5.4 ^b	4.2 ^d	4.2 ^d	0.1	<0.001
L-lactate	0.0 ^f	3.8 ^a	0.4 ^e	1.3 ^c	0.8 ^d	2.9 ^b	2.9 ^b	0.1	<0.001
Acetate	9.0 ^b	11.6 ^{ab}	13.5 ^a	14.3 ^a	8.1 ^b	9.1 ^b	9.9 ^{ab}	3.5	<0.001
Propionate	3.3 ^c	15.5 ^a	17.8 ^a	11.3 ^b	8.6 ^b	10.7 ^b	12.6 ^b	3.6	<0.001
Butyrate	1.5 ^b	0.3 ^b	0.0 ^b	4.2 ^a	0.3 ^b	0.9 ^b	0.7 ^b	1.3	<0.001
Isobutyrate	0.1 ^a	0.3 ^{ab}	0.5 ^{ab}	0.1 ^a	0.2 ^a	1.0 ^b	1.0 ^b	0.8	0.020
Valerate	0.1 ^b	0.0 ^b	0.0 ^b	0.9 ^a	0.0 ^b	0.0 ^b	0.3 ^b	0.3	<0.001
Isovalerate	0.1	0.0	0.0	0.0	0.2	0.9	0.2	0.5	<0.001
Total SCFA	14.1 ^b	27.9 ^a	31.7 ^a	30.8 ^a	17.3 ^b	22.6 ^a	31.0 ^a	6.9	<0.001

FSC = D-fructose/sucrose caramel; FCDC = D-fructose/ β -cyclodextrin caramel; FC = D-fructose caramel; CSC = commercial sucrose caramel. Means (each sample was analysed in triplicate) in the same row with different superscript letters differ ($P < 0.05$).

Table 3 Effect of the type of additive (10 mg/ml, except for FC-2%, which contained 20 mg/ml) on the bacterial \log_{10} number of copies/mg of freeze-dried material after the in vitro fermentation for 24 h within a defined medium

Bacterial group	Additive							Pooled SD	Significance level
	No additive	Raftilose	FSC	FCDC	FC-1%	FC-2%	CSC		
Lactobacilli	6.9 ^a	8.4 ^c	7.4 ^a	7.6 ^b	7.0 ^a	8.1 ^c	8.4 ^c	0.2	<0.001
Bifidobacteria	5.5 ^a	5.6 ^a	6.4 ^b	6.5 ^b	5.4 ^a	5.5 ^a	6.7 ^b	0.2	<0.001
Bacteroides	9.2 ^a	8.4 ^b	8.7 ^{ab}	9.6 ^a	8.9 ^{ab}	8.5 ^b	8.7 ^{ab}	0.5	<0.001
<i>Clostridium coccoides</i> / <i>Eubacterium rectale</i>	8.1 ^a	6.7 ^b	6.8 ^b	7.8 ^a	6.6 ^b	6.7 ^b	7.1 ^b	0.2	<0.001
<i>C. leptum</i>	7.9 ^a	7.4 ^b	7.7 ^{ab}	8.8 ^c	7.7 ^{ab}	7.7 ^{ab}	8.1 ^a	0.3	<0.001
Enterobacteria	8.2 ^a	6.7 ^c	8.0 ^a	8.6 ^a	7.4 ^b	6.2 ^c	7.4 ^b	0.3	<0.001

FSC = D-fructose/sucrose caramel; FCDC = D-fructose/ β -cyclodextrin caramel; FC = D-fructose caramel; CSC = commercial sucrose caramel. Means (each sample was analysed in triplicate) in the same row with different superscript letters differ ($P < 0.01$).




Figure 1 Effects of dietary inclusion of inulin or caramel FC on final BW (a), feed intake (b), feed gain ratio (c) and crop pH (d) of chickens from 1 to 21 days of age. ^{a,b}Bars with different superscripts were significantly different ($P < 0.01$). Values are means (eight replicates of six birds each) with their SD in bars, except for crop pH values, where individual values ($n = 18$ per treatment) were used.

gain values were not different. Feed intake of chickens fed the FC diet was higher ($P < 0.01$) than inulin but not different from controls. Crop pH values were lower

($P < 0.01$) than controls in birds fed the FC diet, with inulin-fed chickens showing values not different from controls or FC-fed birds.

Effects of inulin and caramels on intestinal microbiota composition in vivo

Decreased ($P < 0.05$) lactobacilli \log_{10} number of copies was determined in crop, ileum and caecal contents of birds fed the inulin diet (Table 4). Inulin supplementation also resulted in lower ($P < 0.05$) *C. coccoides*/*E. rectale*, bacteroides and total bacteria in caecal contents. Addition of D-fructose caramel (FC, 2%) to broilers' diets resulted in higher ($P < 0.05$) *C. coccoides*/*E. rectale* number of copies in the crop, and in lower ($P < 0.05$) enterobacteria and *Escherichia-Shigella* \log_{10} number of copies in crop and caecal contents compared with controls. Compared with controls, bacteroides number of copies increased ($P < 0.05$) in the ileum, but decreased ($P < 0.05$) in the caeca, of chickens fed the FC diet.

In vivo intestinal digestibility of substances

In vivo ileal apparent digestibility of DAFs in birds fed FC diet was 0.81 (Table 5). No DAFs were detected in the control diet or in the ileal contents of birds fed the control diet. Inulin digestibility was 14% and 85% at the ileal and fecal levels, respectively. Ileal N digestibility was not affected ($P < 0.05$) by the supplementation with either inulin or FC. On the

Table 4 Bacterial log₁₀ number of copies/mg of freeze-dried contents in the crop, ileum and caeca of broiler chickens fed on control, inulin or DFA-enriched caramel supplemented diets from 1 to 21 days

	Control	Inulin	FC	Pooled SD	Significance level
Crop					
Lactobacilli	6.2 ^a	5.3 ^b	5.8 ^{ab}	0.8	<0.001
Bifidobacteria	2.9	3.0	2.9	0.2	0.092
<i>Clostridium coccoides</i> / <i>Eubacterium rectale</i>	3.8 ^a	3.9 ^{ab}	4.2 ^b	0.3	0.023
<i>C. leptum</i>	4.0	3.9	4.0	0.4	0.648
Enterobacteria	5.5 ^a	5.3 ^{ab}	4.7 ^b	0.7	<0.001
<i>Escherichia-Shigella</i>	5.5 ^a	5.2 ^{ab}	4.8 ^b	0.8	0.014
Bacteroides	4.5	4.7	4.5	0.3	0.027
Total bacteria	6.8	6.9	7.0	0.3	0.317
Ileum					
Lactobacilli	5.2 ^a	4.5 ^b	5.0 ^{ab}	0.7	<0.001
Bifidobacteria	2.7	2.8	2.9	0.4	0.545
<i>C. coccoides</i> / <i>E. rectale</i>	4.7	4.4	4.3	0.5	0.205
<i>C. leptum</i>	4.2	4.1	3.8	0.5	0.100
Enterobacteria	4.4	4.4	4.2	0.6	0.521
<i>Escherichia-Shigella</i>	3.2	3.0	3.0	0.8	0.666
Bacteroides	3.9 ^a	3.9 ^a	4.5 ^b	0.6	0.013
Total bacteria	6.0	5.8	5.9	0.5	0.289
Caecum					
Lactobacilli	7.2 ^a	6.7 ^b	6.9 ^{ab}	0.6	0.004
Bifidobacteria	4.3	4.0	3.9	0.4	0.062
<i>C. coccoides</i> / <i>E. rectale</i>	8.3 ^a	7.8 ^b	7.8 ^b	0.2	<0.001
<i>C. leptum</i>	8.1 ^a	8.0 ^{ab}	7.8 ^b	0.3	<0.001
Enterobacteria	5.9 ^a	5.6 ^a	5.1 ^b	0.6	<0.001
<i>Escherichia-Shigella</i>	5.9 ^a	5.7 ^{ab}	5.2 ^b	0.6	<0.001
Bacteroides	5.7 ^a	5.3 ^b	5.0 ^c	0.3	<0.001
Total bacteria	8.7 ^a	8.3 ^b	8.3 ^b	0.2	<0.001

DFA = D-fructose dianhydride; FC = D-fructose caramel.
Means in the same row with different superscript letters differ ($P < 0.05$).

Table 5 AMEn (cal/g) and ileal and faecal apparent digestibility (%) in growing broiler chickens fed control, inulin or DFA-enriched caramel supplemented diets from 1 to 21 days of age

	Control	Inulin	FC	Pooled SD	Significance level
AMEn	3129	3162	3130	27.2	0.046
Ileal digestibility					
N	83	85	85	2.2	0.059
DFAs	nd	nd	81 (SD 4.1)	–	–
Inulin	nd	14 (SD 9.2)	nd	–	–
Faecal digestibility					
Energy	84 ^a	85 ^b	85 ^b	0.8	0.036
Fat	94 ^a	94 ^a	95 ^b	0.7	0.004
ADF	39 ^a	45 ^b	46 ^b	2.9	<0.001
NDF	66 ^a	68 ^b	69 ^b	1.3	<0.001
NSP	59 ^a	66 ^b	64 ^{ab}	4.3	0.038
Inulin	nd	85 (SD 3.1)	nd	–	–

DFAs = di-D-fructose-dianhydrides; nd = not detected.
Means in each row with different superscript letters differ ($P < 0.05$).

contrary, faecal energy, ADF and NDF apparent digestibilities were greater ($P < 0.05$) than controls in chickens fed diets containing either inulin or FC. Fat faecal apparent digestibility was higher ($P < 0.05$) in FC-fed birds compared with controls or inulin-fed chickens. NSP digestibility was higher ($P < 0.05$) than controls in inulin-fed birds, but not in those fed the FC diet.

Morphology of the ileal mucosa

Inclusion of 20 g of FC/kg in diets for broiler chickens between 1 and 21 days had no significant effect on the histological parameters measured (Table 6). Nevertheless, inulin supplementation with 20 g/kg diet increased ($P < 0.01$) villus height and the ratio of villus height/crypt depth as compared with both control- and FC-fed birds.

Table 6 Morphology of the ileal sections of broiler chickens fed control, inulin or FC diets from 1 to 21 days of age

	Control	Inulin	FC	Pooled SD	Significance level
Villus height (μm)	784 ^a	982 ^b	763 ^a	89	<0.001
Crypt depth (μm)	96	86	87	15	0.222
Villus height/crypt depth	9 ^a	12 ^b	9 ^a	2	<0.001
Villus width (μm)	131	118	116	62	0.536
Villus surface area (μm^2)	325 940	368 549	281 272	183 053	0.175
Mucosal thickness (μm)	47	36	39	16	0.043
Muscular layer thickness (μm)	173	152	177	39	0.144

Means ($n=3$, with five measurements per sample) with different superscripts in each row were significantly different ($P<0.01$).

Discussion

The nutritional effects of sucrose-derived DFA-enriched products and biotechnologically produced individual DFAs have been previously investigated to some extent in laboratory animals (rat), but studies on production animals (pig, broiler chicken) are very scarce. Arribas *et al.* (2010) recently showed that the administration of a DFA-enriched caramel obtained from fructose (containing 70% of an isomeric mixture of 13 DFAs and glycosyl-DFAs, identical to FC in this work) to colitic rats promoted a more favourable intestinal microbiota, increasing lactobacilli and bifidobacteria \log_{10} number of copies, as well as inducing increased concentrations of SCFA in the luminal colonic contents. Previous results on the prebiotic potential of a caramel obtained by pyrolysis of sucrose in the presence of citric acid (Manley-Harris and Richards, 1997) containing ~34% of DFAs and 43% of monosaccharides indicated that DFAs and their glycosylated derivatives might represent promising candidates as prebiotic agents (Orban *et al.*, 1997). It is important to mention here that in the present study fructose-derived caramels were used. These were obtained through a different procedure that involves thermal activation at a much lower temperature (90°C) in the presence of an acid resin approved for its use in the food industry that is then removed from the final product by filtration. The content of DFAs and glycosyl-DFAs, assumed to be the active prebiotic components, in the caramels assayed *in vitro* amounts to 60–80%, being of ~80% in the fructose caramel used for the *in vivo* evaluation.

The DFA-enriched caramels and the commercial sucrose caramel used here increased the numbers of potentially beneficial bacteria *in vitro*. Although numerous studies indicate that inulin and oligofructose selectively stimulate the health-promoting groups of the human intestinal microbiota (Candela *et al.*, 2010), *in vitro* studies on oligosaccharides involving animal microbiota are scarce, and as far as we know there are no references using chicken microbiota. Caramels used here mostly resembled the effects obtained with raftilose. Thus, raftilose and most caramels tested increased lactobacilli, bifidobacteria, bacteroides and clostridia \log number of copies as compared with the non-additive control diet, whereas coliforms and enterobacteria number of colonies were decreased by two of the caramels tested (i.e. FC and FSC) but not by raftilose. This effect is similar to those found with a number of substances

with potential (isomalto-oligosaccharides, lactosucrose, xylo-oligosaccharides) or confirmed (inulin, transgalacto-oligosaccharides, lactulose) prebiotic effects in humans (Candela *et al.*, 2010). Regarding the products of microbial metabolism, it is well established that the principal end products of bacterial fermentation processes are SCFA, with inulin and fructooligosaccharides (FOS) giving rise mainly to acetate (Candela *et al.*, 2010). The results found with raftilose and caramels in our *in vitro* trial (lower pH values, and higher lactate and SCFA, mainly acetate, production) are similar to those reported by Tzortzis *et al.* (2005) using galacto-oligosaccharides and inulin.

Although *in vitro* results provide a preliminary indication, the final demonstration of a prebiotic effect of any food ingredient must be carried out *in vivo* (Roberfroid, 2007). In the current work, the addition to the diet of D-fructose-derived caramel (FC, 2%) did not significantly affect lactobacilli or bifidobacteria number of copies, but led to lower enterobacteria and *Escherichia-Shigella* numbers in crop and caecal contents compared with controls. Compared with controls bacteroides number of copies increased controls in ileal contents but decreased in the caeca of chickens fed the FC diet. As for the inulin-fed birds, lower lactobacilli \log_{10} number of copies was determined in the crop, ileum and caeca compared with controls. Inulin supplementation also resulted in lower bacteroides and total bacteria in caecal contents. These differences in the effects found with inulin and DAFs are likely to be linked to their different intestinal digestibility. Surprisingly, although most of the research interest on prebiotics has focused on their role as modulators of the intestinal microbiota, little effort has been made on their intestinal digestibility, a major criterion to determine their prebiotic potential and mechanism of action (Roberfroid, 2007). However, given that DFAs are highly stable in acidic media and that they are inert to the action of mammalian intestinal glycosidases, it is likely that substantial amounts reach the final sections of the intestinal tract essentially unaltered (Manley-Harris and Richards, 1997). In the current work, ileal apparent digestibility of DAFs in birds fed FC diet was 81%, which means that about 19% of the amounts present in the diet would reach the caeca. The ileal digestibility of inulin was 14% at the ileal level and 85% at the faecal level, which indicates that higher amounts (about 86% of the inulin in the diet) reach the chickens' caeca. These data suggest that the effectiveness of inulin and

caramels is mainly at the caecal and ileal levels, respectively. In fact, this higher digestibility of DFAs in the upper parts of the intestinal tract is likely to explain the lower pH in the crop of birds fed diets containing caramel with respect to control or inulin-fed chickens. In addition, lactobacilli number of copies was not different from controls in the upper gut of caramel-fed birds, but was affected in inulin-fed chickens.

The effectiveness of a given additive in broiler feeding depends on a variety of factors. Variables such as concentration, diet type, animal characteristics, hygiene and husbandry conditions, and environmental stress can influence the response to inulin or FOS in broiler feeding (Patterson and Burkholder, 2003). Dietary supplementation with chito-oligosaccharide (COS) improved the growth rate of broilers, which was likely mediated through the effects of COS on FI and nutrient digestibility. COS may also serve as a growth promoter in broiler production by modulating the concentrations of intestinal microbial flora, as this additive increased the concentrations of cecal lactobacilli and reduced the caecal concentrations of *Eubacterium coli* (Li *et al.*, 2007). In the current work, BW was greater than controls in animals fed the FC diet, and faecal energy, ADF, NDF and NSP apparent digestibilities were greater than controls in chickens fed diets containing inulin or FC. Fat faecal apparent digestibility was higher in FC-fed birds compared with controls or inulin-fed chickens. The mechanism by which NSP exert their anti-nutritive effects in poultry is usually linked to the increased bulk and viscosity of the intestinal contents, which decrease the rate of diffusion of substrates and digestive enzymes, thus hindering their effective interaction at the mucosal surface (Choct *et al.*, 1996). The concentrations of soluble NSP in wheat were inversely correlated with their metabolizable energy (ME_n)-values in broiler chickens (Annisson, 1991). The viscous conditions in the small intestine may interfere with crude fat digestibility owing to draining of bile acids from enterohepatic circulation (Malayoğlu *et al.*, 2010). It has been suggested that an increased digesta viscosity may reduce the amounts of conjugated bile acids, affect fat emulsification negatively and thus decrease fat digestibility (Langhout *et al.*, 1997). Therefore, a higher NSP digestibility as reported here would decrease NSP amounts within the intestine and consequently increase nutrients digestibility.

Although it has been reported that a high level of bird performance may be supported by a range of microbial compositions (Geier *et al.*, 2009), and despite its scientific and practical relevance, the study on the relationship between variations of the microbiota composition and productive/physiological parameters is still at the beginning in poultry production. In a pioneering work, by using T-RFLP analysis together with multivariate statistical methods, Torok *et al.* (2011a) identified and characterized changes in gut microbiota development in chickens in response to enzyme or antimicrobial agents in feed that had performance implications. More specifically, an overgrowth of some microorganisms including enterobacteria in the intestine has been reported (Bourlinoux *et al.*, 2003; Pelicano *et al.*, 2005) to

result in mucosal impairment, villus erosion and damage to the intestinal cells, thus reducing its nutrients' absorptive potential. Fonseca *et al.* (2010) linked a decrease in the quantity of caecal enterobacteria to improved performance in broilers, and Kim *et al.* (2011) proved that lower numbers of certain gut pathogens such as *E. coli* may improve broiler performance. Peinado *et al.* (2012) recently showed that a garlic derivative lowered the intestinal numbers of enteropathogens and improved the ileal histological structure and productive parameters of broilers. On the other hand, in the current work, lactobacilli numbers were lower in the crop, ileum and caeca of chickens fed on the inulin diet and tended to be lower in the FC-fed chickens, which was accompanied by an increase in the bacteroides number of copies in the ileum. Although generally regarded as a beneficial group, higher numbers of lactobacilli have been implicated in broiler growth depression owing to competition for nutrient uptake or impaired fat absorption linked to deconjugation of bile acids (Torok *et al.*, 2011b). Meanwhile, species from the Bacteroidetes family are involved in many important metabolic activities including fermentation of carbohydrates, induction of critical glycolytic enzymes in the enterocytes, utilization of nitrogenous substances, biotransformation of bile acids and prevention of pathogen colonization (Bry *et al.*, 1996; Phillips, 2009). In summary, according to the literature and the current results, it could be concluded that a decrease in the numbers of some bacterial groups such as enterobacteria, and probably lactobacilli, accompanied by an increase in others such as bacteroides in the intestine might be related to improved performance in poultry. However, with the information at present available, it is not possible to rule out other mechanisms such as modifications in the microbiota metabolism, effects on bacterial groups other than those studied here, effects on the immune system, absorption of substances and so on.

Inulin-type fructans are regarded as prebiotics in various intestinal environments of diverse animal genera (mammal, fish, bird, etc.). The specific intestinal fermentation results in a bacterial ecosystem that is less prone to pathogen invasion and in an increased production of SCFA characterized by a higher proportion of butyrate. As a result, the increased absorptive capacity, reflected in a longer intestinal length and increased villus height or crypt depth of the intestine, results in improved feed conversion and better growth in the young animal. The only effect observed in the present work on the histological structure of the chickens' mucosa was that inulin supplementation increased villus height and the ratio between villus height and crypt depth as compared with both control- and FC-fed birds, which is in keeping with previous reports (Rebolé *et al.*, 2010).

In summary, the results presented in this study show that DFAs in DFA-enriched caramels resisted to some extent chickens' small intestinal digestion *in vivo*, were fermented *in vitro* by birds' caecal microbiota and selectively stimulated the growth of bacteria associated with health and well-being *in vitro*. FC supplementation also gave place to lower counts of potentially pathogenic bacteria in the broiler's intestine

in vivo. Faecal energy, fat, ADF, NDF and NSP apparent digestibilities were greater than controls in chickens fed diets containing DFA-enriched caramels.

Acknowledgements

Authors are indebted to Drs L. Lara and A. Nieto and to Mrs M. A. Guzmán for excellent technical assistance, and to the Spanish MICIIN (SAF2008-02616, SAF 2010-15670 and AGL 2009-11925) and the Junta de Andalucía (project AGR395) for financial support. This research has been also partially supported by the FEDER and FSE funds from the European Union. ES-P is a 'Project of Excellence' Fellow (Junta de Andalucía, contract no. P06-AGR-02150) and M. J. P. is a recipient of a JAEpredoc (CSIC) grant.

Supplementary material

To view supplementary material for this article, please visit <http://dx.10.1017/S1751731113001572>

References

- Annisson G 1991. Relationship between the levels of soluble non-starch polysaccharides and the apparent metabolisable energy of wheats assayed in broiler chickens. *Journal of Agricultural and Food Chemistry* 39, 1252–1256.
- Arribas B, Suárez-Pereira E, Ortiz Mellet C, García Fernández JM, Buttersack C, Rodríguez-Cabezas ME, Garrido-Mesa N, Bailon E, Guerra-Hernández E, Zarzuelo A and Gálvez J 2010. Di-D-fructose-enriched caramels: effect on colon microbiota, inflammation, and tissue damage in trinitrobenzenesulfonic acid-induced colitic rats. *Journal of Agricultural and Food Chemistry* 58, 6476–6484.
- Bourlinoux P, Koletzko B and Guarner FV 2003. The intestine and its microflora are partners for the protection of the host. *American Journal of Clinical Nutrition* 78, 675–683.
- Bry L, Falk PG, Midtvedt T and Gordon JI 1996. A model of host-microbial interactions in an open mammalian ecosystem. *Science* 273, 1380–1383.
- Candela M, Maccaferri S, Taroni S, Carnevali P and Brigidi P 2010. Functional intestinal microbiome, new frontiers in prebiotic design. *International Journal of Food Microbiology* 140, 93–101.
- Choct M, Hughes RJ, Wang J, Bedford MR, Morgan AJ, Annison G 1996. Increased small intestinal fermentation responsible for the anti-nutritive activity of non-starch polysaccharides in chickens. *British Poultry Science* 37, 609–621.
- EC 2003. Commission of the European Communities, Commission Regulation (EC) No. 1831/2003. *Official Journal of European Union* L 268, 29–43.
- Englyst HN, Quigley ME, Hudson GJ and Cummings JH 1982. Determination of dietary fiber as non-starch polysaccharides by gas-liquid chromatography. *Analyst* 117, 1707–1714.
- Fenton TW and Fenton M 1979. An improved procedure for the determination of chromic oxide in feed and faeces. *Canadian Journal of Animal Science* 59, 631–634.
- Fonseca BB, Beletti ME, da Silva MS, da Silva PL, Duarte IN and Rossi DA 2010. Microbiota of the cecum, ileum morphology, pH of the crop and performance of broiler chickens supplemented with probiotics. *Revista Brasileira de Zootecnia* 39, 1756–1760.
- Gaggia F, Mattarelli P and Biavati B 2010. Probiotics and prebiotics in animal feeding for safe food production. *International Journal of Food Microbiology* 141, S15–S28.
- Geier MS, Torok VA, Allison GE, Ophel-Keller K and Hughes RJ 2009. Indigestible carbohydrates alter the intestinal microbiota but do not influence the performance of broiler chickens. *Journal of Applied Microbiology* 106, 1540–1548.
- Hill FW and Anderson DL 1958. Comparison of metabolizable energy and productive energy determinations with growing chicks. *Journal of Nutrition* 64, 587–603.
- Huyghebaert G, Ducatelle R and Van Immerseel F 2011. An update on alternatives to antimicrobial growth promoters for broilers. *Veterinary Journal* 187, 182–188.
- Kim G-B, Seo JM, Kim CH and Paik IK 2011. Effect of dietary prebiotic supplementation on the performance, intestinal microflora, and immune response of broilers. *Poultry Science* 90, 75–82.
- Langhout DL, Schutte JB, Geerse C, Kies AK, De Jong J and Verstegen MWA 1997. Effects on chick performance and nutrient digestibility of an endoxylanase added to a wheat- and rye-based diet in relation to fat source. *British Poultry Science* 38, 557–563.
- Li XJ, Piao XS, Kim SW, Liu P, Wang L, Shen YB, Jung SC and Lee HS 2007. Effects of chito-oligosaccharide supplementation on performance, nutrient digestibility, and serum composition in broiler chickens. *Poultry Science* 86, 1107–1114.
- Malayoğlu HB, Baysal S, Misirlioğlu Z, Polat M, Yılmaz H and Turan N 2010. Effects of oregano essential oil with or without feed enzymes on growth performance, digestive enzyme, nutrient digestibility, lipid metabolism and immune response of broilers fed on wheat-soybean meal diets. *British Poultry Science* 51, 67–80.
- Manley-Harris M and Richards GN 1997. Dihexulose dianhydrides. *Advances in Carbohydrate Chemistry and Biochemistry* 52, 207–239.
- Orban JI, Patterson JA, Sutton AL and Richards GN 1997. Effect of sucrose thermal oligosaccharide caramel, dietary vitamin-mineral level, and brooding temperature on growth and intestinal bacterial populations of broiler chickens. *Poultry Science* 76, 482–490.
- Ortiz Mellet C and García Fernández JM 2010. Diffructose dianhydrides (DFAs) and DFA enriched products as functional foods. *Topics in Current Chemistry* 294, 49–77.
- Patterson JA and Burkholder KM 2003. Application of prebiotics and probiotics in poultry production. *Poultry Science* 82, 627–631.
- Peinado MJ, Ruiz R, Echávarri A and Rubio LA 2012. Garlic derivative PTS-O is effective against broiler pathogens *in vivo*. *Poultry Science* 91, 2148–2157.
- Pelicano ERL, Souza PA, Souza HBA, Figueiredo DF, Boiago MM, Carvalho SR and Bordon BF 2005. Intestinal mucosa development in broiler chickens fed natural growth promoters. *Brazilian Journal of Poultry Science* 7, 221–229.
- Phillips ML 2009. Gut reaction: environmental effects on the human microbiota. *Environmental and Health Perspectives* 117, A198–A205.
- Playne MJ 1985. Determination of ethanol, volatile fatty acids, lactic acid and succinic acids in fermentation liquids by gas chromatography. *Journal of the Science of Food and Agriculture* 36, 638–643.
- Rebolé A, Ortiz LT, Rodríguez ML, Alzueta C, Treviño J and Velasco S 2010. Effects of inulin and enzyme complex, individually or in combination, on growth performance, intestinal microflora, cecal fermentation characteristics, and jejunal histomorphology in broiler chickens fed a wheat- and barley-based diet. *Poultry Science* 89, 276–286.
- Roberfroid M 2007. Prebiotics: the concept revisited. *Journal of Nutrition* 137, 830S–837S.
- Rose DJ, Venema K, Keshavarzian A and Hamaker BR 2010. Starch-entrapped microspheres show a beneficial fermentation profile and decrease in potentially harmful bacteria during *in vitro* fermentation in faecal microbiota obtained from patients with inflammatory bowel disease. *British Journal of Nutrition* 103, 1514–1524.
- Ruiz R and Rubio LA 2009. Lyophilization improves the extraction of PCR-quality community DNA from pig faecal samples. *Journal of the Science of Food and Agriculture* 89, 723–727.
- Ruiz R, García MP, Lara A and Rubio LA 2010. Garlic derivatives (PTS and PTS-O) differently affect the ecology of swine faecal microbiota *in vitro*. *Veterinary Microbiology* 144, 110–117.
- Suárez-Pereira E, Rubio EM, Pilard S, Ortiz Mellet C and García-Fernández JM 2010. Di-D-fructose dianhydride (DFA)-enriched caramels by acid ion-exchange resin-promoted caramelization of D-fructose: chemical analysis. *Journal of Agricultural and Food Chemistry* 58, 1777–1787.
- Torok VA, Allison GE, Percy NJ, Ophel-Keller K and Hughes RJ 2011a. Influence of antimicrobial feed additives on broiler commensal posthatch gut microbiota development and performance. *Applied and Environmental Microbiology* 77, 3380–3390.
- Torok VA, Hughes RJ, Mikkelsen LL, Pérez-Maldonado RP, Balding K, MacAlpine R, Percy NJ and Ophel-Keller K 2011b. Identification and characterization of potential performance-related gut microbiotas in broiler chickens

across various feeding trials. *Applied and Environmental Microbiology* 77, 5868–5878.

Tzortzis G, Goulas AK, Gee JM and Gibson GR 2005. A novel galacto-oligosaccharide mixture increases the bifidobacterial population numbers in a continuous *in vitro* fermentation system and in the proximal colonic contents of pigs *in vivo*. *Journal of Nutrition* 135, 1726–1731.

Van Soest PJ, Robertson JB and Lewis BA 1991. Methods for dietary fiber, neutral detergent fiber, and non-starch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583–3597.

Williams BA, Verstegen MW and Tamminga S 2001. Fermentation in the large intestine of single-stomached animals and its relationship to animal health. *Nutrition Research Reviews* 14, 207–228.