



Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra



On solvable Lie and Leibniz superalgebras with maximal codimension of nilradical [☆]

L.M. Camacho ^{a,*}, R.M. Navarro ^b, B.A. Omirov ^c

^a *Dpto. Matemática Aplicada I, Universidad de Sevilla, Sevilla, Spain*

^b *Dpto. de Matemáticas, Universidad de Extremadura, Cáceres, Spain*

^c *National University of Uzbekistan, Institute of Mathematics, Tashkent, Uzbekistan*

ARTICLE INFO

Article history:

Received 23 June 2021

Available online 5 November 2021

Communicated by Alberto Elduque

MSC:

17A32

17A70

17B30

17B40

Keywords:

Solvable Lie superalgebras

Solvable Leibniz superalgebras

Derivations

Nilpotent Lie superalgebras

Nilpotent Leibniz superalgebras

ABSTRACT

Along this paper we show that under certain conditions the method for describing of solvable Lie and Leibniz algebras with maximal codimension of nilradical is also extensible to Lie and Leibniz superalgebras, respectively. In particular, we totally determine the solvable Lie and Leibniz superalgebras with maximal codimension of model filiform and model nilpotent nilradicals. Finally, it is established that the superderivations of the obtained superalgebras are inner.

© 2021 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

[☆] This work has been supported by Agencia Estatal de Investigación (Spain), grant PID2020-115155GB-I00 (European FEDER support included, EU) and by the European Union under the 2014-2020 ERDF Operational Programme and by the Department of Economy, Knowledge, Business and University of the Regional Government of Andalusia. Project reference: FEDER-UCA18-107643. This work has been also supported by Junta de Extremadura and Fondo Europeo de Desarrollo Regional (GR18001 and IB18032).

* Corresponding author.

E-mail addresses: lcamacho@us.es (L.M. Camacho), rnavarro@unex.es (R.M. Navarro), omirovb@mail.ru (B.A. Omirov).

1. Introduction

In 1950 A.I. Malcev proved that a solvable Lie algebra is uniquely determined by its nilradical [19]. A decade later Mubarakzjanov proposed the method of description of solvable Lie algebra in terms of nilradical and its nil-independent derivations [20]. He noted that the codimension of nilradical does not exceed the number of nil-independent derivations and the number of generators of the nilradical. Since then the classification of solvable Lie algebras with the Abelian, Heisenberg, filiform, quasi-filiform nilradicals is obtained, see for instance [5,6,21,25,26,24]. Recently, the topic of research has drawn a lot of attention and in particular in [11] the authors extended Mubarakzjanov's method to Leibniz algebras. Therefore, along the last few years have been obtained classifications with different types of Leibniz nilradicals [10–12,18]. Among all the solvable Lie and Leibniz algebras special mention deserves the maximal (in dimensional sense) solvable Lie and Leibniz algebras with given nilradical, due to its properties, they are in some cases cohomologically rigid [4].

The main goal of this paper is the study of maximal solvable Lie and Leibniz superalgebras with a given nilradical. It should be noted that the structures of solvable Lie and Leibniz superalgebras are more complex than structures of solvable Lie and Leibniz algebras [23]. In particular, Lie's theorem is not true (neither in its general or its reduced forms) for a solvable Leibniz (respectively, Lie) superalgebras L . Moreover, the square of a solvable superalgebra is not necessary to be nilpotent [22]. Nevertheless, throughout this paper we show that under certain conditions Mubarakzjanov's method of description of maximal solvable Lie and Leibniz algebras with given nilradical is also applicable for Lie and Leibniz superalgebras, respectively. In particular, we determine the maximal dimensional solvable Lie and Leibniz superalgebras with model filiform and model nilpotent nilradicals. Additionally, we compute the space of superderivations on the maximal-dimensional solvable Lie and Leibniz superalgebras with filiform and model nilpotent nilradical and show that all of these superderivations are inner. These results constitute an extension of the results obtained for similar Lie and Leibniz algebras. Finally, let us note that all Lie and Leibniz superalgebras obtained in this paper are excellent candidates to cohomologically rigid superalgebras, we leave this study for a further work though.

2. Preliminary results

A vector space V is said to be \mathbb{Z}_2 -graded if it admits a decomposition in direct sum, $V = V_{\bar{0}} \oplus V_{\bar{1}}$, where $\bar{0}, \bar{1} \in \mathbb{Z}_2$. An element $x \in V$ is called *homogeneous of degree \bar{i}* if it is an element of $V_{\bar{i}}$, $\bar{i} \in \mathbb{Z}_2$. In particular, the elements of $V_{\bar{0}}$ (resp. $V_{\bar{1}}$) are also called *even* (resp. *odd*). For a homogeneous element $x \in V$ we denote $|x|$ the degree of x (either $\bar{0}$ or $\bar{1}$).

A *Lie superalgebra* (see [15]) is a \mathbb{Z}_2 -graded vector space $\mathfrak{g} = \mathfrak{g}_{\bar{0}} \oplus \mathfrak{g}_{\bar{1}}$ together with an even bilinear mapping $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ ($[\mathfrak{g}_{\bar{0}}, \mathfrak{g}_{\bar{0}}] \subset \mathfrak{g}_{\bar{0}}$, $[\mathfrak{g}_{\bar{0}}, \mathfrak{g}_{\bar{1}}] \subset \mathfrak{g}_{\bar{1}}$ and $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] \subset \mathfrak{g}_{\bar{0}}$), which for any arbitrary homogeneous elements x, y, z satisfies the conditions:

1. $[x, y] = -(-1)^{|x||y|}[y, x]$,
2. $(-1)^{|z||x|}[x, [y, z]] + (-1)^{|x||y|}[y, [z, x]] + (-1)^{|y||z|}[z, [x, y]] = 0$ (*super Jacobi identity*).

Thus, \mathfrak{g}_0 is an ordinary Lie algebra, and \mathfrak{g}_1 is a module over \mathfrak{g}_0 ; the Lie superalgebra structure also contains the symmetric pairing $S^2\mathfrak{g}_1 \rightarrow \mathfrak{g}_0$.

In general, the *descending central sequence* of a Lie superalgebra $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is defined in the same way as for Lie algebras: $\mathcal{C}^0(\mathfrak{g}) := \mathfrak{g}$, $\mathcal{C}^{k+1}(\mathfrak{g}) := [\mathcal{C}^k(\mathfrak{g}), \mathfrak{g}]$ for all $k \geq 0$. Consequently, if $\mathcal{C}^k(\mathfrak{g}) = \{0\}$ for some k , then the Lie superalgebra is called *nilpotent*. Then, the smallest integer k such that $\mathcal{C}^k(\mathfrak{g}) = \{0\}$ is called the *nilindex* of the Lie superalgebra \mathfrak{g} . Likewise, the *derived sequence* of \mathfrak{g} is defined by $\mathcal{D}^0(\mathfrak{g}) := \mathfrak{g}$, $\mathcal{D}^{k+1}(\mathfrak{g}) := [\mathcal{D}^k(\mathfrak{g}), \mathcal{D}^k(\mathfrak{g})]$ for all $k \geq 0$. If this sequence is stabilized in zero, then the Lie superalgebra is said to be *solvable*. All nilpotent Lie superalgebras are solvable ones. Engel’s theorem and its direct consequences remain valid for Lie superalgebras. In particular, a Lie superalgebra L is nilpotent if and only if $ad_L x$ is nilpotent for every homogeneous element x of L . Moreover, for solvable Lie superalgebras we have that a Lie superalgebra L is solvable if and only if its Lie algebra L_0 is solvable. Nevertheless, we do not have the analog of Lie’s Theorem and neither its corollaries for solvable Lie superalgebras.

At the same time, there are also defined two other crucial sequences denoted by $\mathcal{C}^k(\mathfrak{g}_0)$ and $\mathcal{C}^k(\mathfrak{g}_1)$ which will play an important role in our study. They are defined as follows:

$$\mathcal{C}^0(\mathfrak{g}_i) := \mathfrak{g}_i, \mathcal{C}^{k+1}(\mathfrak{g}_i) := [\mathfrak{g}_0, \mathcal{C}^k(\mathfrak{g}_i)], \quad k \geq 0, i \in \mathbb{Z}_2.$$

This last sequences allow us to introduce filiform Lie superalgebras. Thus, a Lie superalgebra $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is said to be filiform if \mathfrak{g}_0 is a filiform Lie algebra (those algebras with nilindex $\dim(\mathfrak{g}_0) - 1$) and the action of \mathfrak{g}_0 over \mathfrak{g}_1 has the structure of filiform \mathfrak{g}_0 -module, i.e. $\dim(\mathcal{C}^{i-1}(\mathfrak{g}_1)/\mathcal{C}^i(\mathfrak{g}_1)) = 1$ for $1 \leq i \leq \dim(\mathfrak{g}_1)$.

In the study nilpotent Lie algebras we have the invariant called characteristic sequence that can be naturally extended for Lie superalgebras.

Definition 2.1. For an arbitrary element $x \in \mathfrak{g}_0$, the adjoint operator ad_x is a nilpotent endomorphism of the space \mathfrak{g}_i , where $i \in \{\bar{0}, \bar{1}\}$. We denote by $gz_i(x)$ the descending sequence of dimensions of Jordan blocks of ad_x . Then, we define the invariant of a Lie superalgebra \mathfrak{g} as follows:

$$gz(\mathfrak{g}) = \left(\max_{x \in \mathfrak{g}_0 \setminus \{0\}} gz_{\bar{0}}(x) \mid \max_{\tilde{x} \in \mathfrak{g}_0 \setminus \{0\}} gz_{\bar{1}}(\tilde{x}) \right),$$

where gz_i is in lexicographic order.

The couple $gz(\mathfrak{g})$ is called characteristic sequence of the Lie superalgebra \mathfrak{g} .

Let us recall the definition of superderivations for Lie superalgebras [15]. A superderivation of degree s of a Lie superalgebra L , $s \in \mathbb{Z}_2$, is an endomorphism $D \in \text{End}_s L$ with the property

$$D(ab) = D(a)b + (-1)^{s \cdot \text{dega}} aD(b).$$

If we denote $\text{Der}_s L \subset \text{End}_s L$ the space of all superderivations of degree s , then $\text{Der} L = \text{Der}_{\bar{0}} L \oplus \text{Der}_{\bar{1}} L$ is the Lie superalgebra of superderivations of L , with $\text{Der}_{\bar{0}} L$ composed by even superderivations and $\text{Der}_{\bar{1}} L$ by odd ones.

2.1. Preliminaries for Leibniz superalgebras

Remark that many results and definitions of the above section can be extended for Leibniz superalgebras.

Definition 2.2. [2]. A \mathbb{Z}_2 -graded vector space $L = L_{\bar{0}} \oplus L_{\bar{1}}$ is called a *Leibniz superalgebra* if it is equipped with a product $[\cdot, \cdot]$ which for an arbitrary element x and homogeneous elements y, z satisfies the condition

$$[x, [y, z]] = [[x, y], z] - (-1)^{|y||z|} [[x, z], y] \quad (\text{super Leibniz identity}).$$

Note that if a Leibniz superalgebra L satisfies the identity $[x, y] = -(-1)^{|x||y|} [y, x]$ for any homogeneous elements $x, y \in L$, then the super Leibniz identity becomes the super Jacobi identity. Consequently, Leibniz superalgebras are a generalization of Lie superalgebras. Also and in the same way as for Lie superalgebras, isomorphisms are assumed to be consistent with the \mathbb{Z}_2 -graduation.

Let us now denote by R_x the right multiplication operator, i.e., $R_x : L \rightarrow L$ given as $R_x(y) := [y, x]$ for $y \in L$, then the super Leibniz identity can be expressed as $R_{[x,y]} = R_y R_x - (-1)^{|x||y|} R_x R_y$.

If we denote by $R(L)$ the set of all right multiplication operators, then $R(L)$ with respect to the following multiplication

$$\langle R_a, R_b \rangle := R_a R_b - (-1)^{\bar{i}\bar{j}} R_b R_a \tag{2.1}$$

for $R_a \in R(L)_{\bar{i}}$, $R_b \in R(L)_{\bar{j}}$, forms a Lie superalgebra. Note that R_a is a derivation. In fact, the condition for being a derivation of a Leibniz superalgebra (for more details see [17]) is $d([x, y]) = (-1)^{|d||y|} [d(x), y] + [x, d(y)]$. Since the degree of R_z as homomorphism between \mathbb{Z}_2 -graded vector spaces is the same as the degree of the homogeneous element z , that is $|R_z| = |z|$, then the condition for R_z to be a derivation is exactly $R_z([x, y]) = (-1)^{|z||y|} [R_z(x), y] + [x, R_z(y)]$. This last condition can be rewritten $[[x, y], z] = (-1)^{|z||y|} [[x, z], y] + [x, [y, z]]$ which is nothing but the super (graded) Leibniz identity.

Let us note that the concepts of descending central sequence, nilindex, the variety of Leibniz superalgebras and Engel’s theorem are natural extensions from Lie theory.

Let $V = V_{\bar{0}} \oplus V_{\bar{1}}$ be the underlying vector space of L , $L = L_{\bar{0}} \oplus L_{\bar{1}} \in Leib^{n,m}$, being $Leib^{n,m}$ the variety of Leibniz superalgebras, and let $G(V)$ be the group of the invertible linear mappings of the form $f = f_{\bar{0}} + f_{\bar{1}}$, such that $f_{\bar{0}} \in GL(n, \mathbb{C})$ and $f_{\bar{1}} \in GL(m, \mathbb{C})$ (then $G(V) = GL(n, \mathbb{C}) \oplus GL(m, \mathbb{C})$). The action of $G(V)$ on $Leib^{n,m}$ induces an action on the Leibniz superalgebras variety: two laws λ_1, λ_2 are *isomorphic* if there exists a linear mapping $f = f_{\bar{0}} + f_{\bar{1}} \in G(V)$, such that

$$\lambda_2(x, y) = f_{\bar{i}+\bar{j}}^{-1}(\lambda_1(f_{\bar{i}}(x), f_{\bar{j}}(y))), \text{ for any } x \in V_{\bar{i}}, y \in V_{\bar{j}}.$$

Furthermore, the description of the variety of any class of algebras or superalgebras is a difficult problem. Different papers (for example, [3,8,13,14]) are regarding the applications of algebraic groups theory to the description of the variety of Lie and Leibniz algebras.

Definition 2.3. For a Leibniz superalgebra $L = L_{\bar{0}} \oplus L_{\bar{1}}$ we define the *right annihilator* of L as the set $Ann(L) := \{x \in L : [L, x] = 0\}$.

It is easy to see that $Ann(L)$ is a two-sided ideal of L and $[x, x] \in Ann(L)$ for any $x \in L_{\bar{0}}$. This notion is compatible with the right annihilator in Leibniz algebras. If we consider the ideal $I := ideal < [x, y] + (-1)^{|x||y|}[y, x] >$, then $I \subset Ann(L)$.

Let $L = L_{\bar{0}} \oplus L_{\bar{1}}$ be a nilpotent Leibniz superalgebra with $\dim L_{\bar{0}} = n$ and $\dim L_{\bar{1}} = m$. From Equation (2.1) we have that $R(L)$ is a Lie superalgebra, and in particular $R(L_{\bar{0}})$ is a Lie algebra. As $L_{\bar{1}}$ has $L_{\bar{0}}$ -module structure we can consider $R(L_{\bar{0}})$ as a subset of $GL(V_{\bar{1}})$, where $V_{\bar{1}}$ is the underlying vector space of $L_{\bar{1}}$. So, we have a Lie algebra formed by nilpotent endomorphisms of $V_{\bar{1}}$. Applying the Engel’s theorem we have the existence of a sequence of subspaces of $V_{\bar{1}}$, $V_0 \subset V_1 \subset V_2 \subset \dots \subset V_m = V_{\bar{1}}$, with $R(L_{\bar{0}})(V_{\bar{i}+1}) \subset V_{\bar{i}}$. Then, it can be defined the descending sequences $C^k(L_{\bar{0}})$ and $C^k(L_{\bar{1}})$ and the super-nilindex in the same way as for Lie superalgebras. That is, $C^0(L_{\bar{i}}) := L_{\bar{i}}, C^{k+1}(L_{\bar{i}}) := [C^k(L_{\bar{i}}), L_{\bar{0}}], k \geq 0, \bar{i} \in \mathbb{Z}_2$. If $L = L_{\bar{0}} \oplus L_{\bar{1}}$ is a nilpotent Leibniz superalgebra, then L has *super-nilindex* or *s-nilindex* (p, q) if satisfies

$$C^{p-1}(L_{\bar{0}}) \neq 0, \quad C^{q-1}(L_{\bar{1}}) \neq 0, \quad C^p(L_{\bar{0}}) = C^q(L_{\bar{1}}) = 0.$$

We have for Lie superalgebras the invariant called characteristic sequences that can be naturally extended for Leibniz superalgebras. Thus, we have the following definition.

Definition 2.4. For an arbitrary element $x \in L_{\bar{0}}$, the operator R_x is a nilpotent endomorphism of the space L_i , where $i \in \{\bar{0}, \bar{1}\}$. We denote by $gz_i(z)$ the descending sequences of dimensions of Jordan blocks of R_x . Then, we define the invariant of a Leibniz superalgebra L as follows:

$$gz(L) = \left(\max_{x \in L_{\bar{0}} \setminus [L_{\bar{0}}, L_{\bar{0}}]} gz_{\bar{0}}(x) \mid \max_{\tilde{x} \in L_{\bar{0}} \setminus [L_{\bar{0}}, L_{\bar{0}}]} gz_{\bar{1}}(\tilde{x}) \right),$$

where gz_i is in lexicographic order.

The couple $gz(L)$ is called characteristic sequences of a Leibniz superalgebra L .

3. Maximal-dimensional solvable Lie superalgebras with filiform nilradical

Throughout this section we study solvable Lie superalgebras with maximal dimension of the complementary space to nilradical, being the nilradical the model filiform Lie superalgebra $L^{n,m}$. Let us recall that in [9] the authors proved that under the condition of being L^2 nilpotent, any solvable Lie superalgebra over the real or complex numbers field can be obtained by means of outer non-nilpotent superderivations of the nilradical. Therefore, for any solvable Lie superalgebra \mathfrak{r} with \mathfrak{r}^2 nilpotent, we have a decomposition into semidirect sum:

$$\mathfrak{r} = \mathfrak{t} \overrightarrow{\oplus} \mathfrak{n},$$

$$[\mathfrak{t}, \mathfrak{n}] \subset \mathfrak{n}, \quad [\mathfrak{n}, \mathfrak{n}] \subset \mathfrak{n}, \quad [\mathfrak{t}, \mathfrak{t}] \subset \mathfrak{n}.$$

Along the present section we consider as nilradical the model filiform Lie superalgebra $L^{n,m}$, that is, the simplest filiform Lie superalgebra which is defined by the only non-zero products

$$L^{n,m} : \begin{cases} [x_1, x_i] = -[x_i, x_1] = x_{i+1}, & 2 \leq i \leq n - 1 \\ [x_1, y_j] = -[y_j, x_1] = y_{j+1}, & 1 \leq j \leq m - 1 \end{cases}$$

where $\{x_1, \dots, x_n\}$ be a basis of $(L^{n,m})_{\bar{0}}$ and $\{y_1, \dots, y_m\}$ be a basis of $(L^{n,m})_{\bar{1}}$. Note that $L^{n,m}$ is the most important filiform Lie superalgebra, in complete analogy to Lie algebras, since all the other filiform Lie superalgebras can be obtained from it by deformations [7]. These infinitesimal deformations are given by the even 2-cocycles $Z^2_0(L^{n,m}, L^{n,m})$.

On the other hand, $\mathfrak{t} = span\{t_1, t_2, t_3\}$ corresponds with the maximal torus of derivations of $L^{n,m}$, which is composed, in particular, by even superderivations. Then \mathfrak{t} is Abelian ($[\mathfrak{t}, \mathfrak{t}] = 0$) and the operators adt_i ($t_i \in \mathfrak{t}$) are diagonal. A straightforward computation leads to the following action of \mathfrak{t} over $L^{n,m}$:

$$\begin{aligned} [t_1, x_i] &= ix_i, & 1 \leq i \leq n; \\ [t_1, y_j] &= jy_j, & 1 \leq j \leq m; \\ [t_2, x_i] &= x_i, & 2 \leq i \leq n; \\ [t_3, y_j] &= y_j, & 1 \leq j \leq m. \end{aligned}$$

Thus, the solvable Lie superalgebra that we are going to consider, and henceforth named $SL^{n,m}$, is defined in a basis $\{x_1, \dots, x_n, t_1, t_2, t_3, y_1, \dots, y_m\}$ by the only non-zero bracket products

$$SL^{n,m} : \begin{cases} [x_1, x_i] = -[x_i, x_1] = x_{i+1}, & 2 \leq i \leq n - 1; \\ [x_1, y_j] = -[y_j, x_1] = y_{j+1}, & 1 \leq j \leq m - 1; \\ [t_1, x_i] = -[x_i, t_1] = ix_i, & 1 \leq i \leq n; \\ [t_1, y_j] = -[y_j, t_1] = jy_j, & 1 \leq j \leq m; \\ [t_2, x_i] = -[x_i, t_2] = x_i, & 2 \leq i \leq n; \\ [t_3, y_j] = -[y_j, t_3] = y_j, & 1 \leq j \leq m; \end{cases}$$

with $\{x_1, \dots, x_n, t_1, t_2, t_3\}$ a basis of $(SL^{n,m})_{\bar{0}}$ and $\{y_1, \dots, y_m\}$ a basis of $(SL^{n,m})_{\bar{1}}$. The purpose now is to find out whether $SL^{n,m}$ is the unique solvable Lie superalgebra with maximal codimension of nilradical $L^{n,m}$.

Theorem 3.1. *An arbitrary complex maximal-dimensional solvable Lie superalgebra L with L^2 nilpotent and nilradical $L^{n,m}$ is isomorphic to $SL^{n,m}$.*

Proof. It is easy to see that superalgebra $L^{n,m}$ can be considered as nilpotent Lie algebra (a \mathbb{Z}_2 -graded Lie algebra), because of the fact that there is no symmetric bracket products in the law of $L^{n,m}$. Note also, that under the condition of being L^2 nilpotent the techniques used in Lie superalgebras are rather similar to the ones used in Lie algebras, i.e. solvable extension by means of (super)derivations of nilradical, for more details it can be consulted [9]. Considering now $L^{n,m}$ as a Lie algebra, the results of [16] allow us to assert that there is a unique solvable Lie algebra with maximal codimension of nilradical, i.e. maximal dimension of the complementary space to nilradical. Moreover this maximal codimension is equal to the number of generators of the nilradical, 3 in our case. It can be easily seen that this unique solvable Lie algebra described in [16] is isomorphic to $SL^{n,m}$, considered the latter as a Lie algebra. Indeed, following to Theorem 3.2 [16] we have a solvable Lie algebra

$$\begin{cases} [x_1, x_i] = -[x_i, x_1] = x_{i+1}, & 2 \leq i \leq n - 1; \\ [x_1, y_j] = -[y_j, x_1] = y_{j+1}, & 1 \leq j \leq m - 1; \\ [z_1, x_1] = -[x_1, z_1] = x_1, \\ [z_1, x_i] = -[x_i, z_1] = (i - 2)x_i, & 3 \leq i \leq n; \\ [z_1, y_j] = -[y_j, z_1] = (j - 1)y_j, & 2 \leq j \leq m; \\ [z_2, x_i] = -[x_i, z_2] = x_i, & 2 \leq i \leq n; \\ [z_3, y_j] = -[y_j, z_3] = y_j, & 1 \leq j \leq m; \end{cases}$$

and the isomorphism defined by $\{t_1 = z_1 + 2z_2 + z_3, t_2 = z_2, t_3 = z_3\}$ shows that this Lie algebra is isomorphic to $SL^{n,m}$.

Now, we consider $L^{n,m}$ as a Lie superalgebra.

The even superderivations D of $L^{n,m}$ are in particular derivations of the \mathbb{Z}_2 -graded Lie algebra $L^{n,m}$ verifying $D(L_0^{n,m}) \subset L_0^{n,m}$ and $D(L_1^{n,m}) \subset L_1^{n,m}$. The odd superderivations D of $L^{n,m}$, on the other hand, verify the following three conditions:

$$D([x_i, x_j]) = [D(x_i), x_j] + [x_i, D(x_j)], \quad \text{if } x_i, x_j \in L_0^{n,m} \tag{3.1}$$

$$D([x_i, y_j]) = [D(x_i), y_j] + [x_i, D(y_j)], \quad \text{if } x_i \in L_0^{n,m}, y_j \in L_1^{n,m} \tag{3.2}$$

$$D([y_i, y_j]) = [D(y_i), y_j] - [y_i, D(y_j)], \quad \text{if } y_i, y_j \in L_1^{n,m} \tag{3.3}$$

The equations (3.1) and (3.2) correspond with the Lie derivation condition. Thus, it remains to study the equation (3.3). Taking into account now the law of $L^{n,m}$ it can be easily seen that the only possibility for having at least one non-null term in the equation (3.3) corresponds with the existence of y_i such that $x_1 \in D(y_i)$. Next, we explore such possibilities. For $i \geq 2$, we get

$$D(y_i) = D([x_1, y_{i-1}]) = [D(x_1), y_i] + [x_1, D(y_{i-1})] = [x_1, D(y_{i-1})]$$

Since $[x_1, D(y_{i-1})] \in \text{span}\{x_3, \dots, x_n\}$ we can exclude i for $i \geq 2$. Suppose $i = 1$, and then

$$D(y_1) = \alpha_1 x_1 + \sum_{k=2}^n \alpha_k x_k, \quad \text{with } \alpha_1 \neq 0.$$

From equation (3.3) we get in particular

$$D([y_1, y_1]) = 0 = [D(y_1), y_1] - [y_1, D(y_1)] = 2\alpha_1 y_2$$

and then $\alpha_1 = 0$. Therefore the equation (3.3) vanishes over $L^{n,m}$ and consequently all the odd superderivations of $L^{n,m}$ are in particular derivations of the \mathbb{Z}_2 -graded Lie algebra $L^{n,m}$ verifying $D(L_0^{n,m}) \subset L_0^{n,m}$ and $D(L_1^{n,m}) \subset L_0^{n,m}$.

Thus, all the superderivations of the Lie superalgebra $L^{n,m}$ are particular cases of derivations of the \mathbb{Z}_2 -graded Lie algebra $L^{n,m}$. Therefore, we can assert that $SL^{n,m}$ is not only the unique maximal-dimensional solvable Lie algebra with nilradical the Lie algebra $L^{n,m}$, but also is the unique maximal-dimensional solvable Lie superalgebra L with L^2 nilpotent and nilradical the model filiform Lie superalgebra $L^{n,m}$. \square

4. Maximal-dimensional solvable Lie superalgebras with model nilpotent nilradical

Throughout this section firstly we extend the definition of model nilpotent to Lie superalgebras and after that we obtain the description of maximal-dimensional solvable Lie superalgebra with model nilpotent nilradical.

Next we recall the definition of model nilpotent Lie algebra, for more details it can be consulted for instance [4]. Thus, the model nilpotent Lie algebra with arbitrary characteristic sequence $(n_1, n_2, \dots, n_k, 1)$ is composed by the Lie algebras admitting a basis

$\{x_1, \dots, x_{n_1+1}, \dots, x_{n_1+n_2+1}, \dots, x_{n_1+\dots+n_k+1}\}$ such that the only non-null brackets (expanded by skew-symmetry) are exactly the following

$$\begin{aligned} [x_1, x_j] &= x_{j+1}, \quad 2 \leq j \leq n_1, \\ [x_1, x_{n_1+j}] &= x_{n_1+1+j}, \quad 2 \leq j \leq n_2, \\ &\vdots \\ [x_1, x_{n_1+\dots+n_{k-2}+j}] &= x_{n_1+\dots+n_{k-2}+1+j}, \quad 2 \leq j \leq n_{k-1}, \\ [x_1, x_{n_1+\dots+n_{k-1}+j}] &= x_{n_1+\dots+n_{k-1}+1+j}, \quad 2 \leq j \leq n_k. \end{aligned}$$

Therefore, we present the next definition in a natural way.

Definition 4.1. The model nilpotent Lie superalgebra with arbitrary characteristic sequences $(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ is nothing but the Lie superalgebra admitting a basis $\{x_1, \dots, x_{n_1+\dots+n_k+1}, y_1, \dots, y_{m_1+\dots+m_p}\}$ with x_i even basis vectors and y_j odd basis vectors, such that the only non-null brackets are exactly the following

$N(n_1, \dots, n_k, 1|m_1, \dots, m_p)$:

$$\left\{ \begin{aligned} [x_1, x_j] &= -[x_j, x_1] = x_{j+1}, & 2 \leq j \leq n_1, \\ [x_1, x_{n_1+\dots+n_j+i}] &= -[x_{n_1+\dots+n_j+i}, x_1] = x_{n_1+\dots+n_j+i+1}, & 1 \leq j \leq k-1, \quad 2 \leq i \leq n_{j+1}, \\ [x_1, y_j] &= -[y_j, x_1] = y_{j+1}, & 1 \leq j \leq m_1-1, \\ [x_1, y_{m_1+\dots+m_j+i}] &= -[y_{m_1+\dots+m_j+i}, x_1] &= y_{m_1+\dots+m_j+i+1}, & 1 \leq j \leq p-1, \quad 1 \leq i \leq m_{j+1}-1. \end{aligned} \right.$$

Remark 4.1. Note that the model filiform Lie superalgebra $L^{n,m}$ is exactly $N(n-1, 1|m)$.

Let us consider now $\mathfrak{t} = \text{span}\{t_1, \dots, t_{k+1}, t'_1, \dots, t'_p\}$ the maximal torus of derivations of $N(n_1, \dots, n_k, 1|m_1, \dots, m_p)$, which is composed in particular by even superderivations. A straightforward computation leads to the following action of \mathfrak{t} over $N(n_1, \dots, n_k, 1|m_1, \dots, m_p)$:

$$\begin{aligned} [t_1, x_i] &= ix_i, & 1 \leq i \leq n_1 + \dots + n_k + 1; \\ [t_1, y_j] &= jy_j, & 1 \leq j \leq m_1 + \dots + m_p; \\ [t_2, x_i] &= x_i, & 2 \leq i \leq n_1 + 1; \\ [t_{j+2}, x_{n_1+\dots+n_j+i}] &= x_{n_1+\dots+n_j+i}, & 1 \leq j \leq k-1, \quad 2 \leq i \leq n_{j+1} + 1; \\ [t'_1, y_i] &= y_i, & 1 \leq i \leq m_1; \\ [t'_{j+1}, y_{m_1+\dots+m_j+i}] &= y_{m_1+\dots+m_j+i}, & 1 \leq j \leq p-1, \quad 1 \leq i \leq m_{j+1}. \end{aligned}$$

Thus, the solvable Lie superalgebra that we are going to consider and denoted by $SN(n_1, \dots, n_k, 1| m_1, \dots, m_p)$ is defined in a basis $\{x_1, \dots, x_{n_1+\dots+n_k+1}, t_1, \dots, t_{k+1},$

$t'_1, \dots, t'_p, y_1, \dots, y_{m_1+\dots+m_p}$ by the only non-zero bracket products: $SN(n_1, \dots, n_k, 1|m_1, \dots, m_p)$:

$$\left\{ \begin{array}{ll} [x_1, x_j] = -[x_j, x_1] = x_{j+1}, & 2 \leq j \leq n_1; \\ [x_1, x_{n_1+\dots+n_j+i}] = -[x_{n_1+\dots+n_j+i}, x_1] \\ = x_{n_1+\dots+n_j+i+1}, & 1 \leq j \leq k-1, 2 \leq i \leq n_{j+1}; \\ [x_1, y_j] = -[y_j, x_1] = y_{j+1}, & 1 \leq j \leq m_1-1; \\ [x_1, y_{m_1+\dots+m_j+i}] \\ = -[y_{m_1+\dots+m_j+i}, x_1] = y_{m_1+\dots+m_j+i+1}, & 1 \leq j \leq p-1, 1 \leq i \leq m_{j+1}-1; \\ [t_1, x_i] = -[x_i, t_1] = ix_i, & 1 \leq i \leq n_1 + \dots + n_k + 1; \\ [t_1, y_j] = -[y_j, t_1] = jy_j, & 1 \leq j \leq m_1 + \dots + m_p; \\ [t_2, x_i] = -[x_i, t_2] = x_i, & 2 \leq i \leq n_1 + 1; \\ [t_{j+2}, x_{n_1+\dots+n_j+i}] \\ = -[x_{n_1+\dots+n_j+i}, t_{j+2}] = x_{n_1+\dots+n_j+i}, & 1 \leq j \leq k-1, 2 \leq i \leq n_{j+1} + 1; \\ [t'_1, y_i] = -[y_i, t'_1] = y_i, & 1 \leq i \leq m_1; \\ [t'_{j+1}, y_{m_1+\dots+m_j+i}] \\ = -[y_{m_1+\dots+m_j+i}, t'_{j+1}] = y_{m_1+\dots+m_j+i}, & 1 \leq j \leq p-1, 1 \leq i \leq m_{j+1}; \end{array} \right.$$

with $\{x_1, \dots, x_{n_1+\dots+n_k+1}, t_1, \dots, t_{k+1}, t'_1, \dots, t'_p\}$ even basis vectors and $\{y_1, \dots, y_{m_1+\dots+m_p}\}$ odd basis vectors. Next, we show that this solvable Lie superalgebra is the unique with maximal codimension of nilradical $N(n_1, \dots, n_k, 1|m_1, \dots, m_p)$.

Theorem 4.1. *Let L be a complex maximal-dimensional solvable Lie superalgebra with L^2 nilpotent and nilradical isomorphic to $N(n_1, \dots, n_k, 1|m_1, \dots, m_p)$. Then there exists a basis, namely $\{x_1, \dots, x_{n_1+\dots+n_k+1}, t_1, \dots, t_{k+1}, t'_1, \dots, t'_p, y_1, \dots, y_{m_1+\dots+m_p}\}$ with $\{x_1, \dots, x_{n_1+\dots+n_k+1}, t_1, \dots, t_{k+1}, t'_1, \dots, t'_p\}$ even basis vectors and $\{y_1, \dots, y_{m_1+\dots+m_p}\}$ odd basis vectors, in which L is isomorphic to $SN(n_1, \dots, n_k, 1|m_1, \dots, m_p)$.*

Proof. On account of the lack of symmetric bracket products, $N(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ can be regarded as both a nilpotent Lie superalgebra and a nilpotent \mathbb{Z}_2 -graded Lie algebra. Likewise, under the condition of being L^2 nilpotent the techniques used in Lie superalgebras are similar to the ones used in Lie algebras, that is solvable extension by means of (super)derivations of nilradical.

Let us consider now $N(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ as a Lie algebra, the results of [16] allow us to assert that there is only one solvable Lie algebra with maximal codimension of nilradical, which can be expressed in a suitable basis $\{x_1, \dots, x_{n_1+\dots+n_k+1}, z_1, \dots, z_{k+1}, z'_1, \dots, z'_p, y_1, \dots, y_{m_1+\dots+m_p}\}$, by the only non zero bracket products:

$$\left\{ \begin{array}{ll}
 [x_1, x_j] = -[x_j, x_1] = x_{j+1}, & 2 \leq j \leq n_1; \\
 [x_1, x_{n_1+\dots+n_j+i}] \\
 = -[x_{n_1+\dots+n_j+i}, x_1] = x_{n_1+\dots+n_j+i+1}, & 1 \leq j \leq k-1, 2 \leq i \leq n_{j+1}; \\
 [x_1, y_j] = -[y_j, x_1] = y_{j+1}, & 1 \leq j \leq m_1-1; \\
 [x_1, y_{m_1+\dots+m_j+i}] \\
 = -[y_{m_1+\dots+m_j+i}, x_1] = y_{m_1+\dots+m_j+i+1}, & 1 \leq j \leq p-1, 1 \leq i \leq m_{j+1}-1; \\
 [z_1, x_1] = -[x_1, z_1] = x_1; \\
 [z_1, x_i] = -[x_i, z_1] = (i-2)x_i, & 3 \leq i \leq n_1+1; \\
 [z_1, x_{n_1+\dots+n_j+i}] \\
 = -[x_{n_1+\dots+n_j+i}, z_1] = (i-2)x_{n_1+\dots+n_j+i}, & 1 \leq j \leq k-1, 3 \leq i \leq n_{j+1}+1; \\
 [z_1, y_i] = -[y_i, z_1] = (i-1)y_i, & 2 \leq i \leq m_1; \\
 [z_1, y_{m_1+\dots+m_j+i}] \\
 = -[y_{m_1+\dots+m_j+i}, z_1] = (i-1)y_{m_1+\dots+m_j+i}, & 1 \leq j \leq p-1, 2 \leq i \leq m_{j+1}; \\
 [z_2, x_i] = -[x_i, z_2] = x_i, & 2 \leq i \leq n_1+1; \\
 [z_{j+2}, x_{n_1+\dots+n_j+i}] \\
 = -[x_{n_1+\dots+n_j+i}, z_{j+2}] = x_{n_1+\dots+n_j+i}, & 1 \leq j \leq k-1, 2 \leq i \leq n_{j+1}+1; \\
 [z'_1, y_i] = -[y_i, z'_1] = y_i, & 1 \leq i \leq m_1; \\
 [z'_{j+1}, y_{m_1+\dots+m_j+i}] \\
 = -[y_{m_1+\dots+m_j+i}, z'_{j+1}] = y_{m_1+\dots+m_j+i}, & 1 \leq j \leq p-1, 1 \leq i \leq m_{j+1};
 \end{array} \right.$$

and the isomorphism defined by

$$t_1 = z_1 + 2z_2 + \left(\sum_{j=1}^{k-1} (n_1 + \dots + n_j + 2)z_{j+2} \right) + z'_1 + \left(\sum_{j=1}^{p-1} (m_1 + \dots + m_j + 1)z'_{j+1} \right),$$

$$t_i = z_i, \quad 2 \leq i \leq k+1 \quad \text{and} \quad t'_i = z'_i, \quad 1 \leq i \leq p,$$

shows that this Lie algebra is isomorphic to the \mathbb{Z}_2 -graded Lie algebra $SN(n_1, \dots, n_k, 1|m_1, \dots, m_p)$.

Let us consider now $N(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ as a Lie superalgebra. Analogously as it was seen along the proof of Theorem 3.1, the even superderivations are in particular Lie derivations, and for odd superderivations the only condition different from Lie derivation condition is exactly equation (3.3):

$$D([y_i, y_j]) = [D(y_i), y_j] - [y_i, D(y_j)]$$

On account of the law of $N(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ it can be easily seen that the only possibility for having at least one non-null term in the equation (3.3) corresponds with the existence of y_i such that $x_1 \in D(y_i)$. Next, we explore such possibilities. For y_i different from the odd generator vectors, i.e. $i \notin \{1, m_1+1, m_1+m_2+1, \dots, m_1+\dots+m_{p-1}+1\}$, we get

$$D(y_i) = D([x_1, y_{i-1}]) = [D(x_1), y_i] + [x_1, D(y_{i-1})] = [x_1, D(y_{i-1})]$$

As $[x_1, D(y_{i-1})] \in \text{span}\{x_3, \dots, x_n\}$ we can exclude i for $i \notin \{1, m_1 + 1, \dots, m_1 + \dots + m_{p-1} + 1\}$. Suppose now that there exists $i, i \in \{1, m_1 + 1, \dots, m_1 + \dots + m_{p-1} + 1\}$ such that

$$D(y_i) = \alpha_1 x_1 + \sum_{k=2}^{n_1 + \dots + n_k + 1} \alpha_k x_k, \quad \text{with } \alpha_1 \neq 0$$

From equation (3.3) we get in particular

$$D([y_i, y_i]) = 0 = [D(y_i), y_i] - [y_i, D(y_i)] = 2\alpha_1 y_{i+1}$$

and then $\alpha_1 = 0$. Therefore the equation (3.3) vanishes over $N(n_1, \dots, n_k, 1 | m_1, \dots, m_p)$ and consequently all the odd superderivations of $N(n_1, \dots, n_k, 1 | m_1, \dots, m_p)$ are in particular Lie derivations of itself regarded as \mathbb{Z}_2 -graded Lie algebra.

Thus, all the superderivations of the Lie superalgebra $N(n_1, \dots, n_k, 1 | m_1, \dots, m_p)$ are particular cases of Lie derivations of itself regarded as \mathbb{Z}_2 -graded Lie algebra. Therefore, we can assert that $SN(n_1, \dots, n_k, 1 | m_1, \dots, m_p)$ is not only the unique maximal-dimensional solvable Lie algebra with nilradical the Lie algebra $N(n_1, \dots, n_k, 1 | m_1, \dots, m_p)$, but also is the unique maximal-dimensional solvable Lie superalgebra L with L^2 nilpotent and nilradical the model nilpotent Lie superalgebra. \square

5. Maximal-dimensional solvable Leibniz superalgebras with non-Lie filiform nilradical

In this section, we consider Leibniz superalgebra whose nilradical is isomorphic to the filiform (non-Lie) Leibniz superalgebra. This filiform Leibniz superalgebra (denoted by $LP^{n,m}$) can be expressed by the only non-null bracket products that follow:

$$\begin{cases} [x_i, x_1] = x_{i+1}, & 2 \leq i \leq n - 1 \\ [y_j, x_1] = y_{j+1}, & 1 \leq j \leq m - 1 \end{cases}$$

Theorem 5.1. *Let L be a complex maximal-dimensional solvable Leibniz superalgebra with L^2 nilpotent and with nilradical isomorphic to $LP^{n,m}$. Then there exists a basis, namely $\{x_1, \dots, x_n, t_1, t_2, t_3, y_1, \dots, y_m\}$ with $\{x_1, \dots, x_n, t_1, t_2, t_3\}$ a basis of $L_{\overline{0}}$ and $\{y_1, \dots, y_m\}$ a basis of $L_{\overline{1}}$, in which L is isomorphic to the following solvable Leibniz superalgebra:*

$$SLP^{n,m} : \begin{cases} [x_i, x_1] = x_{i+1}, & 2 \leq i \leq n-1; \\ [y_j, x_1] = y_{j+1}, & 1 \leq j \leq m-1; \\ [t_1, x_1] = -x_1, \\ [x_1, t_1] = x_1, \\ [x_i, t_1] = (i-2)x_i, & 3 \leq i \leq n; \\ [y_j, t_1] = (j-1)y_j, & 2 \leq j \leq m; \\ [x_i, t_2] = x_i, & 2 \leq i \leq n; \\ [y_j, t_3] = y_j, & 1 \leq j \leq m; \end{cases}$$

where the omitted products are zero.

Proof. Note that $LP^{n,m}$ can be considered as \mathbb{Z}_2 -graded nilpotent Leibniz algebra. Similar to Lie case, we can use the result of paper [1], which give the description of solvable Leibniz algebras with maximal codimension of nilradical (maximal dimension of the complementary space to nilradical). This maximal codimension is equal to the number of generators of the nilradical, 3 in our case. Then, using Theorem 4 of [1] we have the following products:

$$\begin{aligned} [x_i, x_1] &= x_{i+1}, & 2 \leq i \leq n-1, \\ [y_j, x_1] &= y_{j+1}, & 1 \leq j \leq m-1, \\ [t_1, x_1] &= (b_1 - 1)x_1, \\ [t_2, x_2] &= (b_2 - 1)x_2, \\ [t_3, y_1] &= (b_3 - 1)y_1, \\ [x_1, t_1] &= x_1, \\ [x_i, t_1] &= (i-2)x_i, & 3 \leq i \leq n, \\ [y_j, t_1] &= (j-1)y_j, & 2 \leq j \leq m, \\ [x_i, t_2] &= x_i, & 2 \leq i \leq n, \\ [y_j, t_3] &= y_j, & 1 \leq j \leq m, \end{aligned}$$

with $b_i \in \{0, 1\}$, $1 \leq i \leq 3$. Only rest to determine the following products $[t_k, x_i]$, with $3 \leq i \leq n$, $[t_k, y_j]$, with $2 \leq j \leq m$ and $1 \leq k \leq 3$. Using the Leibniz identity and the induction method we derive the remaining products of $SLP^{n,m}$:

$$\begin{aligned} [t_1, x_i] &= [t_1, y_j] = 0, & 2 \leq i \leq n, 1 \leq j \leq m, \\ [t_2, x_i] &= (b_2 - 1)x_i, & [t_2, y_j] = 0, & 3 \leq i \leq n, 2 \leq j \leq m, \\ [t_3, x_i] &= 0, & [t_3, y_j] &= (b_3 - 1)y_j, & 3 \leq i \leq n, 2 \leq j \leq m. \end{aligned}$$

Finally, from Leibniz identity for the triples $\{x_2, t_1, x_1\}$, $\{t_2, t_1, x_3\}$ and $\{t_3, x_1, y_1\}$ we obtain $b_1 = 0$, $b_2 = 1$ and $b_3 = 1$, respectively.

Now we consider $LP^{n,m}$ as a Leibniz superalgebra. Recall that d is a Leibniz superderivation on $LP^{n,m}$ if d verifies the condition:

$$d([x, y]) = (-1)^{|d||y|}[d(x), y] + [x, d(y)]$$

Analogously to Lie superalgebra, the even superderivations of $LP^{n,m}$ are in particular Leibniz derivations verifying that $d(LP_{\overline{0}}^{n,m}) \subset LP_{\overline{0}}^{n,m}$ and $d(LP_{\overline{1}}^{n,m}) \subset LP_{\overline{1}}^{n,m}$. On the other hand, the odd superderivations d of $LP^{n,m}$ verify the following:

$$d([x_i, x_j]) = [d(x_i), x_j] + [x_i, d(x_j)], \quad x_i, x_j \in LP_{\overline{0}}^{n,m}; \tag{5.1}$$

$$d([x_i, y_j]) = -[d(x_i), y_j] + [x_i, d(y_j)], \quad x_i \in LP_{\overline{0}}^{n,m}, y_j \in LP_{\overline{1}}^{n,m}; \tag{5.2}$$

$$d([y_j, x_i]) = [d(y_j), x_i] + [y_j, d(x_i)], \quad x_i \in LP_{\overline{0}}^{n,m}, y_j \in LP_{\overline{1}}^{n,m}; \tag{5.3}$$

$$d([y_i, y_j]) = -[d(y_i), y_j] + [y_i, d(y_j)], \quad y_i, y_j \in LP_{\overline{1}}^{n,m}. \tag{5.4}$$

Let d be an odd superderivation. Then, we have that

$$d(x_1) = \sum_{k=1}^m a_k y_k, \quad d(x_2) = \sum_{k=1}^m b_k y_k, \quad d(y_1) = \sum_{k=1}^n c_k x_k.$$

Using the equation (5.1) for the pair $[x_1, x_1]$ we get $a_i = 0$ with $1 \leq i \leq m - 1$ and from (5.4) for the pair $[y_1, y_1]$ we obtain $c_1 = 0$. From the equations (5.1) and (5.3) we compute $d(x_i)$ with $3 \leq i \leq n$ and $d(y_j)$ for $2 \leq j \leq m$. Thus, we have:

$$d(x_1) = a_m y_m, \quad d(x_i) = \sum_{k=i-1}^m b_{k-i+2} y_k, \quad 2 \leq i \leq n,$$

$$d(y_j) = \sum_{k=j+1}^n c_{k-j+1} x_k, \quad 1 \leq j \leq m.$$

Finally, from the equation (5.1) for the pair $[x_n, x_1]$ we have that $b_k = 0$ for $1 \leq k \leq m - n + 2$ if $m \geq n - 1$.

It is easy to prove that all odd derivations are Leibniz derivations because the equations (5.2) and (5.4) vanish.

The equations (5.1) and (5.3) correspond with Leibniz derivation condition. Then we can reason as in Theorem 3.1 and we get that all odd superderivations of $LP^{n,m}$ are in particular derivations of the \mathbb{Z}_2 -graded Leibniz algebra $LP^{n,m}$. Then we can assert that $SLP^{n,m}$ is the unique maximal-dimensional solvable Leibniz superalgebra L with L^2 nilpotent and nilradical the Leibniz superalgebra $LP^{n,m}$. \square

6. Maximal-dimensional solvable Leibniz superalgebras with model nilpotent non-Lie nilradical

In this section, we consider as nilradical the equivalent of the model nilpotent Lie superalgebra into (non-Lie) Leibniz superalgebras. This Leibniz superalgebra denoted by $NP(n_1, \dots, n_k, 1 \mid m_1, \dots, m_p)$ can be expressed by the only non-null bracket products that follow:

$$\left\{ \begin{array}{ll} [x_j, x_1] = x_{j+1}, & 2 \leq j \leq n_1, \\ [x_{n_1+\dots+n_j+i}, x_1] = x_{n_1+\dots+n_j+i+1}, & 1 \leq j \leq k-1, 2 \leq i \leq n_{j+1}, \\ [y_j, x_1] = y_{j+1}, & 1 \leq j \leq m_1-1, \\ [y_{m_1+\dots+m_j+i}, x_1] = y_{m_1+\dots+m_j+i+1}, & 1 \leq j \leq p-1, 1 \leq i \leq m_{j+1}-1. \end{array} \right.$$

Theorem 6.1. *Let L be a complex maximal-dimensional solvable Leibniz superalgebra with L^2 nilpotent and nilradical isomorphic to $NP(n_1, \dots, n_k, 1|m_1, \dots, m_p)$. Then there exists a basis, namely $\{x_1, \dots, x_{n_1+\dots+n_k+1}, t_1, \dots, t_{k+1}, t'_1, \dots, t'_p, y_1, \dots, y_{m_1+\dots+m_p}\}$ with $\{x_1, \dots, x_{n_1+\dots+n_k+1}, t_1, \dots, t_{k+1}, t'_1, \dots, t'_p\}$ even basis vectors and $\{y_1, \dots, y_{m_1+\dots+m_p}\}$ odd basis vectors, in which L is isomorphic to $SNP(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ given by:*

$$\left\{ \begin{array}{ll} [x_j, x_1] = x_{j+1}, & 2 \leq j \leq n_1; \\ [x_{n_1+\dots+n_j+i}, x_1] = x_{n_1+\dots+n_j+i+1}, & 1 \leq j \leq k-1, 2 \leq i \leq n_{j+1}; \\ [y_j, x_1] = y_{j+1}, & 1 \leq j \leq m_1-1; \\ [y_{m_1+\dots+m_j+i}, x_1] = y_{m_1+\dots+m_j+i+1}, & 1 \leq j \leq p-1, 1 \leq i \leq m_{j+1}-1; \\ [t_1, x_1] = -x_1, & \\ [x_1, t_1] = x_1, & \\ [x_i, t_1] = (i-2)x_i, & 3 \leq i \leq n_1+1; \\ [x_{n_1+\dots+n_j+i}, t_1] = (i-2)x_{n_1+\dots+n_j+i}, & 1 \leq j \leq k-1, 3 \leq i \leq n_{j+1}+1; \\ [y_j, t_1] = (i-1)y_j, & 2 \leq j \leq m_1; \\ [y_{m_1+\dots+m_j+i}, t_1] = (i-1)y_{m_1+\dots+m_j+i}, & 1 \leq j \leq p-1, 2 \leq i \leq m_{j+1}, \\ [x_i, t_2] = x_i, & 2 \leq i \leq n_1+1; \\ [x_{n_1+\dots+n_j+i}, t_{j+2}] = x_{n_1+\dots+n_j+i}, & 1 \leq j \leq k-1, 2 \leq i \leq n_{j+1}+1; \\ [y_j, t'_1] = y_j, & 1 \leq j \leq m_1; \\ [y_{m_1+\dots+m_j+i}, t'_{j+1}] = y_{m_1+\dots+m_j+i}, & 1 \leq j \leq p-1, 1 \leq i \leq m_{j+1}; \end{array} \right.$$

where the omitted products are zero.

Proof. Similar to Theorem 5.1, we consider the superalgebra $NP(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ as a nilpotent Leibniz algebra (a \mathbb{Z}_2 -graded Leibniz algebra). Thus, we can use the results of the paper [1] and we obtain the following products:

$$\begin{aligned} [t_1, x_1] &= (b_1 - 1)x_1, \\ [x_1, t_1] &= x_1, \\ [x_i, t_1] &= (i - 2)x_i, & 3 \leq i \leq n_1 + 1, \\ [x_{n_1+\dots+n_j+i}, t_1] &= (i - 2)x_{n_1+\dots+n_j+i}, & 1 \leq j \leq k - 1, 3 \leq i \leq n_{j+1} + 1, \\ [y_j, t_1] &= (i - 1)y_j, & 2 \leq j \leq m_1, \\ [y_{m_1+\dots+m_j+i}, t_1] &= (i - 1)y_{m_1+\dots+m_j+i}, & 1 \leq j \leq p - 1, 2 \leq i \leq m_{j+1}, \end{aligned}$$

$$\begin{aligned}
 [t_2, x_2] &= (b_2 - 1)x_2, \\
 [x_2, t_2] &= x_2, \\
 [t_{j+2}, x_{n_1+\dots+n_j+2}] &= (b_{j+2} - 1)x_{n_1+\dots+n_j+2}, \\
 [x_{n_1+\dots+n_j+2}, t_{j+2}] &= x_{n_1+\dots+n_j+2}, \\
 [y_1, t'_1] &= y_1, \\
 [t'_1, y_1] &= (b'_1 - 1)y_1, \\
 [t'_{j+1}, y_{m_1+\dots+m_j+1}] &= (b'_{j+1} - 1)y_{m_1+\dots+m_j+1}, \\
 [y_{m_1+\dots+m_j+1}, t'_{j+1}] &= y_{m_1+\dots+m_j+1}
 \end{aligned}$$

Similar to previous case and applying Leibniz identity we obtain the remaining products and $b_1 = 0$, $b_i = 1$ with $2 \leq i \leq k + 1$ and $b'_j = 1$ with $1 \leq j \leq p$. Thus, we get $SNP(n_1, \dots, n_k, 1|m_1, \dots, m_p)$.

Now, we consider $NP(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ as a superalgebra. The even superderivations of this superalgebra are in particular Leibniz derivations. Then, we go to prove that the odd superderivations are also Leibniz derivations. For that purpose, it is sufficient to verify that the equations (5.2) and (5.4) vanish.

Let d be an odd superderivation. Then, we have

$$\begin{aligned}
 d(x_1) &= \sum_{k=1}^{m_1+\dots+m_p} a_k y_k, & d(x_2) &= \sum_{k=1}^{m_1+\dots+m_p} b_k y_k, & d(x_{n_1+\dots+n_j+2}) \\
 &= \sum_{k=1}^{m_1+\dots+m_p} \alpha_{k j} y_k, & 1 \leq j &\leq k - 1, \\
 d(y_1) &= \sum_{t=1}^{n_1+\dots+n_k+1} c_t x_t, & d(y_{m_1+\dots+m_j+1}) &= \sum_{t=1}^{n_1+\dots+n_k+1} \beta_{t j} x_t, & 1 \leq j \leq p.
 \end{aligned}$$

From the equation (5.1) for the pair $[x_1, x_1]$ we have that

$$d(x_1) = a_{m_1} y_{m_1} + \dots + a_{m_1+\dots+m_p} y_{m_1+\dots+m_p}$$

and for the equation (5.4) for the pair $[y_1, y_1]$ we get to $c_1 = 0$. Analogously, if we consider the pairs $[y_{m_1+\dots+m_j+1}, y_{m_1+\dots+m_j+1}]$ with $1 \leq j \leq p - 1$ we have that $\beta_{1j} = 0$. Thus, it proves that the equation (5.4) is always zero.

On the other hand, we put the equation (5.2). The products $[x_i, d(y_j)]$ are always zero because of $c_1 = \beta_{1j} = 0$ for $1 \leq j \leq p - 1$ in $d(y_j)$. The other products are trivially zero.

Finally, we conclude that the odd superderivations are in particular Leibniz derivations. We can assert that $SNP(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ is the unique maximal-dimensional solvable Leibniz superalgebra L with L^2 nilpotent and nilradical the model nilpotent Leibniz superalgebra $NP(n_1, \dots, n_k, 1|m_1, \dots, m_p)$. \square

7. Superderivations of the maximal-dimensional solvable Lie and Leibniz superalgebras

In this section we establish that the spaces of superderivations for the superalgebras obtained in previous sections (that is, $SL^{n,m}$, $SN(n_1, \dots, n_k, 1|m_1, \dots, m_p)$, $SLP^{n,m}$, $SNP(n_1, \dots, n_k, 1|m_1, \dots, m_p)$) consist of inner superderivations. These results extend analogously results for similar Lie and Leibniz algebras.

Theorem 7.1. *Any superderivation on the Lie superalgebra $SL^{n,m}$ is inner.*

Proof. Our goal is to prove that the following inner superderivations $\{adx_1, \dots, adx_n, adt_1, adt_2, adt_3, ady_1, \dots, ady_m\}$ form a basis of the space of superderivations $Der(SL^{n,m}) = Der_{\bar{0}}(SL^{n,m}) \oplus Der_{\bar{1}}(SL^{n,m})$, with $\{adx_1, \dots, adx_n, adt_1, adt_2, adt_3\}$ a basis of $Der_{\bar{0}}(SL^{n,m})$ and $\{ady_1, \dots, ady_m\}$ a basis of $Der_{\bar{1}}(SL^{n,m})$.

Let D be an even superderivation of $SL^{n,m}$. Then taking into account the embeddings $D(SL_{\bar{0}}^{n,m}) \subset SL_{\bar{0}}^{n,m}$ and $D(SL_{\bar{1}}^{n,m}) \subset SL_{\bar{1}}^{n,m}$ we set

$$\begin{aligned}
 D(x_1) &= \sum_{s=1}^3 \alpha_s t_s + \sum_{k=1}^n a_k x_k, & D(x_2) &= \sum_{s=1}^3 \beta_s t_s + \sum_{k=1}^n b_k x_k, & D(y_1) &= \sum_{r=1}^m p_r y_r, \\
 D(t_1) &= \sum_{s=1}^3 \gamma_s t_s + \sum_{k=1}^n c_k x_k, & D(t_2) &= \sum_{s=1}^3 \delta_s t_s + \sum_{k=1}^n d_k x_k, \\
 D(t_3) &= \sum_{s=1}^3 \nu_s t_s + \sum_{k=1}^n e_k x_k.
 \end{aligned}$$

According to the even superderivation condition, we can summarize the computation in the following table:

Pairs	Constraints
$\{x_1, t_1\}$	$\alpha_i = 0, 1 \leq i \leq 3, \gamma_1 = 0, \alpha_2 = 0, c_k = ka_{k+1}, 2 \leq k \leq n-1$
$\{x_1, t_2\}$	$\delta_1 = 0, d_k = a_{k+1}, 2 \leq k \leq n-1$
$\{x_1, t_3\}$	$\nu_1 = 0, e_k = 0, 2 \leq k \leq n-1$
$\{x_1, y_{j-1}\}$ $2 \leq j \leq m-1$	$d(y_j) = ((j-1)a_1 + p_1)y_j + \sum_{k=j+1}^m p_{k-j+1}y_k, 2 \leq j \leq m$
$\{y_1, x_2\}$	$b_1 = 0, \beta_3 = -\beta_1$
$\{x_1, x_2\}$	$d(x_3) = -\beta_1 x_1 + (a_1 + b_2)x_3 + \sum_{k=4}^n b_{k-1}x_k$
$\{x_3, y_1\}$	$\beta_1 = 0 \Rightarrow \beta_3 = 0$
$\{x_1, x_{i-1}\}$ $4 \leq j \leq n-1$	$d(x_i) = ((i-2)a_1 + b_2)x_i + \sum_{k=i+1}^n b_{k-i+2}x_k, 3 \leq i \leq n$
$\{t_3, t_1\}$	$e_1 = e_n = 0$
$\{t_3, x_2\}$	$\nu_2 = 0$
$\{t_1, x_2\}$	$\beta_2 = \gamma_2 = 0, c_1 = -b_3, b_k = 0, 4 \leq k \leq n$
$\{x_2, t_2\}$	$\delta_2 = d_1 = 0$
$\{t_1, t_2\}$	$c_n = nd_n$
$\{t_1, y_1\}$	$\gamma_3 = 0, p_2 = b_3, p_k = 0, 3 \leq k \leq m$
$\{t_2, y_1\}$	$\delta_3 = 0$
$\{t_3, y_1\}$	$\nu_3 = 0$

Therefore, we get

$$\begin{aligned}
 D(t_1) &= -b_3x_1 + \sum_{k=2}^{n-1} ka_{k+1}x_k + nd_nx_n, & D(t_2) &= \sum_{k=2}^{n-1} a_{k+1}x_k + d_nx_n, & D(t_3) &= 0, \\
 D(x_1) &= a_1x_1 + \sum_{k=3}^n a_kx_k, & D(x_2) &= b_2x_2 + b_3x_3, \\
 D(x_i) &= ((i-2)a_1 + b_2)x_i + b_3x_{i+1}, & & 3 \leq i \leq n, \\
 D(y_1) &= p_1y_1 + b_3y_2, \\
 D(y_j) &= ((j-1)a_1 + p_1)y_j + b_3y_{j+1}, & & 2 \leq j \leq m.
 \end{aligned}$$

Thus, we conclude $\dim(Der_{\bar{0}}(SL^{n,m})) = n+3$. On the other hand, the $(n+3)$ inner superderivations $\{adx_1, \dots, adx_n, adt_1, adt_2, adt_3\}$ are in particular even superderivations. Hence, we obtain a basis of the space $Der_{\bar{0}}(SL^{n,m})$ composed by inner even superderivations. Moreover, D can be expressed via inner as follows

$$D = b_3(adx_1) - \left(\sum_{k=2}^{n-2} a_{k+1}(adx_k) \right) - d_n(adx_n) + a_1(adt_1 - 2adt_2 - adt_3) + b_2(adt_2) + p_1(adt_3).$$

Analogously, we are going to compute the odd superderivations. Let D now be an odd superderivation. We put

$$\begin{aligned}
 D(x_1) &= \sum_{k=1}^m a_ky_k, & D(x_2) &= \sum_{k=1}^m b_ky_k, & D(y_1) &= \sum_{s=1}^3 p_s t_s + \sum_{r=1}^n c_r y_r, \\
 D(t_1) &= \sum_{k=1}^m d_ky_k, & D(t_2) &= \sum_{k=1}^m g_ky_k, & D(t_3) &= \sum_{k=1}^m h_kx_k.
 \end{aligned}$$

According to the odd superderivation conditions we have the following computations:

Pairs	Constraints
$\{x_1, x_{i-1}\}$ $3 \leq j \leq n-1$	$d(x_i) = \sum_{k=i-1}^m b_{k-i+2}y_k, 3 \leq i \leq n$
$\{x_1, y_1\}$	$d(y_2) = -p_1x_1 + \sum_{k=3}^n c_{k-1}x_k$
$\{x_1, y_{j-1}\}$ $3 \leq j \leq m-1$	$d(y_j) = \sum_{k=j+1}^n c_{k-j+1}x_k, 2 \leq j \leq m$
$\{x_2, y_2\}$	$p_1 = 0$
$\{x_2, y_1\}$	$p_2 = c_1 = 0$
$\{t_1, y_1\}$	$p_3 = 0, c_k = 0, 2 \leq k \leq n$
$\{x_1, t_1\}$	$d_k = ka_{k+1}, 1 \leq k \leq m-1$
$\{x_1, t_2\}$	$g_k = 0, 1 \leq k \leq m-1$
$\{x_1, t_3\}$	$h_k = a_{k+1}, 1 \leq k \leq m-1$
$\{x_2, t_1\}$	$b_1 = b_k = 0, 3 \leq k \leq m$
$\{x_2, t_2\}$	$b_2 = 0$
$\{t_1, t_2\}$	$g_m = 0$
$\{t_1, t_3\}$	$d_m = mh_m$

Thus, we get

$$D(t_1) = \sum_{k=1}^{m-1} ka_{k+1}y_k + mh_m y_m, \quad D(t_2) = 0, \quad D(t_3) = \sum_{k=1}^{m-1} a_{k+1}y_k + h_m y_m,$$

$$D(x_1) = \sum_{k=2}^m a_k y_k, \quad D(x_i) = 0, \quad 2 \leq i \leq n, \quad D(y_j) = 0, \quad 1 \leq j \leq m.$$

This implies that $\dim(Der_{\overline{1}}(SL^{n,m})) = m$. On the other hand, we have m odd inner superderivations (they are $\{ady_1, \dots, ady_m\}$). Consequently, a basis of $Der_{\overline{1}}(SL^{n,m})$ form by inner odd superderivations. In particular, D can be expressed via inner superderivations as follows:

$$D = - \left(\sum_{k=1}^{m-1} a_{k+1}(ady_k) \right) - h_m(ady_m). \quad \square$$

Note that all the computations have been duplicated by using the software *Mathematica*.

Consider now the maximal-dimensional solvable Lie superalgebra with model nilpotent nilradical $SN(n_1, \dots, n_k, 1|m_1, \dots, m_p)$.

Theorem 7.2. *Any superderivation of the Lie superalgebra $SN(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ is inner.*

Proof. We are going to prove that the following inner superderivations

$$\{adx_1, \dots, adx_{n_1+\dots+n_k+1}, adt_1, \dots, adt_{k+1}, adt'_1, \dots, adt'_p, ady_1, \dots, ady_{m_1+\dots+m_p}\}$$

form a basis of the space $Der(SN(n_1, \dots, n_k, 1|m_1, \dots, m_p))$ with $\{adx_1, \dots, adx_{n_1+\dots+n_k+1}, adt_1, \dots, adt_{k+1}, adt'_1, \dots, adt'_p\}$ a basis for even superderivations, $Der_{\overline{0}}(SN(n_1, \dots, n_k, 1|m_1, \dots, m_p))$, and $\{ady_1, \dots, ady_{m_1+\dots+m_p}\}$ a basis for the odd ones, $Der_{\overline{1}}(SN(n_1, \dots, n_k, 1|m_1, \dots, m_p))$.

Let D be an even superderivation of $SN(n_1, \dots, n_k, 1|m_1, \dots, m_p)$. Then from superderivation property we derive

$$D(x_1) = \alpha_1 x_1 + \sum_{s=3}^{n_1+1} \alpha_s x_s + \sum_{j=1}^{k-1} \left(\sum_{s=n_1+\dots+n_j+3}^{n_1+\dots+n_{j+1}} \alpha_s x_s \right),$$

$$D(x_i) = ((i - 2)\alpha_2 + \beta)x_i, \quad 2 \leq i \leq n_1 + 1,$$

$$D(x_{n_1+\dots+n_j+i}) = ((i - 2)\alpha_1 + a_j)x_{n_1+\dots+n_j+i}, \quad 1 \leq j \leq k - 1, \quad 2 \leq i \leq n_{j+1} + 1,$$

$$D(t_1) = \sum_{s=2}^{n_1} s\alpha_{s+1}x_s + \sum_{j=1}^{k-1} \left(\sum_{s=n_1+\dots+n_j+2}^{n_1+\dots+n_{j+1}} s\alpha_{s+1}x_s \right) +$$

$$\begin{aligned}
 &+ \sum_{j=1}^k (n_1 + \dots + n_j + 1) b_j x_{n_1 + \dots + n_j + 1}, \\
 D(t_2) &= \sum_{s=2}^{n_1} \alpha_{s+1} x_s + b_1 x_{n_1 + 1}, \\
 D(t_{j+2}) &= \sum_{s=n_1 + \dots + n_j + 2}^{n_1 + \dots + n_{j+1}} \alpha_{s+1} x_s + b_{j+1} x_{n_1 + \dots + n_{j+1} + 1}, \quad 1 \leq j \leq k - 1, \\
 D(t'_j) &= 0, \quad 1 \leq j \leq p, \\
 D(y_i) &= ((i - 1)\alpha_1 + \gamma) y_i, \quad 1 \leq i \leq m_1, \\
 D(y_{m_1 + \dots + m_j + i}) &= ((i - 1)\alpha_1 + q_j) y_{m_1 + \dots + m_j + i}, \quad 1 \leq j \leq p - 1, 1 \leq i \leq m_{j+1}.
 \end{aligned}$$

Let D be an odd superderivation of $SN(n_1, \dots, n_k, 1 | m_1, \dots, m_p)$. Then the superderivation property implies

$$\begin{aligned}
 D(x_1) &= \sum_{s=2}^{m_1} \alpha_s y_s + \sum_{j=1}^{p-1} \left(\sum_{s=m_1 + \dots + m_j + 2}^{m_1 + \dots + m_{j+1}} \alpha_s y_s \right), \\
 D(x_i) &= 0, \quad 2 \leq i \leq n_1 + \dots + n_k + 1, \\
 D(t_1) &= \sum_{s=1}^{m_1 - 1} s \alpha_{s+1} y_s + \sum_{j=1}^{p-1} \left(\sum_{s=m_1 + \dots + m_j + 1}^{m_1 + \dots + m_{j+1} - 1} s \alpha_{s+1} y_s \right) \\
 &\quad + \sum_{j=1}^p (m_1 + \dots + m_j) \delta_j y_{m_1 + \dots + m_j}, \\
 D(t_i) &= 0, \quad 2 \leq i \leq k + 1, \\
 D(t'_1) &= \sum_{s=1}^{m_1 - 1} \alpha_{s+1} y_s + \sum_{j=1}^p \delta_j y_{m_1 + \dots + m_j}, \\
 D(t'_i) &= 0, \quad 2 \leq i \leq p, \\
 D(y_j) &= 0, \quad 1 \leq j \leq m_1 + \dots + m_p.
 \end{aligned}$$

Therefore, we have

$$\begin{aligned}
 \dim(Der_{\overline{0}}(SN(n_1, \dots, n_k, 1 | m_1, \dots, m_p))) &= (n_1 + \dots + n_k + 1) + k + 1 + p, \\
 \dim(Der_{\overline{1}}(SN(n_1, \dots, n_k, 1 | m_1, \dots, m_p))) &= m_1 + \dots + m_p.
 \end{aligned}$$

Now, since both sets of inner superderivations of the statement of the Theorem, even and odd, are linearly independent we conclude that the set

$$\{adx_1, \dots, adx_{n_1 + \dots + n_k + 1}, adt_1, \dots, adt_{k+1}, adt'_1, \dots, adt'_p, ady_1, \dots, ady_{m_1 + \dots + m_p}\}$$

constitutes a basis of the superalgebra of superderivations $Der(SN(n_1, \dots, n_k, 1|m_1, \dots, m_p))$. \square

Since the proofs of the following results based on the application the same arguments and similar computations as in the proofs of Theorems 7.1 and 7.2 we present summaries of their proofs.

Theorem 7.3. *Any superderivation of the Leibniz superalgebra $SLP^{n,m}$ is inner.*

Proof. As a result of computing of the odd and even superderivations properties of the Leibniz superalgebra $SLP^{n,m}$ we obtain $Der_{\bar{1}}(SLP^{n,m}) = \{0\}$ and for an arbitrary $d \in Der_{\bar{0}}(SLP^{n,m})$ we get

$$\begin{aligned} d(x_1) &= \alpha x_1, & d(x_i) &= (\beta + (i - 2)\alpha)x_i - \gamma x_{i+1}, & 2 \leq i \leq n, \\ d(t_1) &= \gamma x_1, & d(t_2) &= d(t_3) = 0, \\ d(y_1) &= \delta y_1 - \gamma y_2, & d(y_j) &= (\delta + (j - 1)\alpha)y_j - \gamma y_{j+1}, & 2 \leq j \leq m, \end{aligned}$$

for some parameters $\alpha, \beta, \gamma, \delta$.

Consequently, $dim(Der(SLP^{n,m})) = 4$ and hence, we obtain $Der(SLP^{n,m}) = span\{R_{x_1}, R_{t_1}, R_{t_2}, R_{t_3}\}$. In particular, d can be expressed via inner derivations as follows:

$$d = -\gamma R_{x_1} + \alpha R_{t_1} + \beta R_{t_2} + \delta R_{t_3}. \quad \square$$

Theorem 7.4. *Any superderivation of the Leibniz superalgebra $SNP(n_1, \dots, n_k, 1|m_1, \dots, m_p)$ is inner.*

Proof. Analogously to the previous superalgebra, we obtain that $Der_{\bar{1}}(SLP(n_1, \dots, n_k, 1|m_1, \dots, m_p)) = \{0\}$ and for an arbitrary even superderivation d we derive the following:

$$\begin{aligned} d(x_1) &= \alpha x_1, \\ d(x_i) &= (\beta + (i - 2)\alpha)x_i - \gamma x_{i+1}, & 3 \leq i \leq n_1, \\ d(x_{n_1+1}) &= (\beta + (n_1 - 1)\alpha)x_{n_1+1}, \\ d(x_{n_1+\dots+n_j+2}) &= \mu_j x_{n_1+\dots+n_j+2} - \gamma x_{n_1+\dots+n_j+3}, & 1 \leq j \leq k - 1, \\ d(x_{n_1+\dots+n_j+i}) &= (\mu_j + (i - 2)\alpha)x_{n_1+\dots+n_j+i} - \gamma x_{n_1+\dots+n_j+i+1}, & 1 \leq j \leq k - 1, 3 \leq j \leq n_{j+1}, \\ d(x_{n_1+\dots+n_{j+1}+1}) &= (\mu_j + (n_{j+1} - 1)\alpha)x_{n_1+\dots+n_{j+1}+1}, & 1 \leq j \leq k - 1, \\ d(t_1) &= \gamma x_1, \\ d(t_i) &= d(t'_j) = 0, & 2 \leq i \leq k + 1, 1 \leq j \leq p, \\ d(y_1) &= \delta y_1 - \gamma y_2, \\ d(y_j) &= (\delta + (j - 1)\alpha)y_j - \gamma y_{j+1}, & 2 \leq j \leq m_1 - 1, \end{aligned}$$

$$\begin{aligned}
d(y_{m_1}) &= (\delta + (m_1 - 1)\alpha)y_{m_1}, \\
d(y_{m_1+\dots+m_s+1}) &= \nu_s y_{m_1+\dots+m_s+1} - \gamma y_{m_1+\dots+m_s+2}, \quad 1 \leq s \leq p-1, \\
d(y_{m_1+\dots+m_s+i}) &= (\nu_s + (i-1)\alpha)y_{m_1+\dots+m_s+i} - \gamma y_{m_1+\dots+m_s+i+1}, \quad 1 \leq s \leq p-1, \quad 2 \leq i \leq m_{s+1}-1, \\
d(y_{m_1+\dots+m_{s+1}}) &= (\nu_s + (m_{s+1}-1)\alpha)y_{m_1+\dots+m_{s+1}}, \quad 1 \leq s \leq p-1.
\end{aligned}$$

Then, $\dim(\text{Der}(\text{SLP}(n_1, \dots, n_k, 1|m_1, \dots, m_p))) = k + p + 2$. On the other hand, we have $k + p + 2$ inner derivations, $\{R_{x_1}, R_{t_1}, R_{t_2}, R_{t_3}, \dots, R_{t_{k+1}}, R_{t'_1}, R_{t'_2}, \dots, R_{t'_p}\}$. In particular, d can be expressed via inner derivations as follows:

$$d = -\gamma R_{x_1} + \alpha R_{t_1} + \beta R_{t_2} + \sum_{j=1}^{k-1} \mu_j R_{t_{j+2}} + \delta R_{t'_1} + \sum_{s=1}^{p-1} \nu_s R_{t'_{s+1}}. \quad \square$$

References

- [1] K.K. Abdurasulov, Maximal solvable extension of nilpotent Leibniz algebras, *Proc. Uzbek. Acad. Sci.* 5 (2019) 3–8.
- [2] S. Albeverio, Sh.A. Ayupov, B.A. Omirov, On nilpotent and simple Leibniz algebras, *Commun. Algebra* 33 (1) (2005) 159–172.
- [3] S. Albeverio, B.A. Omirov, I.S. Rakhimov, Varieties of nilpotent complex Leibniz algebras of dimensions less than five, *Commun. Algebra* 33 (5) (2005) 1575–1585.
- [4] J.M. Ancochea Bermúdez, R. Campoamor-Stursberg, Cohomologically rigid solvable Lie algebras with a nilradical of arbitrary characteristic sequence, *Linear Algebra Appl.* 488 (2016) 135–147.
- [5] J.M. Ancochea Bermúdez, R. Campoamor-Stursberg, L. García Vergnolle, Indecomposable Lie algebras with nontrivial Levi decomposition cannot have filiform radical, *Int. Math. Forum* 1 (2006) 309–316.
- [6] J.M. Ancochea Bermúdez, R. Campoamor-Stursberg, L. García Vergnolle, Classification of Lie algebras with naturally graded quasi-filiform nilradicals, *J. Geom. Phys.* 61 (11) (2011) 2168–2186.
- [7] M. Bordemann, J.R. Gómez, Yu. Khakimjanov, R.M. Navarro, Some deformations of nilpotent Lie superalgebras, *J. Geom. Phys.* 57 (5) (2007) 1391–1403.
- [8] D. Burde, C. Steinhoff, Classification of orbit closures of 4-dimensional complex Lie algebras, *J. Algebra* 214 (1999) 729–739.
- [9] L.M. Camacho, J.M. Fernández-Barroso, R.M. Navarro, Classification of solvable Lie and Leibniz superalgebras with a given nilradical, *Forum Math.* 32 (5) (2020) 1271–1288, <https://doi.org/10.1515/forum-2020-0031>.
- [10] L.M. Camacho, B.A. Omirov, K.K. Masutova, Solvable Leibniz algebras with filiform nilradical, *Bull. Malays. Math. Sci. Soc.* 39 (1) (2016) 283–303.
- [11] J.M. Casas, M. Ladra, B.A. Omirov, I.A. Karimjanov, Classification of solvable Leibniz algebras with null-filiform nilradical, *Linear Multilinear Algebra* 61 (6) (2013) 758–774.
- [12] J.M. Casas, M. Ladra, B.A. Omirov, I.A. Karimjanov, Classification of solvable Leibniz algebras with naturally graded filiform nilradical, *Linear Algebra Appl.* 438 (7) (2013) 2973–3000.
- [13] M. Goze, Yu. Khakimjanov, *Nilpotent Lie Algebras, Mathematics and Its Applications*, vol. 361, Kluwer Academic Publishers Group, Dordrecht, ISBN 0-7923-3932-0, 1996, xvi+336 pp.
- [14] F. Grunewald, J. O’Halloran, Varieties of nilpotent Lie algebras of dimension less than six, *J. Algebra* 112 (2) (1988) 315–326.
- [15] V.G. Kac, A sketch of Lie superalgebra theory, *Commun. Math. Phys.* 53 (1977) 31–64.
- [16] Kh.A. Khalkulova, K.K. Abdurasulov, Solvable Lie algebras with maximal dimension of complementary space to nilradical, *Uzbek Math. J.* 1 (2018) 90–98, <https://doi.org/10.29229/uzmj.2018-1-8>.
- [17] A.Kh. Khudoyberdiyev, B.A. Omirov, Infinitesimal deformations of null-filiform Leibniz superalgebras, *J. Geom. Phys.* 74 (2013) 370–380.
- [18] M. Ladra, K.K. Masutova, B.A. Omirov, Corrigendum to “Classification of solvable Leibniz algebras with naturally graded filiform nilradical” [*Linear Algebra Appl.* 438 (7) (2013) 2973–3000], *Linear Algebra Appl.* 507 (2016) 513–517.

- [19] A.I. Malcev, Solvable Lie algebras, *Transl. Am. Math. Soc.* 27 (1950).
- [20] G.M. Mubarakzjanov, On solvable Lie algebras, *Izv. Vysš. Učebn. Zaved., Mat.* 1 (32) (1963) 114–123 (in Russian).
- [21] J.C. Ndogmo, P. Winternitz, Solvable Lie algebras with Abelian nilradicals, *J. Phys. A* 27 (1994) 405–423.
- [22] Dušan D. Repovš, Mikhail V. Zaicev, Codimension growth of solvable Lie superalgebras, *J. Lie Theory* 28 (4) (2018) 1189–1199.
- [23] M.C. Rodríguez-Vallarte, G. Salgado, On indecomposable solvable Lie superalgebras having a Heisenberg nilradical, *J. Algebra Appl.* 15 (10) (2016) 1650190.
- [24] L. Šnobl, D. Karásek, Classification of solvable Lie algebras with a given nilradical by means of solvable extensions of its subalgebras, *Linear Algebra Appl.* 432 (7) (2010) 1836–1850.
- [25] S. Tremblay, P. Winternitz, Solvable Lie algebras with triangular nilradicals, *J. Phys. A* 31 (1998) 789–806.
- [26] Y. Wang, J. Lin, S. Deng, Solvable Lie algebras with quasifiliform nilradicals, *Commun. Algebra* 36 (2008) 4052–4067.