

Effects of *Chrysochlorum (Aphanizomenon) ovalisporum* extracts containing cylindrospermopsin on growth, photosynthetic capacity, and mineral content of carrots (*Daucus carota*)

Remedios Guzmán-Guillén¹ · Alexandre Campos² · Joana Machado² ·
Marisa Freitas^{2,4} · Joana Azevedo² · Edgar Pinto^{3,4} · Agostinho Almeida³ ·
Ana M. Cameán¹ · Vitor Vasconcelos^{2,5}

Abstract Natural toxins produced by freshwater cyanobacteria, such as cylindrospermopsin, have been regarded as an emergent environmental threat. Despite the risks for food safety, the impact of these water contaminants in agriculture is not yet fully understood. Carrots (*Daucus carota*) are root vegetables, extensively consumed worldwide with great importance for human nourishment and economy. It is, therefore, important to evaluate the possible effects of using water contaminated with cyanotoxins on carrot cultivation. The aim of this work was to investigate cylindrospermopsin effects on *D. carota* grown in soil and irrigated for 30 days, with a *Chrysochlorum ovalisporum* extract containing environmentally relevant concentrations of cylindrospermopsin (10 and 50 µg/L).

The parameters evaluated were plant growth, photosynthetic capacity, and nutritional value (mineral content) in roots of carrots, as these are the edible parts of this plant crop. The results show that, exposure to cylindrospermopsin did not have a clear negative effect on growth or photosynthesis of *D. carota*, even leading to an increase of both parameters. However, alterations in mineral contents were detected after exposure to crude extracts of *C. ovalisporum* containing cylindrospermopsin. A general decline was observed for most minerals (Ca, Mg, Na, Fe, Mn, Zn, Mo, and P), although an increase was shown in the case of K and Cu, pointing to a possible interference of the cyanobacterial extract in mineral uptake. This study is the first to evaluate the effects of *C. ovalisporum* extracts on a root vegetable, however, more research is necessary to understand the effects of this toxin in environmentally relevant scenarios.

Keywords Cylindrospermopsin · *Daucus carota* · Food safety · Photosynthesis · Minerals

✉ Alexandre Campos
acampos@ciimar.up.pt

¹ Area of Toxicology, Department of Pharmacy, University of Sevilla, Sevilla, Spain

² Interdisciplinary Centre of Marine and Environmental Research (CIIMAR/CIMAR), Porto 4050-123, Portugal

³ LAQV/REQUIMTE, Department of Chemical Sciences, Laboratory of Applied Chemistry, Department of Pharmacy, University of Porto, Rua de Jorge Viterbo Ferreira 228, Porto 4050-313, Portugal

⁴ Department of Environmental Health, Escola Superior de Tecnologia da Saúde do Porto, Polytechnic Institute of Porto, CISA/Research Center in Environment and Health, Rua de Valente Perfeito, 322, Gaia 4400-330, Portugal

⁵ Department of Biology, Department of Sciences of the University of Porto, Porto 4169-007, Portugal

Introduction

Toxic cyanobacterial blooms are increasing in aquatic ecosystems, potentially as a consequence of eutrophication and climate change. They can have detrimental effects on the aquatic ecosystems by altering their trophic structure and functionality (Havens 2008), causing mortality to aquatic organisms, and decreasing water quality (Quiblier et al. 2013). Among the cyanobacterial toxins, cylindrospermopsin (CYN) is the most emergent and of increased concern (Elliott 2012). Due to its multiple toxicity targets, severe impacts on health can occur, making CYN a

compound of global concern in water management (De la Cruz et al. 2013).

Currently, 13 species of cyanobacteria are known to produce CYN on four of the five continents (Oceania, America, Asia, and Europe), with *Chrysochloris ovalisporum* being one of the main producers (Moreira et al. 2012). The presence of CYN has been detected predominantly in tropical and subtropical waters, such as lakes, rivers, ponds, and reservoirs (Pearson et al. 2010). Furthermore, the fact that CYN is highly water-soluble because of its zwitterionic structure, leads to significant extracellular levels of this toxin in waters (70–98 %) (Rücker et al. 2007; Van Apeldoorn et al. 2007). Field studies have found a wide range of CYN concentrations in waters of different countries, reaching values from 1.1 to 800 µg/L in Australia (Saker and Eaglesham 1999; Hoeger et al. 2004; Shaw et al. 2009), 97.1 µg/L in Florida (Burns 2008), 12.1 µg/L in Germany (Rücker et al. 2007), 18.4 µg/L in Italy (Bogialli et al. 2006), and 9.4 µg/L in Spain (Quesada et al. 2006). All of these concentrations exceed the guideline value proposed by Humpage and Falconer (2003) for CYN in drinking water (1 µg/L).

The mechanism of toxicity of CYN is mediated by inhibition of protein synthesis, as well as genotoxicity by DNA fragmentation (Frosio et al. 2008; Humpage 2008; Bazin et al. 2010; Żegura et al. 2011). In addition, some studies have reported the induction of oxidative stress in different plant species (Prieto et al. 2011; Freitas et al. 2015). Humans are exposed to cyanobacterial toxins through different routes, including drinking water, recreational contact, food supplements made from cyanobacteria, and through the food chain (Gutiérrez-Praena et al. 2013). Several cyanobacterial toxins have been investigated in regard to their ability to enter the food chain via food (Ibelings and Chorus 2007; Saker et al. 2004; Gutiérrez-Praena et al. 2013; Corbel et al. 2014). The use of cyanotoxin-contaminated water for agricultural purposes may represent a potential risk to food safety and security, and is of particular concern. Indeed, several studies have reported the negative impact of cyanotoxins on the yield, quality and safety of crop plants, as it was recently reviewed by De la Cruz et al. (2013).

To date, the effects of exposure of edible plants to cyanotoxins have been described for rice exposed to microcystins (MC) and CYN (Prieto et al. 2011; Azevedo et al. 2014), tomatoes and carrots exposed to MC alone (Gutiérrez-Praena et al. 2014; Machado 2014), and lettuce contaminated with microcystin-LR (MC-LR), with leucine (L) and arginine (R) as variable L-amino acids in the chemical structure of the toxin, CYN and a mixture of both toxins (Freitas et al. 2015). Although the effects of CYN on plants have been studied to a much lesser extent than other cyanotoxins such as MC, this toxin is expected to become

increasingly recurrent and thus enhancing the knowledge of its impact on crop plants is necessary. Moreover, MC are predominantly intracellular and CYN extracellular due to its high water-solubility; this fact may also make the effects of these toxins very different.

Few studies have indicated so far several effects on plant species exposed to CYN. White mustard (*Sinapis alba*) experienced growth inhibition (Vasas et al. 2002), with reduced fresh mass and the length of cotyledons (M-Hamvas et al. 2010) after exposure to concentrations of CYN between 5 and 10 µg/mL. Alterations in growth were also shown in the macrophyte *Phragmites australis* (Beyer et al. 2009), as well as reduction in pollen germination in tobacco plants (*Nicotiana tabacum*) (Metcalf et al. 2004), and the induction of oxidative stress in rice (*Oryza sativa*) (Prieto et al. 2011). In addition, exposure of *Hydrilla verticillata* to *Cylindrospermopsis raciborskii* whole-cell extracts (25–400 µg/L) resulted in decreases in chlorophyll and changes in the chlorophyll a:b ratio, indicating complex effects that extracts might be exerting on photosynthesis (Kinnear et al. 2008). Moreover, chromatin alterations, chromosomal aberrations and mitotic delay with short-term exposures have been reported in *Vicia faba* cells (Garda et al. 2015).

However, the uptake of toxins by plants may induce not only morphological and physiological changes leading to a loss of productivity, but it could also have considerable implications in their nutritional value, as well as in toxin bioaccumulation. Recently, Kittler et al. (2012) reported that treatment of *Brassica oleracea* var. *sabellica*, *Brassica juncea*, and *S. alba* under varying experimental conditions showed significant CYN uptake, with CYN levels ranging from 10 to 21 % in the leaves compared to the CYN concentration applied to the roots (18–35 µg/L). This work demonstrates that crop plants irrigated with CYN-containing waters may represent an important source of this toxin within the food chain, especially when the cyanotoxins are present in high concentrations in the irrigation water (Pflugmacher et al. 2006). The plant root system is usually more exposed via soil irrigation, and therefore root vegetables are more susceptible to contamination.

Minerals are essential to plant growth and development; they are intrinsic components for normal structure, metabolism and function (Pinto and Ferreira 2015). Interestingly, Saqrane et al. (2009) have reported that exposure of *Triticum durum*, *Zea mays*, *Pisum sativum*, and *Lens esculenta* plants to MC-containing extracts resulted in changes in the mineral content in roots in a concentration-dependent manner. More recently, El Khalloufi et al. (2012) and Lahrouni et al. (2013) demonstrated that cyanobacterial bloom extracts containing MC induced changes in mineral assimilation and content in tomato (*Lycopersicon esculentum*) and faba bean (*V. faba*), respectively. According to

our knowledge, only the study developed by Freitas et al. (2015) reported effects of CYN on the mineral content of plants. Therefore, because carrots are root vegetables used for animal and direct human consumption, and taking into account human health and agricultural economics, it is important to evaluate the possible effects of using water contaminated with CYN for irrigation of this crop.

Regarding the effects of cyanotoxins on plants, most works were based on short-term toxicological studies (5 days) with high toxin concentrations or were conducted with other toxins different from CYN: 1500–20,000 µg/L in Mitrovic et al. (2005), 5000–10,000 µg/L in M-Hamvas et al. (2010), and 2220–22,240 µg/L in El Khalloufi et al. (2011, 2012), and almost all plants were tested in early stages of development. Recently, two studies took into account environmentally relevant cyanotoxin concentrations for carrots exposed to MC (Machado 2014) and lettuce plants exposed to MC-LR and CYN (Freitas et al. 2015). However, the adverse effects of CYN exposure on carrot plants have not been previously investigated.

In this context, the aim of the present study was to assess the effects of CYN on growth, photosynthetic capacity and nutritional value (mineral content) of carrot plants (*Daucus carota*), at the stage of the growth and differentiation of the taproot (carrot), in soil in natural conditions, and irrigated every 2 days with non-contaminated water or with a *C. ovalisporum* extract containing environmentally relevant concentrations of CYN (10 and 50 µg/L).

Materials and methods

Chrysochlorum ovalisporum culture

Chrysochlorum ovalisporum (LEGE X-001) cyanobacterial CYN-producing strain (CYN+) was isolated from Lake Kinneret, Israel (Banker et al. 1997) and was grown for biomass production in the Interdisciplinary Centre of Marine and Environmental Research, CIIMAR (Porto, Portugal) as described in Guzmán-Guillén et al. (2014).

CYN extraction, purification, and quantification

The extraction and purification of CYN from the culture was performed according to Guzmán-Guillén et al. (2012) with modifications. Briefly, the lyophilized biomass (0.5 g) was extracted with 100 mL of MilliQ water, sonicated (15 min), stirred (1 h), and sonicated again (15 min). The resulting mixture was centrifuged (4500 rpm, 10 min), after which the supernatant was collected and 0.1 % TFA was added. The purified CYN fractions were then quantified in an HPLC system on an Atlantis® HILIC phase column (250 × 4.6 mm, 5 µm) from Waters (Milford, USA)

maintained at 40 °C. The PDA range was 210–400 nm with a fixed wavelength of 262 nm. The isocratic elution was also a 5 % MeOH solution containing 2 mM sodium 1-heptanesulfonate monohydrate (99 %) with a flow rate of 0.9 mL/min and an injection volume of 10 µL. The system was calibrated using a set of seven dilutions of the CYN standard (25, 20, 10, 5, 2, 1, and 0.5 µg/mL) in ultrapure water. Each vial was injected in duplicate, and every HPLC run consisting of a series of 10 samples included a blank and two different standard concentrations. The chromatographic purity of CYN was 98 %. The Empower 2 Chromatography Data Software was used for the calculation and reporting the peak information. The retention time of the CYN peak was 7.35 min, and the concentration obtained was 27.2 µg CYN/mL.

Plant material and exposure to CYN

Carrot plants (*Daucus carota* cv. Soprano), approximately 1-month-old and starting the growth and differentiation of the taproots, were acquired from a nursery, taken to the laboratory, transplanted into clear plastic cups (350 mL) with soil, and placed on black trays. Plants were placed in outside conditions, in July 2014, and allowed to acclimate for 1 week before the beginning of the experiment to the natural light, temperature, and air humidity levels. The soil used was Siro® Plant Universal Substrate and its main physicochemical properties were: pH (CaCl₂), 5.5–6.5; soil moisture content, 50–60 %; conductivity, 0.6–1.2 dS/m; N, 150–250 mg/L; P₂O₅, 150–250 mg/L; K₂O, 300–500 mg/L; organic material, >70 %. During the test (30 days, in July), carrots were submitted three times a week to irrigation with 40 mL of the crude extract of *C. ovalisporum* diluted in water to obtain different concentrations of CYN (10 and 50 µg CYN/L), which are considered environmentally relevant. The toxin in the crude extract was quantified on days that it was administered, obtaining an average content of 27.36 µg CYN/mL (levels of CYN in the *C. ovalisporum* extract were maintained generally constant with no noticeable variations, in the range 27.11–27.79 µg CYN/mL). Thus, dilutions were accurately prepared to always achieve the same concentration of 10 and 50 µg CYN/L. Additional irrigation was applied to keep the soil moist during the experiment, with similar volume of clean water (free of toxin) to each plant. A soil moisture sensor (Leroy Merlin, Ref.14785393, Lille, France) with 4 intensity levels (A to D, where A corresponds to dry soil and D to water saturated soil) was used to control soil moisture. In the experiment, moisture levels in the soil were between levels B and C.

There were 9 trays, with 17 carrot plants per tray for a total of 153 plants (51 plants for each of the control group, 10 and 50 µg CYN/L). A 30-days exposure was performed outside under natural light, temperature, and humidity

conditions, allowing plants to develop enlarged and orange taproots (carrots). At the end of the exposure experiment, plants were analyzed with respect to the photosynthetic capacity ($n = 20$ per condition), growth ($n = 37$ per condition), and mineral content ($n = 4$ per condition).

Determination of photosynthetic capacity

Twenty randomly harvested carrots per condition (control, 10, and 50 μg CYN/L) were selected to nondestructively measure the photosynthetic capacity of the leaves by pulse amplitude modulation (PAM) using PAM 2000 (Walz, Effeltrich, Germany) on day 30. Plants were transferred to a dark room where they remained for 30 min. After the adjustment period, the leaves were illuminated with a saturating light pulse and the emitted fluorescence was immediately recorded by the instrument. This procedure allows measuring the maximum efficiency (F_v/F_m), which is directly related to the functional state of photosystem II (PSII) protein complex and photosynthetic efficiency of the plants (Maxwell and Johnson 2000).

Determination of plant growth

After 30 days of CYN exposure, carrot plants were harvested, and roots and leaves separated. The fresh weight (fw) of roots of 37 randomly selected plants per condition (control, 10, and 50 μg CYN/L) was determined on a semi-analytical balance. Root growth was expressed as the mean fresh weight (fw) \pm standard deviation (SD) of $n = 37$.

Determination of mineral content in carrots

At the harvest, four randomly selected plants per exposure condition were taken for determination of mineral content ($n = 4$; 10 g/plant). The root system of each plant, constituted mainly by the differentiated and enlarged taproot (carrot), was harvested and washed with deionized water to remove soil contaminants, weighed, frozen at -80°C and lyophilized (Telstar Lyoquest) for 1 week. Carrot samples were ground and homogenized in a blade homogenizer (GRINDOMIX GM 200, Retsch, Haan, Germany). The microwave-assisted acid digestion and the elemental analysis were performed according to Pinto et al. (2014a). For the microwave-assisted acid digestion, 200 mg of lyophilized sample were weighed in polytetrafluoroethylene (PTFE) vessels and 3 mL of HNO_3 ($\geq 69\%$, w/w) plus 1 mL of H_2O_2 (30%, v/v) were added. Then, the PTFE vessels were placed in the microwave digestion unit (MLS 1200 Mega, Milestone, Sorisole, Italy), which was equipped with a rotor HPR-1000/10 S, and subjected to the following digestion program: 250 W for 1 min, 0 W for 2 min, 250 W for 5 min, 400 W for 5 min, and 600 W for 5 min. Once digested, the sample solutions were

transferred to 25 mL volumetric flasks, and the volume was made up with ultrapure water. This procedure was performed for all samples, as well as for the certified reference material. In each digestion, a blank was included, consisting of the digestion process in the absence of biological material.

Sample solutions were analyzed for their content of Ca, Mg, Na, and K by a 3100 flame (air-acetylene) atomic absorption spectrometer (Perkin Elmer, Überlingen, Germany). Cathode lamps (Perkin Elmer, Überlingen, Germany) were used as a radiation source. The multi-element calibration standards were prepared from 1000 mg/L standard solutions of one element of Ca, Mg, Na, and K (Sigma, St. Louis MO, USA). A lanthanum solution (10% w/v) was used in the determination of Ca and Mg to eliminate chemical interferences.

The inductively coupled plasma-mass spectrometry (ICP-MS) equipment used for the analysis of Fe, Mn, Zn, Mo, Cu, and P was a VG Elemental, PlasmaQuad 3 model (Winsford, UK). A Minipuls 3 peristaltic pump (Gilson, Villiers le Bel, France) was used for sample introduction. Argon of 99.9997% purity (Gasin II, Leça da Palmeira, Portugal) was used as the plasma source. The ICP-MS analysis was performed under the following conditions: argon flow rate (13 L/min); auxiliary argon flow rate (0.7 L/min); nebulizer flow rate (0.8 L/min); RF power (1350 W); scan regions dwell time (100 ms); and detection mode (pulse counting). The elemental isotopes (m/z ratios) ^{31}P , ^{55}Mn , ^{56}Fe , ^{65}Cu , ^{66}Zn and ^{95}Mo were monitored for analytical determination; ^{45}Sc , ^{89}Y , ^{115}In , ^{159}Tb and ^{209}Bi were used as internal standards. The instrument was tuned daily for maximum signal sensitivity and stability using ^{115}In as the target isotope. For this analysis, calibration standards were prepared from the AccuStandard[®] (New Haven, CT) 10 $\mu\text{g}/\text{mL}$ multi-element ICP-MS standard solution (ICP-MS-200.8-CAL1-1) and the 1000 $\mu\text{g}/\text{mL}$ single-element standard solutions (P ion chromatography standard). Results were expressed on a fresh weight basis.

All materials used in the preparation of the samples were decontaminated with HNO_3 (10% v/v) and washed 3 times with ultrapure water. In the preparation of the solutions, all reagents used were analytical grade: HNO_3 ($\geq 69\%$ w/w, TraceSELECT[®] Ultra Fluka, L'Isle d'Abeau Chesnes, France) and H_2O_2 (30% v/v TraceSELECT[®] Fluka, Seelze, Germany). Ultrapure water (18.2 $\text{M}\Omega$ cm, 25°C) was obtained from a Sartorius Arium[®] pro water purification system (Goettingen, Germany). A certified reference material (BCR 679, white cabbage, provided by the EC Institute for Reference Materials and Measurements, Geel, Belgium) was used to verify the precision and accuracy of the analytical procedures. Recovery, calculated as the determined concentration for a given element divided by its certified value, ranged between 97.2 and 103.6% for all the analyzed elements (Online Resource 1).

Statistical analysis

All results were subjected to one-way analysis of variance (ANOVA) (GraphPad Instat Software, La Jolla, USA). Differences in mean values between groups were assessed by the Tukey's test and were considered statistically different at $p < 0.05$.

Results and discussion

Effects of CYN on photosynthetic capacity

Regarding photosynthetic capacity, we observed an increase in the Fv/Fm of the plants irrigated with 10 and 50 μg CYN/L with respect to the control group (Fig. 1). An Fv/Fm between 800 and 830 is considered an optimal value for this parameter and values lower than this may indicate a decrease in the photosynthetic capacity of plants caused by some kind of stress (Maxwell and Johnson 2000). In this work, contrary to what was expected, an increase in Fv/Fm was observed and in the case of 50 μg CYN/L treatment was statistically significant. However, increased Fv/Fm is rarely observed and is poorly characterized in the existing literature at the physiological and plant performance levels (Machado 2014; Madhana Sekhar et al. 2014; Li et al. 2014). This may reflect a hormesis effect, showing an active metabolism in an early stage for low doses of toxicants (such as 10 and 50 μg CYN/L), but it may be possible that for higher CYN concentrations, an inhibition of photosynthesis occurs, as reported in other plant species exposed to different toxic agents (Dai et al. 2010; Capelo et al. 2012; Erofeeva 2014; Jia et al. 2015).

With regard to the relationship between Fv/Fm and photosynthesis, Madhana Sekhar et al. (2014) demonstrated that elevated CO_2 grown mulberry plants showed a persistent stimulation of photosynthesis, avoiding photosynthetic




Fig. 1 Fluorescence measures in leaves from carrots (*Daucus carota*) grown for 30 days with different concentrations of a *Chrysosporium ovalisporum* extract containing CYN (10 or 50 μg CYN/L). Values are expressed as mean \pm SD ($n = 20$). The significance levels observed are ** $p < 0.01$ when comparing the control and CYN-exposed plants (10 or 50 μg CYN/L)

acclimation, based on increased Fv/Fm and other parameters. The authors suggested this is perhaps due to efficient sink capacity, which implies a positive regulation of the photosynthesis by an efficient utilization and/or storage of the photosynthetic assimilates (produced in the source organs, leaves) in the sink organs (roots, flowers, seeds, fruits, etc.). By contrast, ineffective sink capacity would lead to an accumulation of photosynthetic assimilates in the leaves and subsequent inhibition of photosynthesis and production of photosynthetic assimilates. Moreover, photosynthetic activity (Fv/Fm) was improved in *Sedum alfredii* by elevated CO_2 , facilitating the plant to keep normal physiological functions under cadmium stress and therefore alleviating metal toxicity to the plant (Li et al. 2014). These authors also demonstrated a positive correlation between plant growth promotion and increased photosynthetic rate at elevated CO_2 , in agreement with our results. According to these findings, it may be possible that carrot plants are reinforcing their photosynthetic mechanism in order to confront the higher energetic cost needed to fight the toxic effects of CYN.

To date, no studies concerning the effects of CYN on photosynthesis on edible plants have been reported, so there are no data available for comparison. Nevertheless, Kinnear et al. (2008) conducted a study in the aquatic plant *Hydrilla verticillata* exposed to whole-cell extracts of *C. raciborskii* containing 25–400 μg CYN/L, remarking that these might exert some adverse effects on photosynthesis.

In relation to other cyanotoxins, such as MC, contradictory results have been obtained depending on the plant species, concentration and exposure length to the toxin. Similar to our study, the findings by Machado (2014) in carrots exposed to MC-LR for 14–28 days show that this parameter increased from the concentration of 10 μg MC-LR/L and with 50 μg MC-LR/L. However, other studies showed inhibition of the photosynthetic process in wheat, maize, peas, lentils, tomatoes, and rice exposed to MC (Saqrane et al. 2009; El Khalloufi et al. 2012; Gutiérrez-Praena et al. 2014; Azevedo et al. 2014). These studies proved that MC impaired the photosynthetic activity, partially due to a damage of PSII reaction centers, deleterious effects on PSII activity and changes in the function of various proteins involved in the photosynthetic process related to the chloroplast ATP synthesis, carbon fixation, photosynthesis, and metabolism of carbohydrates. However, most of the above-mentioned research was carried out with MC, a cyanotoxin which is chemically and functionally very distinct from CYN. Thus, the effects observed in other works may be not necessarily reproduced in our study with CYN, as can be seen in the present work where the toxin may even have contrasting effects. Taking into account the results obtained in our study, more research will be needed to understand if the high Fv/Fm value observed

in plants exposed to 50 µg CYN/L can be translated into an increased photosynthetic efficiency, or if, conversely, the Fv/Fm value has no physiological significance with CYN.

Effects of CYN on carrot growth

Exposure of carrot plants to 10 and 50 µg CYN/L for 30 days led to a significant increase in the fresh weight of roots (Fig. 2). This was similar to the increase in root growth of *Hydrilla verticillata* observed by Kinnear et al. (2008) after 14 days of exposure to *C. raciborskii* extracts containing 400 µg CYN/L. Moreover, Prieto et al. (2011) found significant increases in the fresh weight of roots from rice plants exposed for 48 h to *C. ovalisporum* cell extracts containing 2.5 µg CYN/L, however, no changes were observed upon a 9-day exposure to the same extracts. The growth of lettuce roots was stimulated after 10 days of exposure to 1, 10, and 100 µg CYN/L, whereas the leaf fresh weight decreased only at 100 µg CYN/L (Freitas et al. 2015). Similarly, Garda et al. (2015) detected transient stimulation of root growth in *V. faba* seedlings at 6 days of treatment with 0.1 µg CYN/mL. According to these authors, growth responses of vascular plants to CYN are variable, showing growth stimulation at low concentrations (0.1–0.4 µg/mL). Kinnear et al. (2008) reported that some benefits could result from an increased root production, such as the potential production of exudates, allelochemicals or phytochelatins, which might scavenge CYN, preventing its uptake by plant cells. By contrast, Vasas et al. (2002) reported the growth inhibition of etiolated *Sinapis alba* seedlings exposed to CYN isolated from *C. ovalisporum*. Moreover, developmental alterations (reduction of fresh mass and length of cotyledons) have been observed in seedlings of *S. alba* by exposure to this toxin (M-Hamvas et al. 2010). The reduction in growth reported by those authors may be attributed to the high toxin concentrations employed (18,200 and 5000–10,000 µg CYN/L, respectively). Beyer et al. (2009)




Fig. 2 Average fresh weight of *Daucus carota* taproots (carrots) grown for 30 days with different concentrations of a *Chrysochloris ovalisporum* extract containing CYN (10 or 50 µg CYN/L). Values are expressed as mean ± SD ($n = 37$). The significance levels observed are * $p < 0.05$ when comparing the control and CYN-exposed plants (10 or 50 µg CYN/L)

showed that concentrations of CYN between 500 and 40,000 µg/L decreases root and shoot elongation in the macrophyte *Phragmites australis*. Regarding carrot plants exposed to MC, Machado (2014) observed a decrease in root growth of carrots exposed to MC-LR (50 µg/L) for 28 days, corroborating the results reported by El Khalloufi et al. (2012) in tomato plants (*Lycopersicon esculentum*). Several factors may explain the differences between those studies and our results, such as the plant species and their stage of development, as well as the time and dose of exposure.

Hence, our results seem to indicate that chronic exposure of carrots to low concentrations of CYN does not impair photosynthesis and plant growth. Among the causes of this tolerance could be the expression of several molecular and specialized tissues that protect adult plants against pathogens and soil-borne toxins. Several of these defense mechanisms are present in roots and are constituted by the root exudates (allelochemicals or phytochelatins) and other secondary metabolites used in defense against pathogenic microorganisms and toxins (Baetz and Martinoia 2014). Other defense mechanisms include the root border cells that engineer the chemical and physical properties of the external environment (Hawes et al. 2000), and the suberized tissues of the epidermis and the endodermis with a casparian band, that function as barriers to the entry of external chemicals and pathogen invasion (Kosma et al. 2015; Enstone et al. 2002). On the other hand, the stimulation of root growth is in line with the stimulation of photosynthesis and the probable increase of photosynthetic assimilates that are translocated and accumulated in storage organs as the carrots. The potential beneficial effects of the cyanobacterial extract nutrients seem unlikely, since these were highly diluted in the irrigation water used in the experiment (dilution rates of around 2700 and 550 to obtain 10 and 50 µg CYN/L, respectively).

The fact that we did not observe serious damages induced by CYN on photosynthetic capacity and growth of carrot plants even after 30 days of exposure, does not exclude the potential hazard of this toxin in other scenarios of prolonged exposure. If we consider that the accumulation of toxin may increase with the length of exposure (Kittler et al. 2012), there is the possibility that toxic CYN concentrations in carrots may be reached after 30 days of exposure, and with the transposition of the defense barriers of the roots. Therefore in terms of risk assessment there is the need to clarify the exposure length effects of CYN in crops (effects on yield, contamination, physiology) and monitor the effects of CYN in the whole life cycle of crops.

Effects of CYN on mineral content in carrot roots

A significant decline in the content of almost all minerals (Ca, Mg, Na, K, Fe, Zn, Mo, and P) in carrot roots exposed

to CYN, especially with 50 µg CYN/L was observed (Figs. 3 and 4). Only Mn and Cu were significantly increased after exposure to CYN, although for Cu this effect was only observed with 10 µg CYN/L (Fig. 4a, b). These results demonstrate that the exposure concentration is an important factor for the toxic effects of CYN on carrots. Contrary to our results, Machado (2014) and Freitas et al. (2015) obtained a general concentration-dependent enhancement of the mineral content in carrot roots exposed to MC and lettuce leaves exposed to CYN, respectively. However, Cu and P followed the same trend (increasing and decreasing, respectively) in the present and the above-mentioned studies, whatever the cyanotoxin was (CYN or MC).

In terms of mineral concentrations, it seems that carrots are more vulnerable to CYN toxicity compared to other plant species, at the conditions assayed. In our study, a general decrease was observed in fully developed carrots (after 30 days of exposure), compared to Freitas et al. (2015), who observed a general increase in mineral content in lettuce leaves after 5 and 10 days of exposure to CYN, indicating the potential tolerance of lettuce plants to this cyanotoxin. One of the reasons for the increased susceptibility of carrots to CYN, in terms of mineral uptake, was the longer time of exposure (30 days) to the toxin, which showed to be critical to mineral uptake, as contrasting to the 5 and 10 days CYN exposure studied in Freitas et al. (2015). In this sense, it seems that the above-mentioned root

defense mechanisms do not protect completely root mineral uptake from the external stressor CYN. Another reason could be that root metabolism is more affected than leaf metabolism, since roots are directly in contact with the toxin, and this is also reflected in terms of mineral content differences in the two organs. Noticeably, CYN had a more severe effect on our carrot roots than cyanotoxin MC-LR, as described by Machado (2014).

The uptake of minerals in the plant, especially micro-nutrients, can be affected by various factors such as the substrate where plants grow, microbial composition of the substrate, environmental factors and the presence of contaminants (Pinto et al. 2014b). In this regard, the differences observed in the mineral uptake in this work vs. Freitas et al. (2015) could also be due to differences in physiology between the hydroponically grown plants of Freitas et al. (2015) compared with our soil-grown plants. Keeping in mind that the present work was carried out by irrigating the plants with water contaminated with *C. ovalisporum* crude extracts, we point out the possible combined action of the toxin CYN and other secondary cyanobacterial metabolites produced by *C. ovalisporum* to explain the differential uptake of minerals observed in this work, compared to Freitas et al. (2015), where purified CYN was used instead. In this sense, cyanobacteria are known to produce a diverse array of secondary metabolites, including toxins,




Fig. 3 Concentration of Ca, Mg, Na, and P **a** and K **b**, (mg/kg F.W.) in *Daucus carota* taproots (carrots) grown for 30 days with different concentrations of a *Chrysochloris ovalisporum* extract containing CYN (10 or 50 µg CYN/L). Values are expressed as mean ± SD ($n = 4$). The significance levels observed are * $p < 0.05$ or ** $p < 0.01$ when comparing the control and CYN-exposed plants (10 or 50 µg CYN/L), and # $p < 0.05$ among both doses of CYN




Fig. 4 Concentration of Fe, Zn, Mn **a**, Mo and Cu **b** (mg/kg F.W.) in *Daucus carota* taproots (carrots) grown for 30 days with different concentrations of a *Chrysochloris ovalisporum* extract containing CYN (10 or 50 µg CYN/L). Values are expressed as mean ± SD ($n = 4$). The significance levels observed are * $p < 0.05$, ** $p < 0.01$ or *** $p < 0.001$ when comparing the control and CYN-exposed plants (10 or 50 µg CYN/L), and # $p < 0.05$ among both doses of CYN

lipopolysaccharides, and other unknown metabolites that, as a whole, are known to contribute to the bioactivity and toxicity of corresponding crude extracts (Berry et al. 2009; Kinnear 2010).

To ensure the proper growth of plants it is essential that nutrients are assimilated at adequate levels, as both a deficiency and an excess of nutrients could affect plant growth by direct effects on cell metabolism (such as the biosynthesis of macromolecules or functioning of the photosynthetic system) (White and Brown 2010). In our study, the general decrease observed in nutrient concentrations does not seem sufficient to impair carrot growth. In this sense, macro and micronutrients contents evaluated in this work are generally within the ranges reported by other authors for several varieties of carrot taproots. The amount of Ca we detected was in the range of 233–298 mg/kg F.W. (for 50 µg/L and control groups, respectively), similarly to Zaccari et al. (2015) who obtained values ranging 290–370 mg/kg F.W. Warman and Havard (1997) reported 1000–1570 mg of Mg/kg F.W. and 1170–4980 mg of Na/kg F.W., compared to our values of 1145–1468 and 1142–2289 mg/kg F.W. for Mg and Na, respectively. The content of Zn in the carrots studied ranges from 3.62 to 5.53 mg/kg F.W. (for 50 µg/L and control groups, respectively), and agrees with the reported by other authors which show different ranges in the literature: 3.1–5.6 mg/kg F.W. (Scherz and Kirchoff 2006), 1.8–3.9 mg/kg F.W. (Nicolle et al. 2004), 7–10 mg/kg F.W. (Zaccari et al. 2015) and 21–26 mg/kg F.W. (Warman and Havard 1997). In the case of Mo, Ensafi and Khaloo (2005) found 1.48 ng/mL, similar to our values of 1.53–1.80 ng Mo/mL. Among all micronutrients, Mo is often present in plants in lower concentrations (Kirkby and Römheld 2007), agreeing with our results. Fe, Mn, and Cu concentrations were lower than those reported by Warman and Havard (1997), even in the control group. However, this is not surprising or alarming because crop plants only need a small amount of micronutrients. These works show that carrots in our work, even after CYN exposure, were generally within the normal amounts of nutrients detected in this species, explaining that growth and photosynthesis were not compromised by this condition.

It would be of interest to further study the role of these minerals in the plant metabolism. Manganese is part of the structure of some photosynthetic proteins, and Zn is essential for the synthesis of chlorophyll and for metalloenzymes. Copper toxicity could affect plant physiological processes and ultimately plant production (Pinto et al. 2014b), being important as cofactor of antioxidant enzymes (Pinto and Ferreira 2015; Kirkby and Römheld 2007).

The absence of mineral content data in leaves makes the link between the nutrient effect and photosynthesis difficult to establish and rather speculative. Taking into account that most minerals decreased in roots, we could expect a similar

effect of CYN in leaves. Since some of these minerals are critical to photosynthesis, the most probable result of such a change in minerals would be the inhibition of photosynthesis. However, this hypothetical situation does not agree with the experimental data on photosynthesis, which leads us to the assumption that, for this particular experiment, mineral, and photosynthesis effects are poorly associated.

Conclusions

In the present study, the effects of ecologically relevant concentrations of CYN (10 and 50 µg/L) were investigated regarding to growth, photosynthetic activity, and mineral content in carrot plants exposed for 30 days to *C. ovalisporum* extracts contained in the irrigation water. Based on the results, we conclude that exposure to CYN did not have a clear negative effect on the growth of *D. carota*, and it even caused an increase in the fresh weights of the exposed plants. Furthermore, exposure of carrots to these CYN concentrations did not appear to affect vital biochemical processes such as photosynthesis. Nevertheless, the decrease in mineral content indicate possible alterations in the nutritional value of carrots after exposure to crude extracts of *C. ovalisporum* containing CYN. This study is the first to evaluate the effects *C. ovalisporum* extracts on a root vegetable. However, more studies are necessary to understand the effect of this toxin in other environmental scenarios.

Funding This work was partially funded by Fundação para a Ciência e Tecnologia (FCT), under the framework of the project UID/Multi/04423/2013, by Ministerio de Ciencia e Innovación of Spain under the project AGL2009-10026, and by Junta de Andalucía under the project P09-AGR-4672. Remedios Guzmán's work is supported by the Spanish Ministerio de Educación fellowship. Alexandre Campos work is supported by a post-doctoral grant (SFRH/BPD/103683/2014) from FCT.

Compliance with ethical standards

Conflict of interest The authors declare that they have no competing interests.

References

- Azevedo CC, Azevedo J, Osório H, Vasconcelos V, Campos A (2014) Early physiological and biochemical responses of rice seedlings to low concentration of microcystin-LR. *Ecotoxicology* 23:107–121
- Baetz U, Martinoia E (2014) Root exudates: The hidden part of plant defense. *Trends Plant Sci* 19:90–98
- Banker R, Carmeli S, Hadas O, Teltsh B, Porat R, Sukenik A (1997) Identification of cylindrospermopsin in *Aphanizomenon ovalisporum* (Cyanophyceae) isolated from lake Kinneret, Israel. *J Phycol* 33:613–616

- Bazin E, Mourot A, Humpage AR, Fessard V (2010) Genotoxicity of a freshwater cyanotoxin, cylindrospermopsin, in two human cell lines: Caco-2 and HepaRG. *Environ Mol Mutagen* 51:251–259
- Berry JP, Gibbs PD, Schmale MC, Saker ML (2009) Toxicity of cylindrospermopsin, and other apparent metabolites from *Cylindrospermopsis raciborskii* and *Aphanizomenon ovalisporum*, to the zebrafish (*Danio rerio*) embryo. *Toxicol* 53:289–299
- Beyer D, Surányi G, Vasas G, Roszik J, Erdodi F, M-Hamvas M, Bácsi I, Batori R, Serfozo Z, Szigeti ZM, Vereb G, Demeter Z, Gonda S, Máthé C (2009) Cylindrospermopsin induces alterations of root histology and microtubule organization in common reed (*Phragmites australis*) plantlets cultured *in vitro*. *Toxicol* 54:440–449
- Boglialli S, Bruno M, Curini R, Di Corcia A, Fanali C, Lagana A (2006) Monitoring algal toxins in lake water by liquid chromatography tandem mass spectrometry. *Environ Sci Technol* 40:2917–2923
- Burns J (2008) Toxic cyanobacteria in Florida waters. *Adv Exp Med Biol* 619:127–137
- Capelo A, Santos C, Loureiro S, Pedrosa MA (2012) Phytotoxicity of lead on *Lactuca sativa*: effects on growth, mineral nutrition, photosynthetic activity and oxidant metabolism. *Fresen Environ Bull* 21:450–459
- Corbel S, Mougín C, Bouaicha N (2014) Cyanobacterial toxins: modes of actions, fate in aquatic and soil ecosystems, phytotoxicity and bioaccumulation in agricultural crops. *Chemosphere* 96:1–5
- Dai HP, Wei Y, Yang TX, Sa WQ, Wei AZ (2010) Influence of cadmium stress on chlorophyll fluorescence characteristics in *Populus 9 canescens*. *J Food Agric Environ* 10:1281–1283
- De la Cruz AA, Hiskia A, Kaloudis T, Chernoff N, Hill D, Antoniou MG, He X, Loftin K, O’Shea K, Zhao C, Pelaez M, Han C, Lynch TJ, Dionysiou DD (2013) A review on cylindrospermopsin: the global occurrence, detection, toxicity and degradation of a potent cyanotoxin. *Environ Sci Process Impacts* 15:1979–2003
- El Khalloufi F, Oufdou K, Lahrouni M, El Ghazali I, Saqrane S, Vasconcelos V, Oudra B (2011) Allelopathic effects of cyanobacteria extracts containing microcystins on *Medicago sativa*-*Rhizobia* symbiosis. *Ecotoxicol Environ Saf* 74:431–438
- El Khalloufi F, El Ghazali I, Saqrane S, Oufdou K, Vasconcelos V, Oudra B (2012) Phytotoxic effects of a natural bloom extract containing microcystins on *Lycopersicon esculentum*. *Ecotoxicol Environ Saf* 79:199–205
- Elliott JA (2012) Is the future blue-green? A review of the current model predictions of how climate change could affect pelagic freshwater cyanobacteria. *Water Res* 46:1364–1371
- Ensafi AA, Khaloo SS (2005) Determination of traces molybdenum by catalytic adsorptive stripping voltammetry. *Talanta* 65:781–788
- Enstone DE, Peterson CA, Ma F (2002) Root endodermis and exodermis: structure, function, and responses to the environment. *J Plant Growth Regul* 21:335–351
- Erofeeva EA (2014) Hormesis and paradoxical effects of wheat seedling (*Triticum Aestivum* L.) parameters upon exposure to different pollutants in a wide range of doses. *Dose Response* 12:121–135
- Freitas M, Azevedo J, Pinto E, Neves J, Campos A, Vasconcelos V (2015) Effects of microcystin-LR, cylindrospermopsin and a microcystin-LR/cylindrospermopsin mixture on growth, oxidative stress and mineral content in lettuce plants (*Lactuca sativa* L.). *Ecotoxicol Environ Saf* 116:59–67
- Froschio SM, Humpage AR, Wickramasinghe W, Shaw G, Falconer IR (2008) Interaction of the cyanobacterial toxin cylindrospermopsin with the eukaryotic protein synthesis system. *Toxicol* 5:191–198
- Garda T, Riba M, Vasas G, Beyer D, M-Hamvas M, Hajdu G, Tándor I, Máthé C (2015) Cytotoxic effects of cylindrospermopsin in mitotic and non-mitotic *Vicia faba* cells. *Chemosphere* 120:145–153
- Gutiérrez-Praena D, Jos A, Pichardo S, Moreno IM, Cameán AM (2013) Presence and bioaccumulation of microcystins and cylindrospermopsin in food and the effectiveness of some cooking techniques at decreasing their concentrations: a review. *Food Chem Toxicol* 53:136–152
- Gutiérrez-Praena D, Campos A, Azevedo J, Neves J, Freitas M, Guzmán-Guillén R, Vasconcelos V (2014) Exposure of *Lycopersicon Esculentum* to microcystin-LR: Effects in the leaf proteome and toxin translocation from water to leaves and fruits. *Toxins* 6:1837–1854
- Guzmán-Guillén R, Prieto AI, González AG, Soria-Díaz ME, Cameán AM (2012) Cylindrospermopsin determination in water by LC–MS/MS: Optimization and validation of the method and application to real samples. *Environ Toxicol Chem* 31:2233–2238
- Guzmán-Guillén R, Prieto AI, Moreno I, Rios V, Vasconcelos VM, Cameán AM (2014) Effects of depuration on oxidative biomarkers in *Tilapia (Oreochromis niloticus)* after subchronic exposure to cyanobacterium producing cylindrospermopsin. *Aquat Toxicol* 149:40–49
- Havens KE (2008) Cyanobacteria blooms: effects on aquatic ecosystems. *Adv Exp Med Biol* 619:733–747
- Hawes MC, Gunawardena U, Miyasaka S, Zhao X (2000) The role of root border cells in plant defense. *Trends Plant Sci* 5:128–133
- Hoeger SJ, Shaw G, Hitzfeld BC, Dietrich DR (2004) Occurrence and elimination of cyanobacterial toxins in two Australian drinking water treatment plants. *Toxicol* 43:639–649
- Humpage A (2008) Toxin types, toxicokinetics and toxicodynamics. In: Hudnel HK (ed) *Cyanobacterial harmful algal blooms state of the science and research needs*. Springer Press, New York, pp 383–415
- Humpage AR, Falconer IR (2003) Oral toxicity of the cyanobacterial toxin cylindrospermopsin in male Swiss albino mice: determination of no observed adverse effect level for deriving a drinking water guideline value. *Environ Toxicol* 18:94–103
- Ibelings BW, Chorus I (2007) Accumulation of cyanobacterial toxins in freshwater “seafood” and its consequences for public health: a review. *Environ Pollut* 150:177–192
- Jia L, Liu Z, Chen W, Ye Y, Yu S, He X (2015) Hormesis effects induced by cadmium on growth and photosynthetic performance in a hyperaccumulator *Lonicera japonica* Thunb. *J Plant Growth Regul* 34:13–21
- Kinnear S, Fabbro L, Duivenvoorden L (2008) Variable growth responses of water thyme (*Hydrilla verticillata*); To whole-cell extracts of *Cylindrospermopsis raciborskii*. *Arch Environ Contam Toxicol* 54:187–194
- Kinnear S (2010) Cylindrospermopsin: a decade of progress on bioaccumulation research. *Mar Drugs* 8:542–564
- Kirkby EA, Römheld V (2007) Micronutrients in Plant Physiology: Functions, Uptake and Mobility. *Proceedings* 543. The International Fertilizer Society, York, United Kingdom.
- Kittler K, Schreiner M, Krumbein A, Manzei S, Koch M, Rohn S, Maul R (2012) Uptake of the cyanobacterial toxin cylindrospermopsin in *Brassica* vegetables. *Food Chem* 133:875–879
- Kosma DK, Rice A, Pollard M (2015) Analysis of aliphatic waxes associated with root periderm or exodermis from eleven plant species. *Phytochemistry* 117:351–362
- Lahrouni M, Oufdou K, El Khalloufi F, Baz M, Lafuente A, Dary M, Pajuelo E, Oudra B (2013) Physiological and biochemical defense reactions of *Vicia faba* L.-*Rhizobium* symbiosis face to chronic exposure to cyanobacterial bloom extract containing microcystins. *Environ Sci Pollut Res* 20:5405–5415
- Li TQ, Tao Q, Di ZZ, Lu F, Yang XE (2014) Effect of elevated CO₂ concentration on photosynthetic characteristics of

- hyperaccumulator *Sedum alfredii* under cadmium stress. *J Integr Plant Biol* 57:653–660
- M-Hamvas M, Máthé C, Vasas G, Jámbrik K, Papp M, Beyer D, Mészáros I, Borbély G (2010) Cylindrospermopsin and microcystin-LR alter the growth, development and peroxidase enzyme activity of white mustard (*Sinapis alba* L.) seedlings, a comparative analysis. *Acta Biol Hung* 61:35–48
- Machado JFF (2014) Implicações do Uso de Água Contaminada com Microcistina na Qualidade de *Daucus Carota* (Cenoura). Dissertação de Mestrado. Universidade do Porto, Portugal
- Madhana Sekhar K, Rachapudi VS, Mudalkar S, Reddy AR (2014) Persistent stimulation of photosynthesis in short rotation coppice mulberry under elevated CO₂ atmosphere. *J Photochem Photobiol B*: 137:21–30
- Maxwell K, Johnson N (2000) Chlorophyll fluorescence – A practical guide. *J Exp Bot* 51:659–668
- Metcalf JS, Barakate A, Codd GA (2004) Inhibition of plant protein synthesis by the cyanobacterial hepatotoxin, cylindrospermopsin. *FEMS Microbiol Lett* 235:125–129
- Mitrovic SM, Allis O, Furey A, James KJ (2005) Bioaccumulation and harmful effects of microcystin-LR in the aquatic plants *Lemna minor* and *Wolffia arrhiza* and the filamentous alga *Chlorella vulgaris*. *Ecotoxicol Environ Saf* 61:345–352
- Moreira C, Azevedo J, Antunes A, Vasconcelos V (2012) Cylindrospermopsin: occurrence, methods of detection and toxicology. *J Appl Microbiol* 114:605–620
- Nicolle C, Simon G, Rock E, Amouroux P, Révész C (2004) Genetic variability influences carotenoid, vitamin, phenolic, and mineral content in white, yellow, purple, orange and dark orange carrot cultivars. *J Am Soc Hortic Sci* 129:523–529
- Pearson L, Mihali T, Moffitt M, Kellmann R, Neilan B (2010) On the chemistry, toxicology and genetics of the cyanobacterial toxins, microcystin, nodularin, saxitoxin and cylindrospermopsin. *Mar Drugs* 8:1650–1680
- Pflugmacher S, Jung K, Lundvall L, Neumann S, Peuthert A (2006) Effects of cyanobacterial toxins and cyanobacterial cell-free crude extract on germination of alfalfa (*Medicago sativa*) and induction of oxidative stress. *Environ Toxicol Chem* 25:2381–2387
- Pinto E, Almeida AA, Aguiar AA, Ferreira IM (2014a) Changes in macrominerals, trace elements and pigments content during lettuce (*Lactuca sativa* L.) growth: influence of soil composition. *Food Chem* 152:603–611
- Pinto E, Aguiar AA, Ferreira IM (2014b) Influence of soil chemistry and plant physiology in the phytoremediation of Cu, Mn, and Zn. *Crit Rev Plant Sci* 33:351–373
- Pinto E, Ferreira IMPLVO (2015) Cation transporters/channels in plants: tools for nutrient biofortification. *J Plant Physiol* 179:64–82
- Prieto A, Campos A, Cameán A, Vasconcelos V (2011) Effects on growth and oxidative stress status of rice plants (*Oryza sativa*) exposed to two extracts of toxin-producing cyanobacteria (*Aphanizomenon ovalisporum* and *Microcystis aeruginosa*). *Ecotoxicol Environ Saf* 74:1973–1980
- Quesada A, Moreno E, Carrasco D, Paniagua T, Wörmer L, De Hoyos C, Sukenik A (2006) Toxicity of *Aphanizomenon ovalisporum* (Cyanobacteria) in a Spanish water reservoir. *Eur J Phycol* 41:39–45
- Quiblier C, Susanna W, Isidora ES, Mark H, Aurélie V, Jean-François H (2013) A review of current knowledge on toxic benthic freshwater cyanobacteria - Ecology, toxin production and risk management. *Water Res* 47:5464–5479
- Rücker J, Stüken A, Nixdorf B, Fastner J, Chorus I, Wiedne C (2007) Concentrations of particulate and dissolved cylindrospermopsin in 21 *Aphanizomenon*-dominated temperate lakes. *Toxicol* 50:800–809
- Saker ML, Eaglesham GK (1999) The accumulation of cylindrospermopsin from the cyanobacterium *Cylindrospermopsis raciborskii* in tissues of the Redclaw crayfish *Cherax quadricarinatus*. *Toxicol* 37:1065–1077
- Saker ML, Metcalf JS, Codd GA, Vasconcelos VM (2004) Accumulation and depuration of the cyanobacterial toxin cylindrospermopsin in the freshwater mussel *Anodonta cygnea*. *Toxicol* 43:185–194
- Saqrane S, Ouahid Y, El Ghazali I, Oudra B, Bouarab L, del Campo FF (2009) Physiological changes in *Triticum durum*, *Zea mays*, *Pisum sativum* and *Lens esculenta* cultivars, caused by irrigation with water contaminated with microcystins: a laboratory experimental approach. *Toxicol* 53:786–796
- Scherz H, Kirchhoff E (2006) Trace elements in foods: zinc contents of raw foods—A comparison of data originating from different geographical regions of the world. *Critical Review. J Food Compos Anal* 19:420–433
- Shaw GR, Seawright AA, Moore MR, Lam PKS (2009) Cylindrospermopsin, a cyanobacterial alkaloid: evaluation of its toxicologic activity. *Ther Drug Monit* 22:89–92
- Van Apeldoorn ME, Van Egmond HP, Speijers GJA, Bakker GJJ (2007) Toxins of cyanobacteria. *Mol Nutr Food Res* 51:7–60
- Vasas G, Gáspár A, Surányi G, Batta G, Gyémánt G, M-Hamvas M, Máthé C, Grigorszky I, Molnár E, Borbély G (2002) Capillary electrophoretic assay and purification of cylindrospermopsin, a cyanobacterial toxin from *Aphanizomenon ovalisporum*, by plant test (blue-green sinapis test). *Anal Biochem* 302:95–103
- Warman PR, Havard KA (1997) Yield, vitamin and mineral contents of organically and conventionally grown carrots and cabbage. *Agr Ecosyst Environ* 61:155–162
- White PJ, Brown PH (2010) Plant nutrition for sustainable development and global health. *Ann Bot* 105:1073–1080
- Zaccari F, Cabrera MC, Ramos A, Saadoun A (2015) *In vitro* bioaccessibility of β -carotene, Ca, Mg and Zn in landrace carrots (*Daucus carota*, L.). *Food Chem* 166:365–371
- Žegura B, Gajski G, Štraser A, Garaj-Vrhovac V (2011) Cylindrospermopsin induced DNA damage and alteration in the expression of genes involved in the response to DNA damage, apoptosis and oxidative stress. *Toxicol* 58:471–479