



SHARP ESTIMATES FOR BIORTHOGONAL FAMILIES TO EXPONENTIAL FUNCTIONS ASSOCIATED TO COMPLEX SEQUENCES WITHOUT GAP CONDITIONS

MANUEL GONZALEZ-BURGOS✉*¹ AND LYDIA OUAILI✉²

¹Dpto. Ecuaciones Diferenciales y Análisis Numérico
and Instituto de Matemáticas de la Universidad de Sevilla (IMUS)
Facultad de Matemáticas, Universidad de Sevilla, C/ Tarfia S/N
41012 Sevilla, SPAIN

²Aix Marseille Université, CNRS, Centrale Marseille, I2M
UMR 7373, 39 rue F. Joliot Curie
13453 Marseille, FRANCE

(Communicated by Handling Editor)

ABSTRACT. The general goal of this work is to obtain upper and lower bounds for the L^2 -norm of biorthogonal families to complex exponential functions associated to sequences $\{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$ which satisfy appropriate assumptions but without imposing a gap condition on the elements of the sequence. As a consequence, we also present new results on the cost of the boundary null controllability of two parabolic systems at time $T > 0$: a phase-field system and a parabolic system whose generator has eigenvalues that accumulate. In the latter case, the behavior of the control cost when T goes to zero depends strongly on the accumulation parameter of the eigenvalue sequence.

1. Introduction and main results. In the last years, an increasing number of authors have addressed the problem of the null controllability of coupled parabolic systems with less controls than equations (see [3], [19], [4], [31],...). One of the most important problems in this framework is obtaining necessary and sufficient conditions that allow the system to be controlled with a reduced number of distributed or boundary controls.

Another important problem is the study of the dependence of the so-called control cost with respect to the final observation time $T > 0$, when T is small enough and the corresponding null controllability result holds at any time $T > 0$. Regarding this latter problem, we highlight the works [17], [18], [40], [23], [24], [20], [33], [43], [7], [15], [30], [13], [14], [12], etc., where the authors study an estimate of the control cost $\mathcal{K}(T)$ (for the definition, see (9)) in the case of scalar parabolic problems (problems that, under general assumptions, are null controllable for any $T > 0$). Most of the previous works use the moment method to obtain an estimate of the control cost.

2020 Mathematics Subject Classification. 35K40, 35Q93, 93B05, 93B60, 93C20.

Key words and phrases. Non scalar parabolic systems, null controllability, moment problem, biorthogonal families, no gap conditions, control cost.

The first author is supported by grant PID2020-114976GB-I00, Ministry of Science and Innovation (Spain).

*Corresponding author: Manuel González-Burgos.

In order to solve both problems, a classical tool in Control Theory is the use of biorthogonal families to appropriate sequences of exponentials in $L^2(0, T; \mathbb{C})$ and, to be precise, sharp estimates on the L^2 -norm of the elements of the biorthogonal family. We will provide more details in what follows.

Given $\{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$, a complex sequence of pairwise distinct elements, we will use the following notation:

$$e_k(t) = e^{-\Lambda_k t}, \quad \forall t \in (0, T), \quad (1)$$

where $T > 0$ is fixed. With this notation, we define

Definition 1.1. Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$ be a complex sequence and $T > 0$. We say that the family of functions $\{q_k\}_{k \geq 1} \subset L^2(0, T; \mathbb{C})$ is a biorthogonal family to the sequence of complex exponentials $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$, if for every $k, n \in \mathbb{N}$, one has

$$\int_0^T e_k(t) \bar{q}_n(t) dt = \delta_{kn},$$

where the function e_k is given in (1).

1.1. State of the art. In what follows we will give a non-exhaustive state of the art on conditions of the sequence Λ that ensure the existence of biorthogonal families to the corresponding complex exponential sequence.

1.1.1. Existence of biorthogonal families and bounds without explicit dependence on T . The existence of biorthogonal families $\{q_k\}_{k \geq 1}$ to sequences of exponentials $\{e_k\}_{k \geq 1}$ (e_k is the function given in (1)) strongly depends on the properties of the sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$. When $\Lambda = \{\Lambda_k\}_{k \geq 1}$ is a positive real sequence, it is well known (see [39]) that the existence of a biorthogonal family to the exponentials $\{e_k\}_{k \geq 1}$ in $L^2(0, T)$ is equivalent to the condition

$$\sum_{k \geq 1} \frac{1}{\Lambda_k} < \infty.$$

Let us provide some general properties for real or complex sequences Λ appearing in the literature which imply the existence of sequences $\{q_k\}_{k \geq 1}$ biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$ ($T > 0$) satisfying appropriate estimates.

The first results on existence and estimates of families $\{q_k\}_{k \geq 1}$ biorthogonal to sequences of exponentials $\{e_k\}_{k \geq 1}$ was proved in [17], [18] and [23] (see also [27], [41], [33], [42], [28] and [29]) for increasing real sequences that satisfy

$$\Lambda_k \in (0, \infty), \quad \Lambda_k = A(k + \omega)^2 + o(k), \quad \forall k \geq 1, \quad (2)$$

with $A > 0$ and $\omega \in \mathbb{R}$.

The previous results has been extended to the complex case in [24], [4], [5] and [7]. In [5], the authors prove the existence of biorthogonal sequences $\{q_k\}_{k \geq 1}$ under general assumptions on the sequence Λ and prove appropriate estimates of $\|q_k\|_{L^2(0, T; \mathbb{C})}$. Assume that the sequence $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$ satisfies

$$\left\{ \begin{array}{l} \Lambda_i \neq \Lambda_k, \quad \forall i, k \in \mathbb{N} \text{ with } i \neq k, \\ \Re(\Lambda_k) \geq \delta |\Lambda_k| > 0, \quad \forall k \geq 1, \quad \text{and} \quad \sum_{k \geq 1} \frac{1}{|\Lambda_k|} < \infty, \end{array} \right. \quad (3)$$

for a positive constant δ . Then, the family of exponentials $\{e_k\}_{k \geq 1}$ is minimal¹ in $L^2(0, T; \mathbb{C})$ for any $T > 0$ and, therefore, there exists a biorthogonal family $\{\tilde{q}_k\}_{k \geq 1}$ to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$ (see for instance [39], [37], [4], Theorem 4.1 in [5],...). In addition, in [5], the authors prove that there exist two positive constants C_1 and C_2 (only depending on Λ and T) such that

$$C_1 \frac{|1 + \Lambda_k|^2}{|\Lambda_k|} W_k \leq \|\tilde{q}_k\|_{L^2(0, T; \mathbb{C})} \leq C_2 \frac{|1 + \Lambda_k|^2}{|\Lambda_k|} W_k, \quad (4)$$

where C_1 and C_2 are positive constants depending on T and W_k is the infinite Blaschke product given by

$$W_k = \frac{1}{2\Re(\lambda_k)} \prod_{\substack{n \geq 1 \\ n \neq k}} \left| \frac{\Lambda_n + \Lambda_k}{\Lambda_n - \Lambda_k} \right|.$$

Nevertheless, the authors do not provide an explicit dependence of the constants C_1 and C_2 in (4) with respect to the final time $T > 0$. This is due to the method used by the authors to prove (4): these inequalities are first obtained in $L^2(0, \infty; \mathbb{C})$ ($T = \infty$) and, then, proved in $L^2(0, T; \mathbb{C})$ ($T \in (0, \infty)$) after a contradiction argument (see [5] for the details).

Observe that, in general, a sequence Λ satisfying (3) does not fulfill the so-called gap condition:

$$\inf_{k, n \geq 1: k \neq n} |\Lambda_k - \Lambda_n| > 0, \quad (5)$$

and, therefore, the elements of Λ could condensate.

From inequality (4), among other properties, in [5], the authors prove a general result of null controllability for abstract parabolic problems that develop a minimal time $T_0 \in [0, \infty]$ of controllability: the system is null-controllable at any time $T > T_0$ and not null-controllable for $T < T_0$. This minimal time is related to the Bernstein's condensation index of the sequence of eigenvalues $\Lambda = \{\Lambda_k\}_{k \geq 1}$ of the generator of the semigroup (see [5] and [8] for further details).

1.1.2. *Existence of biorthogonal families and bounds with explicit dependence on T .* In [17] the authors provide an approach that allows to construct biorthogonal families $\{q_k\}_{k \geq 1}$ to the sequence $\{e_k\}_{k \geq 1}$ in $L^2(0, T)$ ($T > 0$) with explicit bounds of the L^2 -norm of q_k with respect to the final time T . To be precise, for increasing sequences $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{R}$ satisfying (2), with $A > 0$ and $\omega \in \mathbb{R}$, there exist $\mathcal{C}_0, \tau_0 \in (0, \infty)$ and a family $\{q_k\}_{k \geq 1}$ biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T)$ such that

$$\|q_k\|_{L^2(0, T)} \leq \mathcal{C}_0 e^{\mathcal{C}_0(\sqrt{\Lambda_k} + \frac{1}{T})}, \quad \forall T \in (0, \tau_0), \quad \forall k \geq 1, \quad (6)$$

(see for instance [17] and [33]).

Estimate (6) is known to be optimal with respect the factor $\exp(C/T)$ thanks to the work [23]: under assumption (2), there exists a positive constant \mathcal{C}_1 such that for any sequence $\{q_k\}_{k \geq 1} \subset L^2(0, T)$ biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T)$, one has

$$\|q_k\|_{L^2(0, T)} \geq \frac{M(k)}{\sqrt{T}} e^{\frac{\mathcal{C}_1}{T}}, \quad \forall T > 0, \quad \forall k \geq 1, \quad (7)$$

where $M(k)$ is a positive constant only depending on k and L .

¹A sequence $\{x_k\}_{k \geq 1}$ in a Hilbert space H is said to be minimal if it satisfies $x_n \notin \overline{\text{span}}\{x_k : k \neq n\}$ for any $n \geq 1$.

The existence of biorthogonal families to real exponentials that satisfy (6) and (7) has been also treated by some authors with assumptions on the sequence Λ different from (2) but always under assumptions on Λ that imply the gap condition (5) (see [13], [14] and [15]). We will analyse these assumptions in Subsection 2.3.

It is also interesting to remark that in [12] the authors also obtain an upper bound of the L^2 -norm of biorthogonal families to real exponentials with explicit time dependence when the corresponding real sequence does not satisfy the gap condition (5). In [2], the authors prove similar results when the sequence does not satisfy (5) but without proving an explicit dependence of the constants on T .

Inequality (6) has also been generalized to the case of complex sequences. Let us describe the result on existence and estimates of biorthogonal families to complex exponentials proved in [7]. One has:

Theorem 1.2 ([7]). *Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$ be a sequence satisfying assumptions (H1)–(H5), in Definition 1.4, the gap condition (5) and*

$$|p\sqrt{r} - \mathcal{N}(r)| \leq \alpha, \quad \forall r > 0,$$

(\mathcal{N} is the counting function associated with the sequence Λ , defined in (21)), for some parameters $\beta \in [0, \infty)$, $\rho, p, \alpha \in (0, \infty)$ and $q \in \mathbb{N}$. Then, there exists $T_0 > 0$ such that for every $T \in (0, T_0)$, there exists a sequence of \mathbb{C} -valued functions

$$\{q_k\}_{k \geq 1} \subset L^2(0, T; \mathbb{C})$$

biorthogonal to the exponentials $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$, e_k given in (1), which, in addition, satisfies (6) for a positive constant C_0 independent of k and T .

To the best of our knowledge, Theorem 1.2 is the most general result that provides existence of biorthogonal families $\{q_k\}_{k \geq 1}$ to complex exponentials with explicit estimates of $\|q_k\|_{L^2(0, T; \mathbb{C})}$ with respect to k and T . It is interesting to note that the sequences under the hypotheses of Theorem 1.2 fulfill assumptions in (3) (see [7]) and, of course, the gap condition (5).

1.1.3. *The cost of fast controls.* Biorthogonal families play a crucial role in the moment method. This method was developed by Fattorini and Russell (see [17] and [18]) to study the boundary null controllability of one-dimensional scalar parabolic problems with second order elliptic self adjoint generator. This method uses in a key way the existence and estimates of biorthogonal families to $\{e_k\}_{k \geq 1}$. As a consequence of inequality (6), in [17], the authors prove that the one-dimensional heat equation

$$\begin{cases} \partial_t y - \partial_{xx} y = 0 & \text{in } (0, T) \times (0, L), \\ y(\cdot, 0) = v, \quad y(\cdot, L) = 0 & \text{on } (0, T), \\ y(0, \cdot) = y_0 & \text{in } (0, L), \end{cases} \quad (8)$$

($L > 0$) is null controllable in $H^{-1}(0, L)$ at any time $T > 0$ with controls $v \in L^2(0, T)$. In fact, they prove the existence of a constant C_0 (only depending on L) such that for any $y_0 \in H^{-1}(0, L)$ there exists a control $v \in L^2(0, T)$ satisfying

$$\|v\|_{L^2(0, T)} \leq C_0 e^{\frac{C_0}{T}} \|y_0\|_{H^{-1}(0, L)},$$

and such that the solution to (8) satisfies $y(T, \cdot) = 0$ in $(0, L)$. Thus, the set

$$\mathcal{C}_T(y_0) := \{v \in L^2(0, T) : y(T, \cdot) = 0 \text{ in } (0, L), y \text{ solution of (8)}\},$$

is non empty and we can define the so-called *control cost* of system (8) at time T :

$$\mathcal{K}(T) := \sup_{\|y_0\|_{H^{-1}}=1} \inf_{v \in \mathcal{C}_T(y_0)} \|v\|_{L^2(0,T)}. \quad (9)$$

Therefore, for system (8), one has

$$\mathcal{K}(T) \leq C_0 e^{\frac{C_0}{T}}, \quad \forall T > 0, \quad (10)$$

for a positive constant C_0 only depending on L .

Again, estimate (10) is optimal with respect to the factor $\exp(C/T)$ thanks to the work [23]: inequality (7) implies the existence of new positive constants C_1 and τ_1 (only depending on L) such that the control cost for system (8) satisfies

$$\mathcal{K}(T) \geq C_1 e^{\frac{C_1}{T}}, \quad \forall T \in (0, \tau_1). \quad (11)$$

Observe that the assumptions on the sequence Λ in Theorem 1.2 are more general than condition (2). Therefore, Theorem 1.2 can be applied to a large range of scalar and coupled parabolic problems. In particular, it assures that the system under consideration is null controllable at any time $T > 0$ (thanks to the gap condition (5)). In addition, Theorem 1.2 provides the inequality (10) for the control cost $\mathcal{K}(T)$ as in the case of scalar parabolic problems (C_0 is a positive constant).

In the framework of N -dimensional scalar parabolic problems, [40] and [20] give an estimate of the cost $\mathcal{K}(T)$ similar to (10) using different approaches: In [40] the authors use the exact controllability of the wave equation to prove inequality (10) for the null-controllability of the heat equation. In [20], inequality (10) is deduced from appropriate global Carleman inequalities for general parabolic operators.

The work [30] is of special relevance because in it, the author studies the cost of the controllability of the one-dimensional heat equation with a pointwise control at point x_0 and, in this framework, there might exist a positive minimal time of null-controllability $T_0 \in [0, \infty]$ (which depends on x_0 and could take any arbitrary value in $[0, \infty]$, see [16]). In this work the eigenvalues satisfy (5) and the minimal time comes from the action of the control. In particular, the author proves that, if $T_0 > 0$, the cost of the controllability at time $T > T_0$ when T is close to T_0 , may explode in any arbitrary way.

1.2. Three examples of sequences of eigenvalues without gap condition.

In the framework of the controllability of non-scalar parabolic problems, new phenomena associated with the vectorial nature of the problem arise (hyperbolic phenomena): minimal time of null controllability and dependence of the controllability result on the position of the control domain (see [5], [6], [38], [34],...). This minimal time may come from the control action itself (as in [16] and [30]) or from the condensation index of the sequence of eigenvalues of the generator of the semigroup associated to the system (see [5]). In this latter case, the sequence Λ , in general, does not satisfy the gap condition (5). Let us provide more details in the case of systems with a minimal time which comes from the condensation index of the sequence.

We consider a boundary controllability problem for the generic 2×2 system

$$\begin{cases} \partial_t y + Ly = 0 & \text{in } (0, T) \times (0, \pi), \\ y(\cdot, 0) = Bv, \quad y(\cdot, \pi) = 0 & \text{on } (0, T), \\ y(0, \cdot) = y_0 & \text{in } (0, \pi), \end{cases} \quad (12)$$

where L is a second order elliptic operator, with domain $D(L) = H^2(0, \pi; \mathbb{R}^2) \cap H_0^1(0, \pi; \mathbb{R}^2)$, $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$ is the initial datum, $B \in \mathbb{R}^2$ is the control vector and $v \in L^2(0, T)$ is a scalar control.

The null controllability properties of the first example has been analyzed in [5]. We consider system (12) with $L = L_1 = -(D_1 \partial_{xx} + A_1)$, with domain $D(L_1) = H^2(0, \pi; \mathbb{R}^2) \cap H_0^1(0, \pi; \mathbb{R}^2)$, and

$$D_1 := \text{diag}(1, d), \quad d > 0, \quad d \neq 1, \quad \text{and} \quad A_1 := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix},$$

(see system (48)). Observe that the sequence of eigenvalues associated to the operator L_1 is $\Lambda^{(1)} = \{k^2\}_{k \geq 1} \cup \{dk^2\}_{k \geq 1}$. If $\sqrt{d} \notin \mathbb{Q}$ (and this condition is necessary for the approximate controllability at time $T > 0$ of the system (12) with the previous data, i.e., system (48)), the sequence $\Lambda^{(1)}$ can be rearranged as an increasing sequence $\Lambda^{(1)} = \{\lambda_k^{(1)}\}_{k \geq 1} \subset \mathbb{R}$ that fulfills property (3). It is clear that $\Lambda^{(1)}$ does not satisfy, in general, the gap condition (5). As a consequence, system (48) has a minimal time $T_0 = T_0(d) \in [0, \infty]$ which, for some d , with $\sqrt{d} \notin \mathbb{Q}$, is positive. Therefore, the system is not null controllable at time T when $T < T_0$ (see [5] for the details).

The controllability properties of our second example has been analyzed in [34]. Let us consider system (12) with

$$L = L_2 := \begin{pmatrix} -\partial_{xx} & 0 \\ 0 & -\partial_{xx} + Q \end{pmatrix}, \quad D(L_2) = H^2(0, \pi; \mathbb{R}^2) \cap H_0^1(0, \pi; \mathbb{R}^2), \quad (13)$$

with $Q \in L^2(0, \pi)$. In this case, the sequence of eigenvalues of the vectorial operator L_2 is given by $\Lambda^{(2)} = \{k^2\}_{k \geq 1} \cup \{\lambda_k^{(2)}\}_{k \geq 1} \subset \mathbb{R}$, where $\{\lambda_k^{(2)}\}_{k \geq 1}$ is the sequence of eigenvalues of the operator $-\partial_{xx} + Q$ with domain $H^2(0, \pi) \cap H_0^1(0, \pi)$. When $Q \in L^2(0, \pi)$ satisfies

$$\int_0^\pi Q(x) dx = 0, \quad (14)$$

then

$$\lambda_k^{(2)} = k^2 + \varepsilon_k, \quad \forall k \geq 1,$$

with $\{\varepsilon_k\}_{k \geq 1} \in \ell^2$. In particular, $\lim \varepsilon_k = 0$ and $\Lambda^{(2)}$ does not fulfill the gap condition (5). Assume that $\lambda_k^{(2)} \neq n^2$ for any $k, n \geq 1$ (that, in fact, is a necessary condition for the approximate controllability of system (12) with $L = L_2$, see [34] and Section 5). In this case, the sequence $\Lambda^{(2)}$ satisfies property (3). Again, system (12) has a minimal time $T_0 = T_0(Q) \in [0, \infty]$ and there exists coefficients $Q \in L^2(0, \pi)$ such that $T_0(Q) > 0$. Thus, the system is not null controllable at time T when $T < T_0$ (see [34] and Section 5 for the details).

Let us consider a third example of non-scalar parabolic system. In [22] the authors study the boundary null controllability of a phase field system of Caginalp type which is a model describing the transition between the solid and liquid phases in solidification/melting processes of a material occupying the interval $(0, \pi)$. For

that purpose, they consider the nonlinear system

$$\begin{cases} \theta_t - \xi\theta_{xx} + \frac{1}{2}\rho\xi\phi_{xx} + \frac{\rho}{\tau}\theta = f(\phi) & \text{in } (0, T) \times (0, \pi), \\ \phi_t - \xi\phi_{xx} - \frac{2}{\tau}\theta = -\frac{2}{\rho}f(\phi) & \text{in } (0, T) \times (0, \pi), \\ \theta(\cdot, 0) = v, \phi(\cdot, 0) = c, \theta(\cdot, \pi) = 0, \phi(\cdot, \pi) = c & \text{on } (0, T), \\ \theta(0, \cdot) = \theta_0, \phi(0, \cdot) = \phi_0 & \text{in } (0, \pi), \end{cases} \quad (15)$$

where: $\theta = \theta(t, x)$ is the temperature of the material; $\phi = \phi(t, x)$ is the phase-field function used to identify the solidification level of the material; $c \in \{-1, 0, 1\}$; f is the nonlinear term which comes from the derivative of the classical regular double-well potential W :

$$f(\phi) = -\frac{\rho}{4\tau}(\phi - \phi^3).$$

On the other hand, $\rho > 0$, $\tau > 0$ and $\xi > 0$ are, resp., the latent heat, a relaxation time and the thermal diffusivity. Finally, $v \in L^2(0, T)$ is the control function, and θ_0, ϕ_0 are the initial data.

The null controllability property of the nonlinear system (15) depends on the coefficients ρ , τ and ξ . This property is obtained from the corresponding one of the linear version of (15) around the constant trajectory $(0, c)$ (see [22] for more details). This linear system is as system (12) with $y = (\theta, \phi)$ and $L = L_3$ given by

$$\begin{cases} L = L_3 := -D_2\partial_{xx} + A_2, & \text{with} \\ D = D_2 := \begin{pmatrix} \xi & -\frac{1}{2}\rho\xi \\ 0 & \xi \end{pmatrix}, A = A_2 := \begin{pmatrix} \frac{\rho}{\tau} & -\frac{\rho}{2\tau} \\ -\frac{2}{\tau} & \frac{1}{\tau} \end{pmatrix}, B = \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \end{cases} \quad (16)$$

In this case the sequence of eigenvalues of the operator L_3 , with domain $D(L_3) = H^2(0, \pi; \mathbb{R}^2) \cap H_0^1(0, \pi; \mathbb{R}^2)$, is given by $\Lambda^{(3)} = \{\lambda_k^{(3,1)}, \lambda_k^{(3,2)}\}_{k \geq 1}$ with

$$\lambda_k^{(3,1)} = \xi k^2 + \frac{\rho+1}{2\tau} - r_k, \quad \lambda_k^{(3,2)} = \xi k^2 + \frac{\rho+1}{2\tau} + r_k, \quad \forall k \geq 1, \quad (17)$$

where r_k is given by

$$r_k := \sqrt{\frac{\xi\rho}{\tau}k^2 + \left(\frac{\rho+1}{2\tau}\right)^2}, \quad \forall k \geq 1. \quad (18)$$

If $\lambda_k^{(3,1)} \neq \lambda_n^{(3,2)}$ for any $k, n \geq 1$ (which in fact is a condition equivalent to the approximate controllability of the linear system (12) with $L = L_3$), the sequence $\Lambda^{(3)}$ can be rearranged in such a way that $\Lambda^{(3)} = \{\Lambda_k^{(3)}\}_{k \geq 1}$ is an increasing sequence that satisfies (3) for $\delta = 1$. However, if for some integer $j \geq 1$ one has

$$\xi = \frac{1}{j^2} \frac{\rho}{\tau}, \quad (19)$$

then, the eigenvalues of L_3 concentrate and one has

$$\inf_{k \geq 1} \left(\Lambda_{k+1}^{(3)} - \Lambda_k^{(3)} \right) = 0,$$

and condition (5) does not hold (see [22] and Section 5 for the details). Therefore, we have another system where the associated sequence of eigenvalues does not satisfy the gap condition (5).

Remark 1.3. The objective of the work [22] is to study the exact boundary controllability to constant trajectories at time T , $T > 0$ arbitrary, of the nonlinear system (15). To this end, the authors follow a technique developed in [32]. This methodology consists of obtaining a null controllability result at time T for system (12), with $L = L_3$, and an estimate of the cost of fast controls like (10). In order to obtain inequality (10) for the linear version of system (15), the authors assume the condition

$$\xi \neq \frac{1}{j^2} \frac{\rho}{\tau}, \quad \forall j \in \mathbb{N}.$$

This condition is crucial in [22] because it assures that the sequence $\Lambda^{(3)}$ satisfies (5) and the conditions in Theorem 1.2. Thus, system (12), with $L = L_3$, is null controllable at time T for any $T > 0$ and the control cost $\mathcal{K}(T)$ satisfies (10) for a positive constant C_0 only depending on ρ , τ and ξ . \square

1.3. Objective. We have seen three examples of sequences of eigenvalues satisfying (3) and for which the gap condition (5) fails. The corresponding parabolic systems could have a positive minimal time of null controllability T_0 and the system would not be null controllable at time T when $T \in (0, T_0)$. Even if $T_0 = 0$, it is not clear that the control cost of the associated system fulfills inequality (10) or inequality (11) and this is an open problem.

In order to obtain sharp estimates of the control cost $\mathcal{K}(T)$ associated to non-scalar parabolic systems, it is very important to prove sharp estimates for biorthogonal families to the exponentials associated to the corresponding sequence of eigenvalues of the generator when this sequence does not satisfy a gap condition. This is the objective of this work: Given a complex sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ satisfying appropriate assumptions and such that inequality (5) does not hold, is there a biorthogonal family $\{q_k\}_{k \geq 0}$ to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$ (e_k is given in (1)) satisfying an appropriate estimate for $\|q_k\|_{L^2(0, T; \mathbb{C})}$ which, in particular, provides an estimate of the control cost $\mathcal{K}(T)$? Understanding the behavior of the control cost $\mathcal{K}(T)$ for general systems as (12) would allow us to extend the null controllability result in the one-dimensional case to some parabolic systems in any dimension (see for instance [7, 1]) and to some nonlinear parabolic equations using the method of Liu, Takahashi and Tucsnak introduced in [32] (see for instance, [22] and [35]).

Summarizing, in this paper we will consider a class of complex sequences $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$ that satisfy (3) and for which condition (5) fails. To this class of sequences we have as our main objective to prove the existence of families $\{q_k\}_{k \geq 1}$ biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$ that satisfy sharp and explicit estimates of $\|q_k\|_{L^2(0, T; \mathbb{C})}$ with respect to T , Λ_k and some appropriate parameters associated to the class of sequences under consideration. As a second objective, we will apply the previous results to system (12) when $L = L_2$ (see (13)) and $L = L_3$ (see (16)) in order to obtain new results on the cost of the boundary null controllability of these systems at time $T > 0$. In this sense, we will see that the accumulation of the eigenvalues of the operator L_2 implies that the corresponding control cost $\mathcal{K}(T)$ has a more explosive behavior when $T \rightarrow 0$ than in the scalar parabolic case.

1.4. Main results. Let us now present the main results of this work. To this aim, let us first introduce the class of complex sequences we will work with throughout this work:

Definition 1.4. Let $\Lambda = \{\Lambda_k\}_{k \geq 1}$ be a complex sequence and let us fix constants

$$\beta \in [0, \infty), \quad \rho, p_0, p_1, p_2, \alpha \in (0, \infty) \quad \text{and} \quad q \in \mathbb{N}. \quad (20)$$

We say that the sequence Λ is in the class $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$, if the following properties hold:

- (H1) $\Lambda_k \neq \Lambda_n$ for all $n, k \in \mathbb{N}^*$ with $n \neq k$;
- (H2) $\Re(\Lambda_n) > 0$ for every $n \geq 1$;
- (H3) $|\Im(\Lambda_n)| \leq \beta \sqrt{\Re(\Lambda_n)}$, for any $n \geq 1$;
- (H4) $\{\Lambda_n\}_{n \geq 1}$ is nondecreasing in modulus, i.e., $|\Lambda_n| \leq |\Lambda_{n+1}|$, for any $n \geq 1$;
- (H5) $\rho |k^2 - n^2| \leq |\Lambda_k - \Lambda_n|$ for any $n, k \geq 1 : |k - n| \geq q$;
- (H6) $p_1, p_2 \geq p_0$ and one has

$$-\alpha + p_1 \sqrt{r} \leq \mathcal{N}(r) \leq \alpha + p_2 \sqrt{r}, \quad \forall r > 0,$$

where \mathcal{N} is the counting function associated with the sequence Λ , defined by

$$\mathcal{N}(r) = \#\{k : |\Lambda_k| \leq r\}, \quad \forall r > 0. \quad (21)$$

Remark 1.5. Observe that from the definition of the counting function \mathcal{N} (see (21)) associated with the sequence $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ (the parameters are given and satisfy (20)), we deduce the following properties:

1. For any $r > 0$, one has

$$\mathcal{N}(r) = k \iff |\Lambda_k| \leq r \quad \text{and} \quad |\Lambda_{k+1}| > r.$$

2. If for some $k_1, k_2 \geq 1$ and $r_1, r_2 > 0$ one has $|\Lambda_{k_1}| \leq r_1$ and $|\Lambda_{k_2}| > r_2$, then

$$k_1 \leq \mathcal{N}(r_1) \quad \text{and} \quad k_2 \geq \mathcal{N}(r_2) + 1.$$

We will use these properties throughout this work. \square

Remark 1.6. The parameter $q \in \mathbb{N}$ in Definition 1.4 plays an important role in this paper. Observe that in this work we are dealing with sequences Λ that, in general, do not satisfy condition (5) and whose terms could condense. With condition (H5) and the parameter q we “measure” the maximal cardinal of the condensation groupings of the sequence Λ , that is to say, the maximal number of elements in Λ around the term Λ_k that do not satisfy (H5) and could condense. In fact, the parameter q plays an important role even in the case of increasing real sequences that satisfy (5). At the end of Section 2 we will see an example of real sequence Λ that satisfies the gap condition (5) and for which the parameters p_1 and p_2 increasingly depend on q .

Sequences not satisfying the gap condition (5) have also been considered in [8] in the real case. Motivated by extending previous results to sequences satisfying (H5), the authors introduce the so-called block-moment method. In particular this method introduces block-biorthogonal families using a similar approach as in [5]. As a consequence, the authors do not obtain the explicit dependence with respect to T of the L^2 -norm of the elements of the biorthogonal sequence. \square

We will see in Section 2 that the class $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ includes sequences $\Lambda = \{\Lambda_k\}_{k \geq 1}$ satisfying (5) (e.g., sequences satisfying condition (2)), and also sequences where the gap condition (5) fails (for instance, $\Lambda = \{k^2\}_{k \geq 1} \cup \{dk^2\}_{k \geq 1}$, with $\sqrt{d} \notin \mathbb{Q}$, or $\Lambda = \{k^2\} \cup \{k^2 + \varepsilon_k\}$, with $\{\varepsilon_k\}_{k \geq 1} \in \ell^2$, or the sequence considered in [22], see Remark 1.3).

We are now in a position to establish the first main result of this work. It reads as follows:

Theorem 1.7. *Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ be a complex sequence with parameters satisfying (20). Then, given $T > 0$, there exists a family of complex functions $\{q_k\}_{k \geq 1} \subset L^2(0, T; \mathbb{C})$, biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$ (e_k is given in (1)) which, in addition, satisfies*

$$\|q_k\|_{L^2(0, T; \mathbb{C})} \leq \mathcal{H}_1 \exp \left[C \left(1 + \mathcal{H}_2 \sqrt{|\Lambda_k|} + \frac{(1 + p_2)^2}{T} \right) \right] \mathcal{P}_k, \quad (22)$$

for every $k \geq 1$. In (22), C is a positive constant only depending on $|\Lambda_1|$, β , p_0 and α (increasing with respect to α), and \mathcal{P}_k , $\mathcal{H}_1 = \mathcal{H}_1(\rho, q, p_1, p_2)$ and $\mathcal{H}_2 = \mathcal{H}_2(\rho, q, p_1, p_2, T)$ are respectively given by

$$\mathcal{P}_k := \prod_{\{n \geq 1: 1 \leq |k-n| < q\}} |\Lambda_k - \Lambda_n|^{-1}, \quad \forall k \geq 1, \quad \text{if } q \geq 2, \quad (23)$$

$\mathcal{P}_k := 1$, for every $k \geq 1$, if $q = 1$,

$$\begin{cases} \mathcal{H}_1(\rho, q, p_1, p_2) = \left(\frac{1 + \rho p_2^2 + q^2}{\rho^2 p_1^4} \right)^{2q-2}, \\ \mathcal{H}_1(\rho, q, p_1, p_2) = \left(\frac{1 + \rho p_2^2}{\rho^2 p_1^4} \right)^{2q-2}, \quad \text{when } \Lambda \text{ is real} \end{cases}, \quad (24)$$

and

$$\begin{cases} \mathcal{H}_2(\rho, q, p_1, p_2, T) = 1 + q + \sqrt{T} + \frac{1 + q}{\rho^2 p_1^2} + p_2, \\ \mathcal{H}_2(\rho, q, p_1, p_2, T) = 1 + q + \sqrt{T} + \frac{1}{\rho^2 p_1^2} + p_2, \quad \text{when } \Lambda \text{ is real.} \end{cases} \quad (25)$$

Remark 1.8. It is clear that if $\Lambda = \{\Lambda_k\}_{k \geq 1}$ is a sequence satisfying the assumptions in Theorem 1.2 for some parameters $\beta \in [0, \infty)$, $\rho, p, \alpha \in (0, \infty)$ and $q \in \mathbb{N}$, then Λ belongs to $\mathcal{L}(\beta, \rho, q, p, p, \alpha)$, and satisfies

$$|\Lambda_k - \Lambda_n| \geq \gamma > 0, \quad \forall k, n \geq 1 : k \neq n,$$

for a positive constant γ . As a consequence, we can apply Theorem 1.7 and deduce the existence of $\{q_k\}_{k \geq 1} \subset L^2(0, T; \mathbb{C})$, a biorthogonal family to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$, satisfying (22). Thanks to the previous gap condition, we get $\mathcal{P}_k = 1$, if $q = 1$, or

$$\mathcal{P}_k \leq \gamma^{2-2q}, \quad \forall k \geq 1, \quad \text{if } q \geq 2.$$

Combining this inequality and (22) we deduce (6) for a positive constant \mathcal{C}_0 independent of k and T . Therefore, Theorem 1.7 is a generalization of Theorem 1.2 to the case of complex sequences that do not satisfy the gap condition (5).

We will also see in Section 2 that Theorem 1.7 generalizes the results on bounds of biorthogonal families to exponentials proved in [17], [33], [15], [13] and [14]. \square

The quantity \mathcal{P}_k in Theorem 1.7 provides a measure of the condensation of the sequence Λ . When condition (5) holds, then, there exists a constant $\mathcal{C} > 0$ such that $|\mathcal{P}_k| \leq \mathcal{C}$ for any positive integer k . But in general, \mathcal{P}_k could have any explosive behavior with respect to k (see for instance Remark 5.8).

In the next result we will prove that inequality (22) is optimal with respect to \mathcal{P}_k . This is our second main result:

Theorem 1.9. Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$ be a complex sequence satisfying

$$|\Lambda_k - \Lambda_n| \leq \nu |k^2 - n^2|, \quad \forall k, n \geq 1, \quad (26)$$

for $\nu > 0$, and $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$, for parameters satisfying (20). Then, for any sequence $\{q_k\}_{k \geq 1} \subset L^2(0, T; \mathbb{C})$ biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$ (e_k is given in (1)), one has

$$\|q_k\|_{L^2(0, T; \mathbb{C})} \geq \max \left\{ \frac{6}{\pi^2} \mathcal{B}_k e^{\frac{1}{T\nu}} \sqrt{\delta |\Lambda_1| + \frac{1}{2T}}, \mathcal{E}_k \right\} \mathcal{P}_k, \quad \forall k \geq 3, \quad (27)$$

where \mathcal{P}_k is given in (23),

$$\mathcal{B}_k = \begin{cases} \nu^{k+q-2} \frac{(q-1)!}{(q+3)!} (k+q)! \frac{(\nu T)^{k+1}}{(1+\nu T)^{2k+q+1}} \frac{(2k+q-1)!}{(2k+q+1)!}, & \text{if } k < q, \\ \nu^{2(q-1)} \frac{[(q-1)!]^2}{(q+3)!} \frac{(k+q)!k}{(2k-q)!} \frac{(\nu T)^{k+1}}{(1+\nu T)^{2k+q+1}} \frac{(2k+q-1)!}{(2k+q+1)!}, & \text{if } k \geq q, \end{cases} \quad (28)$$

$$\mathcal{E}_k = \begin{cases} \frac{(k+q-2)!}{T^{k+q-2}} \left(\frac{2(k+q)-3}{2T} + \delta |\Lambda_1| \right)^{1/2}, & \text{if } 1 \leq k < q, \\ \frac{(2q-2)!}{T^{2(q-1)}} \left(\frac{4q-3}{2T} + \delta |\Lambda_{k+1-q}| \right)^{1/2}, & \text{if } k \geq q, \end{cases} \quad (29)$$

and δ is a positive constant only depending on β ($\delta = 1$ when $\beta = 0$).

Remark 1.10. It is important to note that the main estimates of $\|q_k\|_{L^2(0, T; \mathbb{C})}$ in Theorems 1.7 and 1.9 are a combination of two phenomena. The first one comes from the fact of having an infinite family of exponential functions associated to the sequence Λ . This introduces the terms

$$\mathcal{H}_1(\rho, q, p_1, p_2) \exp \left[C \left(1 + \mathcal{H}_2(\rho, q, p_1, p_2, T) \sqrt{|\Lambda_k| + \frac{(1+p_2)^2}{T}} \right) \right]$$

in inequality (22) of Theorem 1.7, and

$$\max \left\{ \frac{6}{\pi^2} \mathcal{B}_k e^{\frac{1}{T\nu}}, \mathcal{E}_k \right\}$$

in inequality (27) of Theorem 1.9. This kind of terms also appears in the case of sequences satisfying the gap condition (5). The second phenomenon is new. It is the contribution of the condensation of a finite number of elements of the sequence Λ . This condensation introduces the factor \mathcal{P}_k in inequalities (22) and (27). Of course, this new term does not appear when the sequence Λ satisfies the gap condition (5) (see Remark 1.8). Observe that even if this term involves a finite number of elements of Λ , its contribution to the corresponding control cost $\mathcal{K}(T)$ of the associated parabolic control problem could be more explosive than the contribution of the first one (see Remark 5.8). To our knowledge, this is the first time where this fact has been highlighted. \square

Remark 1.11. Theorem 1.9 generalizes the results proved in [23], [13] and [14] to general complex sequences that might not satisfy the gap condition (5). \square

Remark 1.12. In [11], the authors revisit the block moment method and obtain similar results to those of Theorem 1.7 under similar conditions on the sequence Λ using a different approach. To be precise, the authors use the Laplace transform

and an explicit estimate of the norm of the restriction operator between the space generated by the exponentials in $L^2(0, \infty; \mathbb{C})$ and the space generated in $L^2(0, T; \mathbb{C})$.

□

As an application of Theorems 1.7 and 1.9, we will study the cost of fast controls $\mathcal{K}(T)$ for system (12) in two situations in which condition (5) does not hold:

1. First, we will analyze system (12) when the operator $L = L_2$ is given by (13) with $Q \in L^2(0, \pi)$ such that the sequence of eigenvalues of L_2 is given by

$$\Lambda^{(2)} = \left\{ k^2, k^2 + e^{-k^{2\gamma}} \right\}_{k \geq 1}$$

and $\gamma \in (0, 1)$. In this example the minimal time associated to system (12) with $L = L_2$ is $T_0(Q) = 0$. Observe that the sequence $\Lambda^{(2)}$ does not satisfy (5) and, therefore, Theorem 1.2 cannot be applied. We will see that the sequence $\Lambda^{(2)}$ fulfills the assumptions in Theorems 1.7 and 1.9 and, as a consequence, we will obtain new estimates from above and from below for the control cost $\mathcal{K}(T)$ associated to system (12) for $L = L_2$ (see Theorems 5.9 and 5.10). These estimates show that the fast controls for system (12) with $L = L_2$ are more violent than those of the heat equation. This violent behavior comes from the condensation of the eigenvalues of the elliptic operator L_2 .

2. We will also study system (12) with $L = L_3$ (see (16)), and ρ, τ and ξ positive constant satisfying (19) for an integer $j \geq 1$. In this case we will check that system (12) is null controllable for any $T > 0$ and the corresponding control cost $\mathcal{K}(T)$ satisfies (10) for a constant $C_0 = C_0(\rho, \tau, \xi) > 0$. With this example we generalize the null controllability result obtained in [22] for the linear version of (15).

In a forthcoming paper (see [9]) we will carry out a more in-depth analysis of the cost of fast controls $\mathcal{K}(T)$ of parabolic systems with a positive minimal time T_0 which comes from the condensation index associated to the sequence of eigenvalues of the generator of the corresponding C_0 -semigroup.

1.5. Plan of the paper. The plan of the paper is the following: In Section 2, we will study some general properties of the sequences Λ in $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$, with parameters satisfying (20). We will also provide in this section some examples of sequences Λ in the literature that satisfy the conditions in Definition 1.4. Sections 3 and 4 will be respectively devoted to the proofs of the main results of this work, namely, Theorem 1.7 and Theorem 1.9. Finally, in Section 5 we will apply the results on general bounds of biorthogonal families to complex sequences that do not satisfy the gap condition (5) to system (12) when $L = L_2$ (see (13)) and

$$\sigma(L_2) = \left\{ k^2, k^2 + e^{-k^{2\gamma}} \right\}_{k \geq 1}$$

with $\gamma \in (0, 1)$, and when $L = L_3$ (see (16)) is such that $\rho, \tau, \xi \in (0, \infty)$ satisfy (19) for an integer $j \geq 1$. Some results presented in this fifth section have been announced in [21].

2. Some general properties of sequences under the assumptions of Definition 1.4. Some examples. We will devote this section to prove some general properties of sequences Λ in the class of Definition 1.4. These properties will be used in the proof of Theorems 1.7 and 1.9. We also complete this section with some examples of sequences Λ that fulfill assumptions in Definition 1.4.

2.1. Some relations between the conditions in Definition 1.4. Real sequences. Let us first analyze the conditions which appear in Definition 1.4 and condition (26) because in some particular cases they are redundant. To be precise, let us first check that the properties (H1)–(H5) and (26) imply property (H6) for some p_0, p_1, p_2 and α . One has:

Proposition 2.1. *Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$ be a complex sequence. Then,*

1. *If Λ satisfies properties (H1), (H4), (H5) and (26) for some $\rho, \nu > 0$ and $q \geq 1$, then, (H6) holds, with*

$$p_0 = \frac{1}{\sqrt{\nu}}, \quad p_1 = \frac{1}{\sqrt{\nu}}, \quad p_2 = \frac{1}{\sqrt{\rho}}, \quad \alpha = \max \left\{ q - \sqrt{\frac{|\Lambda_1|}{\rho}}, \sqrt{\frac{|\Lambda_1|}{\rho}} + 1, \sqrt{\frac{|\Lambda_1|}{\nu}} + 1 \right\}.$$

2. *Assume now that Λ fulfills (H1), (H4) and (H6) for some positive constants p_0, p_1, p_2 . Then,*
 - (a) *If Λ satisfies property (H5), then*

$$p_1 \leq \frac{1}{\sqrt{\rho}}. \quad (30)$$

- (b) *If (26) holds, then*

$$\frac{1}{\sqrt{\nu}} \leq p_2.$$

Proof. Let us first assume that $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$ satisfies the assumption in item 1 of the proposition, and let us prove that (H6) holds for appropriate parameters p_0, p_1, p_2 and α .

From (H5) and (26), we have $\rho(k^2 - n^2) \leq |\Lambda_k| + |\Lambda_n|$, for any $k, n : k \geq n + q$, and $|\Lambda_k| - |\Lambda_n| \leq \nu(k^2 - n^2)$, for any $k, n : k \geq n$. In particular,

$$\begin{cases} |\Lambda_k| \geq \rho(k^2 - 1) - |\Lambda_1|, & \forall k \geq q + 1, \\ |\Lambda_k| \leq \nu(k^2 - 1) + |\Lambda_1|, & \forall k \geq 1. \end{cases} \quad (31)$$

Let us consider $r \geq |\Lambda_{q+1}|$. Taking into account the first item in Remark 1.5, if $\mathcal{N}(r) = k$, then $k \geq q + 1$, $|\Lambda_k| \leq r$ and $|\Lambda_{k+1}| > r$. The first inequality in (31) gives $r \geq \rho(k^2 - 1) - |\Lambda_1|$, i.e.,

$$\mathcal{N}(r) = k \leq \sqrt{\frac{1}{\rho}r + \frac{|\Lambda_1|}{\rho}} + 1 \leq \frac{1}{\sqrt{\rho}}\sqrt{r} + \sqrt{\frac{|\Lambda_1|}{\rho}} + 1, \quad \forall r \geq |\Lambda_{q+1}|. \quad (32)$$

On the other hand, the second inequality in (31) also provides $r < \nu[(k+1)^2 - 1] + |\Lambda_1|$ and

$$\mathcal{N}(r) = k > -1 + \sqrt{\frac{1}{\nu}r - \frac{|\Lambda_1|}{\nu}} + 1 > \sqrt{\frac{1}{\nu}r - \frac{|\Lambda_1|}{\nu}} - 1 \geq \frac{1}{\sqrt{\nu}}\sqrt{r} - \sqrt{\frac{|\Lambda_1|}{\nu}} - 1. \quad (33)$$

Observe that this inequality is also valid when $0 < r < |\Lambda_{q+1}|$. In the previous reasoning we have used the inequalities

$$\begin{cases} \sqrt{a+b} \leq \sqrt{a} + \sqrt{b}, & \forall a, b \in [0, \infty), \\ \sqrt{a-b} \geq \sqrt{a} - \sqrt{b}, & \forall a, b \in [0, \infty), \quad a \geq b. \end{cases} \quad (34)$$

Let us now take r such that $|\Lambda_1| \leq r < |\Lambda_{q+1}|$. In this case,

$$\mathcal{N}(r) \leq q \leq \frac{1}{\sqrt{\rho}}\sqrt{r} + q - \frac{1}{\sqrt{\rho}}\sqrt{|\Lambda_1|}.$$

Finally, when r is such that $0 < r < |\Lambda_1|$, $\mathcal{N}(r) = 0 \leq \sqrt{r}/\sqrt{\rho}$. We deduce then that Λ satisfies (H6) with p_0, p_1, p_2 and α given in the statement. This proves the first item.

The second item in the statement can be deduced from (32) and (33). Indeed, if Λ satisfies (H5) and (H6), one has (32) and $p_1\sqrt{r} - \alpha \leq \mathcal{N}(r)$, for any $r > 0$. This clearly implies (30). On the other hand, if the sequence Λ satisfies (26), then (33) holds. Combining this inequality with the second inequality in (H6), we get the second point in item. This proves the result. \square

Remark 2.2. Property (H5) does not imply, in general, (H6), even for increasing positive real sequences. Indeed, $\Lambda = \{k^3\}_{k \geq 1}$ is an increasing positive real sequence that satisfies (H5), with $\rho = 1$ and $q = 1$, and does not satisfy (H6).

Something similar can be said for property (26): $\Lambda = \{k\}_{k \geq 1}$ is an increasing positive real sequence that satisfies (26) with $\nu = 1$ and does not satisfy (H6).

On the other hand, sequences Λ satisfying (H1)–(H5) and (26) for $\beta \geq 0, \rho, \nu > 0$ and $q \geq 1$, also satisfy condition (H6) with parameters p_0, p_1, p_2 and α given in the statement of Proposition 2.1. In conclusion, $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$. \square

Remark 2.3. Observe that, if Λ is a sequence under the conditions of item 2 in Proposition 2.1, from inequality (30) we also deduce

$$0 < \rho \leq \frac{1}{p_0^2}, \quad \rho p_1^2 \leq 1, \quad \text{and} \quad \rho p_1 \leq \sqrt{\rho} \leq \frac{1}{p_0}. \quad (35)$$

These estimates will be used later. \square

Let us now analyze the case of increasing positive real sequences $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset (0, \infty)$. This case is specially interesting because some assumptions in Definition 1.4 are direct. For instance, Λ satisfies (H1)–(H4) for $\beta = 0$. In addition, one has:

Proposition 2.4. *Let $\Lambda = \{\Lambda_k\}_{k \geq 1}$ be a positive real sequence satisfying (H1), (H4) and (H6) for some $p_0, \alpha \in (0, \infty)$, with $p_1 = p_2 = p \geq p_0$. Then, $\Lambda \in \mathcal{L}(0, \rho, q, p_0, p, p, \alpha)$ and (26) holds, with*

$$q = 3\alpha, \quad \rho = \frac{1}{3p^2} \quad \text{and} \quad \nu = \frac{1}{3} \left(\frac{2 + \alpha}{p} \right)^2. \quad (36)$$

Proof. Let us take $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset (0, \infty)$, a sequence satisfying (H1), (H4) and (H6) for some $p_0, \alpha \in (0, \infty)$, with $p_1 = p_2 = p \geq p_0$. It is clear that Λ satisfies (H2) and (H3) for $\beta = 0$.

Let us see that Λ also satisfies (H5) for appropriate constants ρ and q . Indeed, using (H1) and (H4) we infer that Λ is an increasing positive real sequence. Thus, $\mathcal{N}(\Lambda_k) = k$, for any $k \geq 1$, (see (21)) and, from (H6) ($p_1 = p_2 = p$), we deduce

$$k - \alpha \leq p\sqrt{\Lambda_k} \leq k + \alpha, \quad \forall k \geq 1. \quad (37)$$

If $k, n \in \mathbb{N}$ are such that $k - n \geq 3\alpha$, then, $k \geq \alpha$ and inequality (37) provides

$$\frac{p^2(\Lambda_k - \Lambda_n)}{k^2 - n^2} \geq \frac{(k - \alpha)^2 - (n + \alpha)^2}{k^2 - n^2} = \frac{k - n - 2\alpha}{k - n} = 1 - \frac{2\alpha}{k - n} \geq \frac{1}{3}.$$

Therefore, sequence Λ satisfies assumption (H5) for q and ρ as in the statement of the proposition.

Let us now check property (26). To this end, we will again use (37). Without loss of generality, we can assume that $\alpha \geq 1$. Thus, if $\alpha < n < k$, one has

$$\frac{p^2(\Lambda_k - \Lambda_n)}{k^2 - n^2} \leq \frac{(k + \alpha)^2 - (n - \alpha)^2}{k^2 - n^2} = 1 + \frac{2\alpha}{k - n} \leq 1 + 2\alpha \leq \frac{1}{3}(2 + \alpha)^2.$$

On the other hand, if $n \leq \alpha < k$, i.e., if $n \leq \lfloor \alpha \rfloor < \lfloor \alpha \rfloor + 1 \leq k$ ($\lfloor \cdot \rfloor$ is the floor function: given $x \in \mathbb{R}$, $\lfloor x \rfloor$ is the greatest integer less than or equal to x), we also deduce

$$\frac{p^2(\Lambda_k - \Lambda_n)}{k^2 - n^2} \leq \frac{(k + \alpha)^2}{k^2 - \lfloor \alpha \rfloor^2} \leq \frac{(\lfloor \alpha \rfloor + \alpha + 1)^2}{2\lfloor \alpha \rfloor + 1} \leq \frac{1}{3}(2 + \alpha)^2.$$

In the previous inequality we have used that $\alpha \geq 1$.

Finally, let us assume that $\alpha \geq 2$ and take $n < k \leq \alpha$. We can write

$$\frac{p^2(\Lambda_k - \Lambda_n)}{k^2 - n^2} \leq \frac{(k + \alpha)^2}{2k - 1} \leq \frac{1}{3}(2 + \alpha)^2.$$

Summarizing, property (H5) holds for ν given in (36). This ends the proof. \square

Remark 2.5. Let us consider $\Lambda = \{\Lambda_k\}_{k \geq 1}$, an increasing positive sequence, satisfying property (H6) with $p_1 = p_2 = p > 0$. In this case, this condition can be written under the equivalent form

$$\Lambda_k = \frac{1}{p^2}k^2 + O(k), \quad \forall k \geq 1. \quad (38)$$

Indeed, from (H6) with $p_1 = p_2 = p$, we infer (37), i.e., $p\sqrt{\Lambda_k} = k + O(1)$ for any $k \geq 1$. So, (38) holds. On the other hand, from (38) we deduce

$$\frac{1}{p^2}k^2 - \alpha_1 k \leq \Lambda_k \leq \frac{1}{p^2}k^2 + \alpha_1 k, \quad \forall k \geq 1,$$

with $\alpha_1 \geq 0$. Thus, given $r > 0$, if $\mathcal{N}(r) = k$, then, (see Remark 1.5) we also have

$$\frac{1}{p^2}k^2 - \alpha_1 k \leq \Lambda_k \leq r \quad \text{and} \quad r < \Lambda_{k+1} \leq \frac{1}{p^2}(k+1)^2 + \alpha_1(k+1),$$

i.e.,

$$\begin{cases} \mathcal{N}(r) = k \leq \frac{1}{2} \left(p^2 \alpha_1 + p \sqrt{p^2 \alpha_1^2 + 4r} \right) \leq p\sqrt{r} + p^2 \alpha_1, \\ \mathcal{N}(r) + 1 = k + 1 > \frac{1}{2} \left(-p^2 \alpha_1 + p \sqrt{p^2 \alpha_1^2 + 4r} \right) > p\sqrt{r} - \frac{1}{2} p^2 \alpha_1. \end{cases}$$

Therefore, (H6) holds with $p_0 = p_1 = p_2 = p$ and

$$\alpha = \max \left\{ p^2 \alpha_1, \frac{1}{2} p^2 \alpha_1 + 1 \right\}.$$

Observe that, in particular, if $\Lambda = \{\Lambda_k\}_{k \geq 1}$ is an increasing real sequence such that (2) holds, then Λ also satisfies (H6) with $A = 1/p^2$. Proposition 2.4 implies that $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ and (26) holds for $\beta = 0$, $p_0 = p_1 = p_2 = p = 1/\sqrt{A}$, $\alpha \in (0, \infty)$ and q, ρ and ν as in (36). Therefore, Theorems 1.7 and 1.9 generalize the results on estimates of biorthogonal families established in [17] and [23]. \square

2.2. Some additional properties of sequences in the class \mathcal{L} . Let us continue showing some properties for sequences Λ in the class $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$. One has:

Lemma 2.6. *Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ be a complex sequence with parameters satisfying (20). Then,*

$$\sum_{k \geq 1} \frac{1}{|\Lambda_k|} < \infty \quad \text{and} \quad |\Lambda_k| \leq \Re(\Lambda_k) + \beta \sqrt{\Re(\Lambda_k)}, \quad \forall k \geq 1. \quad (39)$$

On the other hand, there exists a positive constant C , only depending on $|\Lambda_1|$, β , p_0 and α (increasing with respect to α), such that

$$\frac{1}{p_2}(k - \alpha) \leq \sqrt{|\Lambda_k|} \leq \frac{1}{p_1}k + \frac{C(1+q)}{\rho p_1^2}, \quad \forall k \geq 1. \quad (40)$$

Proof. Let us take a sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ under assumptions of the lemma. From items (H4) and (H6) of Definition 1.4, we have that:

$$\begin{aligned} \sum_{k \geq 1} \frac{1}{|\Lambda_k|} &= \int_{|\Lambda_1|}^{\infty} \frac{1}{r} d\mathcal{N}(r) = \int_{|\Lambda_1|}^{\infty} \frac{1}{r^2} \mathcal{N}(r) dr \leq \int_{|\Lambda_1|}^{\infty} \frac{\alpha + p_2 \sqrt{r}}{r^2} dr \\ &= \frac{\alpha}{|\Lambda_1|} + \frac{2p_2}{\sqrt{|\Lambda_1|}} < \infty. \end{aligned}$$

On the other hand, using assumption (H3), we deduce that

$$|\Lambda_k|^2 = \Re(\Lambda_k)^2 + \Im(\Lambda_k)^2 \leq \Re(\Lambda_k)^2 + \beta^2 \Re(\Lambda_k) \leq \left(\Re(\Lambda_k) + \beta \sqrt{\Re(\Lambda_k)} \right)^2.$$

Therefore, we have the proof of (39).

Let us now prove property (40). Let us first assume that Λ is a positive real sequence in $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ ($\beta = 0$). We have that $\mathcal{N}(\Lambda_k) = k$, for any $k \geq 1$. In particular, taking $r = \Lambda_k$ in assumption (H6), we deduce

$$\frac{k - \alpha}{p_2} \leq \sqrt{\Lambda_k} \leq \frac{k + \alpha}{p_1} = \frac{k}{p_1} + \frac{\alpha}{p_1} \leq \frac{k}{p_1} + \frac{\alpha}{p_0} \frac{1}{\rho p_1^2} \leq \frac{k}{p_1} + \frac{\alpha}{p_0} \frac{1}{\rho p_1^2}, \quad \forall k \geq 1.$$

In the previous inequality we have used (35). This shows inequality (40) in the real case.

Let us now assume that the sequence $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ is complex, i.e., $\beta > 0$. As before, we are going to work with property (H6) with $r = |\Lambda_k|$ ($k \geq 1$). From Remark 1.5, (H4) and (H6) (see Definition 1.4), we can write that, if $n = \mathcal{N}(|\Lambda_k|)$, then $k \leq n$, $|\Lambda_k| = |\Lambda_n|$ and

$$\frac{-\alpha + n}{p_2} = \frac{-\alpha + \mathcal{N}(|\Lambda_k|)}{p_2} \leq \sqrt{|\Lambda_k|} \leq \frac{\alpha + \mathcal{N}(|\Lambda_k|)}{p_1} = \frac{\alpha + n}{p_1}, \quad \forall k \geq 1. \quad (41)$$

In particular, $k \leq n$ and

$$-\alpha + k \leq -\alpha + n \leq p_2 \sqrt{|\Lambda_k|}, \quad \forall k \geq 1.$$

This proves the first inequality in (40) in the complex case.

In order to show the second inequality in (40), let us estimate $n = \mathcal{N}(|\Lambda_k|)$. As $|\Lambda_k| = |\Lambda_n|$, using property (H3), we infer

$$\left| \Re(\Lambda_n)^2 - \Re(\Lambda_k)^2 \right| = \left| \Im(\Lambda_n)^2 - \Im(\Lambda_k)^2 \right| \leq \beta^2 (\Re(\Lambda_k) + \Re(\Lambda_n)),$$

that is to say,

$$|\Re(\Lambda_k) - \Re(\Lambda_n)| \leq \beta^2.$$

Again, assumption (H3) also provides the inequality

$$|\Lambda_k - \Lambda_n| \leq |\Re(\Lambda_k) - \Re(\Lambda_n)| + |\Im(\Lambda_k) - \Im(\Lambda_n)| \leq \beta^2 + 2\beta\sqrt{|\Lambda_k|}.$$

If $|k - n| \geq q$, combining the previous inequality and assumption (H5) we obtain

$$\rho|k - n|(k + n) = \rho|k^2 - n^2| \leq |\Lambda_k - \Lambda_n| \leq \beta^2 + 2\beta\sqrt{|\Lambda_k|}.$$

Thus,

$$n - k = |k - n| \leq \max \left\{ q, \frac{\beta^2 + 2\beta\sqrt{|\Lambda_k|}}{\rho(k + n)} \right\},$$

i.e.,

$$n \leq k + \max \left\{ q, \frac{\beta^2 + 2\beta\sqrt{|\Lambda_k|}}{\rho(k + \mathcal{N}(|\Lambda_k|))} \right\},$$

and, from (41)

$$p_1\sqrt{|\Lambda_k|} \leq \alpha + k + \max \left\{ q, \frac{\beta^2 + 2\beta\sqrt{|\Lambda_k|}}{\rho(k + \mathcal{N}(|\Lambda_k|))} \right\}. \quad (42)$$

If the maximum in (42) is q , in particular,

$$p_1\sqrt{|\Lambda_k|} \leq k + \alpha + q.$$

Taking into account inequalities (30) and (35), we also deduce

$$p_1\sqrt{|\Lambda_k|} \leq k + \frac{\alpha + q}{\rho p_1} \rho p_1 \leq k + \frac{(\alpha + q)/p_0}{\rho p_1}.$$

Thus, we get the second inequality in (40) for a positive constant C only depending on α and p_0 and increasing with respect to α .

Let us now assume that the maximum in (42) is given by the second term. Using again (H6) and (35), for $k \geq \alpha$, we can write

$$\begin{aligned} p_1\sqrt{|\Lambda_k|} &\leq k + \alpha + \frac{\beta^2 + 2\beta\sqrt{|\Lambda_k|}}{\rho(k + \mathcal{N}(|\Lambda_k|))} \leq k + \alpha + \frac{\beta^2 + 2\beta\sqrt{|\Lambda_k|}}{\rho(k - \alpha + p_1\sqrt{|\Lambda_k|})} \\ &\leq k + \frac{\alpha}{\rho p_1} \frac{1}{p_0} + \frac{\beta^2 + 2\beta\sqrt{|\Lambda_k|}}{\rho p_1\sqrt{|\Lambda_k|}}. \end{aligned}$$

This inequality provides the second inequality in (40) when $k \geq \alpha$ for a positive constant C only depending on $|\Lambda_1|$, β , p_0 and α (of course, increasing with respect to α).

Finally, let us consider the case $k < \alpha$. Thus, there exists a positive constant C (only depending on α and increasing with respect to α) such that

$$\sqrt{|\Lambda_k|} \leq C \leq \frac{k}{p_1} + \frac{C}{\rho p_1^2}.$$

In the previous inequality we have used (35). This ends the proof. \square

Remark 2.7. Analyzing the proof of Lemma 2.6 we deduce that, in fact, if the sequence Λ is real, then the second inequality in (40) can be written as follows: there exists a positive constant C , only depending on p_0 and α (increasing with respect to α) such that

$$\frac{1}{p_2}(k - \alpha) \leq \sqrt{|\Lambda_k|} \leq \frac{1}{p_1}k + \frac{C}{\rho p_1^2}, \quad \forall k \geq 1. \quad (43)$$

In particular the previous inequalities are independent of q . We will use this property for real sequences Λ throughout the paper. \square

Remark 2.8. From Lemma 2.6 we deduce that, if the sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ is in the class $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$, then one also has (3) for some $\delta > 0$, only depending on β ($\delta = 1$ when $\beta = 0$). As said before, property (3) implies that the family of exponentials $\{e_k\}_{k \geq 1}$, e_k is given in (1), is minimal in $L^2(0, T; \mathbb{C})$ for any $T > 0$. Thus, there exists a biorthogonal family $\{\tilde{e}_k\}_{k \geq 1}$ to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$ (see for instance [39], [37], [4], Theorem 4.1 in [5],...). \square

2.3. Some examples of sequences in $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$. Let us complete this section providing some examples of sequences $\Lambda = \{\Lambda_k\}_{k \geq 1}$ such that $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ for some parameters satisfying (20). In order to have a clearer exposition, we will present the results and we will include the corresponding proofs in an appendix, at the end of this paper.

In [13] and [14], the authors consider increasing real positive sequences $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{R}$ satisfying a ‘‘global gap condition’’:

$$\gamma_0 \leq \sqrt{\Lambda_{k+1}} - \sqrt{\Lambda_k} \leq \gamma_1, \quad \forall k \geq 1, \quad (44)$$

and an ‘‘asymptotic gap condition’’:

$$\gamma_0^* \leq \sqrt{\Lambda_{k+1}} - \sqrt{\Lambda_k} \leq \gamma_1^*, \quad \forall k \geq N,$$

where $N \geq 1$ and $\gamma_0, \gamma_1, \gamma_0^*, \gamma_1^* \in (0, \infty)$ are such that $0 < \gamma_1^* - \gamma_0^* < \gamma_1 - \gamma_0$. Under these assumptions on Λ the authors obtain general and precise upper and lower bounds for biorthogonal families as (6) or (7), paying attention to the dependence of the constant \mathcal{C}_0 and \mathcal{C}_1 with respect to the parameters $\gamma_0, \gamma_1, \gamma_0^*$ and γ_1^* .

One has:

Proposition 2.9. *Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset (0, \infty)$ be a real sequence satisfying (44) for two constants $\gamma_0, \gamma_1 > 0$. Then, $\Lambda \in \mathcal{L}(0, \rho, q, p_0, p_1, p_2, \alpha)$ and (26) holds with*

$$\begin{cases} p_0 = p_1 = \frac{1}{\gamma_1}, & p_2 = \frac{1}{\gamma_0}, & \alpha = \max \left\{ 1 - \frac{\sqrt{\Lambda_1}}{\gamma_0}, \frac{\sqrt{\Lambda_1}}{\gamma_1} \right\}, \\ q = 1, & \rho = \min \left\{ \gamma_0^2, \frac{1}{3}\gamma_0^2 + \frac{2}{3}\gamma_0\sqrt{\Lambda_1} \right\} & \text{and } \nu = \max \left\{ \gamma_1^2, \frac{1}{3}\gamma_1^2 + \frac{2}{3}\gamma_1\sqrt{\Lambda_1} \right\}. \end{cases}$$

In particular, the gap condition (5) holds.

For the proof, see A.1.

Remark 2.10. Sequences $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset (0, \infty)$ under the assumptions of Proposition 2.9 satisfy the general assumptions in Theorem 1.7 and Theorem 1.9 with parameters given in the statement of the proposition. Observe, in particular, that $q = 1$ and $\mathcal{P}_k = 1$. Thus, Theorems 1.7 and 1.9 cover the results in [13] and [14]. \square

We continue our analysis of real sequences that fulfill general assumptions previously discussed in the literature. In [15], the authors consider a real increasing sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ that is given as

$$\Lambda = \left\{ \lambda_k^{(1)} \right\}_{k \geq 1} \cup \left\{ \lambda_k^{(2)} \right\}_{k \geq 1},$$

with $\{\lambda_k^{(1)}\}_{k \geq 1}$ and $\{\lambda_k^{(2)}\}_{k \geq 1}$ two increasing sequences of positive real numbers satisfying

$$\begin{cases} \left| \lambda_k^{(1)} - \frac{1}{\pi_1^2} k^2 \right| \leq c_1 k, & \left| \lambda_k^{(2)} - \frac{1}{\pi_2^2} k^2 \right| \leq c_1 k, & \forall k \geq 1, \\ \inf_{n \geq 1} \left| \sqrt{\lambda_k^{(2)}} - \sqrt{\lambda_n^{(1)}} \right| \geq \frac{r}{k}, & \forall k \geq 1, \end{cases} \quad (45)$$

and the strong gap condition

$$\sqrt{\lambda_{k+1}^{(1)}} - \sqrt{\lambda_k^{(1)}} \geq c_2, \quad \sqrt{\lambda_{k+1}^{(2)}} - \sqrt{\lambda_k^{(2)}} \geq c_2, \quad \forall k \geq 1, \quad (46)$$

for some positive constants π_1, π_2, c_1, c_2 and r . For this class of sequences, the authors prove the existence of a sequence $\{q_k\}_{k \geq 1} \subset L^2(0, T)$ ($T > 0$ is given) biorthogonal to $\{e_k\}_{k \geq 1}$ (e_k given in (1)) in $L^2(0, T)$ which satisfies (6) for a positive constant \mathcal{C}_0 independent of k and T and uniform for the class of sequences Λ satisfying the previous assumptions.

One has:

Proposition 2.11. *Let us consider two increasing sequences of positive real numbers $\Lambda_1 = \{\lambda_k^{(1)}\}_{k \geq 1}$ and $\Lambda_2 = \{\lambda_k^{(2)}\}_{k \geq 1}$ satisfying (45) and*

$$\lambda_{k+1}^{(1)} - \lambda_k^{(1)} \geq c_0 \quad \text{and} \quad \lambda_{k+1}^{(2)} - \lambda_k^{(2)} \geq c_0, \quad \forall k \geq 1, \quad (47)$$

for some positive constants π_1, π_2, c_0, c_1 and r . Then, the sequence $\Lambda = \Lambda_1 \cup \Lambda_2$ can be rearranged as an increasing sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ satisfying the gap condition (5), $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ and (26), with $\beta = 0$, $p_0 = \min\{\pi_1, \pi_2\}$, $p_1 = p_2 = p = \pi_1 + \pi_2$,

$$\alpha = \max \left\{ 2 + \frac{1}{2} c_1 (\pi_1^2 + \pi_2^2), c_1 (\pi_1^2 + \pi_2^2) \right\}$$

and q, ρ and ν given in (36).

The proof of this result can be seen in A.2.

Remark 2.12. Observe that the sequences $\{\lambda_k^{(1)}\}_{k \geq 1}$ and $\{\lambda_k^{(2)}\}_{k \geq 1}$ satisfying (45), for some positive constants π_1, π_2, c_1 and r , and the strong gap condition (46), with c_2 a positive constant, in particular, fulfill assumptions (45) and (47) in Proposition 2.11 (and, therefore, the general hypotheses imposed to general complex sequences $\{\Lambda_k\}_{k \geq 1}$ in [7]; see Theorem 1.2). Thus, the results on existence and sharp estimates of biorthogonal families established in [15] can be deduced from the corresponding results proved in [7]. Of course, Theorem 1.7 generalizes the results in [15] and [7] to complex sequences that do not satisfy the gap condition (5). \square

In [5] the authors prove the existence of a minimal time of controllability for some parabolic problems. This minimal time is related to the condensation index of the sequence of eigenvalues of the corresponding operator. In order to illustrate the existence of this minimal time, the authors consider the system

$$\begin{cases} \partial_t y - (D_1 \partial_{xx} + A_1) y = 0 & \text{in } (0, T) \times (0, \pi), \\ y(\cdot, 0) = Bv, \quad y(\cdot, \pi) = 0 & \text{on } (0, T), \\ y(0, \cdot) = y_0 & \text{in } (0, \pi), \end{cases} \quad (48)$$

where $B \in \mathbb{R}^2$, $v \in L^2(0, T)$ is the control,

$$D_1 := \text{diag}(1, d), \quad d > 0, \quad d \neq 1, \quad \text{and} \quad A_1 := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

The sequence of eigenvalues associated to the operator $L_1 = -(D_1 \partial_{xx} + A_1)$, with domain $D(L_1) = H^2(0, \pi; \mathbb{R}^2) \cap H_0^1(0, \pi; \mathbb{R}^2)$, is given by $\Lambda = \{k^2\}_{k \geq 1} \cup \{dk^2\}_{k \geq 1}$. Recall that the condition $\sqrt{d} \notin \mathbb{Q}$ is necessary for the approximate controllability of the system (48) at time $T > 0$. On the other hand, under this assumption, there exists a minimal time $T_0 = T_0(d) \in [0, \infty]$ such that the system is not null controllable at time T when $T < T_0$ (see [5] for the details). In our second example we will consider the sequence of eigenvalues associated to this system:

Proposition 2.13. *Let us consider $d \in (0, \infty)$ such that $\sqrt{d} \notin \mathbb{Q}$. Then, the sequence $\Lambda = \{k^2\}_{k \geq 1} \cup \{dk^2\}_{k \geq 1}$ can be rearranged as an increasing sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ satisfying $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ and condition (26) with $\beta = 0$, $p_0 = 1$,*

$$p_1 = p_2 = p = 1 + \frac{1}{\sqrt{d}}, \quad \alpha = 2, \quad q = 2, \quad \rho = \frac{5}{8} \frac{1}{p^2} \quad \text{and} \quad \nu = \frac{8}{3} \frac{1}{p^2}. \quad (49)$$

The proof of Proposition 2.13 can be found in A.3.

Let us now analyze a fourth example of sequence Λ which satisfies (26) and the general conditions appearing in Definition 1.4. With this example we cover the kind of sequences associated to some parabolic problems studied in [34]:

Proposition 2.14. *Let us consider two real positive sequences $\Lambda_1 = \{\lambda_k^{(1)}\}_{k \geq 1}$ and $\Lambda_2 = \{\lambda_k^{(2)}\}_{k \geq 1} = \{\lambda_k^{(1)} + \varepsilon_k\}_{k \geq 1}$ where $\{\varepsilon_k\}_{k \geq 1}$ is a real bounded sequence. Assume that Λ_1 satisfies $\Lambda_1 \in \mathcal{L}(0, \rho_1, 1, \pi_0, \pi_1, \pi_2, \alpha_1)$, for $\rho_1, \pi_0, \pi_1, \pi_2, \alpha_1 \in (0, \infty)$, and (26), for $\nu = \nu_1 \in (0, \infty)$. On the other hand, assume*

$$\lambda_k^{(2)} \neq \lambda_n^{(2)}, \quad \forall k, n \geq 1, \quad \text{with } k \neq n, \quad \text{and} \quad \lambda_k^{(1)} \neq \lambda_n^{(2)}, \quad \forall k, n \geq 1.$$

Let us take $\varepsilon_0 = \sup_{k \geq 1} |\varepsilon_k|$. Then, the sequence

$$\{\lambda_k^{(1)}\}_{k \geq 1} \cup \{\lambda_k^{(2)}\}_{k \geq 1}$$

can be rearranged as a positive increasing sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ satisfying $\Lambda \in \mathcal{L}(0, \rho, q, \pi_0, p_1, p_2, \alpha)$ and (26), with $\beta = 0$, $p_1 = 2\pi_1$, $p_2 = 2\pi_2$, $\alpha = \pi_2 \sqrt{\varepsilon_0} + 2\alpha_1$, $q = 2$ and ρ and ν positive constants only depending, resp., on ρ_1 and ε_0 and on ρ_1, ν_1 and ε_0 .

For the proof, see A.4.

Remark 2.15. Proposition 2.14 covers the sequence of eigenvalues of operator L in system (12) when $L = L_2$ (see (13)). We will use this proposition in Section 5. \square

Remark 2.16. Under assumptions of Proposition 2.14, observe that the sequence Λ satisfies the gap condition (5) if and only if

$$\liminf |\varepsilon_k| > 0.$$

On the other hand, analyzing the proof of Proposition 2.14, it is possible to provide some additional information about parameters ρ and α in Proposition 2.14

when the sequence $\{\varepsilon_k\}_{k \geq 1}$ satisfies appropriate properties. Indeed, when the bounded sequence $\{\varepsilon_k\}_{k \geq 1}$ is such that $\varepsilon_0 = \sup_{k \geq 1} |\varepsilon_k|$ satisfies

$$|\varepsilon_k| \leq \varepsilon_0 \leq \frac{\rho_1}{4}, \quad \forall k \geq 1,$$

then, the sequence Λ can be explicitly defined by (see (133)):

$$\Lambda_k = \begin{cases} \min \left\{ \lambda_\ell^{(1)}, \lambda_\ell^{(1)} + \varepsilon_\ell \right\}, & \text{if } k = 2\ell - 1, \\ \max \left\{ \lambda_\ell^{(1)}, \lambda_\ell^{(1)} + \varepsilon_\ell \right\}, & \text{if } k = 2\ell, \end{cases} \quad (50)$$

for any $k \geq 1$. In addition, from the proof of Proposition 2.14, we can deduce

$$\begin{cases} \Lambda_k - \Lambda_n \geq \frac{\rho_1}{16} (k^2 - n^2), & \forall k, n \in \mathbb{N} : k \geq n + 2, \\ \Lambda_k - \Lambda_n \leq \frac{\nu_1 + \varepsilon_0}{2} (k^2 - n^2), & \forall k, n \in \mathbb{N}, \end{cases}$$

i.e., we can take $\rho = \rho_1/16$ and $\nu = (\nu_1 + \varepsilon_0)/2$ in Proposition 2.14. We will use this observation in the proof of Proposition 5.6. \square

As said in Remark 1.6, let us finalize this section with an academic example of a positive sequence Λ in the class $\mathcal{L}(0, \rho, q, p_0, p_1, p_2, \alpha)$ with a parameter q which can be chosen as large as we want. With this example will see that the parameters p_1 and p_2 are increasing with respect to q . To this end, let us fix a positive integer $m \geq 2$ and define

$$\Lambda = \left\{ k^2 + \frac{\ell - 1}{m} : k \geq 1, \quad 1 \leq \ell \leq m \right\}. \quad (51)$$

It is clear that the set Λ can be written as an increasing positive sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ that satisfies the gap condition (5). Let us see that it also satisfies $\Lambda \in \mathcal{L}(0, \rho, q, p_0, p_1, p_2, \alpha)$, for appropriate parameters, and condition (26), for $\nu > 0$. One has:

Proposition 2.17. *Let us take a positive integer $m \geq 2$ and consider the sequence Λ defined in (51). Then,*

1. $\Lambda \in \mathcal{L}(0, \rho, q, p_0, p_1, p_2, \alpha)$, with $q = m$, $p_0 = 2$, $p_1 = p_2 = m$, $\alpha = m$ and

$$\rho = \frac{2}{(2m - 1)(2m + 1)}.$$

In fact, property (H5) does not hold if $q \leq m - 1$.

2. The sequence Λ satisfies (26) with

$$\nu = \frac{4m - 1}{m(2m + 1)}.$$

Proof. If $m \geq 2$, it is clear that the sequence Λ is an increasing sequence that satisfies items (H1)–(H4), with $\beta = 0$. Let us check (H5), (H6) and condition (26):

- (a) Let us prove item (H6) for the sequence Λ . To be precise, let us see

$$-m + m\sqrt{r} < \mathcal{N}(r) \leq m\sqrt{r}, \quad \forall r > 0, \quad (52)$$

where $\mathcal{N}(r)$ is defined in (21). First, if $r \in (0, 1)$, $\mathcal{N}(r) = 0$ and it is clear that (52) holds.

When $r \geq 1$ the function $\mathcal{N}(r)$ is given by

$$\mathcal{N}(r) = \sum_{\ell=1}^m \# \left\{ k : k^2 + \frac{\ell-1}{m} \leq r \right\} = \sum_{\ell=1}^m \left\lfloor \sqrt{r - \frac{\ell-1}{m}} \right\rfloor \leq \sum_{\ell=1}^m \lfloor \sqrt{r} \rfloor = m\sqrt{r}.$$

On the other hand, we can explicitly calculate $\mathcal{N}(r)$: Given $r \geq 1$, there exists an integer $k \geq 1$ such that $r \in [k^2, (k+1)^2)$. In this case,

$$\mathcal{N}(r) = \begin{cases} mk - m + \tilde{\ell}, & \text{if } r \in \left[k^2 + \frac{\tilde{\ell}-1}{m}, k^2 + \frac{\tilde{\ell}}{m} \right), \text{ with } 1 \leq \tilde{\ell} \leq m, \\ mk, & \text{if } r \in [k^2 + 1, (k+1)^2). \end{cases} \quad (53)$$

Indeed, if $r \in \left[k^2 + \frac{\tilde{\ell}-1}{m}, k^2 + \frac{\tilde{\ell}}{m} \right)$, with $\tilde{\ell} \in \mathbb{N} : \tilde{\ell} \leq m$, then, for any $\ell : 1 \leq \ell \leq \tilde{\ell}$,

$$r - \frac{\ell-1}{m} \in \left[k^2 + \frac{\tilde{\ell}-\ell}{m}, k^2 + \frac{\tilde{\ell}-\ell+1}{m} \right) \subset [k^2, (k+1)^2),$$

and $\left\lfloor \sqrt{r - \frac{\ell-1}{m}} \right\rfloor = k = \lfloor \sqrt{r} \rfloor$. Also, if $\ell : \tilde{\ell} + 1 \leq \ell \leq m$, one has

$$r - \frac{\ell-1}{m} \in \left[k^2 - \frac{\ell-\tilde{\ell}}{m}, k^2 - \frac{\ell-\tilde{\ell}-1}{m} \right) \subset [(k-1)^2, k^2),$$

and $\left\lfloor \sqrt{r - \frac{\ell-1}{m}} \right\rfloor = k - 1$. We deduce in this case

$$\mathcal{N}(r) = \sum_{\ell=1}^m \left\lfloor \sqrt{r - \frac{\ell-1}{m}} \right\rfloor = mk - m + \tilde{\ell} = m\lfloor \sqrt{r} \rfloor - m + \tilde{\ell},$$

and the first equality in (53).

Now, if $r \in [k^2 + 1, (k+1)^2)$, we can apply the same reasoning as before and deduce the second equality of (53).

Let us now prove the first inequality in (52) for $r \geq 1$ (the second one is a direct consequence of (53)). As before, assume $r \in [k^2, (k+1)^2)$ with $k \geq 1$ an appropriate integer. Thus, if $r \in \left[k^2 + \frac{\tilde{\ell}-1}{m}, k^2 + \frac{\tilde{\ell}}{m} \right)$, with $\tilde{\ell} \in \mathbb{N} : \tilde{\ell} \leq m$, then

$$\begin{aligned} \mathcal{N}(r) + m - m\sqrt{r} &= mk + \tilde{\ell} - m\sqrt{r} > mk - m\sqrt{k^2 + \frac{\tilde{\ell}}{m} + \tilde{\ell}} \\ &= \frac{(mk + \tilde{\ell})^2 - (m^2k^2 + m\tilde{\ell})}{mk + \tilde{\ell} + m\sqrt{k^2 + \frac{\tilde{\ell}}{m}}} = \frac{\tilde{\ell}^2 + m\tilde{\ell}(2k-1)}{mk + \tilde{\ell} + m\sqrt{k^2 + \frac{\tilde{\ell}}{m}}} > 0. \end{aligned}$$

Finally, if $r \in [k^2 + 1, (k+1)^2)$, we can write

$$\mathcal{N}(r) = mk > m(\sqrt{r} - 1).$$

This proves (52) and property (H6) for the sequence Λ with $p_0 = 2$, $p_1 = p_2 = m$ and $\alpha = m$.

(b) Let us now see that property (H5) holds for $q = m$ (and an appropriate parameter $\rho > 0$) and is not valid if $q < m$. To this end, let us first provide the expression of the terms of the sequence Λ . It is not difficult to see that, given an

integer $k \geq 1$, this can be written as $k = m\tilde{k} + \ell$, with $\tilde{k} \geq 0$ and $\ell \in \mathbb{N}$ with $1 \leq \ell \leq m$. Thus,

$$\Lambda_k = (\tilde{k} + 1)^2 + \frac{\ell - 1}{m}.$$

Negative part: Fix $q \in \mathbb{N}$, with $1 \leq q \leq m - 1$, and take $n = m\tilde{k} + 1$ and $k = m\tilde{k} + q + 1$, with $\tilde{k} \geq 0$, an arbitrary integer. It is clear that $k - n = q \leq m - 1$ and

$$\frac{\Lambda_k - \Lambda_n}{k^2 - n^2} = \frac{(\tilde{k} + 1)^2 + \frac{q}{m} - (\tilde{k} + 1)^2}{(m\tilde{k} + q + 1)^2 - (m\tilde{k} + 1)^2} = \frac{1}{m(2m\tilde{k} + 2 + q)} \rightarrow 0, \text{ when } \tilde{k} \rightarrow \infty.$$

We deduce that property (H5) is not valid when $q \leq m - 1$.

Positive part: Let us take $q = m$ and $n, k \geq 1$ with $k - n \geq q$. In this case,

$$k = m\tilde{k} + \ell_2, \quad n = m\tilde{n} + \ell_1, \quad \text{with } \tilde{n}, \ell_1, \tilde{k}, \ell_2 \in \mathbb{Z}, \quad 1 \leq \ell_1, \ell_2 \leq m \text{ and } \tilde{k}, \tilde{n} \geq 0.$$

Observe that, using $k - n \geq q = m$, we can conclude $\tilde{k} - \tilde{n} \geq 1$. So,

$$\begin{aligned} \frac{\Lambda_k - \Lambda_n}{k^2 - n^2} &= \frac{(\tilde{k} - \tilde{n})(\tilde{k} + \tilde{n} + 1) + (\tilde{k} - \tilde{n}) + \frac{\ell_2 - \ell_1}{m}}{(m\tilde{k} + \ell_2)^2 - (m\tilde{n} + \ell_1)^2} \\ &\geq \frac{(\tilde{k} - \tilde{n})(\tilde{k} + \tilde{n} + 1) + \tilde{k} - \tilde{n} - 1 + \frac{1}{m}}{(m\tilde{k} + m)^2 - (m\tilde{n} + 1)^2} \\ &> \frac{\tilde{k} - \tilde{n}}{m(\tilde{k} - \tilde{n}) + m - 1} \cdot \frac{\tilde{k} + \tilde{n} + 1}{m(\tilde{k} + \tilde{n} + 1) + 1} \geq \frac{1}{2m - 1} \cdot \frac{2}{2m + 1}. \end{aligned}$$

This shows property (H5) for the sequence Λ with $q = m$ and ρ given in the statement.

(c) In order to finish the proof of this result, let us show property (26). Again, let us take $k, n \in \mathbb{N}$ with $k > n$. As before, $k = m\tilde{k} + \ell_2$ and $n = m\tilde{n} + \ell_1$, with $\tilde{n}, \ell_1, \tilde{k}, \ell_2 \in \mathbb{Z}$, $1 \leq \ell_1, \ell_2 \leq m$, and $\tilde{k}, \tilde{n} \geq 0$ with $\tilde{k} \geq \tilde{n} \geq 0$.

Let us first analyze the case $\tilde{k} = \tilde{n} = \tilde{k} \geq 0$ and, of course, $1 \leq \ell_1 < \ell_2 \leq m$. We deduce,

$$\frac{\Lambda_k - \Lambda_n}{k^2 - n^2} = \frac{(\hat{k} + 1)^2 + \frac{\ell_2 - 1}{m} - (\hat{k} + 1)^2 - \frac{\ell_1 - 1}{m}}{(m\hat{k} + \ell_2)^2 - (m\hat{k} + \ell_1)^2} = \frac{1}{m} \frac{1}{2m\hat{k} + \ell_2 + \ell_1} \leq \frac{1}{3m}.$$

Now, if $\tilde{k} > \tilde{n}$ and $1 \leq \ell_1, \ell_2 \leq m$, one gets

$$\begin{aligned} \frac{\Lambda_k - \Lambda_n}{k^2 - n^2} &= \frac{(\tilde{k} - \tilde{n})(\tilde{k} + \tilde{n} + 2) + \frac{\ell_2 - \ell_1}{m}}{(m\tilde{k} + \ell_2)^2 - (m\tilde{n} + \ell_1)^2} \leq \frac{(\tilde{k} - \tilde{n})(\tilde{k} + \tilde{n} + 2) + 1 - \frac{1}{m}}{(m\tilde{k} + 1)^2 - (m\tilde{n} + m)^2} \\ &= \frac{\tilde{k} - \tilde{n}}{m(\tilde{k} - \tilde{n}) + 1 - m} \cdot \frac{\tilde{k} + \tilde{n} + 2}{m(\tilde{k} + \tilde{n} + 1) + 1} + \frac{1 - \frac{1}{m}}{(m\tilde{k} + 1)^2 - (m\tilde{n} + m)^2} \\ &\leq \frac{3}{2m + 1} + \frac{m - 1}{m(2m + 1)} = \frac{4m - 1}{m(2m + 1)}. \end{aligned}$$

Taking into account that $m \geq 2$, we can infer that

$$\frac{1}{3m} \leq \frac{4m-1}{m(2m+1)}$$

and, therefore, the sequence Λ fulfills inequality (26) with ν given in the statement. This ends the proof of the proposition. \square

Remark 2.18. It is interesting to point out that, thanks to Proposition 2.4, once property (H6) is proved for the sequence Λ with $p_0 = 2$, $p_1 = p_2 = m$ and $\alpha = m$, we can conclude that $\Lambda \in \mathcal{L}(0, \tilde{\rho}, \tilde{q}, 2, m, m, m)$ and (26) holds, with (see (36))

$$\tilde{q} = 3m, \quad \tilde{\rho} = \frac{1}{3m^2} \quad \text{and} \quad \tilde{\nu} = \frac{1}{3} \left(\frac{2+m}{m} \right)^2.$$

The parameters provided by Proposition 2.17 are better than the previous values. Indeed, taking into account that $m \geq 2$, it is clear that $q = m < \tilde{q} = 3m$,

$$\rho = \frac{2}{(2m-1)(2m+1)} > \tilde{\rho} = \frac{1}{3m^2} \quad \text{and} \quad \nu = \frac{4m-1}{m(2m+1)} < \tilde{\nu} = \frac{1}{3} \left(\frac{2+m}{m} \right)^2. \quad \square$$

Remark 2.19. We can apply Theorems 1.7 and 1.9 to the sequence Λ given by (51) and conclude the existence of a sequence $\{q_k\}_{k \geq 1} \subset L^2(0, T)$, biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$ (e_k is given in (1)), which satisfies (22) and (27). If we make use of Proposition 2.17, these two inequalities can be written under the form

$$\mathcal{A}_k^{(1)}(m) \mathcal{P}_k \leq \|q_k\|_{L^2(0, T)} \leq \mathcal{A}_k^{(2)}(m) \mathcal{P}_k, \quad \forall k \geq 3, \quad (54)$$

where $\mathcal{A}_k^{(1)}(m) := \mathcal{E}_k$ (see (29)) and

$$\mathcal{A}_k^{(2)}(m) := \mathcal{H}_1(\rho, q, p_1, p_2) \exp \left[C \left(1 + \mathcal{H}_2(\rho, q, p_1, p_2, T) \sqrt{|\Lambda_k|} + \frac{(1+p_2)^2}{T} \right) \right],$$

(see (24) and (25) in the real case) with ρ, q, p_1, p_2 and ν given in Proposition 2.17 (recall that the parameter m is the maximal cardinal of the condensation groupings of the sequence Λ , that is to say the maximal number of elements in Λ that do not satisfy (H5)).

Observe that, taking into account Remark 2.5, the elements of the sequence Λ satisfy

$$\Lambda_k = \frac{1}{m^2} k^2 + O(k), \quad \forall k \geq 1$$

and, therefore, one has

$$\lim_{m \rightarrow \infty} \mathcal{S}(m) = \infty \quad \text{where} \quad \mathcal{S}(m) = \sum_{k \geq 1} \frac{1}{\Lambda_k}, \quad \forall m \geq 2.$$

In some sense, the family of exponentials $\{e_k\}_{k \geq 1}$ (e_k given in (1)) “loses” its property of minimality in $L^2(0, T)$ when m tends to infinity. Thus, it is natural that the constants $\mathcal{A}_k^{(1)}(m)$ and $\mathcal{A}_k^{(2)}(m)$ in (54) satisfy

$$\lim_{m \rightarrow \infty} \mathcal{A}_k^{(1)}(m) = \lim_{m \rightarrow \infty} \mathcal{A}_k^{(2)}(m) = \infty, \quad \forall k \geq 1. \quad (55)$$

Let us see that (55) holds. To this end, we will analyze the asymptotic behavior of $\mathcal{A}_k^{(1)}(m)$ and $\mathcal{A}_k^{(2)}(m)$ when $m \rightarrow \infty$. In what follows, we will provide an explicit expression of these constants when $3 \leq k \leq m$.

1. Let us first analyze $\mathcal{A}_k^{(1)}(m)$. From the expression of \mathcal{E}_k for $3 \leq k \leq m$ (see (29)) and Proposition 2.17, we can write

$$\mathcal{A}_k^{(1)}(m) := \mathcal{E}_k = \frac{(m+k-2)!}{T^{m+k-2}} \left(\frac{2(m+k)-3}{2T} + 1 \right)^{1/2}.$$

Observe that Stirling's formula implies the existence of a positive constant $c_0 > 0$ such that

$$n! \geq c_0 \sqrt{2\pi n} \left(\frac{n}{e} \right)^n, \quad \forall n \in \mathbb{N}.$$

In particular, for a new positive constant c (independent of m), we deduce

$$\mathcal{A}_k^{(1)}(m) \geq c \sqrt{(m+k-2) \left[\frac{2(m+k)-3}{2T} + 1 \right]} \left(\frac{m+k-2}{eT} \right)^{m+k-2},$$

which is valid for any $m \geq 2$ and any $k : 3 \leq k \leq m$. One has the first equality in (55).

2. We continue with the analysis of $\mathcal{A}_k^{(2)}(m)$. Let us start with $\mathcal{H}_1(\rho, q, p_1, p_2)$ (see (24) in the real case). From Proposition 2.17, this constant only depends on m and has the expression:

$$\mathcal{H}_1(\rho, q, p_1, p_2) \equiv \mathcal{H}_1(m) = \left[\frac{(6m^2-1)(4m^2-1)}{4m^4} \right]^{2(m-1)}, \quad \forall m \geq 2.$$

It is not difficult to see that

$$\lim_{m \rightarrow \infty} \frac{\mathcal{H}_1(m)}{6^{2(m-1)}} = 1,$$

and, then

$$c_1 6^{2(m-1)} \leq \mathcal{H}_1(m) \leq c_2 6^{2(m-1)}, \quad \forall m \geq 2.$$

for two positive constants c_1 and c_2 , independent of m .

On the other hand, from the expression of $\mathcal{H}_2(\rho, q, p_1, p_2)$ (see (25) in the real case), we can write

$$\mathcal{H}_2(\rho, q, p_1, p_2, T) \equiv \mathcal{H}_2(m, T) = 4m^2 + 2m - 1 + \frac{1}{4m^2} + \sqrt{T}.$$

Observe that in our case $\alpha = m$. We can conclude that $\mathcal{A}_k^{(2)}(m)$ is given by

$$\mathcal{A}_k^{(2)}(m) := \mathcal{H}_1(m) \exp \left[C(m) \left(1 + \mathcal{H}_2(m, T) \sqrt{\Lambda_k} + \frac{(1+m)^2}{T} \right) \right], \quad m \geq 2,$$

with $C(m)$ a positive constant only depending on m and increasing with respect to m (see Theorem 1.7 with $\alpha = m$). Clearly, $\mathcal{A}_k^{(1)}(m)$ has an exponential behavior with respect to m and we can write

$$\mathcal{A}_k^{(2)}(m) \geq \exp \left[C \left(1 + m^2 \left(\sqrt{\Lambda_k} + \frac{1}{T} \right) + \sqrt{T \Lambda_k} \right) \right], \quad \forall m \geq 2, \quad \forall k : 3 \leq k \leq m.$$

We can conclude that $\mathcal{A}_k^{(2)}(m)$ has an exponential behavior with respect to m and satisfies the second equality in (55). \square

3. Proof of the first main result. This section is devoted to prove Theorem 1.7. To this end, we have to take into account the lack of the gap condition (5) for the sequence Λ and the fact that we cannot assume that T is small (see [5] and [34]). As we will see, this introduces new difficulties.

We will use the Fourier transform together with the Paley-Wiener Theorem:

Theorem 3.1. *Let f be an entire function such that*

$$|f(z)| \leq B e^{A|z|}, \quad \forall z \in \mathbb{C},$$

for two positive constants A, B , and $\|f\|_{L^2(\mathbb{R})} < \infty$. Then, there exists a function $\phi \in L^2(-A, A; \mathbb{C})$ such that

$$f(z) = \frac{1}{\sqrt{2\pi}} \int_{-A}^A \phi(t) e^{izt} dt.$$

Moreover, the Plancherel theorem gives

$$\|\phi\|_{L^2(-A, A; \mathbb{C})} = \|f\|_{L^2(\mathbb{R})}.$$

For the proof of Theorem 3.1 we refer to [44, Theorem 18. p. 101].

Remark 3.2. In what follows, C will denote a positive constant independent of T , $k \in \mathbb{N}$, ρ , q , p_1 and p_2 , which may change from one line to another (C may depend on $|\Lambda_1|$, β , p_0 and α , and is increasing with respect to α). In this work, the dependence of the constants with respect to the parameters ρ , q , p_1 and p_2 (see assumptions (H5) and (H6)) will be explicitly given. \square

Let us begin with a result of existence of entire functions satisfying appropriate properties. Our first main result will be a consequence of this theorem. One has:

Theorem 3.3. *Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ be a complex sequence with parameters satisfying (20). Then, for all $T > 0$, there exists a sequence of entire functions $\{G_k\}_{k \geq 1}$, with the following properties:*

1. For any $k \geq 1$ and $\varepsilon > 0$, there exists a positive constant $C'_{T, k, \varepsilon}$ such that

$$\left| e^{-iz\frac{T}{2}} G_k(z) \right| \leq C'_{T, k, \varepsilon} e^{(\frac{T}{2} + \varepsilon)|z|}, \quad \forall z \in \mathbb{C}; \quad (56)$$

2. $G_k(i\bar{\Lambda}_n) = \frac{1}{\sqrt{2\pi}} \delta_{kn}$, for all $k, n \geq 1$;
3. G_k belongs to $L^2(\mathbb{R})$, for any $k \geq 1$, and there exists a positive constant $C > 0$, only depending on $|\Lambda_1|$, β , p_0 and α (increasing with respect to α), such that

$$\|G_k\|_{L^2(\mathbb{R})} \leq \mathcal{H}_1 \exp \left[C \left(1 + \mathcal{H}_2 \sqrt{|\Lambda_k|} + \frac{(1 + p_2)^2}{T} \right) \right] \mathcal{P}_k, \quad (57)$$

for any $k \geq 1$, where \mathcal{P}_k , $\mathcal{H}_1 = \mathcal{H}_1(\rho, q, p_1, p_2)$ and $\mathcal{H}_2 = \mathcal{H}_2(\rho, q, p_1, p_2, T)$ are respectively given in (23), (24) and (25).

Theorem 1.7 is a direct consequence of Theorem 3.3. Therefore, before providing the proof of the technical result established in Theorem 3.3, we will complete the proof of Theorem 1.7.

Proof of Theorem 1.7. Let us consider $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ with parameters satisfying (20). On the other hand, let us fix $T > 0$. With the previous data, let us consider the function

$$F_k(z) := G_k(z) e^{-iz\frac{T}{2}}, \quad z \in \mathbb{C}, \quad k \geq 1,$$

where $\{G_k\}_{k \geq 1}$ is the sequence provided by Theorem 3.3. Let us see some properties of the function F_k . First, F_k is, for any $k \geq 1$, an entire function over \mathbb{C} . In fact, $F_k \in L^2(\mathbb{R})$ with $\|F_k\|_{L^2(\mathbb{R})} = \|G_k\|_{L^2(\mathbb{R})}$ for any $k \geq 1$.

Secondly, for any $\varepsilon > 0$ and $k \geq 1$, F_k is an entire function of exponential type $T/2 + \varepsilon$ (see (56)). So, we can apply Payley-Wiener Theorem (see Theorem 3.1) and deduce that there exists $\psi_k \in L^2(-T/2 - \varepsilon, T/2 + \varepsilon; \mathbb{C})$ such that

$$F_k(z) = e^{-iz\frac{T}{2}} G_k(z) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi_k(t) e^{izt} dt, \quad \forall z \in \mathbb{C}, \quad \forall k \geq 1.$$

Observe that the support of the function ψ_k is contained in $[-T/2 - \varepsilon, T/2 + \varepsilon]$, for any $k \geq 1$ and $\varepsilon > 0$. We conclude that, in fact, $\psi_k \in L^2(-T/2, T/2; \mathbb{C})$ and

$$F_k(z) = e^{-iz\frac{T}{2}} G_k(z) = \frac{1}{\sqrt{2\pi}} \int_{-\frac{T}{2}}^{\frac{T}{2}} \psi_k(t) e^{izt} dt, \quad \forall z \in \mathbb{C}, \quad \forall k \geq 1. \quad (58)$$

Let us now consider the function

$$q_k(t) := \psi_k \left(t - \frac{T}{2} \right), \quad t \in [0, T], \quad k \geq 1. \quad (59)$$

It is clear that q_k is well defined and $q_k \in L^2(0, T; \mathbb{C})$ for any $k \geq 1$. The objective now is to prove that the sequence $\{q_k\}_{k \geq 1} \subset L^2(0, T; \mathbb{C})$ satisfies Theorem 1.7. Let us first see that $\{q_k\}_{k \geq 1}$ is biorthogonal to $\{e^{-\Lambda_k t}\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$. Indeed, for any $k, n \geq 1$ and thanks to (58) and item 2 in Theorem 3.3, we can write,

$$\begin{aligned} \int_0^T q_k(t) e^{-\bar{\Lambda}_n t} dt &= \int_0^T \psi_k \left(t - \frac{T}{2} \right) e^{-\bar{\Lambda}_n t} dt = e^{-\bar{\Lambda}_n \frac{T}{2}} \int_{-\frac{T}{2}}^{\frac{T}{2}} \psi_k(t) e^{-\bar{\Lambda}_n t} dt \\ &= e^{-\bar{\Lambda}_n \frac{T}{2}} \sqrt{2\pi} e^{\bar{\Lambda}_n \frac{T}{2}} G(i\bar{\Lambda}_n) = \delta_{kn}. \end{aligned}$$

Let us now estimate $\|q_k\|_{L^2(0, T)}$. To this aim, we will use Plancherel Theorem and estimate (57). From the expression of q_k (see (59)), one has

$$\|q_k\|_{L^2(0, T; \mathbb{C})} = \|\psi_k\|_{L^2(-\frac{T}{2}, \frac{T}{2}; \mathbb{C})} = \|F_k\|_{L^2(\mathbb{R})} = \|G_k\|_{L^2(\mathbb{R})}.$$

Combining the previous inequality and inequality (57) we deduce (22). This completes the proof of Theorem 1.7. \square

Once Theorem 1.7 is proved, our next objective will be to show Theorem 3.3. The proof of this result is very technical. In order to make it clearer, we will divide it in two subsections:

1. In the first subsection (see Subsection 3.1) we will introduce an entire function $f_k(z)$ ($k \geq 1$) with simple zeros at Λ_n with $n \geq 1$ and $n \neq k$. To this end, we will use the natural infinite product that satisfies the condition $f_k(\Lambda_n) = 0$ for any $n \neq k$. We will show some properties of this function that, in particular, will imply item 2 in Theorem 3.3.
2. In the second subsection (see Subsection 3.2) we will introduce a ‘‘mollifier’’ function that we will use in the definition of the entire function G_k ($k \geq 1$) in Theorem 3.3. We will prove some properties of this function (which, in particular, will provide the property of item 3 in Theorem 3.3) and we will complete the proof of Theorem 3.3.

Remark 3.4. Let us remark that, if $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$, with parameters satisfying (20) (see Definition 1.4), then

$$\bar{\Lambda} := \{\bar{\Lambda}_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha).$$

We will prove Theorem 3.3 for the sequence $\overline{\Lambda}$ instead of Λ . \square

3.1. An infinite product. In this section we will consider again a complex sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ in $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ (the parameters satisfy (20)). Thus, for each $k \geq 1$ and $z \in \mathbb{C}$, we define

$$f_k(z) := \prod_{\substack{n \geq 1 \\ n \neq k}} \left(1 - \frac{z}{\Lambda_n}\right), \quad z \in \mathbb{C}. \quad (60)$$

The objective of this section is to prove some interesting properties satisfied by the function f_k .

First, observe that, by property (39), the previous product is uniformly convergent on compact sets of \mathbb{C} . Therefore, f_k is, for any $k \geq 1$, an entire function over \mathbb{C} (see for instance [25, p. 457]). Moreover, $f_k(\Lambda_n) = 0$, for any $n \neq k$. In fact, the zeros of f_k are exactly the elements of the sequence $\{\Lambda_n\}_{n \geq 1, n \neq k}$ and they are zeros of multiplicity 1.

We have the following property of function f_k :

Lemma 3.5 ([7]). *Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ be a complex sequence, with parameters satisfying (20). Then, for every $z \in \mathbb{C}$ and $k \geq 1$, we have*

$$\log |f_k(z)| \leq (p_2\pi + 1)\sqrt{|z|} + C, \quad (61)$$

where p_2 is given in assumption (H6) and C is a positive constant only depending on α and $|\Lambda_1|$ and increasing with respect to α . \square

The proof of this result can be found in [7].

Recall that our objective is to construct a sequence $\{G_k\}_{k \geq 1}$ of entire functions over \mathbb{C} satisfying items 1–3 in Theorem 3.3. This construction will use the function $f_k(z)$ and an estimate from below of the non-zero quantity $|f_k(\Lambda_k)|$. This is one of the key points of this work and is established in the next

Lemma 3.6. *Let $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ be a complex sequence, with parameters satisfying (20). Then,*

$$|f_k(\Lambda_k)| \geq \mathcal{H}_1(\rho, q, p_1, p_2)^{-1} e^{-C \mathcal{H}_3(\rho, q, p_1, p_2) \sqrt{|\Lambda_k|}} \mathcal{P}_k^{-1}, \quad \forall k \geq 1, \quad (62)$$

where C is a positive constant, only depending on $|\Lambda_1|$, β , p_0 and α (increasing with respect to α), $\mathcal{H}_1(\rho, q, p_1, p_2)$, f_k and \mathcal{P}_k are respectively given in (24), (60) and (23), and \mathcal{H}_3 is defined by

$$\begin{cases} \mathcal{H}_3(\rho, q, p_1, p_2) = 1 + q + \frac{1+q}{\rho^2 p_1^2} + p_2, \\ \mathcal{H}_3(\rho, q, p_1, p_2) = 1 + q + \frac{1}{\rho^2 p_1^2} + p_2, \quad \text{when } \Lambda \text{ is real.} \end{cases}$$

Proof. As said before, if $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$, then f_k (see (60)) is an entire function over \mathbb{C} with simple zeros at the points $\{\Lambda_n\}_{n \geq 1, n \neq k}$. Moreover, from assumption (H1), we have

$$|f_k(\Lambda_k)| = \prod_{\substack{n \geq 1 \\ n \neq k}} \left| \frac{\Lambda_n - \Lambda_k}{\Lambda_n} \right| \neq 0.$$

In order to obtain lower estimates of $|f_k(\Lambda_k)|$ let us decompose the set $\{n \geq 1 : n \neq k\}$ into the following sets:

$$\begin{cases} S_1(k) := \{n \geq 1 : 1 \leq |n - k| < q\}, \\ S_2(k) := \{n \geq 1 : |n - k| \geq q, |\Lambda_n| \leq 2|\Lambda_k|\}, \\ S_3(k) := \{n \geq 1 : |n - k| \geq q, |\Lambda_n| > 2|\Lambda_k|\}. \end{cases}$$

Then,

$$|f_k(\Lambda_k)| = \prod_{n \in S_1(k)} \left| 1 - \frac{\Lambda_k}{\Lambda_n} \right| \prod_{n \in S_2(k)} \left| 1 - \frac{\Lambda_k}{\Lambda_n} \right| \prod_{n \in S_3(k)} \left| 1 - \frac{\Lambda_k}{\Lambda_n} \right| := \prod_{i=1}^3 P_i^{(k)}. \quad (63)$$

Let us estimate each term in (63) and, to this aim, let us take $n \in S_1(k)$. In particular, $n < k + q$ and, from (H4) and (40) (or (43) in the real case), we deduce

$$\begin{aligned} |\Lambda_n| &\leq |\Lambda_{k+q}| \leq \frac{2}{p_1^2} |k+q|^2 + \frac{2C(1+q)^2}{\rho^2 p_1^4} \leq \frac{2}{p_1^2} \left| p_2 \sqrt{|\Lambda_k|} + \alpha + q \right|^2 + \frac{2C(1+q)^2}{\rho^2 p_1^4} \\ &\leq 4 \frac{p_2^2}{p_1^2} |\Lambda_k| + \frac{2C(1+q)^2}{p_1^2} + \frac{2C(1+q)^2}{\rho^2 p_1^4} := 4 \frac{p_2^2}{p_1^2} |\Lambda_k| + A, \quad \forall n \in S_1(k), \end{aligned}$$

(or

$$|\Lambda_n| \leq 4 \frac{p_2^2}{p_1^2} |\Lambda_k| + \frac{2C}{p_1^2} + \frac{2C}{\rho^2 p_1^4} := 4 \frac{p_2^2}{p_1^2} |\Lambda_k| + A, \quad \forall n \in S_1(k),$$

when Λ is a real sequence). In the previous inequalities, C is a positive constant independent of ρ , q , p_1 and p_2 .

If $x \geq |\Lambda_1|$, one has

$$\log \left(4 \frac{p_2^2}{p_1^2} x + A \right) = \log(x) + \log \left(4 \frac{p_2^2}{p_1^2} + \frac{A}{x} \right) \leq \sqrt{x} + \log \left(4 \frac{p_2^2}{p_1^2} + \frac{A}{|\Lambda_1|} \right).$$

On the other hand, thanks to (30) and (35), we also deduce

$$\begin{aligned} 4 \frac{p_2^2}{p_1^2} + \frac{A}{|\Lambda_1|} &= 4 \frac{p_2^2}{p_1^2} + \frac{C(1+q^2)}{|\Lambda_1|} \left(\frac{1}{p_1^2} + \frac{1}{\rho^2 p_1^4} \right) \\ &= \frac{4|\Lambda_1| \rho^2 p_1^2 p_2^2 + C(1+q^2)(\rho^2 p_1^2 + 1)}{|\Lambda_1| \rho^2 p_1^4} \leq \frac{C(1 + \rho p_2^2 + q^2)}{\rho^2 p_1^4}, \end{aligned}$$

(or $4 \frac{p_2^2}{p_1^2} + \frac{A}{|\Lambda_1|} \leq \frac{C(1 + \rho p_2^2)}{\rho^2 p_1^4}$ when Λ is real).

Thus,

$$\begin{aligned} P_1^{(k)} &= \prod_{n \in S_1(k)} \frac{|\Lambda_n - \Lambda_k|}{|\Lambda_n|} \geq \prod_{n \in S_1(k)} \frac{|\Lambda_n - \Lambda_k|}{\frac{4p_2^2}{p_1^2} |\Lambda_k| + A} = \frac{\mathcal{P}_k^{-1}}{\left(\frac{4p_2^2}{p_1^2} |\Lambda_k| + A \right)^{2q-2}} \\ &\geq \frac{e^{-(2q-2)\sqrt{|\Lambda_k|}}}{\left(\frac{4p_2^2}{p_1^2} + \frac{A}{|\Lambda_1|} \right)^{2q-2}} \mathcal{P}_k^{-1} \geq C \left(\frac{\rho^2 p_1^4}{1 + \rho p_2^2 + q^2} \right)^{2q-2} e^{-(2q-2)\sqrt{|\Lambda_k|}} \mathcal{P}_k^{-1}, \end{aligned} \quad (64)$$

where \mathcal{P}_k is given in (23) and C is a new a positive constant independent of ρ , q , p_1 and p_2 . In the real case, we deduce the following inequality for $P_1^{(k)}$:

$$P_1^{(k)} \geq C \left(\frac{\rho^2 p_1^4}{1 + \rho p_2^2} \right)^{2q-2} e^{-(2q-2)\sqrt{|\Lambda_k|}} \mathcal{P}_k^{-1}. \quad (65)$$

Let us now estimate the product $P_2^{(k)}$. At this point we will use the gap condition assumed in hypothesis (H5) when $|n - k| \geq q$. We will follow some ideas from [7]. Using again Lemma 2.6 (or inequality (43) in the real case), we deduce

$$\frac{k+n}{\sqrt{|\Lambda_k|}} \geq \frac{k}{\frac{k}{p_1} + \frac{C(1+q)}{\rho p_1^2}} = \frac{\rho p_1^2 k}{\rho p_1 k + C(1+q)} \geq \frac{\rho p_1^2}{\rho p_1 + C(1+q)} := B, \quad \forall n, k \geq 1. \quad (66)$$

(or

$$\frac{k+n}{\sqrt{|\Lambda_k|}} \geq \frac{k}{\frac{k}{p_1} + \frac{C}{\rho p_1^2}} = \frac{\rho p_1^2 k}{\rho p_1 k + C} \geq \frac{\rho p_1^2}{\rho p_1 + C} := B, \quad \forall n, k \geq 1. \quad (67)$$

when Λ is real). Then, if $n \in S_2(k)$,

$$P_2^{(k)} = \prod_{n \in S_2(k)} \left| \frac{\Lambda_n - \Lambda_k}{\Lambda_n} \right| \geq \prod_{n \in S_2(k)} \frac{\rho |k-n||k+n|}{2 |\Lambda_k|} \geq \prod_{n \in S_2(k)} \frac{B\rho |k-n|}{2 \sqrt{|\Lambda_k|}},$$

where B is given in (66) (or (67) in the real case).

Let $r_k := \#\{n \in S_2(k) : n \leq k - q\}$ and $s_k := \#\{n \in S_2(k) : n \geq q + k\}$. Then, from the previous estimate, one has

$$P_2^{(k)} \geq r_k! \left(\frac{B\rho}{2\sqrt{|\Lambda_k|}} \right)^{r_k} s_k! \left(\frac{B\rho}{2\sqrt{|\Lambda_k|}} \right)^{s_k} := P_{2,1}^{(k)} P_{2,2}^{(k)}.$$

Observe that Stirling's formula implies the existence of a positive constant $c_0 > 0$ such that

$$n! \geq c_0 \sqrt{2\pi n} \left(\frac{n}{e} \right)^n, \quad \forall n \in \mathbb{N}.$$

On the other hand, for $c_1 = e^{-1}$ one has

$$x \log x \geq -c_1, \quad \forall x \in (0, \infty).$$

Thus,

$$\begin{aligned} P_{2,1}^{(k)} &= r_k! \left(\frac{B\rho}{2\sqrt{|\Lambda_k|}} \right)^{r_k} \geq c_0 \left(\frac{B\rho r_k}{2e\sqrt{|\Lambda_k|}} \right)^{r_k} \\ &= c_0 \exp \left(\frac{2e\sqrt{\Lambda_k}}{B\rho} \frac{B\rho r_k}{2e\sqrt{|\Lambda_k|}} \log \left(\frac{B\rho r_k}{2e\sqrt{|\Lambda_k|}} \right) \right) \geq c_0 \exp \left(\frac{-2c_1 e}{B\rho} \sqrt{|\Lambda_k|} \right). \end{aligned}$$

Taking into account the expression of B (see (66), resp., (67) in the real case) and inequalities (30) and (35), we can conclude the existence of positive constant C_1 and C_2 (independent of ρ , q , p_1 , p_2 and T) such that

$$\frac{-C_1(1+q)}{\rho^2 p_1^2} \leq \frac{-1}{B\rho} \leq \frac{-C_2(1+q)}{\rho^2 p_1^2}, \quad (\text{resp.}, \frac{-C_1}{\rho^2 p_1^2} \leq \frac{-1}{B\rho} \leq \frac{-C_2}{\rho^2 p_1^2}, \text{ in the real case}).$$

Therefore, $P_{2,1}^{(k)} \geq C \exp \left(\frac{-C(1+q)}{\rho^2 p_1^2} \sqrt{|\Lambda_k|} \right)$, (resp., $P_{2,1}^{(k)} \geq C \exp \left(\frac{-C}{\rho^2 p_1^2} \sqrt{|\Lambda_k|} \right)$, in the real case).

A similar reasoning can be applied to $P_{2,2}^{(k)}$. Therefore, we have proved:

$$\begin{aligned} P_2^{(k)} &\geq C \exp \left(\frac{-C(1+q)}{\rho^2 p_1^2} \sqrt{|\Lambda_k|} \right), \\ (\text{resp.}, P_2^{(k)} &\geq C \exp \left(\frac{-C}{\rho^2 p_1^2} \sqrt{|\Lambda_k|} \right), \text{ in the real case}), \end{aligned} \quad (68)$$

for any $k \geq 1$. Again, C is a positive constant independent of ρ , q , p_1 , p_2 and T .

In order to finish, let us analyze the third product in (63). To this aim, we will use the inequalities

$$\log(1-x) \geq -2x, \quad \forall x \in \left[0, \frac{1}{2}\right], \quad \text{and} \quad \frac{|\Lambda_k|}{|\Lambda_n|} < \frac{1}{2}, \quad \forall n \in S_3(k).$$

From these inequalities and (H6), we can write $\mathcal{N}(r) \leq \alpha + p_2\sqrt{r}$, for any $r > 0$, and

$$\begin{aligned} \log P_3^{(k)} &\geq \sum_{n \in S_3(k)} \log \left(1 - \frac{|\Lambda_k|}{|\Lambda_n|}\right) \geq -2|\Lambda_k| \sum_{n \in S_3(k)} \frac{1}{|\Lambda_n|} \geq -2|\Lambda_k| \int_{2|\Lambda_k|}^{\infty} \frac{1}{r} d\mathcal{N}(r) \\ &= -2|\Lambda_k| \left(-\frac{\mathcal{N}(2|\Lambda_k|)}{2|\Lambda_k|} + \int_{2|\Lambda_k|}^{\infty} \frac{\mathcal{N}(r)}{r^2} dr \right) \geq -2|\Lambda_k| \int_{2|\Lambda_k|}^{\infty} \frac{\mathcal{N}(r)}{r^2} dr \\ &\geq -2|\Lambda_k| \int_{2|\Lambda_k|}^{\infty} \frac{\alpha + p_2\sqrt{r}}{r^2} dr = -2|\Lambda_k| \left(\frac{\alpha}{2|\Lambda_k|} + \frac{2p_2}{\sqrt{2|\Lambda_k|}} \right) \\ &= -\alpha - 2\sqrt{2}p_2\sqrt{|\Lambda_k|}. \end{aligned}$$

Coming back to (63) and putting together the previous inequality and inequalities (64) (or (65) in the real case) and (68), we conclude that inequality (62) holds. This ends the proof. \square

3.2. Additional properties and proof of Theorem 3.3. In this paragraph we will prove some additional properties that we will use in the proof of Theorem 3.3. To this end, we will introduce a ‘‘mollifier’’ function and we will construct the entire function G_k ($k \geq 1$) in Theorem 3.3 by means of this function and function f_k (see (60)). In order to construct this ‘‘mollifier’’ function, we follow the strategy of [41, 14].

Let us take $T > 0$ and a sequence $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$, with parameters satisfying (20) (see Definition 1.4). With all these data, we fix an integer $N \geq 2$ and we define the sequence $\{a_k\}_{k \geq 1} \subset (0, \infty)$ given by

$$a_k := \frac{C_{N,T}}{k^2}, \quad \text{where} \quad C_{N,T} := \frac{T}{2 \sum_{k \geq N} \frac{1}{k^2}}, \quad (69)$$

in order to have

$$\sum_{k \geq N} a_k = \frac{T}{2}.$$

Observe that this choice implies

$$\frac{1}{N} = \int_N^{\infty} \frac{1}{y^2} dy \leq \sum_{k \geq N} \frac{1}{k^2} \leq \int_{N-1}^{+\infty} \frac{1}{y^2} dy = \frac{1}{N-1},$$

and the estimate

$$\left(\frac{N-1}{2}\right)T \leq C_{N,T} \leq \frac{N}{2}T. \quad (70)$$

Consider now the function

$$P_{N,T}(z) := e^{iz\frac{T}{2}} \prod_{k \geq N} \cos(a_k z), \quad z \in \mathbb{C}.$$

With the previous data, one has:

Lemma 3.7. *Under the previous conditions, the following properties hold:*

1. *The function $P_{N,T}$ is entire over \mathbb{C} and satisfies*

$$\begin{cases} P_{N,T}(0) = 1, \\ |P_{N,T}(z)| \leq 1, \quad \forall z \in \mathbb{C} \text{ such that } \Im(z) \geq 0, \\ \left| e^{-iz\frac{T}{2}} P_{N,T}(z) \right| \leq e^{|z|\frac{T}{2}}, \quad \forall z \in \mathbb{C}. \end{cases}$$

2. *There exist positive constants $\theta_0 > 0$ and $\theta_1 > 0$ (independent of T and N) such that*

$$\begin{cases} \left(\frac{C_{N,T}}{\theta_0} \right)^{\frac{1}{2}} \sqrt{|x|} + 1 \geq N \implies \log |P_{N,T}(x)| \leq \frac{-\theta_1}{2^3} \left(\frac{C_{N,T}}{\theta_0} \right)^{\frac{1}{2}} \sqrt{|x|}, \\ \left(\frac{C_{N,T}}{\theta_0} \right)^{\frac{1}{2}} \sqrt{|x|} + 1 \leq N \implies \log |P_{N,T}(x)| \leq \frac{-\theta_1}{N^3} \left(\frac{C_{N,T}}{\theta_0} \right)^2 |x|^2. \end{cases} \quad (71)$$

3. *There exists a positive constant $\theta_2 > 0$ (independent of T and N) such that*

$$P_{N,T}(ix) \geq e^{-\theta_2 \sqrt{C_{N,T}x}}, \quad \forall x \geq 0. \quad (72)$$

For the proof of Lemma 3.7, see [41, 13, 14].

We are ready to prove the fundamental result stated in Theorem 3.3.

Proof of Theorem 3.3. Recall that $T > 0$ is given and $\Lambda = \{\Lambda_k\}_{k \geq 1}$ is a complex sequence in $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$. Let us define the function

$$G_k(z) := \frac{1}{\sqrt{2\pi}} \frac{f_k(-iz) P_{N,T}(z + \Im(\Lambda_k))}{f_k(\Lambda_k) P_{N,T}(i\Re(\Lambda_k))}; \quad (73)$$

(the function f_k is given in (60)). From the properties of the functions $P_{N,T}$ (see Lemma 3.7) and f_k we deduce that the function G_k is well defined and is an entire function over \mathbb{C} . In addition,

$$G_k(i\Lambda_n) = \frac{1}{\sqrt{2\pi}} \delta_{kn}, \quad \forall k, n \geq 1.$$

Observe that the function $P_{N,T}$ only has real zeros ($\{a_n\}_{n \geq 1}$ is a real sequence) and, then, the sequence $\{\Lambda_n\}_{n \geq 1, n \neq k}$ are zeros of G_k of multiplicity 1. This proves item 2 in Theorem 3.3.

Let us now see that $e^{-iz\frac{T}{2}} G_k$ satisfies inequality (56). From Lemmas 3.5 and 3.7, one has

$$\left| e^{-iz\frac{T}{2}} G_k(z) \right| \leq \frac{e^{(p_2\pi+1)\sqrt{|z|}+C} e^{|z+\Im(\Lambda_k)|\frac{T}{2}}}{\sqrt{2\pi} |f_k(\Lambda_k)| |P_{N,T}(i\Re(\Lambda_k))|} \leq \frac{e^{|\Im(\Lambda_k)|\frac{T}{2}+(p_2\pi+1)\sqrt{|z|}+C}}{\sqrt{2\pi} |f_k(\Lambda_k)| |P_{N,T}(i\Re(\Lambda_k))|} e^{|z|\frac{T}{2}},$$

for any $k \geq 1$. If we combine the previous inequality with

$$(p_2\pi + 1) \sqrt{|z|} \leq \frac{1}{4\varepsilon} (p_2\pi + 1)^2 + \varepsilon|z|,$$

valid for any $\varepsilon > 0$, we conclude that there exists a positive constant $C'_{N,T,k,\varepsilon}$ such that one has (56). This proves item 1 in Theorem 3.3.

Let us prove that G_k belongs to $L^2(\mathbb{R})$ and satisfies estimate (57). To this end, we will make the following choice of N :

$$2 + \gamma (p_2\pi + 1)^2 \frac{1}{T} \leq N \leq 4 + \gamma (p_2\pi + 1)^2 \frac{1}{T} \quad \text{with} \quad \gamma = \frac{2^7 \theta_0}{\theta_1^2}; \quad (74)$$

(p_2 is given in assumption (H6)).

Using (61) and (71), we have that for $|x|$ large enough one has

$$|G_k(x)| \leq \frac{e^{(p_2\pi+1)\sqrt{|x|+C} - \frac{\theta_1}{2^3} \left(\frac{C_{N,T}}{\theta_0}\right)^{\frac{1}{2}} \sqrt{|x+\Im(\Lambda_k)|}}}{\sqrt{2\pi} |f_k(\Lambda_k)| |P_{N,T}(i\Re(\Lambda_k))|}.$$

Observe that if

$$p_2\pi + 1 < \frac{\theta_1}{2^3} \left(\frac{C_{N,T}}{\theta_0}\right)^{\frac{1}{2}},$$

then $G_k \in L^2(\mathbb{R})$. In fact, thanks to assumption (74) the previous estimate is satisfied. Indeed, recall that a_k and $C_{N,T}$ are given in (69) and satisfy (70). So, from (74)

$$\frac{\theta_1}{2^3} \left(\frac{C_{N,T}}{\theta_0}\right)^{\frac{1}{2}} \geq \frac{\theta_1}{2^3} \left(\frac{(N-1)T}{2\theta_0}\right)^{\frac{1}{2}} > p_2\pi + 1.$$

This proves $G_k \in L^2(\mathbb{R})$.

In what follows, we will estimate $\|G_k\|_{L^2(\mathbb{R})}$. First, from the expression of G_k (see (73)) and using (61), (70), (72) and (74), one has

$$\begin{aligned} \|G_k\|_{L^2(\mathbb{R})}^2 &\leq \frac{e^{2\theta_2\sqrt{C_{N,T}\Re(\Lambda_k)+2C}}}{2\pi |f_k(\Lambda_k)|^2} \int_{-\infty}^{\infty} e^{2(p_2\pi+1)\sqrt{|x|}} |P_{N,T}(x + \Im(\Lambda_k))|^2 \\ &\leq \frac{e^{2\theta_2\sqrt{[2T+\gamma(p_2\pi+1)^2/2]\Re(\Lambda_k)+2C}}}{2\pi |f_k(\Lambda_k)|^2} \int_{-\infty}^{\infty} e^{2(p_2\pi+1)\sqrt{|x|}} |P_{N,T}(x + \Im(\Lambda_k))|^2 \\ &:= \frac{e^{2\theta_2\sqrt{[2T+\gamma(p_2\pi+1)^2/2]\Re(\Lambda_k)+2C}}}{2\pi |f_k(\Lambda_k)|^2} I. \end{aligned} \tag{75}$$

Denote

$$\mathcal{A}_1 := \{x \in \mathbb{R} : |x + \Im(\Lambda_k)| < X_{N,T}\}, \quad \mathcal{A}_2 := \{x \in \mathbb{R} : |x + \Im(\Lambda_k)| \geq X_{N,T}\},$$

where

$$X_{N,T} := \frac{\theta_0(N-1)^2}{C_{N,T}}.$$

Let us first observe that, thanks to inequalities (70) and (74), it is not difficult to see the property

$$\frac{1}{2}\theta_0 \left(\frac{1}{T} + \gamma \frac{(p_2\pi+1)^2}{T^2}\right) \leq X_{N,T} \leq 18\theta_0 \left(\frac{1}{T} + \gamma \frac{(p_2\pi+1)^2}{T^2}\right), \tag{76}$$

with γ given in (74).

With the previous notations, we can write

$$\begin{aligned} I &= \int_{\mathcal{A}_1} e^{2(p_2\pi+1)\sqrt{|x|}} |P_{N,T}(x + \Im(\Lambda_k))|^2 dx \\ &\quad + \int_{\mathcal{A}_2} e^{2(p_2\pi+1)\sqrt{|x|}} |P_{N,T}(x + \Im(\Lambda_k))|^2 dx := I_1 + I_2. \end{aligned}$$

The next objective is to provide an estimate of I_1 and I_2 . To this end, we will use property (71) of Lemma 3.7. Firstly, we estimate I_1 . We have:

$$\begin{aligned} I_1 &\leq e^{2(p_2\pi+1)\sqrt{|\Im(\Lambda_k)|}} \int_{\mathcal{A}_1} e^{2(p_2\pi+1)\sqrt{|x+\Im(\Lambda_k)|}} e^{\frac{-2\theta_1}{N^3} \frac{C_{N,T}}{\theta_0} |x+\Im(\Lambda_k)|^2} dx \\ &\leq e^{2(p_2\pi+1)(\sqrt{|\Im(\Lambda_k)|}+\sqrt{X_{N,T}})} \int_{\mathcal{A}_1} e^{\frac{-2\theta_1}{N^3} \frac{C_{N,T}}{\theta_0} |x+\Im(\Lambda_k)|^2} dx \\ &\leq e^{2(p_2\pi+1)(\sqrt{|\Im(\Lambda_k)|}+\sqrt{X_{N,T}})} |\mathcal{A}_1| = 2e^{2(p_2\pi+1)(\sqrt{|\Im(\Lambda_k)|}+\sqrt{X_{N,T}})} X_{N,T}. \end{aligned} \quad (77)$$

Let us now estimate I_2 . If we denote

$$L := \frac{\theta_1}{2^2} \sqrt{\frac{C_{N,T}}{\theta_0}} - 2(p_2\pi + 1),$$

and we use again (75), we get

$$\begin{aligned} I_2 &\leq e^{2(p_2\pi+1)\sqrt{|\Im(\Lambda_k)|}} \int_{\mathcal{A}_2} e^{2(p_2\pi+1)\sqrt{|x+\Im(\Lambda_k)|}} |P_{N,T}(x + \Im(\Lambda_k))|^2 dx, \\ &\leq e^{2(p_2\pi+1)\sqrt{|\Im(\Lambda_k)|}} \int_{\mathcal{A}_2} e^{-L\sqrt{|x+\Im(\Lambda_k)|}} dx \\ &\leq 2e^{2(p_2\pi+1)\sqrt{|\Im(\Lambda_k)|}} \int_0^\infty e^{-L\sqrt{x}} dx = 4e^{2(p_2\pi+1)\sqrt{|\Im(\Lambda_k)|}} \frac{1}{L^2}. \end{aligned} \quad (78)$$

As before and in order to bound L , we use again (70) and (74). Thus,

$$\begin{aligned} L &\geq \sqrt{\frac{\theta_1^2}{2^5\theta_0} (N-1)T - 2(p_2\pi+1)} \geq \sqrt{\frac{\theta_1^2 T}{2^5\theta_0} + 2^2(p_2\pi+1)^2 - 2(p_2\pi+1)} \\ &= \frac{\frac{\theta_1^2 T}{2^5\theta_0}}{\sqrt{\frac{\theta_1^2 T}{2^5\theta_0} + 2^2(p_2\pi+1)^2 + 2(p_2\pi+1)}} \geq \frac{\frac{\theta_1^2 T}{2^5\theta_0}}{2\sqrt{\frac{\theta_1^2 T}{2^5\theta_0} + 2^2(p_2\pi+1)^2}} > 0, \end{aligned}$$

and

$$\frac{1}{L^2} \leq \gamma \left(\frac{1}{T} + \gamma(p_2\pi+1)^2 \frac{1}{T^2} \right), \quad (79)$$

with γ given in (74).

Coming back to (75) and taking into account the inequality

$$x \leq e^{\sqrt{x}}, \quad \forall x \geq 0,$$

assumption (H3), (77) and (78), with $X_{N,T}$ and L satisfying (76) and (79), we deduce

$$\begin{aligned} \|G\|_{L^2(\mathbb{R})} &\leq \frac{e^{C(1+\sqrt{(1+T+p_2)\Re(\Lambda_k)})}}{|f_k(\Lambda_k)|} e^{C(1+p_2)(\sqrt{|\Im(\Lambda_k)|}+\sqrt{X_{N,T}})} \left(\frac{1}{T} + \frac{\gamma}{T^2} (p_2\pi+1)^2 \right) \\ &\leq \frac{e^{C(1+\sqrt{(1+T+p_2)\Re(\Lambda_k)})}}{|f_k(\Lambda_k)|} e^{C(1+p)(1+\Im(\Lambda_k)+\frac{1}{\sqrt{T}}+\frac{p_2+1}{T})} \\ &\leq \frac{e^{C(1+p_2+\sqrt{(1+T+p_2)|\Lambda_k|})}}{|f_k(\Lambda_k)|} e^{C(1+p_2)(\sqrt{|\Lambda_k|}+\frac{p_2+1}{T})}. \end{aligned}$$

Finally, the previous inequality and (62) provide estimate (57), for $G_k(x)$. This ends the proof. \square

4. A lower bound for the norm of arbitrary biorthogonal families: Proof of Theorem 1.9. This section will be devoted to prove the second main result, Theorem 1.9, of this paper. Observe that in inequality (27) two important factors appear. The first one, $e^{\frac{1}{T\nu}}$ ($\nu > 0$), is due to the fact we are dealing with infinite sequences. In order to obtain this factor, we will follow some ideas developed by Güichal in [23] (see also [14]). The second factor, \mathcal{P}_k (see (23)), comes from the eventual condensation of the elements of the sequence Λ . The corresponding part of the proof is completely new.

Let us consider a sequence $\Lambda = \{\Lambda_k\}_{k \geq 1} \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ (the parameters are given and satisfy (20)) such that property (26) holds, for $\nu > 0$. On the other hand, let us also consider $\{q_k\}_{k \geq 1} \subset L^2(0, T; \mathbb{C})$, a biorthogonal family to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$ (e_k is the function given by (1)).

Associated to the sequence Λ we introduce the spaces:

$$E(\Lambda, T) := \overline{\text{span}\{e_n : n \geq 1\}}^{L^2(0, T; \mathbb{C})},$$

$$E_k(\Lambda, T) := \overline{\text{span}\{e_n : n \geq 1, n \neq k\}}^{L^2(0, T; \mathbb{C})}, \quad \forall k \geq 1.$$

With this notation, one has:

Lemma 4.1. *Assume that $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset \mathbb{C}$ is a complex sequence satisfying (3) for a positive constant δ . Then, the closed space $E(\Lambda, T)$ is a proper subspace of $L^2(0, T; \mathbb{C})$. Moreover, the family of exponentials $\{e_k\}_{k \geq 1}$ is minimal in $L^2(0, T; \mathbb{C})$, that is to say, for every $k \geq 1$, one has*

$$e_k \notin E_k(\Lambda, T).$$

The previous lemma is a well-known result for sequences that satisfy (3) (see for instance [39], [37], [4], [5] and Remark 2.8).

As a consequence of Lemma 4.1, we can consider $d_{T,k} > 0$, the distance between the function e_k and $E_k(\Lambda, T)$, i.e.,

$$d_{T,k}^2 = \inf_{p \in E_k(\Lambda, T)} \|e_k - p\|_{L^2(0, T; \mathbb{C})}^2 = \int_0^T |e^{-\Lambda_k t} - p_k(t)|^2 dt, \quad \forall k \geq 1,$$

where $p_k \in E_k(\Lambda, T)$ is the orthogonal projection of the function $e_k(t) = e^{-\Lambda_k t}$ on $E_k(\Lambda, T)$. Observe that the function p_k is characterized by: $p_k \in E_k(\Lambda, T)$ and

$$(e_k - p_k, e_n)_{L^2(0, T; \mathbb{C})} = 0, \quad \forall n \geq 1 : n \neq k.$$

Thus, if we consider the function s_k given by

$$s_k(t) := \frac{e_k(t) - p_k(t)}{d_{T,k}^2} = \frac{e^{-\Lambda_k t} - p_k(t)}{d_{T,k}^2}, \quad t \in (0, T), \quad \forall k \geq 1,$$

we deduce that the sequence $\{s_k\}_{k \geq 1} \subset E(\Lambda, T)$ is biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$. In fact, this family is the unique sequence biorthogonal to the exponentials $\{e_k\}_{k \geq 1}$ whose elements belong to $E(\Lambda, T)$. Moreover, it is optimal in the following sense: if we consider another biorthogonal family $\{\tilde{q}_k\}_{k \geq 1}$ to $\{e_k\}_{k \geq 1}$ in $L^2(0, T; \mathbb{C})$, then $\tilde{q}_k - s_k \in E(\Lambda, T)^\perp$. Since $s_k \in E(\Lambda, T)$, we deduce

$$\|\tilde{q}_k\|_{L^2(0, T; \mathbb{C})}^2 = \|s_k\|_{L^2(0, T; \mathbb{C})}^2 + \|\tilde{q}_k - s_k\|_{L^2(0, T; \mathbb{C})}^2 \geq \|s_k\|_{L^2(0, T; \mathbb{C})}^2 = \frac{1}{d_{T,k}^2}, \quad \forall k \geq 1.$$

In particular,

$$\|q_k\|_{L^2(0, T; \mathbb{C})}^2 \geq \frac{1}{d_{T,k}^2}, \quad \forall k \geq 1.$$

The goal now is to obtain an upper bound of $d_{T,k}$, for any $k \geq 1$. From the definition of $d_{T,k}$ we clearly have

$$d_{T,k} \leq \|e_k - p\|_{L^2(0,T;\mathbb{C})}, \quad \forall p \in E_k(\Lambda, T), \quad \forall k \geq 1.$$

Then,

$$\|q_k\|_{L^2(0,T;\mathbb{C})} \geq \frac{1}{d_{T,k}} \geq \frac{1}{\|e_k - p\|_{L^2(0,T;\mathbb{C})}}, \quad \forall p \in E_k(\Lambda, T), \quad \forall k \geq 1. \quad (80)$$

In order to obtain (27), we are going to apply the previous inequality to two appropriate functions $p \in E_k(\Lambda, T)$. Inequality (27) will be a direct consequence of inequality (80), written for these two functions.

4.1. A lower bound for the norm of arbitrary biorthogonal families. First part. Let us prove that, for any $k \geq 3$, one has

$$\|q_k\|_{L^2(0,T;\mathbb{C})} \geq \frac{6}{\pi^2} \mathcal{B}_k \mathcal{P}_k e^{\frac{1}{T\nu}}, \quad (81)$$

where \mathcal{P}_k and \mathcal{B}_k are respectively given in (23) and (28).

Following [23], the idea is to construct a particular function p in $E_k(\Lambda, T)$. To this end, let us fix a positive integer $M \geq q+k$, where q is given in assumption (H5). On the other hand, let us take

$$f_1(t) := \sum_{n=1}^{M+1} A_n e_n(t) = \sum_{n=1}^{M+1} A_n e^{-\Lambda_n t}, \quad t \in (0, T), \quad (82)$$

with coefficients $A_1, A_2, \dots, A_{M+1} \in \mathbb{C}$. Observe that $f_1 \in E_k(\Lambda, T)$ if and only if $A_k = 0$ and, when $A_k \neq 0$ then

$$\frac{1}{A_k} f_1(t) = e^{-\Lambda_k t} + \sum_{n=1}^{k-1} \frac{A_n}{A_k} e^{-\Lambda_n t} + \sum_{n=k+1}^{M+1} \frac{A_n}{A_k} e^{-\Lambda_n t} = e_k(t) - p(t), \quad t \in (0, T).$$

Therefore,

$$d_{T,k} \leq \left\| \frac{1}{A_k} f_1 \right\|_{L^2(0,T;\mathbb{C})}, \quad \forall k \geq 1. \quad (83)$$

Let us consider the coefficients $A_1, A_2, \dots, A_{M+1} \in \mathbb{C}$ given by

$$A_n := \prod_{\substack{i=1 \\ i \neq n}}^{M+1} (\Lambda_i - \Lambda_n)^{-1}, \quad 1 \leq n \leq M+1. \quad (84)$$

The next task will be to estimate $\|f_1\|_{L^2(0,T;\mathbb{C})}$, with f_1 and A_n , $1 \leq n \leq M+1$, respectively given in (82) and (84). To this aim, we recall the following results:

Lemma 4.2. *Let $B := \{a_n\}_{1 \leq n \leq r+1} \subset \mathbb{C}$ be a set of distinct points, $r \geq 1$, and let us fix g an analytic function in a convex domain $\Omega \subset \mathbb{C}$ such that $B \subset \Omega$. Then, there exists $\theta \in [-1, 1]$ and $\xi \in \text{Conv}(B)$, the convex hull of B , such that*

$$\sum_{n=1}^{r+1} \frac{g(a_n)}{\prod_{\substack{a_i \in B \\ i \neq n}} (a_n - a_i)} = \frac{\theta}{r!} \frac{\partial^r g}{\partial z^r}(\xi).$$

Lemma 4.3. *The following properties hold:*

1. $\int_0^T t^N e^{-\lambda t} dt \leq \frac{2T^{N+1}}{N+1+\lambda T}$, for any $N \geq 1$ and $\lambda > 0$.

$$2. \frac{1}{N!} \left(\frac{x}{1+x} \right)^N e^x \leq \sum_{n=N}^{\infty} \frac{x^n}{n!}, \text{ for any } x \geq 0 \text{ and } N \geq 1.$$

Lemma 4.2 is a formula due to Jensen. On the other hand, the proof of Lemma 4.3 can be found in [14, Lemma 4.2, Lemma 4.3].

Now, using assumption (26), we can provide an estimate of $|A_k|^{-1}$. One has:

Lemma 4.4. *Let us fix $k \geq 1$ and $M \geq q + k$. Then, under the assumptions of Theorem 1.9, we have*

$$|A_k|^{-1} \leq \begin{cases} \frac{\nu^{M+2-q-k}}{(q-1)!(2k+q-1)!} (M+1-k)! (M+1+k)! \mathcal{P}_k^{-1}, & \text{if } k \leq q, \\ \frac{\nu^{M-2(q-1)}(2k-q)!}{k[(q-1)!]^2(2k+q-1)!} (M+1-k)! (M+1+k)! \mathcal{P}_k^{-1}, & \text{if } k > q, \end{cases} \quad (85)$$

where A_k and \mathcal{P}_k are respectively given in (84) and (23).

Proof. The proof is a direct consequence of assumption (26). Indeed, let us first assume that $k > q$. From the expression of A_k (see (84)), we obtain,

$$|A_k|^{-1} = \prod_{\substack{n=1 \\ n \neq k}}^{M+1} |\Lambda_k - \Lambda_n| = \mathcal{P}_k^{-1} \prod_{n=1}^{k-q} |\Lambda_k - \Lambda_n| \prod_{n=q+k}^{M+1} |\Lambda_k - \Lambda_n| := \mathcal{P}_k^{-1} S_{1,k} S_{2,k},$$

where \mathcal{P}_k is given in (23). On the other hand, assumption (26) provides the following estimate

$$\begin{aligned} S_{1,k} S_{2,k} &\leq \prod_{n=1}^{k-q} (\nu |k^2 - n^2|) \prod_{n=q+k}^{M+1} (\nu |k^2 - n^2|) \\ &= \nu^{M-2(q-1)} \prod_{n=1}^{k-q} (k+n) \prod_{n=1}^{k-q} (k-n) \prod_{n=q+k}^{M+1} (k+n) \prod_{n=q+k}^{M+1} (n-k) \\ &= \nu^{M-2(q-1)} \frac{(2k-q)!}{k!} \frac{(k-1)!}{(q-1)!} \frac{(k-1)!}{(q-1)!} \frac{(M+1+k)!}{(2k+q-1)!} \frac{(M+1-k)!}{(q-1)!} \\ &= \nu^{M-2(q-1)} \frac{(2k-q)!}{k[(q-1)!]^2(2k+q-1)!} (M+1-k)! (M+1+k)!. \end{aligned}$$

Putting both inequalities together we deduce (85) in the case $k > q$.

We can reason as before in the case $k \leq q$. In this case, the first product $S_{1,k}$ in the expression of $|A_k|^{-1}$ does not appear. It is not difficult to deduce the following estimate:

$$S_{2,k} \leq \prod_{n=q+k}^{M+1} (\nu |k^2 - n^2|) = \frac{\nu^{M+2-q-k}}{(q-1)!(2k+q-1)!} (M+1-k)! (M+1+k)!.$$

The previous inequality implies (85) for $k \leq q$. This ends the proof. \square

Let us continue with the proof of inequality (81) when $k \geq 3$. Observe that we can apply Lemma 4.2 to f_1 with coefficients A_n given in (84), $r = M$, $B =$

$\{\Lambda_n\}_{1 \leq n \leq M+1}$ and $g(z) = e^{-tz}$ ($t \in [0, T]$ is fixed). We obtain,

$$f_1(t) = \sum_{n=1}^{M+1} \frac{(-1)^M}{M+1} e^{-\Lambda_n t} = (-1)^M \frac{\theta}{M!} \frac{\partial^M g}{\partial z^M}(\xi) = \frac{\theta t^M}{M!} e^{-t\xi},$$

where $\theta = \theta(t)$ satisfies $|\theta| \leq 1$ and

$$\xi = \sum_{n=1}^{M+1} \alpha_n \Lambda_n, \quad \text{with } \alpha_n \geq 0 \quad \text{and} \quad \sum_{n=1}^{M+1} \alpha_n = 1.$$

Using assumption (H4) and property (3), we can write:

$$\Re(\xi) = \sum_{n=1}^{M+1} \alpha_n \Re(\Lambda_n) \geq \delta \sum_{n=1}^{M+1} \alpha_n |\Lambda_n| \geq \delta |\Lambda_1|,$$

where $\delta > 0$ is a constant only depending on β ($\delta = 1$ when $\beta = 0$). Thus,

$$|f_1(t)| = \left| \frac{\theta t^M e^{-t\xi}}{M!} \right| \leq \frac{t^M}{M!} e^{-t\Re(\xi)} \leq \frac{t^M}{M!} e^{-\delta |\Lambda_1| t}, \quad \forall t \in [0, T].$$

Coming back to (83) with A_k given in (84), we deduce that

$$d_{T,k} \leq \frac{1}{|A_k|} \|f_1\|_{L^2(0,T;\mathbb{C})} \leq \frac{1}{M!} |A_k|^{-1} \left(\int_0^T t^{2M} e^{-2\delta |\Lambda_1| t} dt \right)^{1/2}, \quad \forall k \geq 1.$$

Let us introduce the quantity

$$\mathcal{D}_k = \begin{cases} \nu^{k+q-2} (q-1)! (2k+q-1)! \sqrt{\delta |\Lambda_1| + \frac{1}{2T}} \mathcal{P}_k, & \text{if } k \leq q, \\ \nu^{2(q-1)} [(q-1)!]^2 \frac{k(2k+q-1)!}{(2k-q)!} \sqrt{\delta |\Lambda_1| + \frac{1}{2T}} \mathcal{P}_k, & \text{if } k \geq q. \end{cases} \quad (86)$$

Let us first work with $k \geq \max\{3, q\}$. If we use Lemma 4.4 and item 1 of Lemma 4.3, we deduce

$$\begin{aligned} d_{T,k} &\leq \frac{\nu^{M-2(q-1)} (2k-q)! (M+k+1)! (M-k+1)!}{M!} \frac{T^M \sqrt{2T}}{\sqrt{2M+1+2\delta T |\Lambda_1|}} \mathcal{P}_k^{-1} \\ &\leq \nu^{-2(q-1)} \sqrt{\frac{2T}{1+2\delta T |\Lambda_1|}} \frac{(2k-q)! \mathcal{P}_k^{-1} (M-k+1)!}{k [(q-1)!]^2 (2k+q-1)! M!} (M+k+1)! (\nu T)^M \\ &= \mathcal{D}_k^{-1} \frac{(M+k+1)!}{M(M-1) \cdots (M-k+4)(M-k+3)(M-k+2)} (\nu T)^M \\ &\leq \mathcal{D}_k^{-1} \frac{1}{(k+q)(k+q-1) \cdots (q+4)} \frac{(M+k+1)!}{(M+1-k-q)^2} (\nu T)^M \\ &= \mathcal{D}_k^{-1} \frac{(q+3)!}{(k+q)!} \frac{(M+k+1)!}{(M+1-k-q)^2} (\nu T)^M, \quad \forall k \geq \max\{3, q\}, \end{aligned}$$

where \mathcal{D}_k is given in (86). In the previous inequalities we have used that $k \geq 3$ and $M \geq k+q$.

Now, if k is such that $3 \leq k \leq q$, we can argue as before and deduce the same inequality. Summarizing, for any $k \geq 3$, one has

$$d_{T,k} \leq \mathcal{D}_k^{-1} \frac{(q+3)!}{(k+q)!} \frac{(M+k+1)!}{(M+1-k-q)^2} (\nu T)^M, \quad \forall k \geq 3,$$

where \mathcal{D}_k is given in (86).

Let us finalize the proof of inequality (81) when $k \geq 3$. The previous estimate of $d_{T,k}$ and item 2 of Lemma 4.3 allow us to write:

$$\begin{aligned} \frac{\pi^2}{6} \frac{1}{d_{T,k}} &= \sum_{M=k+q}^{\infty} \frac{1}{(M+1-k-q)^2} \frac{1}{d_{T,k}} \geq \mathcal{D}_k \frac{(k+q)!}{(q+3)!} \sum_{M=k+q}^{\infty} \frac{1}{(M+k+1)!} \frac{1}{(\nu T)^M} \\ &= (\nu T)^{k+1} \mathcal{D}_k \frac{(k+q)!}{(q+3)!} \sum_{n=2k+q+1}^{\infty} \frac{1}{n!} \frac{1}{(\nu T)^n} \\ &\geq (\nu T)^{k+1} \mathcal{D}_k \frac{(k+q)!}{(q+3)!} \frac{1}{(2k+q+1)!} \frac{1}{(1+\nu T)^{2k+q+1}} e^{\frac{1}{\nu T}}, \end{aligned}$$

where \mathcal{D}_k is given in (86). Coming back to (80), the previous inequality proves (81).

4.2. A lower bound for the norm of arbitrary biorthogonal families. Second part. In order to finish the proof of Theorem 1.9, let us now show that, for any $k \geq 1$, one has

$$\|q_k\|_{L^2(0,T;\mathbb{C})} \geq \mathcal{E}_k \mathcal{P}_k, \quad (87)$$

where \mathcal{E}_k and \mathcal{P}_k are respectively given in (29) and (23).

Let us introduce the function

$$f_2(t) = \sum_{\{n \geq 1: |k-n| < q\}} \tilde{A}_n e_n(t) = \sum_{\{n \geq 1: |k-n| < q\}} \tilde{A}_n e^{-\Lambda_n t}, \quad t \in (0, T), \quad (88)$$

with coefficients $\tilde{A}_n \in \mathbb{C}$ given by

$$\tilde{A}_n := \prod_{\substack{\{i \geq 1: |k-i| < q\} \\ i \neq n}} (\Lambda_i - \Lambda_n)^{-1}, \quad n \geq 1: |k-n| < q. \quad (89)$$

Observe that $|\tilde{A}_k| = \mathcal{P}_k \neq 0$ (\mathcal{P}_k is given in (23)). As in the previous subsection, we can write

$$\frac{1}{\tilde{A}_k} f_2(t) = e^{-\Lambda_k t} + \sum_{\{n \geq 1: 0 < |k-n| < q\}} \frac{\tilde{A}_n}{\tilde{A}_k} e^{-\Lambda_n t} = e_k(t) - \tilde{p}(t), \quad t \in (0, T).$$

Therefore,

$$d_{T,k} \leq \|e_k - \tilde{p}\|_{L^2(0,T;\mathbb{C})} = \mathcal{P}_k^{-1} \|f_2\|_{L^2(0,T;\mathbb{C})}, \quad \forall k \geq 1. \quad (90)$$

Given $k \geq 1$, we consider the set

$$B = \{\Lambda_n : |k-n| < q\}.$$

and the number $r+1 = \#B$. It is not difficult to see that

$$r = \begin{cases} k+q-2, & \text{if } 1 \leq k < q, \\ 2(q-1), & \text{if } k \geq q, \end{cases}$$

and, therefore $r \geq 1$ ($q \geq 2$). Now, if we apply Lemma 4.2 to f_2 with coefficients \tilde{A}_n given by (89), the set B , r and $g(z) = e^{-tz}$ ($t \in [0, T]$ is fixed), we deduce

$$f_2(t) = \frac{\tilde{\theta}}{r!} t^r e^{-t\tilde{\xi}},$$

where $\tilde{\theta} = \tilde{\theta}(t)$ is such that $|\tilde{\theta}| \leq 1$ and $\tilde{\xi} \in \text{Conv}(B)$, i.e.,

$$\tilde{\xi} = \sum_{\{n \geq 1: |k-n| < q\}} \tilde{\alpha}_n \Lambda_n \quad \text{with } \tilde{\alpha}_n \geq 0 \quad \text{and} \quad \sum_{\{n \geq 1: |k-n| < q\}} \tilde{\alpha}_n = 1.$$

The previous expression of $\tilde{\xi}$ and assumption (H4) also allow us to deduce

$$\begin{aligned} \Re(\tilde{\xi}) &= \sum_{\{n \geq 1: |k-n| < q\}} \tilde{\alpha}_n \Re(\Lambda_n) \geq \delta \sum_{\{n \geq 1: |k-n| < q\}} \tilde{\alpha}_n |\Lambda_n| \geq \delta \min_{\Lambda_n \in B} |\Lambda_n| \\ &= \delta |\Lambda_{\max\{1, k+1-q\}}|, \end{aligned}$$

with $\delta > 0$ as in (3) ($\delta = 1$ when $\beta = 0$). Summarizing, we have proved

$$|f_2(t)| \leq \frac{1}{r!} |t|^r e^{-\delta |\Lambda_{k+1-q}| t}, \quad \forall t \in [0, T].$$

Let us finalize the proof of (87). To this end, we work with the previous inequality, inequality (90), item 1 of Lemma 4.3 and the expression of r . Thus, if $1 \leq k < q$, we obtain

$$\begin{aligned} d_{T,k} &\leq \frac{1}{(k+q-2)!} \left(\int_0^T |t|^{2(k+q-2)} e^{-2\delta |\Lambda_1| t} dt \right)^{1/2} \mathcal{P}_k^{-1} \\ &\leq \frac{1}{(k+q-2)!} \frac{T^{k+q-2} \sqrt{2T}}{\sqrt{2(k+q-2) + 1 + 2\delta |\Lambda_1| T}} \mathcal{P}_k^{-1}. \end{aligned}$$

Now, if $k \geq q$, $r = 2(q-1)$ and a similar computation provides

$$d_{T,k} \leq \frac{1}{(2q-2)!} \frac{T^{2(q-1)} \sqrt{2T}}{\sqrt{4(q-1) + 1 + 2\delta |\Lambda_{k+1-q}| T}} \mathcal{P}_k^{-1}.$$

Of course, inequality (87) is a direct consequence of these inequalities and inequality (80). This finally ends the proof of Theorem 1.9. \square

5. Application to the boundary controllability problem for some parabolic systems. This section will be devoted to apply Theorems 1.7 and 1.9 to two particular parabolic systems in order to provide some new results on the control cost for the boundary controllability problem associated to these systems. To be precise, we will revisit the controllability problems analyzed in [34] and [22] and we will prove new estimates of the control cost with respect to the final time $T > 0$. Some results in this section have been previously announced in [21].

5.1. A 2×2 linear coupled parabolic system. Let us consider the one-dimensional Dirichlet-Laplace operator $\tilde{L}_1 := -\partial_{xx}$ with domain $D(\tilde{L}_1) = H^2(0, \pi) \cap H_0^1(0, \pi)$. It is well-known that $(\tilde{L}_1, D(\tilde{L}_1))$ is self-adjoint and admits a sequence of eigenvalues $\Lambda_1 = \{\lambda_k^{(1)}\}_{k \geq 1} = \{k^2\}_{k \geq 1}$ and normalized eigenfunctions given by

$$\varphi_k^{(1)}(x) := \sqrt{\frac{2}{\pi}} \sin(kx), \quad \forall k \geq 1, \quad x \in (0, \pi).$$

On the other hand, let $Q \in L^2(0, \pi)$ be a given function and consider the operator $\tilde{L}_2 := -\partial_{xx} + Q$ with domain $D(\tilde{L}_2) = D(\tilde{L}_1)$. Again, $(\tilde{L}_2, D(\tilde{L}_2))$ is self-adjoint, admits a sequence of increasing eigenvalues $\Lambda_2 = \{\lambda_k^{(2)}\}_{k \geq 1}$ and a sequence of normalized eigenfunctions $\{\varphi_k^{(2)}\}_{k \geq 1}$ which is an orthonormal basis of $L^2(0, \pi)$.

In this section we will revisit the boundary controllability problem for the system

$$\begin{cases} \partial_t y + L_2 y = 0 & \text{in } Q_T := (0, T) \times (0, \pi), \\ y(\cdot, 0) = Bv, \quad y(\cdot, \pi) = 0 & \text{on } (0, T), \\ y(0, \cdot) = y_0 & \text{in } (0, \pi), \end{cases} \quad (91)$$

where $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$ is the initial datum, $v \in L^2(0, T)$ is a scalar control, the operator $(L_2, D(L_2))$ is given by (13) and $B \in \mathbb{R}^2$ is the control vector. It is interesting to observe that we want to control system (91), two variables, with a unique control function $v \in L^2(0, T)$.

For every $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$, system (91) admits a unique solution defined by transposition, y , which satisfies $y \in L^2(Q_T; \mathbb{R}^2) \cap C^0([0, T]; H^{-1}(0, \pi; \mathbb{R}^2))$.

It is well-known that, when $Q \in L^2(0, \pi)$ satisfies (14), one has

$$\lambda_k^{(2)} = \lambda_k^{(1)} + \varepsilon_k = k^2 + \varepsilon_k, \quad \forall k \geq 1,$$

with $\{\varepsilon_k\}_{k \geq 1} \in \ell^2$. In particular, $\lim \varepsilon_k = 0$ (see for instance [26]). Observe that in this case, the eigenvalues of the operator L_2 do not fulfill the gap condition (5) and the null controllability of system (91) has a minimal time T_0 of null controllability which is defined as:

$$T_0 = \limsup \frac{-\log |\varepsilon_k|}{k^2} \in [0, \infty]. \quad (92)$$

To be precise, one has:

Theorem 5.1. *Let us consider $Q \in L^2(0, \pi)$, a function satisfying $Q \not\equiv 0$ and (14). Given $T > 0$ and $B = (b_1, b_2)^t$, one has*

1. *System (91) is approximately controllable at time $T > 0$ if and only if*

$$b_1 b_2 \neq 0 \quad \text{and} \quad \lambda_k^{(1)} \neq \lambda_n^{(2)} \quad \forall k, n \geq 1. \quad (93)$$

2. *Assume that (93) holds and consider T_0 given in (92). Then*

(a) *If $T > T_0$, system (91) is null controllable at time T .*

(b) *If $T < T_0$, system (91) is not null controllable at time T .* \square

The previous result has been proved in [34]. In this reference, the author also shows that T_0 depends on $Q \in L^2(0, \pi)$ and satisfies this property: given $\tau \in [0, \infty]$, there exists $Q \in L^2(0, \pi)$ satisfying (14) such that $T_0 = \tau$. Thus, the minimal time T_0 associated to system (91) could reach any value in the interval $[0, \infty]$. Therefore, there exist coefficients $Q \in L^2(0, \pi)$ such that the corresponding minimal time of system (91) satisfies $T_0 > 0$.

Remark 5.2. The study of the controllability of system (91) is easier when $Q \in L^2(0, \pi)$ does not satisfy condition (14). In fact, we have the following property: system (91) is null controllable at time $T > 0$ if and only if the system is approximately controllable at this time, i.e., if and only if (93) holds. In this case, we have that $T_0 = 0$ and the null controllability of the system is valid for any $T > 0$ (see [34]). On the other hand, it is not difficult to check that we can apply Theorem 1.2 to the sequence Λ . As a consequence, the associated control cost $\mathcal{K}(T)$ for

system (91) satisfies (10) and (11) for appropriate positive constants C_0, C_1, τ_0 and τ_1 independent of T . \square

From now on, let us suppose that (14) and (93) hold. Then, when $T > T_0$, we deduce that system (91) is null controllable at time T . So, for any $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$, the set

$$\mathcal{C}_T(y_0) := \{v \in L^2(0, T) : y(T, \cdot) = 0 \text{ in } (0, \pi), y \text{ solution of (91)}\},$$

is non empty and therefore, we can define the control cost of system (91) in time T , $\mathcal{K}(T)$, when $T > T_0$ (see (9)).

The positive part of the null controllability result for system (91) at time $T > 0$ stated in Theorem 5.1 is proved in [34] by using the moment method. Let us briefly describe this method for system (91).

From the previous assumptions, we deduce that $(L_2, D(L_2))$ is a self-adjoint operator. Its spectrum is given by

$$\sigma(L_2) := \Lambda = \Lambda_1 \cup \Lambda_2 = \left\{ \lambda_k^{(1)}, \lambda_k^{(2)} \right\}_{k \geq 1} = \{k^2, k^2 + \varepsilon_k\}_{k \geq 1}, \quad (94)$$

and the eigenspaces of L_2 associated to $\lambda_k^{(1)}$ and $\lambda_k^{(2)}$ are respectively generated by

$$\phi_k^{(1)} = \begin{pmatrix} \varphi_k^{(1)} \\ 0 \end{pmatrix} \quad \text{and} \quad \phi_k^{(2)} = \begin{pmatrix} 0 \\ \varphi_k^{(2)} \end{pmatrix}, \quad \forall k \geq 1. \quad (95)$$

Moreover, the sequence $\{\phi_k^{(1)}, \phi_k^{(2)}\}_{k \geq 1}$ is an orthonormal basis of $L^2(0, \pi; \mathbb{R}^2)$ and an orthogonal basis of $H_0^1(0, \pi; \mathbb{R}^2)$ and $H^{-1}(0, \pi; \mathbb{R}^2)$.

Using the spectral properties of the operator L_2 (see (13)) we can rewrite the null controllability problem for system (91) at time T as a moment problem. To be precise, one has:

Proposition 5.3. *Under the previous assumptions, given $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$, the control $v \in L^2(0, T)$ is such that the corresponding solution of (91) satisfies $y(T, \cdot) = 0$ in $(0, \pi)$ if and only if $v \in L^2(0, T)$ satisfies*

$$\begin{cases} b_1 \varphi_{k,x}^{(1)}(0) \int_0^T v(T-t) e^{-\lambda_k^{(1)} t} dt = -e^{-\lambda_k^{(1)} T} \langle y_0, \phi_k^{(1)} \rangle_{H^{-1}, H_0^1}, \\ b_2 \varphi_{k,x}^{(2)}(0) \int_0^T v(T-t) e^{-\lambda_k^{(2)} t} dt = -e^{-\lambda_k^{(2)} T} \langle y_0, \phi_k^{(2)} \rangle_{H^{-1}, H_0^1}. \end{cases} \quad (96)$$

for any $k \geq 1$, where $\lambda_k^{(i)}$ and $\phi_k^{(i)}$ are respectively given in (94) and (95). \square

For a proof of the previous property, see [34].

In fact, when (93) holds and $T > T_0$, T_0 given in (92), the corresponding null controllability problem at time T for system (91) (or equivalently, the moment problem stated in Proposition 5.3) can be explicitly solved as follows (see [34] for the details): The sequence Λ given in (94) satisfies (3). Therefore, Lemma 4.1 can be applied to deduce the existence of a sequence $\{q_k^{(1)}, q_k^{(2)}\}_{k \geq 1} \subset L^2(0, T)$ biorthogonal to $\{e_k^{(1)}, e_k^{(2)}\}_{k \geq 1} \subset L^2(0, T)$, where

$$e_k^{(i)}(t) = e^{-\lambda_k^{(i)} t}, \quad \forall t \in (0, T), \quad i = 1, 2. \quad (97)$$

Thus, a formal solution of the moment problem (96) is:

$$v(t) = \sum_{k \geq 1} \left(e^{-\lambda_k^{(1)} T} m_k^{(1)} q_k^{(1)}(T-t) + e^{-\lambda_k^{(2)} T} m_k^{(2)} q_k^{(2)}(T-t) \right), \quad \forall t \in (0, T), \quad (98)$$

where

$$m_k^{(i)} = \frac{-1}{b_i \varphi_{k,x}^{(i)}(0)} \langle y_0, \phi_k^{(i)} \rangle_{H^{-1}, H_0^1}, \quad \forall k \geq 1, \quad i = 1, 2. \quad (99)$$

Furthermore, when $T > T_0$, with T_0 given in (92), the series (98) converges absolutely in $L^2(0, T)$ and provides a null control $v \in L^2(0, T)$ which in fact is a solution of the moment problem (96).

Let us see that we can conveniently choose the sequence $\{q_k^{(1)}, q_k^{(2)}\}_{k \geq 1}$ in order to select a null control for system (91) associated to $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$ with minimal norm in $L^2(0, T)$. For that purpose, we define (see Section 4)

$$\begin{aligned} E(\Lambda, T) &:= \overline{\text{span} \left\{ e_n^{(i)} : n \geq 1, i = 1, 2 \right\}}^{L^2(0, T)}, \\ E_k^{(1)}(\Lambda, T) &:= \overline{\text{span} \left\{ e_n^{(1)}, e_l^{(2)} : n \geq 1, n \neq k, l \geq 1 \right\}}^{L^2(0, T)}, \quad \forall k \geq 1 \\ E_k^{(2)}(\Lambda, T) &:= \overline{\text{span} \left\{ e_n^{(1)}, e_l^{(2)} : n \geq 1, l \geq 1, l \neq k \right\}}^{L^2(0, T)}, \quad \forall k \geq 1. \end{aligned}$$

We have:

Proposition 5.4. *Under the previous assumptions, let us suppose that (93) holds. Let us also consider $T > T_0$ and the sequence of functions*

$$s_k^{(i)}(t) := \frac{e^{-\lambda_k^{(i)} t} - p_k^{(i)}(t)}{d_{T,k,i}^2}, \quad t \in (0, T), \quad \forall k \geq 1, \quad i = 1, 2,$$

where $d_{T,k,i}$ and $p_k^{(i)} \in E_k^{(i)}(\Lambda, T)$ are defined by

$$d_{T,k,i}^2 = \inf_{p \in E_k^{(i)}(\Lambda, T)} \left\| e_k^{(i)} - p \right\|_{L^2(0, T)}^2 = \int_0^T \left| e^{-\lambda_k^{(i)} t} - p_k^{(i)}(t) \right|^2 dt, \quad \forall k \geq 1, \quad i = 1, 2.$$

Then, the sequence $\{s_k^{(1)}, s_k^{(2)}\}_{k \geq 1} \subset E(\Lambda, T)$ is biorthogonal to $\{e_k^{(1)}, e_k^{(2)}\}_{k \geq 1}$ in $L^2(0, T)$ (the function $e_k^{(i)}$ is given in (97)). Moreover, given $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$, the control $u \in L^2(0, T)$ given by

$$u(t) = \sum_{k \geq 1} \left(e^{-\lambda_k^{(1)} T} m_k^{(1)} s_k^{(1)}(T-t) + e^{-\lambda_k^{(2)} T} m_k^{(2)} s_k^{(2)}(T-t) \right), \quad \forall t \in (0, T), \quad (100)$$

where $m_k^{(i)}$ is given in (99), satisfies $u \in \mathcal{C}_T(y_0)$, $\hat{u} \in E(\Lambda, T)$ (\hat{u} is the function $\hat{u}(t) = u(T-t)$, $t \in (0, T)$) and

$$\|u\|_{L^2(0, T)} = \inf_{v \in \mathcal{C}_T(y_0)} \|v\|_{L^2(0, T)}.$$

Proof. As said before, under assumption (93), the sequence Λ satisfies (3). Then, the family $\{e_k^{(1)}, e_k^{(2)}\}_{k \geq 1}$ is minimal in $L^2(0, T)$. In particular, we deduce that the functions $s_k^{(i)}$ are well defined, live in $E(\Lambda, T)$, for any $k \geq 1$ and $i = 1, 2$, and are biorthogonal to $\{e_k^{(1)}, e_k^{(2)}\}_{k \geq 1}$. These properties together with $T > T_0$ imply that the function u defined in (100) satisfies $\hat{u} \in E(\Lambda, T)$ and solves the null controllability problem at time T for system (91) and $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$, i.e., $u \in \mathcal{C}_T(y_0)$.

Let us now consider another null control $v \in \mathcal{C}_T(y_0)$. Using the equivalence stated in Proposition 5.3 we infer that v satisfies the moment problem (96). Therefore,

$$\int_0^T [v(T-t) - u(T-t)] e^{-\lambda_k^{(i)} t} dt = 0, \quad \forall k \geq 1 \text{ and } i = 1, 2,$$

that is to say, $\widehat{v} - \widehat{u} \in E(\Lambda, T)^\perp$ (\widehat{v} and \widehat{u} are defined as $\widehat{v}(t) = v(T-t)$ and $\widehat{u}(t) = u(T-t)$, $t \in (0, T)$). Using that $\widehat{u} \in E(\Lambda, T)$, we deduce

$$\begin{aligned} \|v\|_{L^2(0,T)}^2 &= \|\widehat{v}\|_{L^2(0,T)}^2 = \|(\widehat{v} - \widehat{u}) + \widehat{u}\|_{L^2(0,T)}^2 = \|\widehat{v} - \widehat{u}\|_{L^2(0,T)}^2 + \|\widehat{u}\|_{L^2(0,T)}^2 \\ &\geq \|\widehat{u}\|_{L^2(0,T)}^2 = \|u\|_{L^2(0,T)}^2. \end{aligned}$$

The previous inequalities prove the result. This finalizes the proof. \square

Our objective is to apply Theorems 1.7 and 1.9 to system (91) in a particular case. To this end, let us state a technical result of inverse spectral theory whose proof can be found in [36] (see also [34]):

Lemma 5.5. *Let us consider $\{\varepsilon_k\}_{k \geq 1}$, a sequence in ℓ^2 . Then, there exists a function $Q \in L^2(0, \pi)$ satisfying (14) such that*

$$\sigma(\widetilde{L}_2) = \Lambda_2 = \{k^2 + \varepsilon_k\}_{k \geq 1},$$

where $\widetilde{L}_2 := -\partial_{xx} + Q$ with domain $D(\widetilde{L}_2) = H^2(0, \pi) \cap H_0^1(0, \pi)$. \square

From now on, we will take

$$\varepsilon_k = e^{-k^{2\gamma}}, \quad k \geq 1,$$

with $\gamma \in (0, \infty)$, and $B = (b_1, b_2)^t$ with $b_1 b_2 \neq 0$. Clearly $\{\varepsilon_k\}_{k \geq 1} \in \ell^2$ and we can apply Lemma 5.5. We will work with the function Q_γ associated to the previous sequence provided by Lemma 5.5 and the corresponding sequences of eigenvalues and orthogonal basis Λ_1, Λ_2 and $\{\varphi_k^{(1)}\}_{k \geq 1}$ and $\{\varphi_k^{(2)}\}_{k \geq 1}$ associated to the operators \widetilde{L}_1 and \widetilde{L}_2 . With this choice, we consider the parabolic control system (91) with L_2 given in (13).

Observe that the sequence Λ of eigenvalues of the operator L_2 can be rearranged as an increasing sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ ($\gamma \in (0, \infty)$) doing:

$$\Lambda_{2k-1} = k^2, \quad \Lambda_{2k} = k^2 + e^{-k^{2\gamma}}, \quad \forall k \geq 1. \quad (101)$$

It is clear that the functions

$$\phi_{2k-1} = \begin{pmatrix} \varphi_k^{(1)} \\ 0 \end{pmatrix} \quad \text{and} \quad \phi_{2k} = \begin{pmatrix} 0 \\ \varphi_k^{(2)} \end{pmatrix}, \quad \forall k \geq 1. \quad (102)$$

are an orthonormal basis of eigenfunctions of the operator L_2 in $L^2(0, \pi; \mathbb{R}^2)$ and an orthogonal basis of $H_0^1(0, \pi; \mathbb{R}^2)$ and $H^{-1}(0, \pi; \mathbb{R}^2)$.

The controllability properties of system (91) at time $T > 0$ can be deduced from Theorem 5.1. In this case, system (91) is approximately controllable for any final time $T > 0$. The expression of the minimal time is (see (92))

$$T_0 = \lim_{k \rightarrow \infty} \frac{-\log(e^{-k^{2\gamma}})}{k^2} = \begin{cases} 0 & \text{if } \gamma \in (0, 1), \\ 1 & \text{if } \gamma = 1, \\ \infty & \text{if } \gamma > 1. \end{cases}$$

We deduce then

1. If $\gamma \in (0, 1)$, system (91) is null controllable at any final time $T > 0$.

2. If $\gamma = 1$, system (91) is null controllable at any final time $T > 1$ and is not controllable at time T when $T < 1$.
3. When $\gamma > 1$, system (91) is never null controllable at any final time $T > 0$.

Observe that, when $\gamma \in (0, 1)$ and $Q_\gamma \in L^2(0, \pi)$ is the function provided by Lemma 5.5 associated to $\varepsilon_k = e^{-k^{2\gamma}}$, system (91) is null controllable at time T , for any $T > 0$. We can introduce the control cost $\mathcal{K}(T)$ associated to this system (see (9)). Our objective is to analyze the dependence of $\mathcal{K}(T)$ with respect to T and $\gamma \in (0, 1)$.

First, let us see that the sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ (see (101)) of eigenvalues of the operator L_2 is in the class $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ (see Definition 1.4) for appropriate parameters satisfying (20). One has:

Proposition 5.6. *Let us fix $\gamma \in (0, 1)$ and consider the sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ with Λ_k given in (101), $k \geq 1$. Then, the sequence Λ satisfies $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ and (26), with $\beta = 0$, $q = 2$,*

$$\rho = \frac{1}{16}, \quad p_0 = 1, \quad p_1 = p_2 = 2, \quad \alpha = 2 + \frac{1}{\sqrt{e}} \quad \text{and} \quad \nu = \frac{1}{2} \left(1 + \frac{1}{e}\right).$$

Proof. The proof of this result is a direct consequence of Proposition 2.14 and Remark 2.16. Indeed, the sequence Λ can be written as $\Lambda = \Lambda_1 \cup \Lambda_2$ with

$$\Lambda_1 = \left\{ \lambda_k^{(1)} \right\}_{k \geq 1} = \{k^2\}_{k \geq 1} \quad \text{and} \quad \Lambda_2 = \left\{ \lambda_k^{(2)} \right\}_{k \geq 1} = \left\{ k^2 + e^{-k^{2\gamma}} \right\}_{k \geq 1}.$$

It is easy to see that $\Lambda_1 \in \mathcal{L}(\beta, \rho_1, q, \pi_0, \pi_1, \pi_2, \alpha_1)$ and satisfies (26) with $\beta = 0$, $\rho_1 = 1$, $q = 1$, $\pi_0 = \pi_1 = \pi_2 = 1$, $\alpha_1 = 1$ and $\nu = \rho_1 = 1$. On the other hand,

$$\varepsilon_0 = \sup_{k \geq 1} |\varepsilon_k| = \sup_{k \geq 1} e^{-k^{2\gamma}} = e^{-1}.$$

In addition, the sequence Λ can be explicitly defined by (101) (see (50)). So, from Proposition 2.14 and Remark 2.16, we deduce that $\Lambda \in \mathcal{L}(\beta, \rho, q, \pi_0, p_1, p_2, \alpha)$ and satisfies (26) with parameters given in the statement of the result. This finalizes the proof. \square

With the previous choice, the sequence Λ satisfies property (H5) for $q = 2$. In this case, let us see how the term \mathcal{P}_k (see (23)) can be estimated. One has:

Proposition 5.7. *Let us fix $\gamma \in (0, 1)$ and consider the sequence Λ given by (101). Then, $\mathcal{P}_1 = e$ and*

$$\begin{cases} \frac{1}{(2n-1)} e^{n^{2\gamma}} \leq \mathcal{P}_{2n-1} \leq \frac{1}{(2n-1) - e^{-1}} e^{n^{2\gamma}}, & \forall n \geq 2, \\ \frac{1}{(2n+1)} e^{n^{2\gamma}} \leq \mathcal{P}_{2n} \leq \frac{1}{(2n+1) - e^{-1}} e^{n^{2\gamma}}, & \forall n \geq 1, \end{cases}$$

where \mathcal{P}_k and the sequence Λ are respectively given in (23) and (101).

Proof. Let us prove the result when $k = 2n$, with $n \geq 1$. The case $k = 2n - 1$, with $n \geq 1$, is similar. From (23) and (101), we deduce

$$\mathcal{P}_{2n}^{-1} = (\Lambda_{2n} - \Lambda_{2n-1})(\Lambda_{2n+1} - \Lambda_{2n}) = e^{-n^{2\gamma}} \left[(2n+1) - e^{-n^{2\gamma}} \right], \quad \forall n \geq 1.$$

The previous formula provides the proof of the result. \square

Remark 5.8. Let us take a sequence $\{\varepsilon_k\}_{k \geq 1}$ in ℓ^2 such that $0 < \varepsilon_k < 1$ for any $k \geq 1$. From Lemma 5.5, there exists a function $Q \in L^2(0, \pi)$ such that $\sigma(\tilde{L}_2) = \{k^2 + \varepsilon_k\}_{k \geq 1}$. As before, we can consider the operator L_2 associated to system (91) (see (13)) and the corresponding sequence of eigenvalues Λ given by (94). In this case, we can repeat the computations in Proposition 5.7 and deduce $\mathcal{P}_1 = \varepsilon_1^{-1}$ and

$$\begin{cases} \frac{1}{(2n-1)\varepsilon_n} \leq \mathcal{P}_{2n-1} \leq \frac{1}{(2n-2)\varepsilon_n}, & \forall n \geq 2, \\ \frac{1}{(2n+1)\varepsilon_n} \leq \mathcal{P}_{2n} \leq \frac{1}{2n\varepsilon_n}, & \forall n \geq 1, \end{cases}$$

(\mathcal{P}_k is given in (23) with $q = 2$). The previous estimates prove that we can construct functions $Q \in L^2(0, \pi)$ such that the sequence $\{\mathcal{P}_k\}_{k \geq 1}$ associated to the sequence $\Lambda = \sigma(L_2)$ can have any arbitrary explosive behavior. \square

The main results of this section concern the control cost $\mathcal{K}(T)$ associated to system (91). First, let us state a bound from above of the control cost:

Theorem 5.9. *Let us fix $\gamma \in (0, 1)$ and take the function $Q_\gamma \in L^2(0, \pi)$ provided by Lemma 5.5 associated to $\varepsilon_k = e^{-k^{2\gamma}}$. If we denote $\mathcal{K}_\gamma(T)$ the control cost of system (91) in $L^2(0, T)$ at time $T > 0$, then, there exists a positive constant \mathcal{C} , independent of γ , such that*

$$\mathcal{K}_\gamma(T) \leq \exp \left[\mathcal{C} \left(1 + \frac{1}{T} \right) + \frac{\mathcal{C}}{(1-\gamma)T} + \frac{1-\gamma}{T^{1-\gamma}} \right], \quad \forall T > 0. \quad (103)$$

Proof. Under the assumptions of the theorem, we can apply Proposition 5.6 and deduce that the sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ (Λ_k given in (101)) of eigenvalues of the operator L_2 (see (13)) satisfies $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ and (26), with $\beta, \rho, q, p_0, p_1, p_2$ and α given in the statement of this result.

Let us now take $T > 0$. Recall that the minimal time associated to system (91) is $T_0 = 0$. Therefore, without loss of generality, we can assume that $T \in (0, 1)$. Thus, Theorem 1.7 can be applied to Λ and we deduce the existence of a family of functions $\{q_k\}_{k \geq 1} \subset L^2(0, T)$, biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T)$ (for the expression of e_k , see (1)) which satisfies (22). In particular, there exists a positive constant \mathcal{C} , independent of γ , such that

$$\|q_k\|_{L^2(0, T)} \leq \exp \left[\mathcal{C} \left(1 + \sqrt{|\Lambda_k|} + \frac{1}{T} \right) \right] \mathcal{P}_k, \quad \forall k \geq 1.$$

If we combine the previous inequality with Proposition 5.7 and (101), we get

$$\begin{cases} \|q_{2k-1}\|_{L^2(0, T)} \leq \exp \left[\mathcal{C} \left(1 + k + \frac{1}{T} \right) + k^{2\gamma} \right], & \forall k \geq 1, \\ \|q_{2k}\|_{L^2(0, T)} \leq \exp \left[\mathcal{C} \left(1 + k + \frac{1}{T} \right) + k^{2\gamma} \right], & \forall k \geq 1, \end{cases} \quad (104)$$

for a new positive constant \mathcal{C} , independent of γ .

Let us prove the result. To this end, we consider $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$ with

$$\|y_0\|_{H^{-1}(0, \pi; \mathbb{R}^2)} \leq 1.$$

Using the moment method, in [34], the author proves that, taking

$$v(t) = \sum_{k \geq 1} \left(e^{-k^2 T} m_k^{(1)} q_{2k-1}(T-t) + e^{-(k^2 + e^{-k^{2\gamma}})T} m_k^{(2)} q_{2k}(T-t) \right), \quad (105)$$

$t \in (0, T)$, where $m_k^{(i)}$ is given in (99), one has $v \in L^2(0, T)$ and the corresponding solution of system (91), $y \in C^0([0, T]; H^{-1}(0, \pi; \mathbb{R}^2))$, satisfies $y(T, \cdot) = 0$ in $(0, \pi)$ ($\phi_k^{(i)}$ is given in (95)). In [34] the author also shows that there exists a positive constant \mathcal{C} (independent of k) such that

$$\left| m_k^{(i)} \right| \leq \mathcal{C} \|y_0\|_{H^{-1}(0, \pi; \mathbb{R}^2)} \leq \mathcal{C}, \quad \forall k \geq 1, \quad i = 1, 2.$$

Coming back to (105) and taking into account (104) and the previous estimate, we deduce

$$\|v\|_{L^2(0, T)} \leq e^{\mathcal{C}(1 + \frac{1}{T})} \sum_{k \geq 1} e^{-k^2 T + \mathcal{C}k + k^{2\gamma}},$$

for a new positive constant \mathcal{C} , independent of γ . Let us now take $\varepsilon \in (0, 1/2)$, which will be fixed later. Observe that Young inequality implies

$$\mathcal{C}k \leq \varepsilon k^2 T + \frac{\mathcal{C}^2}{4\varepsilon T}, \quad \forall k \geq 1,$$

and therefore, we can write

$$-k^2 T + \mathcal{C}k + k^{2\gamma} \leq -k^2 T + \varepsilon k^2 T + \frac{\mathcal{C}^2}{4\varepsilon T} + k^{2\gamma} = h_\varepsilon(k^2) + \frac{\mathcal{C}^2}{4\varepsilon T} - \varepsilon k^2 T, \quad \forall k \geq 1,$$

where the function h_ε is given by

$$h_\varepsilon(x) = -(1 - 2\varepsilon)Tx + x^\gamma, \quad x \in (0, \infty).$$

To summarize, the previous calculations provide the following estimate:

$$\|v\|_{L^2(0, T)} \leq e^{\mathcal{C}(1 + \frac{1}{T})} e^{\frac{\mathcal{C}^2}{4\varepsilon T}} \sum_{k \geq 1} e^{h_\varepsilon(k^2)} e^{-\varepsilon k^2 T}, \quad (106)$$

for any $\varepsilon > 0$.

It is easy to see that h_ε possesses an absolute maximum in $(0, \infty)$ at point

$$x^* = \left(\frac{\gamma}{1 - 2\varepsilon} \frac{1}{T} \right)^{\frac{1}{1-\gamma}}.$$

Thus, if we take $\varepsilon = (1 - \gamma)/2$, we can write

$$h_\varepsilon(x) \leq h_\varepsilon(x^*) = (1 - \gamma) \left(\frac{\gamma}{1 - 2\varepsilon} \right)^{\frac{\gamma}{1-\gamma}} \frac{1}{T^{\frac{\gamma}{1-\gamma}}} = \frac{1 - \gamma}{T^{\frac{\gamma}{1-\gamma}}}, \quad \forall x \in (0, \infty).$$

Going back to the formula (106), we deduce

$$\|v\|_{L^2(0, T)} \leq \exp \left[\mathcal{C} \left(1 + \frac{1}{T} \right) + \frac{\mathcal{C}^2}{2(1 - \gamma)T} + \frac{1 - \gamma}{T^{\frac{\gamma}{1-\gamma}}} \right] \sum_{k \geq 1} e^{-\frac{1}{2}(1 - \gamma)k^2 T}.$$

Finally, a comparison with Gauss integral gives

$$\sum_{k \geq 1} e^{-\frac{1}{2}(1 - \gamma)k^2 T} \leq \int_0^\infty e^{-\frac{1}{2}(1 - \gamma)Tx^2} dx = \frac{\sqrt{2\pi}}{2} \frac{1}{\sqrt{(1 - \gamma)T}} \leq e^{\frac{1}{2(1 - \gamma)T}},$$

and then,

$$\|v\|_{L^2(0, T)} \leq \exp \left[\mathcal{C} \left(1 + \frac{1}{T} \right) + \frac{\mathcal{C}^2 + 1}{2(1 - \gamma)T} + \frac{1 - \gamma}{T^{\frac{\gamma}{1-\gamma}}} \right].$$

It is clear that, from the previous inequality, we can deduce (103) for a new positive constant \mathcal{C} , independent of γ . This completes the proof. \square

Our second result provides an estimate from below of the control cost $\mathcal{K}_\gamma(T)$ for system (91) in $L^2(0, T)$ at the final time $T > 0$. As before, we are going to fix $\gamma \in (0, 1)$ and take the function $Q_\gamma \in L^2(0, \pi)$ provided by Lemma 5.5 associated to $\varepsilon_k = e^{-k^{2\gamma}}$. One has:

Theorem 5.10. *Under the assumptions of Theorem 5.9, there exists two positive constants τ_0 and \mathcal{C} , independent of γ , such that*

$$\mathcal{K}_\gamma(T) \geq \mathcal{C} \exp\left(\frac{\mathcal{C}}{T} + \frac{\mathcal{C}(1-\gamma)}{T^{1-\gamma}}\right), \quad \forall T \in (0, \tau_0). \quad (107)$$

Before starting the proof of Theorem 5.10, we will show a technical result that we will use in its proof:

Lemma 5.11. *Let us consider $T > 0$ and $\gamma \in (0, 1)$ and define the function*

$$\tilde{h}(x) = -Tx + x^\gamma, \quad \forall x \in (0, \infty).$$

Let us assume that

$$T < \gamma \left(\frac{\sqrt{2}-1}{\sqrt{2}}\right)^{2(1-\gamma)}. \quad (108)$$

Then, there exists $k_0 \geq 1$ such that

$$\tilde{h}(k_0^2) \geq \frac{(1 + \log 2)(1-\gamma)}{2e} \frac{(1-\gamma)}{T^{1-\gamma}}.$$

Proof. Under the assumptions of the lemma, it is easy to see that the function \tilde{h} is increasing in $(0, \tilde{x})$ and decreasing in (\tilde{x}, ∞) , where

$$\tilde{x} = \left(\frac{\gamma}{T}\right)^{\frac{1}{1-\gamma}}.$$

Thus, if $k_0 \geq 1$ is such that

$$\frac{1}{2}\tilde{x} \leq k_0^2 \leq \tilde{x}, \quad (109)$$

then

$$\begin{aligned} \tilde{h}(k_0^2) &\geq \tilde{h}(\tilde{x}/2) = \left(\frac{1}{2^\gamma} - \frac{\gamma}{2}\right) \left(\frac{\gamma}{T}\right)^{\frac{\gamma}{1-\gamma}} > \frac{1}{2}(1 + \log 2)(1-\gamma) \left(\frac{\gamma}{T}\right)^{\frac{\gamma}{1-\gamma}} \\ &\geq \frac{(1 + \log 2)(1-\gamma)}{2e} \frac{(1-\gamma)}{T^{1-\gamma}}, \end{aligned}$$

and we would have the proof of the result.

In order to finish the proof, let us check that there exists $k_0 \geq 1$ such that (109) holds. Indeed, (109) is equivalent to

$$\frac{1}{\sqrt{2}}\sqrt{\tilde{x}} \leq k_0 \leq \sqrt{\tilde{x}}.$$

Observe that this property occurs if

$$\sqrt{\tilde{x}} - \frac{1}{\sqrt{2}}\sqrt{\tilde{x}} > 1,$$

i.e., if T satisfies (108). This ends the proof. \square

Proof of Theorem 5.10. As before, under the assumptions of the theorem, we know that the sequence Λ of eigenvalues of the operator L_2 (see (13)) satisfies $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ and (26), with $\beta, \rho, q, p_0, p_1, p_2$ and α given in the statement of Proposition 5.6.

Let us fix $T > 0$. The minimal time T_0 for system (91) associated to the function Q_γ is $T_0 = 0$. In addition, we can apply Proposition 5.4 and Theorem 1.9. We deduce that the optimal family $\{s_k\}_{k \geq 1} \subset E(\Lambda, T)$ biorthogonal to $\{e_k\}_{k \geq 1}$ in $L^2(0, T)$ satisfies (27) (e_k is given in (1)).

We will divide the proof of the result into two parts:

1. Assume that $\gamma \in (0, 1/2]$. In this case, it is easy to check that, for any $\tau_0 \in (0, 1]$, one has

$$\frac{1}{T} \geq \frac{1-\gamma}{T^{1-\gamma}}, \quad \forall T < \tau_0.$$

Therefore, inequality (107) is equivalent to prove the existence of a positive constant \mathcal{C}_0 , independent of γ , such that

$$\mathcal{K}_\gamma(T) \geq \mathcal{C}_0 \exp\left(\frac{\mathcal{C}_0}{T}\right), \quad \forall T \in (0, \tau_0). \quad (110)$$

Our objective is to find $\mathcal{C}_0 > 0$ and $\tau_0 \in (0, 1]$, independent of γ , such that one has inequality (110).

From inequality (27) written for the function s_3 , we deduce ($\nu = \frac{1}{2}(1 + \frac{1}{e})$):

$$\|s_3\|_{L^2(0, T)} \geq \frac{6}{\pi^2} \mathcal{B}_3 \mathcal{P}_3 e^{\frac{1}{T\nu}}$$

where (see (28) for $q = 2$)

$$\mathcal{B}_3 = \mathcal{C} \frac{(\nu T)^4}{(1 + \nu T)^9} \sqrt{|\Lambda_1| + \frac{1}{2T}},$$

and \mathcal{C} is a positive constant ($\beta = 0$ and then $\delta = 1$). From the previous expression, it is not difficult to see that there exist $\mathcal{C}_0 > 0$ and $\tau_0 \in (0, 1]$, independent of γ , such that

$$\mathcal{B}_3 \mathcal{P}_3 \geq \mathcal{C}_0 e^{-\frac{1}{2T\nu}}, \quad \forall T \in (0, \tau_0).$$

Coming back to the expression of $\|s_3\|_{L^2(0, T)}$, we finally deduce:

$$\|s_3\|_{L^2(0, T)} \geq \mathcal{C} e^{\frac{1}{2T\nu}}, \quad \forall T \in (0, \tau_0). \quad (111)$$

Let us take $y_0 = \phi_3 / \|\phi_3\|_{H^{-1}}$ (see (102)). Then, applying Proposition 5.4 to y_0 , it is possible to construct the null control with minimal L^2 -norm for system (91) associated to y_0 (see (100)):

$$u(t) = e^{-4T} \frac{1}{b_1 \varphi_{2,x}^{(1)}(0)} \frac{1}{\|\phi_3\|_{H^{-1}}} s_3(T-t), \quad \forall t \in (0, T).$$

From (111), we also have

$$\mathcal{K}_\gamma(T) \geq \inf_{v \in \mathcal{C}_T(y_0)} \|v\|_{L^2(0, T)} = \|u\|_{L^2(0, T)} = \mathcal{C} \|s_3\|_{L^2(0, T)} \geq \mathcal{C} e^{\frac{1}{2T\nu}}, \quad \forall T \in (0, \tau_0).$$

This proves inequality (110) and inequality (107) when $\gamma \in (0, 1/2]$.

2. Let us now assume that $\gamma \in (1/2, 1)$. In this case, inequality (107) is equivalent to

$$\mathcal{K}_\gamma(T) \geq \mathcal{C}_0 \exp\left(\frac{\mathcal{C}_0}{T^{1-\gamma}}\right), \quad \forall T \in (0, \tau_0). \quad (112)$$

and therefore, our goal is to prove that there exist two positive constants \mathcal{C}_0 and τ_0 , independent of γ , in such a way that the previous inequality holds. As before, we are going to work with an appropriate element s_{k_0} of the optimal biorthogonal family $\{s_k\}_{k \geq 1} \subset E(\Lambda, T)$ provided by Proposition 5.4.

Let us define τ_0 as

$$\tau_0 = \frac{1}{2} \left(\frac{\sqrt{2} - 1}{\sqrt{2}} \right).$$

Observe that if $T \in (0, \tau_0)$, then inequality (108) is valid for any $\gamma \in (1/2, 1)$. From Lemma 5.11, we can infer the existence of $k_0 \geq 1$ such that

$$\tilde{h}(k_0^2) = -k_0^2 T + (k_0)^{2\gamma} \geq \frac{(1 + \log 2)(1 - \gamma)}{2e} \frac{(1 - \gamma)}{T^{1-\gamma}} = \frac{\mathcal{C}(1 - \gamma)}{T^{1-\gamma}}. \quad (113)$$

Consider $y_0 = \phi_{2k_0-1} / \|\phi_{2k_0-1}\|_{H^{-1}}$, i.e. (see (102)),

$$y_0(x) = k_0 \sqrt{\frac{2}{\pi}} \begin{pmatrix} \sin(k_0 x) \\ 0 \end{pmatrix}.$$

On the other hand, let us also consider the null control for system (91) associated to y_0 provided by Proposition 5.4:

$$u(\cdot) = e^{-k_0^2 T} \frac{1}{b_1 \varphi_{k_0, x}^{(1)}(0)} \langle y_0, \phi_{2k_0-1} \rangle_{H^{-1}, H_0^1} s_{2k_0-1}(T \cdot) = \frac{1}{b_1} \sqrt{\frac{2}{\pi}} e^{-k_0^2 T} s_{2k_0-1}(T \cdot).$$

Using inequality (27), written for the function s_{2k_0-1} , and taking into account Proposition 5.7 ($q = 2$ and $\delta = 1$) and (113), we deduce

$$\begin{aligned} \|u\|_{L^2(0, T)} &= \mathcal{C} e^{-k_0^2 T} \|s_{2k_0-1}\|_{L^2(0, T)} \geq \frac{\mathcal{C}}{T^2} \left(\frac{5}{2T} + \Lambda_{2k_0-2} \right)^{1/2} e^{-k_0^2 T} \mathcal{P}_{2k_0-1} \\ &\geq \frac{\mathcal{C}}{T^2} \left(\frac{5}{2T} + \Lambda_{2k_0-2} \right)^{1/2} \frac{1}{2k_0 - 1} e^{-k_0^2 T + (k_0)^{2\gamma}} \\ &= \frac{\mathcal{C}}{T^2} \left(\frac{5}{2T} + \Lambda_{2k_0-2} \right)^{1/2} \frac{e^{\tilde{h}(k_0^2)}}{2k_0 - 1} \geq \frac{\mathcal{C}}{T^2} \exp \left(\frac{\mathcal{C}(1 - \gamma)}{T^{1-\gamma}} \right). \end{aligned}$$

where \mathcal{C} is a constant independent of γ and k_0 .

As before,

$$\mathcal{K}_\gamma(T) \geq \inf_{v \in \mathcal{C}_T(y_0)} \|v\|_{L^2(0, T)} = \|u\|_{L^2(0, T)} \geq \frac{\mathcal{C}}{T^2} \exp \left(\frac{\mathcal{C}(1 - \gamma)}{T^{1-\gamma}} \right), \quad \forall T \in (0, \tau_0).$$

Thus, we can conclude that inequality (112) holds. This ends the proof of Theorem 5.10. \square

Remark 5.12. Observe that inequalities (103) and (107) are valid when $\gamma \in (0, 1)$. In fact, these inequalities are equivalent to:

1. If $\gamma \in (0, 1/2]$, then, there exist three positive constants τ_0 , \mathcal{C}_0 and \mathcal{C}_1 (independent of γ) such that

$$\exp \left[\mathcal{C}_0 \left(1 + \frac{1}{T} \right) \right] \leq \mathcal{K}_\gamma(T) \leq \exp \left[\mathcal{C}_1 \left(1 + \frac{1}{T} \right) \right], \quad \forall T \in (0, \tau_0).$$

Observe that the previous estimates for the control cost of system (91) are similar to those obtained for the control cost of the heat equation (see for instance [23] and [20]).

2. If $\gamma \in (1/2, 1)$, again, there exist three positive constants τ_0 , \mathcal{C}_0 and \mathcal{C}_1 (independent of γ) such that

$$\exp \left[\mathcal{C}_0 \left(1 + \frac{1}{T^{1-\gamma}} \right) \right] \leq \mathcal{K}_\gamma(T) \leq \exp \left[\mathcal{C}_0 \left(1 + \frac{1}{T^{1-\gamma}} \right) \right], \quad \forall T \in (0, \tau_0).$$

The previous expressions prove that the control cost blows up when $\gamma \rightarrow 1^-$. This is natural because the minimal time for system (91) when $\gamma = 1$ is $T_0 = 1$ and the system is not null controllable at time T when $T < 1$. \square

5.2. The linear phase-field system. Let us now apply Theorem 1.7 and Theorem 1.9 to the linear version of (15) around the constant trajectory $(0, c)$ with $c = 1$ or $c = -1$. To be precise, we will work with the linear system (12) with $L = L_3$ (see (16)) and $\rho, \tau, \xi \in (0, \infty)$. As said above, the controllability properties of this system has been analyzed in [22] under the condition $\xi \neq \frac{1}{j^2} \frac{\rho}{\tau}$, for any $j \in \mathbb{N}$. The approximate controllability of this system is given by the next result:

Theorem 5.13 (Approximate controllability). *Fix $T > 0$. Then, system (12) with $L = L_3$ (see (16)) is approximately controllable in $H^{-1}(0, \pi; \mathbb{R}^2)$ at time $T > 0$ if and only if $\lambda_k^{(3,1)} \neq \lambda_n^{(3,2)}$ for any $k, n \geq 1$ (see (17)), that is to say, if and only if*

$$\xi^2 \tau^2 (\ell^2 - k^2)^2 - 2\xi \rho \tau (\ell^2 + k^2) - 2\rho - 1 \neq 0, \quad \forall k, \ell \geq 1, \quad \ell > k. \quad (114)$$

The proof of this result can be found in [22].

Now, our objective is to give a null controllability result at time $T > 0$ for this system when (114) holds (which, in fact, is a necessary condition for the null controllability at time T of system (12) with $L = L_3$) and obtain a bound for the corresponding control cost $\mathcal{K}(T)$. This problem has analyzed in [22] under additional assumptions on the parameters ξ , ρ and τ . To be precise, in [22] the authors prove:

Theorem 5.14. *Let us us fix $T > 0$ and consider ξ , ρ and τ , positive real numbers satisfying (114) and*

$$\xi \neq \frac{1}{j^2} \frac{\rho}{\tau}, \quad \forall j \geq 1. \quad (115)$$

Then, system (12) with $L = L_3$ (see (16)) is exactly controllable to zero at time $T > 0$ in $H^{-1}(0, \pi; \mathbb{R}^2)$. Moreover, there exist two positive constants C and M , only depending on ξ , ρ and τ , such that

$$\mathcal{K}(T) \leq C e^{M/T}, \quad \forall T > 0,$$

where $\mathcal{K}(T)$ is the control cost for system (12) with $L = L_3$:

$$\mathcal{K}(T) = \sup_{\|y_0\|_{H^{-1}(0, \pi; \mathbb{R}^2)} = 1} \left(\inf_{v \in \mathcal{Z}_T(y_0)} \|v\|_{L^2(0, T)} \right), \quad \forall T > 0.$$

and

$$\mathcal{Z}_T(y_0) := \{v \in L^2(0, T) : y(\cdot, T) = 0 \text{ in } (0, \pi), \text{ } y \text{ solution of (12) for } L = L_3\}.$$

Conditions (114) and (115) implies that the sequence $\Lambda^{(3)} = \{\lambda_k^{(3,1)}, \lambda_k^{(3,2)}\}_{k \geq 1}$ (see (17)) satisfies the conditions in Theorem 1.2 (see Remark 1.3). In fact, condition (115) provides the gap condition (5) for the sequence Λ_3 . Therefore, Theorem 5.14 is a consequence of Theorem 1.2.

As said before, our objective is to analyze the null controllability of system (12) with $L = L_3$ without imposing condition (115) to the sequence Λ_3 of eigenvalues of

the operator L_3 . Let us first see that this sequence is in $\mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ with $\beta = 0$ and appropriate parameters $\rho, p_0, p_1, p_2, \alpha \in (0, \infty)$ (see Definition 1.4):

Proposition 5.15. *Let us consider ξ, ρ and τ , positive real numbers satisfying (114). Then, the sequence $\Lambda^{(3)} = \{\lambda_k^{(3,1)}, \lambda_k^{(3,2)}\}_{k \geq 1}$, with $\lambda_k^{(3,i)}$ given in (17), can be rearranged as a positive increasing sequence $\Lambda^{(3)} = \{\Lambda_k\}_{k \geq 1}$ satisfying $\Lambda^{(3)} \in \mathcal{L}(0, \rho, q, p_0, p_1, p_2, \alpha)$ and (26), with*

$$p_0 = p_1 = p_2 = \frac{2}{\sqrt{\xi}} \quad \text{and} \quad \alpha = \frac{1}{2\sqrt{\xi}} \left(\sqrt{\frac{\rho}{\tau}} + \sqrt{\frac{3\rho+4}{\tau}} \right) + 2,$$

and $q \geq 2$, ρ and ν positive constants only depending on ξ, ρ and τ .

Proof. The proof of this result is a direct consequence of the results in [22]. Indeed, from Proposition 3.2 of [22] one has,

$$0 < \lambda_k^{(3,1)} < \lambda_k^{(3,2)}, \quad \forall k \geq 1.$$

Secondly, as a consequence of assumption (114) and Theorem 5.13, we deduce that the elements of the sequence $\Lambda^{(3)}$ are pairwise different. Thus, this sequence can be rearranged into a positive increasing sequence $\Lambda^{(3)} = \{\Lambda_k\}_{k \geq 1}$ that satisfies (H1) and, of course, (H2), (H3) ($\beta = 0$) and (H4).

On the other hand, taking into account the proof of Proposition 3.3 in [22], we also have that $\Lambda^{(3)}$ satisfies condition (H6) in Definition 1.4 with parameters p_0, p_1, p_2 and α as in the statement of the proposition.

Finally, we deduce properties (H5) and (26) from Proposition 2.4 with q, ρ and ν given in (36). This ends the proof of the proposition. \square

In the next result we will provide further properties of the sequence $\Lambda^{(3)}$ that will be used later. Again, we will use some properties that has been proved in [22]. One has:

Proposition 5.16. *Let us consider ξ, ρ and τ , positive real numbers. Then,*

$$\lambda_k^{(3,2)} - \lambda_{k+i}^{(3,1)} = \xi \left(\sqrt{\frac{\rho}{\xi\tau}} - i \right) (2k+i) + \left(\frac{\epsilon_{k+i}}{k+i} + \frac{\epsilon_k}{k} \right), \quad \forall k, i \geq 1, \quad (116)$$

where $\lambda_k^{(3,i)}$ is given in (17) and $\{\epsilon_k\}_{k \geq 1}$ is the increasing sequence given by

$$\epsilon_k = \left(\frac{\rho+1}{2\tau} \right)^2 \frac{1}{\sqrt{\frac{\xi\rho}{\tau} + \frac{1}{k^2} \left(\frac{\rho+1}{2\tau} \right)^2} + \sqrt{\frac{\xi\rho}{\tau}}}, \quad \forall k \geq 1. \quad (117)$$

Proof. The proof of the result can be found in [22]. \square

Let us now analyze the control cost for the linear phase-field system, i.e., the control cost for system (12) with $L = L_3$. One has:

Theorem 5.17. *Let us consider ξ, ρ and τ , positive real numbers satisfying (114). Then, system (12) with $L = L_3$ (see (16)) is exactly controllable to zero at any time $T > 0$. Moreover, there exist positive constants C_0, C_1, M_0 and M_1 (only depending on ξ, ρ and τ) such that*

$$C_0 e^{M_0/T} \leq \mathcal{K}(T) \leq C_1 e^{M_1/T}, \quad \forall T \in (0, 1], \quad (118)$$

where $\mathcal{K}(T)$ is the control cost for system (12) with $L = L_3$ defined in the statement of Theorem 5.14.

Proof. The result is proved in [22] when the coefficients ξ , ρ and τ satisfy conditions (114) and (115). Thus, let us prove the result when these coefficients do not satisfy (115), that is to say, when one has

$$\xi = \frac{1}{j_0^2} \frac{\rho}{\tau},$$

for some integer $j_0 \geq 1$. In this case, (116) becomes

$$\lambda_k^{(3,2)} - \lambda_{k+i}^{(3,1)} = \xi(j_0 - i)(2k + i) + \left(\frac{\epsilon_{k+i}}{k+i} + \frac{\epsilon_k}{k} \right), \quad \forall k, i \geq 1, \quad (119)$$

where $\lambda_k^{(3,i)}$ is given in (17) and $\{\epsilon_k\}_{k \geq 1}$ is the increasing sequence given by (117). In particular, we can estimate the terms ϵ_k of the sequence as follows:

$$\left(\frac{\rho+1}{2\tau} \right)^2 \frac{1}{\sqrt{\frac{\xi\rho}{\tau} + \left(\frac{\rho+1}{2\tau}\right)^2} + \sqrt{\frac{\xi\rho}{\tau}}} = \epsilon_1 \leq \epsilon_k < \lim_{k \rightarrow \infty} \epsilon_k = \left(\frac{\rho+1}{2\tau} \right)^2 \frac{\sqrt{\tau}}{2\sqrt{\xi\rho}} := L,$$

for all $k \geq 1$. We will use the previous inequalities in what follows.

If we choose i such that $1 \leq i \leq j_0 - 1$, from (119), we infer

$$\begin{cases} \lambda_k^{(3,2)} - \lambda_{k+i}^{(3,1)} > \xi(j_0 - i)(2k + i), \\ \lambda_k^{(3,2)} - \lambda_{k+i}^{(3,1)} < \xi(j_0 - i)(2k + i) + \frac{2L}{k} \leq \xi(j_0 - 1)(2k + j_0 - 1) + 2L, \end{cases} \quad (120)$$

for any $k \geq 1$. Now, if we take $i = j_0$, using again (119), we deduce

$$\begin{cases} \lambda_k^{(3,2)} - \lambda_{k+j_0}^{(3,1)} = \frac{\epsilon_{k+j_0}}{k+j_0} + \frac{\epsilon_k}{k} > \frac{2\epsilon_1}{k+j_0}, \quad \forall k \geq 1, \\ \lambda_k^{(3,2)} - \lambda_{k+j_0}^{(3,1)} = \frac{\epsilon_{k+j_0}}{k+j_0} + \frac{\epsilon_k}{k} < \frac{2L}{k}, \quad \forall k \geq 1. \end{cases} \quad (121)$$

Finally, if $i \geq j_0 + 1$, equality (119) provides the formula

$$\lambda_{k+i}^{(3,1)} - \lambda_k^{(3,2)} = \xi(i - j_0)(2k + i) - \left(\frac{\epsilon_{k+i}}{k+i} + \frac{\epsilon_k}{k} \right), \quad \forall k \geq 1, \quad \forall i \geq j_0 + 1.$$

If we take $k_0 \geq 1$ (only depending on ξ , ρ and τ) such that

$$\frac{2L}{k_0} \leq \frac{\xi}{2} (2k_0 + j_0 + 1),$$

in particular, for any $k \geq k_0$ and $i \geq j_0 + 1$, one has

$$\frac{\epsilon_{k+i}}{k+i} + \frac{\epsilon_k}{k} < \frac{2L}{k} \leq \frac{2L}{k_0} \leq \frac{\xi}{2} (2k_0 + j_0 + 1) \leq \frac{\xi}{2} (i - j_0)(2k + i),$$

and

$$\begin{cases} \lambda_{k+i}^{(3,1)} - \lambda_k^{(3,2)} > \frac{\xi}{2} (i - j_0)(2k + i) \geq \frac{\xi}{2} (2k + j_0 + 1), \quad \forall k \geq k_0, \\ \lambda_{k+i}^{(3,1)} - \lambda_k^{(3,2)} < \xi(i - j_0)(2k + i). \end{cases} \quad (122)$$

The first consequence that we can obtain from (120)–(122) is the following one: for any $k \geq k_0$, we can write

$$\lambda_{k+j_0}^{(3,1)} < \lambda_k^{(3,2)} < \lambda_{k+1+j_0}^{(3,1)} < \lambda_{k+1}^{(3,2)} < \dots, \quad \forall k \geq k_0,$$

($\lambda_k^{(3,i)}$ is given in (17)). Thus, we can give an explicit expression of the elements of the increasing sequence $\Lambda^{(3)} = \{\Lambda_k\}_{k \geq 1}$ (see Proposition 5.15): if $1 \leq k \leq 2k_0 + j_0 - 2$, we define Λ_k such that

$$\{\Lambda_k\}_{1 \leq k \leq 2k_0 + j_0 - 2} \equiv \left\{ \lambda_k^{(3,1)} \right\}_{1 \leq k \leq k_0 + j_0 - 1} \cup \left\{ \lambda_k^{(3,2)} \right\}_{1 \leq k \leq k_0 - 1},$$

and $\Lambda_k < \Lambda_{k+1}$, for any $k : 1 \leq k \leq 2k_0 + j_0 - 3$. From the $(2k_0 + j_0 - 1)$ -th term, we define

$$\Lambda_{2k_0 + j_0 + 2s - 1} = \lambda_{k_0 + j_0 + s}^{(3,1)} \quad \text{and} \quad \Lambda_{2k_0 + j_0 + 2s} = \lambda_{k_0 + s}^{(3,2)}, \quad \forall s \geq 0.$$

Equivalently, in the case $k \geq 2k_0 + j_0 - 1$, we have

$$\begin{cases} \Lambda_k = \lambda_{\frac{1}{2}(k+j_0+1)}^{(3,1)}, & \text{if } k \geq 2k_0 + j_0 - 1 \text{ and } k + j_0 \text{ is odd,} \\ \Lambda_k = \lambda_{\frac{1}{2}(k-j_0)}^{(3,2)}, & \text{if } k \geq 2k_0 + j_0 - 1 \text{ and } k + j_0 \text{ is even.} \end{cases} \quad (123)$$

Our next objective will be to obtain appropriate estimates of the products \mathcal{P}_k (see (23)) for the sequence $\Lambda^{(3)}$. Recall that $\Lambda^{(3)} \in \mathcal{L}(0, \rho, q, p_0, p_1, p_2, \alpha)$ and satisfies (26), with p_0, p_1 and p_2 given in Proposition 5.15, and $q \geq 2, \rho$ and ν positive constants only depending on ξ, ρ and τ . We will reason for arbitrary $k \geq 2k_0 + j_0 + q - 2$ because if k is such that $1 \leq k < 2k_0 + j_0 + q - 2$, taking into account that $|\Lambda_k - \Lambda_n| > 0$ for any $k \neq n$ (assumption (114)), we deduce the existence of two positive constants c_0 and c_1 (only depending on ξ, ρ and τ) such that

$$0 < c_0 \leq \mathcal{P}_k \leq c_1, \quad \forall k : 1 \leq k < 2k_0 + j_0 + q - 2. \quad (124)$$

Let us then take $k \geq 2k_0 + j_0 + q - 2$ and $n \geq 1$ such that $1 \leq |k - n| < q$. In particular, $n \geq 2k_0 + j_0 - 1$ and we can use formulae (123) for the expression of Λ_k and Λ_n , and inequalities (120)–(122) for the corresponding indexes.

We will reason when $k + j_0$ is odd. A similar argument will provide the proof when $k + j_0$ is even. Indeed, if $k + j_0$ is odd, from (123), one has $\Lambda_k = \lambda_{\tilde{k}}^{(3,1)}$ and $\Lambda_{k+1} = \lambda_{\tilde{k}-j_0}^{(3,2)}$ with $\tilde{k} = \frac{1}{2}(k + j_0 + 1)$. Thus, we can apply (121) for $\tilde{k} - j_0$ and write

$$\frac{2\epsilon_1}{\frac{1}{2}(k + j_0 + 1)} \leq \Lambda_{k+1} - \Lambda_k \leq \frac{2L}{\frac{1}{2}(k - j_0 + 1)}.$$

On the other hand, let us take $n \neq k+1$ with $1 \leq |k - n| < q$. Using properties (120) and (122) and the expression of $\lambda_k^{(3,i)}$ (see (17) and (18)) and Λ_n (see (123)), it is not difficult to check the existence of positive constants c_0 and c_1 (as before, only depending on ξ, ρ and τ) such that

$$c_0 k \leq |\Lambda_k - \Lambda_n| \leq c_1 k, \quad \forall n \neq k+1 \quad \text{with} \quad 1 \leq |k - n| < q.$$

As a consequence of the previous inequalities, again, we deduce the existence of positive constants c_0 and c_1 (only depending on ξ, ρ and τ) such that

$$c_0 k^{2q-4} \leq \prod_{\{n \geq 1: 1 \leq |k-n| < q\}} |\Lambda_k - \Lambda_n| \leq c_1 k^{2q-4},$$

or, equivalently (see (23)),

$$\frac{c_0}{k^{2q-4}} \leq \mathcal{P}_k \leq \frac{c_1}{k^{2q-4}}, \quad \forall k \geq 2k_0 + j_0 + q - 2, \quad (125)$$

(c_0 and c_1 are new positive constants only depending on ξ, ρ and τ). We will use this inequality later.

Let us now revisit some properties on null controllability of system (12) with $L = L_3$ proved in [22]: Given $T > 0$ and $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$, there exists a control $v \in L^2(0, T)$ such that the solution of (12) with $L = L_3$ satisfies $y(\cdot, T) = 0$ in $(0, \pi)$ if and only if $v \in L^2(0, T)$ solves the moment problem

$$\int_0^T e^{-\Lambda_k t} v(T-t) dt = e^{-\Lambda_k T} m_k, \quad \forall k \geq 1. \quad (126)$$

In the previous equality $\Lambda^{(3)} = \{\Lambda_k\}_{k \geq 1} = \{\lambda_k^{(3,1)}, \lambda_k^{(3,2)}\}_{k \geq 1}$ ($\lambda_k^{(3,i)}$ is given in (17)) and m_k only depends on y_0 and satisfies

$$|m_k| \leq Ck \|y_0\|_{H^{-1}}, \quad \forall k \geq 1, \quad (127)$$

with $C > 0$ only depending on ξ , ρ and τ .

On the other hand, the sequence $\Lambda^{(3)}$ belongs to $\mathcal{L}(0, \rho, q, p_0, p_1, p_2, \alpha)$ and satisfies (26) (p_0 , p_1 and p_2 are given in Proposition 5.15, $q \geq 2$, and ρ and ν are positive constants only depending on ξ , ρ and τ). Then, we can apply Theorems 1.7 and 1.9 to the sequence $\Lambda^{(3)}$. We deduce the existence of a biorthogonal family $\{q_k\}_{k \geq 1}$ to the exponentials $\{e_k\}_{k \geq 1}$ (see (1)) associated to the sequence $\Lambda^{(3)}$ satisfying (22) and (27).

Let us first prove that, under the assumptions of Theorem 5.17, system (12) with $L = L_3$ is null controllable at any time $T > 0$ and satisfies the second inequality in (118). To this end, we will solve the previous moment problem for any $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$. An explicit solution of this problem is

$$v(t) = \sum_{k \geq 1} e^{-\Lambda_k T} m_k q_k(T-t), \quad \forall t \in (0, T).$$

Since q_k , \mathcal{P}_k and m_k respectively satisfy (22), (124) or (125), and (127), we can prove that the previous series is absolutely convergent in $L^2(0, T)$ and provide an estimate of the L^2 -norm of v . Indeed,

$$\begin{cases} |m_k| \|q_k\|_{L^2(0, T)} \leq Ck e^{C\sqrt{\Lambda_k}} e^{C/T} \mathcal{P}_k \leq C e^{C\sqrt{\Lambda_k}} e^{C/T} \|y_0\|_{H^{-1}} \\ \leq C e^{C/T} e^{\frac{C^2}{2T} + \frac{\tau}{2}\Lambda_k} = C e^{C/T} e^{\frac{\tau}{2}\Lambda_k}, \quad \forall k \geq 1, \end{cases}$$

for a new positive constant C , only depending on ξ , ρ and τ . If we use (43) (p_2 and α are given in the statement of Proposition 5.15), we deduce that $v \in L^2(0, T)$ and

$$\begin{cases} \|v\|_{L^2(0, T)} \leq C e^{C/T} \sum_{k=1}^{\infty} e^{-\frac{\tau}{2}\Lambda_k} \leq C e^{C/T} \sum_{k=1}^{\alpha} e^{-\frac{\tau}{2}\Lambda_k} \sum_{k > \alpha} e^{-\frac{\tau}{8}\xi(k-\alpha)^2} \\ \leq C e^{C/T} \int_{\mathbb{R}} e^{-\frac{\tau}{8}\xi(x-\alpha)^2} dx = C \sqrt{\frac{8\pi}{\xi T}} e^{C/T}. \end{cases}$$

From this inequality we deduce the estimate from above of $\mathcal{K}(T)$ in (118).

Let us now prove the first inequality in (118). To this end, we will reason as in Subsection 5.1 and, to be precise, as in Proposition 5.4 and the first point of the proof of Theorem 5.10. We first construct the sequence $\{s_k\}_{k \geq 1}$ biorthogonal to the exponentials $\{e_k\}_{k \geq 1}$ associated to the sequence $\Lambda^{(3)}$. Given $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$, we know that the null control with minimal L^2 -norm for system (12) with $L = L_3$ (see (16)) associated to $y_0 \in H^{-1}(0, \pi; \mathbb{R}^2)$ is

$$u(t) = \sum_{k \geq 1} e^{-\Lambda_k T} m_k s_k(T-t), \quad \forall t \in (0, T),$$

where m_k depends on y_0 and appears in the corresponding moment problem (126).

Let us take $\ell = \max\{3, q\}$ and $y_0 = \Psi_\ell$, with Ψ_ℓ the eigenvector of L_3 associated to Λ_ℓ with $\|\Psi_\ell\|_{H^{-1}} = 1$ (for the expression of Ψ_ℓ see Proposition 3.1 in [22]). In this case, the corresponding null control with minimal L^2 -norm is

$$u(t) = e^{-\Lambda_\ell t} m_\ell s_\ell(T-t), \quad \forall t \in (0, T),$$

and $\mathcal{K}(T) \geq \|u\|_{L^2(0, T)} = e^{-\Lambda_\ell T} |m_\ell| \|s_\ell\|$ (m_ℓ only depends on ρ , ξ and τ). If we use inequalities (27), for the function s_ℓ , and (124) or (125) for $k = \ell$, we deduce the existence of a positive constant C , only depending on ρ , ξ and τ , such that

$$\mathcal{K}(T) \geq C \mathcal{B}_\ell e^{\frac{1}{T^v}} = C \frac{(\nu T)^{\ell+1}}{(1 + \nu T)^{2\ell+q+1}} \sqrt{|\Lambda_1| + \frac{1}{2T}} e^{\frac{1}{T^v}}, \quad \forall T > 0.$$

Finally, there exist $C > 0$, only depending on ρ , ξ and τ , such that

$$\frac{(\nu T)^{\ell+1}}{(1 + \nu T)^{2\ell+q+1}} \sqrt{|\Lambda_1| + \frac{1}{2T}} \geq C e^{\frac{-1}{2T^v}}, \quad \forall T \in (0, 1].$$

Therefore,

$$\mathcal{K}(T) \geq C e^{\frac{1}{2T^v}}, \quad \forall T \in (0, 1],$$

for a new constant $C > 0$ only depending on ρ , ξ and τ . This proves (118) and finalizes the proof of Theorem 5.17 \square

Theorem 5.17 in particular provides a local boundary exact controllability result to the trajectory $(0, c)$ ($c = \pm 1$) for the nonlinear system (15) under assumption (114). One has:

Theorem 5.18. *Let us consider ξ, τ and ρ three positive numbers satisfying (114), and let us fix $T > 0$ and $c = -1$ or $c = 1$. Then, there exists $\varepsilon > 0$ such that, for any $(\theta_0, \phi_0) \in H^{-1}(0, \pi) \times (c + H_0^1(0, \pi))$ fulfilling*

$$\|\theta_0\|_{H^{-1}} + \|\tilde{\phi}_0 - c\|_{H_0^1} \leq \varepsilon,$$

there exists $v \in L^2(0, T)$ for which system (15) has a unique solution

$$(\theta, \phi) \in [L^2(Q_T) \cap C^0([0, T]; H^{-1}(0, \pi; \mathbb{R}^2))] \times C^0(\overline{Q}_T)$$

which satisfies

$$\theta(\cdot, T) = 0 \quad \text{and} \quad \phi(\cdot, T) = c \quad \text{in } (0, \pi).$$

In order to obtain the proof of the previous local controllability result for system (15), it is enough to follow the reasoning of the reference [22] that combines inequality (118) with the general methodology developed in [32]. For further details, see [22].

Remark 5.19. Theorem 5.18 is valid under the only assumption (114). In this sense, Theorem 5.18 generalizes the local controllability result for system (15) stated in [22] where the authors prove the same result under assumptions (114) and (115). \square

Acknowledgments. Part of this work has been carried out when the second author was developing the research period of the Doc-Course ‘‘Partial Differential Equations: Analysis, Numerics and Control’’ in the Institute of Mathematics of the University of Seville (IMUS). The authors want to thank the IMUS for providing a great working framework. We would also like to thank Prof. Assia Benabdallah for her fruitful discussions that have helped to improve this work.

Appendix A. Proof of Propositions 2.9, 2.11, 2.13 and 2.14.

A.1. Proof of Proposition 2.9. Let us take $\Lambda = \{\Lambda_k\}_{k \geq 1} \subset (0, \infty)$, a sequence under the assumptions of the proposition. It is clear that the sequence Λ satisfies (H1)–(H4) for $\beta = 0$.

Let us first see that property (44) implies property (H6). Indeed, given $r > 0$, one has $\mathcal{N}(r) = k$ if and only if $\Lambda_k \leq r$ and $\Lambda_{k+1} > r$. Since the sequence Λ satisfies

$$\gamma_0 k + \sqrt{\Lambda_1} - \gamma_0 \leq \sqrt{\Lambda_k} \leq \gamma_1 k + \sqrt{\Lambda_1} - \gamma_1, \quad \forall k \geq 1, \quad (128)$$

we can write

$$\gamma_0 k + \sqrt{\Lambda_1} - \gamma_0 \leq \sqrt{r} \quad \text{and} \quad \sqrt{r} < \gamma_1(k+1) + \sqrt{\Lambda_1} - \gamma_1,$$

i.e., condition (H6) holds with p_0, p_1, p_2 and α as in the statement of the proposition.

Let us now see that we can deduce (H5) from property (44). First, one has

$$\sqrt{\Lambda_k} - \sqrt{\Lambda_n} \geq \gamma_0(k-n), \quad \forall k, n : k \geq n.$$

As a direct consequence, one also has

$$\Lambda_k - \Lambda_n \geq \gamma_0(k-n) \left(\sqrt{\Lambda_k} + \sqrt{\Lambda_n} \right), \quad \forall k, n : k \geq n,$$

that together with (128) provides

$$\Lambda_k - \Lambda_n \geq \gamma_0^2(k^2 - n^2) + 2\gamma_0(k-n) \left(\sqrt{\Lambda_1} - \gamma_0 \right),$$

for any $k, n : k \geq n$. If $\sqrt{\Lambda_1} \geq \gamma_0$, clearly one gets (H5) with ρ as in the statement. Otherwise, $\sqrt{\Lambda_1} < \gamma_0$ and, from the previous inequality, we deduce

$$\frac{\Lambda_k - \Lambda_n}{k^2 - n^2} \geq \gamma_0^2 - \frac{2\gamma_0}{k+n} \left(\gamma_0 - \sqrt{\Lambda_1} \right) \geq \gamma_0^2 - \frac{2}{3}\gamma_0 \left(\gamma_0 - \sqrt{\Lambda_1} \right),$$

for any $k, n : k \geq n$. In this case we also deduce (H5) with ρ given in the statement.

Finally, let us prove (26). Reasoning as before, we can write

$$\Lambda_k - \Lambda_n = \left(\sqrt{\Lambda_k} - \sqrt{\Lambda_n} \right) \left(\sqrt{\Lambda_k} + \sqrt{\Lambda_n} \right) \leq \gamma_1(k-n) \left(\sqrt{\Lambda_k} + \sqrt{\Lambda_n} \right),$$

for any $k, n : k \geq n$, that together with (128) gives

$$\Lambda_k - \Lambda_n \leq \gamma_1^2(k^2 - n^2) + 2\gamma_1(k-n) \left(\sqrt{\Lambda_1} - \gamma_1 \right), \quad \forall k, n : k \geq n.$$

In the case in which $\sqrt{\Lambda_1} \leq \gamma_1$, we deduce (26) with $\rho = \gamma_1^2$. Otherwise,

$$\frac{\Lambda_k - \Lambda_n}{k^2 - n^2} \leq \gamma_1^2 + \frac{2\gamma_1}{k^2 - n^2} \left(\sqrt{\Lambda_1} - \gamma_1 \right) \leq \gamma_1^2 + \frac{2}{3}\gamma_1 \left(\sqrt{\Lambda_1} - \gamma_1 \right),$$

for any $k, n : k \geq n$. We also obtain (26) in this case with ρ given in the statement.

This finalizes the proof of Proposition 2.9. \square

A.2. Proof of Proposition 2.11. Let us consider $\{\lambda_k^{(1)}\}_{k \geq 1}$ and $\{\lambda_k^{(2)}\}_{k \geq 1}$, two sequences satisfying (45) and (47). It is clear that, from (47) and the third condition in (45), the sequence $\{\lambda_k^{(1)}\}_{k \geq 1} \cup \{\lambda_k^{(2)}\}_{k \geq 1}$ can be rearranged as an increasing sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$.

First, let us see that (26) holds and $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ for appropriate positive constants $\rho, q, p_0, p_1, p_2, \alpha$ and ν . It is clear that Λ satisfies (H1)–(H4).

On the other hand, using that $\lambda_k^{(1)} \neq \lambda_n^{(2)}$ for any $k, n \geq 1$, we also have

$$\mathcal{N}(r) = \# \left\{ k : \lambda_k^{(1)} \leq r \right\} + \# \left\{ k : \lambda_k^{(2)} \leq r \right\} := \mathcal{N}_1(r) + \mathcal{N}_2(r), \quad \forall r > 0,$$

where $\mathcal{N}(r)$ is given in (21). Using the first property in (45) we infer

$$\frac{1}{\pi_i^2}k^2 - c_1k \leq \lambda_k^{(i)} \leq \frac{1}{\pi_i^2}k^2 + c_1k, \quad \forall k \geq 1, \quad i = 1, 2.$$

Therefore, we can follow the arguments in Remark 2.5 and deduce

$$-1 - \frac{1}{2}\pi_i^2c_1 + \pi_i\sqrt{r} < \mathcal{N}_i(r) \leq \pi_i\sqrt{r} + c_1\pi_i^2, \quad i = 1, 2.$$

Coming back to the expression of $\mathcal{N}(r)$, we finally obtain

$$-2 - \frac{1}{2}c_1(\pi_1^2 + \pi_2^2) + (\pi_1 + \pi_2)\sqrt{r} < \mathcal{N}(r) \leq (\pi_1 + \pi_2)\sqrt{r} + c_1(\pi_1^2 + \pi_2^2), \quad \forall r > 0.$$

Thus, condition (H6) holds with p_0, p_1, p_2 and α as in the statement of Proposition 2.11. Finally, applying Proposition 2.4, we also have that the sequence Λ satisfies (26) and $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$ with the parameters $\rho, q, p_0, p_1, p_2, \alpha$ and ν given in the statement of Proposition 2.11.

Let us now check the gap condition (5). Taking into account property (47), we just have to check the following property

$$\left| \lambda_k^{(1)} - \lambda_n^{(2)} \right| \geq c_2 > 0, \quad \forall k, n \geq 1,$$

and this will be deduced from the third condition in (45). Indeed, this condition implies

$$\left| \lambda_k^{(1)} - \lambda_n^{(2)} \right| \geq \frac{r}{k} \left(\sqrt{\lambda_k^{(1)}} + \sqrt{\lambda_n^{(2)}} \right) \geq \frac{r}{k} \sqrt{\lambda_k^{(1)}}, \quad \forall k \geq 1.$$

If $k \leq 2c_1p_1^2$, from the previous inequality we deduce the existence of a constant $c > 0$ such that

$$\left| \lambda_k^{(1)} - \lambda_n^{(2)} \right| \geq c, \quad \forall n \in \mathbb{N}.$$

If $k > 2c_1p_1^2$, then we can apply the first assumption in (45) and deduce

$$\left| \lambda_k^{(1)} - \lambda_n^{(2)} \right| \geq \frac{r}{k} \sqrt{\lambda_k^{(1)}} \geq \frac{r}{k} \sqrt{\frac{k^2}{\pi_1^2} - c_1k} = \frac{r}{k} \sqrt{\frac{k}{\pi_1^2} (k - c_1\pi_1^2)} \geq \frac{r}{k} \sqrt{\frac{k^2}{2\pi_1^2}} = \frac{r}{\sqrt{2}\pi_1}.$$

This proves (5) and ends the proof of the result. \square

A.3. Proof of Proposition 2.13. Let us consider the sequence $\Lambda = \{k^2\}_{k \geq 1} \cup \{dk^2\}_{k \geq 1}$ with $d > 0$. Thanks to assumption $\sqrt{d} \notin \mathbb{Q}$, it is clear that $k^2 \neq dn^2$ for any $k, n \geq 1$. So, the sequence $\Lambda = \{k^2\}_{k \geq 1} \cup \{dk^2\}_{k \geq 1}$ can be rearranged as an increasing sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$ that satisfies (H1)–(H4) with $\beta = 0$. On the other hand,

$$\mathcal{N}(r) = \# \{k : k^2 \leq r\} + \# \{k : dk^2 \leq r\} = \lfloor \sqrt{r} \rfloor + \left\lfloor \frac{\sqrt{r}}{\sqrt{d}} \right\rfloor, \quad \forall r > 0,$$

i.e.,

$$-2 + \left(1 + \frac{1}{\sqrt{d}}\right) \sqrt{r} \leq \mathcal{N}(r) \leq \left(1 + \frac{1}{\sqrt{d}}\right) \sqrt{r}, \quad \forall r > 0.$$

Thus, condition (H6) holds with $p_1 = p_2 = p$ and α given in (49).

As a direct consequence of Proposition 2.4 we can deduce (26) and

$$\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha),$$

(q , ρ and ν given in (36)). We will provide better values of these parameters using the expression of Λ . Indeed, if we take $r = \Lambda_k$, $k \geq 1$, one has $k = \mathcal{N}(\Lambda_k)$ and

$$k = \mathcal{N}(\Lambda_k) = \left\lfloor \sqrt{\Lambda_k} \right\rfloor + \left\lfloor \frac{\sqrt{\Lambda_k}}{\sqrt{d}} \right\rfloor, \quad \forall k \geq 1.$$

Observe that if $\Lambda_k = n_k^2$ for some $n_k \geq 1$, from the previous inequality we deduce

$$k = n_k + \left\lfloor \frac{\sqrt{\Lambda_k}}{\sqrt{d}} \right\rfloor.$$

Using that $x - 1 \leq \lfloor x \rfloor \leq x$, for any $x > 0$, the previous inequality provides,

$$\sqrt{\Lambda_k} + \frac{\sqrt{\Lambda_k}}{\sqrt{d}} - 1 = n_k + \frac{\sqrt{\Lambda_k}}{\sqrt{d}} - 1 \leq n_k + \left\lfloor \frac{\sqrt{\Lambda_k}}{\sqrt{d}} \right\rfloor = k \leq n_k + \frac{\sqrt{\Lambda_k}}{\sqrt{d}} = \sqrt{\Lambda_k} + \frac{\sqrt{\Lambda_k}}{\sqrt{d}},$$

and

$$k \leq \left(1 + \frac{1}{\sqrt{d}}\right) \sqrt{\Lambda_k} \leq k + 1, \quad \forall k \geq 1. \quad (129)$$

The same property can be proved in the case in which $\Lambda_k = dn_k^2$ for some $n_k \geq 1$.

Let us now prove conditions (H5), with $q = 2$, and (26). If $k - n \geq 2$, from (129), one has

$$\left(1 + \frac{1}{\sqrt{d}}\right)^2 \frac{\Lambda_k - \Lambda_n}{k^2 - n^2} \geq \frac{k^2 - (n+1)^2}{k^2 - n^2} = \left(1 + \frac{1}{k+n}\right) \left(1 - \frac{1}{k-n}\right) \geq \frac{5}{8}.$$

Thus, (H5) holds with ρ given in (49). On the other hand, if $k > n$, we deduce (see (129))

$$\left(1 + \frac{1}{\sqrt{d}}\right)^2 \frac{\Lambda_k - \Lambda_n}{k^2 - n^2} \leq \frac{(k+1)^2 - n^2}{k^2 - n^2} = \left(1 + \frac{1}{k+n}\right) \left(1 + \frac{1}{k-n}\right) \leq \frac{8}{3},$$

and property (26) with ν given in (49). This ends the proof of Proposition 2.13. \square

A.4. Proof of Proposition 2.14. Let us consider two sequences $\Lambda_1 = \{\lambda_k^{(1)}\}_{k \geq 1}$ and $\Lambda_2 = \{\lambda_k^{(2)}\}_{k \geq 1}$ under the conditions of Proposition 2.14. In particular, the sequence $\Lambda_1 \cup \Lambda_2$ can be rearranged as a positive increasing sequence $\Lambda = \{\Lambda_k\}_{k \geq 1}$. Let us see that $\Lambda \in \mathcal{L}(\beta, \rho, q, p_0, p_1, p_2, \alpha)$, for $\beta = 0$ and appropriate positive constants ρ, q, p_0, p_1, p_2 and α , and (26) holds for $\nu > 0$.

First, it is clear that Λ satisfies (H1)–(H4) ($\beta = 0$). As above, using that $\lambda_k^{(1)} \neq \lambda_n^{(2)}$ for any $k, n \geq 1$, we also have

$$\mathcal{N}(r) = \#\left\{k : \lambda_k^{(1)} \leq r\right\} + \#\left\{k : \lambda_k^{(2)} \leq r\right\} := \mathcal{N}_1(r) + \mathcal{N}_2(r), \quad \forall r > 0.$$

From Remark 1.5 we deduce the following property:

$$\mathcal{N}_1(r - \varepsilon_0) \leq \mathcal{N}_2(r) \leq \mathcal{N}_1(r + \varepsilon_0), \quad \forall r > 0, \quad (130)$$

(in the previous inequality we have taken $\mathcal{N}_1(r - \varepsilon_0) = 0$ when $r \leq \varepsilon_0$). Indeed, given $r > 0$, if $k_2 = \mathcal{N}_2(r)$, then $\lambda_{k_2}^{(2)} \leq r$ and $\lambda_{k_2+1}^{(2)} > r$. In particular,

$$\lambda_{k_2}^{(1)} - \varepsilon_0 \leq \lambda_{k_2}^{(2)} \leq r \quad \text{and} \quad r < \lambda_{k_2+1}^{(2)} \leq \lambda_{k_2+1}^{(1)} + \varepsilon_0,$$

($\varepsilon_0 = \sup_{k \geq 1} |\varepsilon_k|$) and $\lambda_{k_2}^{(1)} \leq r + \varepsilon_0$ and $r - \varepsilon_0 < \lambda_{k_2+1}^{(1)}$. Applying item 2 of Remark 1.5, property (130) can be easily deduced.

Recall that $\Lambda_1 = \left\{ \lambda_k^{(1)} \right\}_{k \geq 1} \in \mathcal{L}(0, \rho_1, 1, \pi_0, \pi_1, \pi_2, \alpha_1)$. Thus, from (H6), we deduce $\pi_1 \sqrt{r} - \alpha_1 \leq \mathcal{N}_1(r) \leq \pi_2 \sqrt{r} + \alpha_1$ for any $r > 0$. Combining this inequality and the expression of $\mathcal{N}(r)$ with (130), we obtain

$$\begin{cases} \pi_1 \sqrt{r} - \alpha_1 \leq \mathcal{N}(r) \leq \pi_2 \sqrt{r} + \pi_2 \sqrt{r + \varepsilon_0} + 2\alpha_1, & \text{if } r \leq \varepsilon_0, \\ \pi_1 \sqrt{r} + \pi_1 \sqrt{r - \varepsilon_0} - 2\alpha_1 \leq \mathcal{N}(r) \leq \pi_2 \sqrt{r} + \pi_2 \sqrt{r + \varepsilon_0} + 2\alpha_1, & \text{if } r > \varepsilon_0. \end{cases}$$

Now, from the previous property and taking into account inequalities (34), it is easy to deduce that $\mathcal{N}(r)$ satisfies

$$\begin{cases} \pi_1 \sqrt{r} - \alpha_1 \leq \mathcal{N}(r) \leq 2\pi_2 \sqrt{r} + \pi_2 \sqrt{\varepsilon_0} + 2\alpha_1, & \text{if } r \leq \varepsilon_0, \\ 2\pi_1 \sqrt{r} - \pi_1 \sqrt{\varepsilon_0} - 2\alpha_1 \leq \mathcal{N}(r) \leq 2\pi_2 \sqrt{r} + \pi_2 \sqrt{\varepsilon_0} + 2\alpha_1, & \text{if } r > \varepsilon_0. \end{cases}$$

In particular,

$$2\pi_1 \sqrt{r} - \pi_1 \sqrt{\varepsilon_0} - 2\alpha_1 \leq \mathcal{N}(r) \leq 2\pi_2 \sqrt{r} + \pi_2 \sqrt{\varepsilon_0} + 2\alpha_1, \quad \forall r > 0.$$

Therefore, condition (H6) holds with p_1 , p_2 and α as in the statement of Proposition 2.14.

Let us now see that the sequence Λ satisfies (H5) and (26) with $q = 2$ and appropriate positive parameters ρ and ν . To this end, we will use that Λ_1 is in $\mathcal{L}(0, \rho_1, 1, \pi_0, \pi_1, \pi_2, \alpha_1)$ ($q = 1$) and satisfies (26), for $\nu_1 \in (0, \infty)$ or, more precisely, we will use

$$\rho_1 |k^2 - n^2| \leq \left| \lambda_k^{(1)} - \lambda_n^{(1)} \right| \leq \nu_1 |k^2 - n^2|, \quad \forall k, n \in \mathbb{N}. \quad (131)$$

The sequence $\{\varepsilon_k\}_{k \geq 1}$ is bounded. So, there exists $k_0 \geq 1$, depending on ρ_1 and ε_0 , such that

$$|\varepsilon_k| \leq \varepsilon_0 \leq \frac{\rho_1}{4} (2k - 1) \leq \frac{\rho_1}{4} |k^2 - n^2|, \quad \forall k, n \geq 1 : k \geq k_0, n \neq k.$$

With this value of k_0 and (131), written for k and n , we obtain

$$\begin{aligned} \left| \lambda_k^{(1)} - \lambda_n^{(1)} \right| &\geq \rho_1 |k^2 - n^2| \geq \frac{\rho_1}{2} |k^2 - n^2|, \\ \left| \lambda_k^{(1)} - \lambda_n^{(2)} \right| &\geq \left| \lambda_k^{(1)} - \lambda_n^{(1)} \right| - \varepsilon_0 \geq \rho_1 |k^2 - n^2| - \frac{\rho_1}{4} |k^2 - n^2| \geq \frac{\rho_1}{2} |k^2 - n^2|, \\ \left| \lambda_k^{(2)} - \lambda_n^{(2)} \right| &\geq \left| \lambda_k^{(1)} - \lambda_n^{(1)} \right| - 2\varepsilon_0 \geq \rho_1 |k^2 - n^2| - \frac{\rho_1}{2} |k^2 - n^2| \geq \frac{\rho_1}{2} |k^2 - n^2|, \end{aligned}$$

with $k \geq k_0$ and $n \neq k$, i.e.,

$$\left| \lambda_k^{(i)} - \lambda_n^{(j)} \right| \geq \frac{\rho_1}{2} |k^2 - n^2|, \quad \forall k, n \geq 1 : k \geq k_0, n \neq k, \quad \forall i, j \in \{1, 2\}. \quad (132)$$

As a consequence of (132), we also obtain $\lambda_k^{(1)} < \lambda_{k+1}^{(2)}$ and $\lambda_k^{(2)} < \lambda_{k+1}^{(1)}$, for any $k \geq k_0$. This provides the following explicit formula for the terms of the increasing sequence Λ when $k \geq 2k_0 - 1$:

$$\Lambda_k = \begin{cases} \min \left\{ \lambda_\ell^{(1)}, \lambda_\ell^{(2)} \right\}, & \text{if } k = 2\ell - 1, \\ \max \left\{ \lambda_\ell^{(1)}, \lambda_\ell^{(2)} \right\}, & \text{if } k = 2\ell. \end{cases} \quad (133)$$

We are going to use the previous expression of the terms Λ_k in order to prove condition (H5) with $q = 2$. Recall that the sequence Λ is real and increasing. Then,

$$\frac{\Lambda_k - \Lambda_n}{k^2 - n^2} > 0, \quad \forall k, n \geq 1 : k \geq n + 1.$$

Assume that, for every $n \in \{1, \dots, 2k_0 - 2\}$ fixed, one has

$$\liminf_{k \rightarrow \infty} \frac{\Lambda_k - \Lambda_n}{k^2 - n^2} \geq \frac{\rho_1}{4} \quad \text{and} \quad \limsup_{k \rightarrow \infty} \frac{\Lambda_k - \Lambda_n}{k^2 - n^2} \leq \frac{\nu_1}{4}. \quad (134)$$

Then, there exists a positive constant $\tilde{\rho}$, only depending on k_0 and ρ_1 or, equivalently, on ρ_1 and ε_0 , such that

$$\Lambda_k - \Lambda_n \geq \tilde{\rho}(k^2 - n^2), \quad \forall k, n \in \mathbb{N} : 1 \leq n \leq 2k_0 - 2 \text{ and } n \leq k.$$

In this way, we have proved condition (H5) for $q = 1$ and $k, n \in \mathbb{N}$ such that $1 \leq n \leq 2k_0 - 2$ and $n \leq k$. We will prove (134) below.

Let us now see that the sequence Λ satisfies (H5), with $q = 2$ and an appropriate value of the parameter ρ , when $k, n \geq 1$ with $k \geq n + 2$ and $n \geq 2k_0 - 1$. We divide the proof into four cases:

1. Assume that $k = 2\ell - 1$ and $n = 2m - 1$, with $\ell, m \geq k_0$ and $k - n \geq 2$. In particular, $\ell - m \geq 1$, $\Lambda_k = \lambda_\ell^{(i)}$ and $\Lambda_n = \lambda_m^{(j)}$, with $i, j \in \{1, 2\}$. Thus, from (132)

$$\begin{aligned} \Lambda_k - \Lambda_n &= \lambda_\ell^{(i)} - \lambda_m^{(j)} \geq \frac{\rho_1}{2} (\ell^2 - m^2) = \frac{\rho_1}{8} \left[(k+1)^2 - (n+1)^2 \right] \\ &= \frac{\rho_1}{8} (k+n+2)(k-n) \geq \frac{\rho_1}{8} (k^2 - n^2). \end{aligned}$$

2. Assume now that $k = 2\ell - 1$ and $n = 2m$, with $\ell, m \geq k_0$ and $k - n \geq 2$. In particular, $\ell - m \geq 3/2$, $\Lambda_k = \lambda_\ell^{(i)}$ and $\Lambda_n = \lambda_m^{(j)}$, with $i, j \in \{1, 2\}$, and we can apply (132):

$$\Lambda_k - \Lambda_n = \lambda_\ell^{(i)} - \lambda_m^{(j)} \geq \frac{\rho_1}{2} (\ell^2 - m^2) = \frac{\rho_1}{8} \left[(k+1)^2 - n^2 \right] \geq \frac{\rho_1}{8} (k^2 - n^2).$$

3. If $k = 2\ell$ and $n = 2m$, with $\ell, m \geq k_0$ and $k - n \geq 2$, then $\ell - m \geq 1$, $\Lambda_k = \lambda_\ell^{(i)}$ and $\Lambda_n = \lambda_m^{(j)}$, with $i, j \in \{1, 2\}$. Applying again (132), we get

$$\Lambda_k - \Lambda_n = \lambda_\ell^{(i)} - \lambda_m^{(j)} \geq \frac{\rho_1}{2} (\ell^2 - m^2) = \frac{\rho_1}{8} (k^2 - n^2).$$

4. In the case $k = 2\ell$ and $n = 2m - 1$, with $\ell, m \geq k_0$ and $k - n \geq 2$ we will use the inequality

$$k^2 - (n+1)^2 \geq \frac{1}{2} (k^2 - n^2)$$

which is valid for any $k, n \geq 1$ such that $k \geq n + 2$. Also, $\ell - m \geq 1/2$, i.e., $\ell - m \geq 1$ and we can apply (132). As before, $\Lambda_k = \lambda_\ell^{(i)}$ and $\Lambda_n = \lambda_m^{(j)}$, with $i, j \in \{1, 2\}$, and

$$\Lambda_k - \Lambda_n = \lambda_\ell^{(i)} - \lambda_m^{(j)} \geq \frac{\rho_1}{2} (\ell^2 - m^2) = \frac{\rho_1}{8} (k^2 - (n+1)^2) \geq \frac{\rho_1}{16} (k^2 - n^2).$$

We can conclude that property (H5) holds for the sequence Λ with $q = 2$ and

$$\rho = \min \left\{ \tilde{\rho}, \frac{\rho_1}{16} \right\}.$$

Recall that the constant $\tilde{\rho}$ only depends on ρ_1 and ε_0 . Therefore, ρ only depends on ρ_1 and ε_0 .

The next task will be the proof of (134). To this end, let us fix n such that $1 \leq n \leq 2k_0 - 2$ and $k \geq 2k_0 - 1$. Then $k = 2\ell$ or $k = 2\ell - 1$ with $\ell \geq k_0$. In both cases, $\Lambda_k = \lambda_\ell^{(i)}$, with $i \in \{1, 2\}$, and we can write (see (131)):

$$\left\{ \begin{array}{l} \Lambda_k - \Lambda_n \geq \lambda_\ell^{(1)} - \varepsilon_0 - \Lambda_{2k_0-2} \geq \rho_1 (\ell^2 - 1) + \lambda_1^{(1)} - \varepsilon_0 - \Lambda_{2k_0-2} \\ \geq \rho_1 \left(\frac{k^2}{4} - 1 \right) - \varepsilon_0 - \Lambda_{2k_0-2}, \\ \Lambda_k - \Lambda_n \leq \lambda_\ell^{(1)} + \varepsilon_0 - \Lambda_1 \leq \nu_1 (\ell^2 - 1) + \lambda_1^{(1)} + \varepsilon_0 - \Lambda_1 \\ \leq \nu_1 \left(\frac{(k+1)^2}{4} - 1 \right) + \lambda_1^{(1)} + \varepsilon_0 - \Lambda_1. \end{array} \right.$$

This proves (134).

In order to finish the proof of Proposition 2.14, let us check that the sequence Λ fulfills condition (26) for an appropriate $\nu > 0$. The proof is very close to that of condition (H5). First, one has

$$|\varepsilon_k| \leq \varepsilon_0 \leq \varepsilon_0 (2k - 1) \leq \varepsilon_0 |k^2 - n^2|, \quad \forall k, n \geq 1 : n \neq k.$$

From this inequality and (26), for $\nu_1 \in (0, \infty)$, we deduce

$$\left\{ \begin{array}{l} |\lambda_k^{(1)} - \lambda_n^{(1)}| \leq \nu_1 |k^2 - n^2| \leq (\nu_1 + \varepsilon_0) |k^2 - n^2|, \\ |\lambda_k^{(1)} - \lambda_n^{(2)}| \leq |\lambda_k^{(1)} - \lambda_n^{(1)}| + \varepsilon_0 \leq (\nu_1 + \varepsilon_0) |k^2 - n^2|, \\ |\lambda_k^{(2)} - \lambda_n^{(2)}| \leq (\nu_1 + 2\varepsilon_0) |k^2 - n^2|, \end{array} \right. \quad (135)$$

for any $k, n \geq 1$ with $k \neq n$.

Let us now prove condition (26) for the sequence Λ . As before, from the second property in (134) we deduce the existence of a positive constant $\tilde{\nu}$, only depending on k_0 and ν_1 , such that

$$\Lambda_k - \Lambda_n \leq \tilde{\nu} (k^2 - n^2), \quad \forall k, n \in \mathbb{N} : 1 \leq n \leq 2k_0 - 2 \text{ and } n \leq k.$$

Let us now see inequality (26) when $k, n \in \mathbb{N}$ are such that $2k_0 - 1 \leq n \leq k$. Recall that, in this case, we have an explicit formula of the terms of the sequence Λ (see (133)). Let us first consider the case $n \geq 2k_0 - 1$ and $k = n + 1$. Thus,

$$\left\{ \begin{array}{l} \Lambda_{n+1} - \Lambda_n = \Lambda_{2\ell} - \Lambda_{2\ell-1} = |\varepsilon_\ell| \leq \varepsilon_0 \left((n+1)^2 - n^2 \right), \\ \Lambda_{n+1} - \Lambda_n = \Lambda_{2\ell+1} - \Lambda_{2\ell} \leq \lambda_{\ell+1}^{(1)} - \lambda_\ell^{(1)} \leq \nu_1 (2\ell + 1) \leq \nu_1 \left((n+1)^2 - n^2 \right). \end{array} \right.$$

In the general case, i.e., when $k, n \in \mathbb{N}$ are such that $2k_0 - 1 \leq n \leq k$ with $k \geq n + 2$, we can repeat the arguments above and deduce inequality (26). Indeed, as a consequence of (135), we deduce

1. If $k = 2\ell - 1$ and $n = 2m - 1$, with $\ell, m \geq k_0$ and $k - n \geq 2$, then, $\ell - m \geq 1$, $\Lambda_k = \lambda_\ell^{(i)}$ and $\Lambda_n = \lambda_m^{(j)}$, with $i, j \in \{1, 2\}$. (135) implies

$$\begin{aligned} \Lambda_k - \Lambda_n &= \lambda_\ell^{(i)} - \lambda_m^{(j)} \leq (\nu_1 + 2\varepsilon_0) (\ell^2 - m^2) = \frac{\nu_1 + 2\varepsilon_0}{4} \left[(k+1)^2 - (n+1)^2 \right] \\ &= \frac{\nu_1 + 2\varepsilon_0}{4} (k+n+2)(k-n) \leq \frac{\nu_1 + 2\varepsilon_0}{2} (k^2 - n^2). \end{aligned}$$

2. Assume now that $k = 2\ell - 1$ and $n = 2m$, with $\ell, m \geq k_0$ and $k - n \geq 2$. In this case, $\ell - m \geq 3/2$, $\Lambda_k = \lambda_\ell^{(i)}$ and $\Lambda_n = \lambda_m^{(j)}$, with $i, j \in \{1, 2\}$. On the

other hand, it is not difficult to check that

$$(k+1)^2 - n^2 \leq 2(k^2 - n^2), \quad \forall k, n \geq 1 : k \geq n+2.$$

Thus, from (135) we get:

$$\begin{aligned} \Lambda_k - \Lambda_n &= \lambda_\ell^{(i)} - \lambda_m^{(j)} \leq (\nu_1 + 2\varepsilon_0)(\ell^2 - m^2) = \frac{\nu_1 + 2\varepsilon_0}{4} [(k+1)^2 - n^2] \\ &\leq \frac{\nu_1 + 2\varepsilon_0}{2} (k^2 - n^2). \end{aligned}$$

3. When $k = 2\ell$ and $n = 2m$, with $\ell, m \geq k_0$ and $k - n \geq 2$, one has $\ell - m \geq 1$, $\Lambda_k = \lambda_\ell^{(i)}$ and $\Lambda_n = \lambda_m^{(j)}$, with $i, j \in \{1, 2\}$. Applying again (135), we get

$$\Lambda_k - \Lambda_n = \lambda_\ell^{(i)} - \lambda_m^{(j)} \leq (\nu_1 + 2\varepsilon_0)(\ell^2 - m^2) = \frac{\nu_1 + 2\varepsilon_0}{4} (k^2 - n^2).$$

4. Finally, let us take $k = 2\ell$ and $n = 2m - 1$, with $\ell, m \geq k_0$ and $k - n \geq 2$. Again, $\ell - m \geq 1/2$, i.e., $\ell - m \geq 1$ and we can apply (135). As in the previous cases, $\Lambda_k = \lambda_\ell^{(i)}$ and $\Lambda_n = \lambda_m^{(j)}$, with $i, j \in \{1, 2\}$, and

$$\begin{aligned} \Lambda_k - \Lambda_n &= \lambda_\ell^{(i)} - \lambda_m^{(j)} \leq (\nu_1 + 2\varepsilon_0)(\ell^2 - m^2) = \frac{\nu_1 + 2\varepsilon_0}{4} (k^2 - (n+1)^2) \\ &\leq \frac{\nu_1 + 2\varepsilon_0}{4} (k^2 - n^2). \end{aligned}$$

Summarizing, we have prove property (26) for the sequence Λ with

$$\nu = \max \left\{ \tilde{\nu}, \frac{\nu_1 + 2\varepsilon_0}{2} \right\}.$$

Recall again that the constant $\tilde{\nu}$ only depends on k_0 and ν_1 , that is to say, on ρ_1 , ε_0 and ν_1 . Therefore, the parameter ν only depends on ρ_1 , ν_1 and ε_0 .

With the proof of property (26) we end the proof of Proposition 2.14. \square

REFERENCES

- [1] D. Allonsius and F. Boyer, Boundary null-controllability of semi-discrete coupled parabolic systems in some multi-dimensional geometries, *Math. Control Relat. Fields* **10** (2020), 217–256.
- [2] D. Allonsius, F. Boyer and M. Morancey, Analysis of the null controllability of degenerate parabolic systems of Grushin type via the moments method, *J. Evol. Equ.* **21** (2021), 4799–4843.
- [3] F. Ammar Khodja, A. Benabdallah, C. Dupaix and M. González-Burgos, A Kalman rank condition for the localized distributed controllability of a class of linear parabolic systems, *J. Evol. Equ.* **9** (2009), 267–291.
- [4] F. Ammar Khodja, A. Benabdallah, M. González-Burgos and L. de Teresa, The Kalman condition for the boundary controllability of coupled parabolic systems. Bounds on biorthogonal families to complex matrix exponentials, *J. Math. Pures Appl. (9)* **96** (2011), 555–590.
- [5] F. Ammar Khodja, A. Benabdallah, M. González-Burgos and L. de Teresa, Minimal time for the null controllability of parabolic systems: the effect of the condensation index of complex sequences, *J. Funct. Anal.* **267** (2014), 2077–2151.
- [6] F. Ammar Khodja, A. Benabdallah, M. González-Burgos and L. de Teresa, New phenomena for the null controllability of parabolic systems: minimal time and geometrical dependence, *J. Math. Anal. Appl.* **444** (2016), 1071–1113.
- [7] A. Benabdallah, F. Boyer, M. González-Burgos and G. Olive, Sharp estimates of the one-dimensional boundary control cost for parabolic systems and application to the N -dimensional boundary null controllability in cylindrical domains, *SIAM J. Control Optim.* **52** (2014), 2970–3001.

- [8] A. Benabdallah, F. Boyer and M. Morancey, A block moment method to handle spectral condensation phenomenon in parabolic control problems, *Ann. H. Lebesgue* **3** (2020), 717–793.
- [9] A. Benabdallah and M. González-Burgos, The cost of the control in the case of parabolic systems with a positive minimal time coming from the condensation index of the spectrum, work in progress.
- [10] V. Bernstein, *Leçons sur les Progrès Récents de la Théorie des Séries de Dirichlet*, Gauthier-Villars, Paris, 1933.
- [11] F. Boyer and M. Morancey, Analysis of non scalar control problems for parabolic systems by the block moment method, to appear, *Comptes Rendus. Mathématique*.
- [12] P. Cannarsa, A. Duca and C. Urbani, Exact controllability to eigensolutions of the bilinear heat equation on compact networks, *Discrete Contin. Dyn. Syst. Ser. S* **15** (2022), 1377–1401.
- [13] P. Cannarsa, P. Martinez and J. Vancostenoble, The cost of controlling weakly degenerate parabolic equations by boundary controls, *Math. Control Relat. Fields* **7** (2017), 171–211.
- [14] P. Cannarsa, P. Martinez and J. Vancostenoble, Precise estimates for biorthogonal families under asymptotic gap conditions, *Discrete Contin. Dyn. Syst. Ser. S* **13** (2020), 1441–1472.
- [15] N. Cîndea, S. Micu, I. Roventa and M. Tucsnak, Particle supported control of a fluid-particle system, *J. Math. Pures Appl. (9)* **104** (2015), 311–353.
- [16] S. Dolecki, Observability for the one-dimensional heat equation, *Studia Math.* **48** (1973), 291–305.
- [17] H.O. Fattorini and D. L. Russell, Exact controllability theorems for linear parabolic equations in one space dimension, *Arch. Rational Mech. Anal.* **43** (1971), 272–292.
- [18] H.O. Fattorini and D. L. Russell, Uniform bounds on biorthogonal functions for real exponentials with an application to the control theory of parabolic equations, *Quart. Appl. Math.* **32** (1974/75), 45–69.
- [19] E. Fernández-Cara, M. González-Burgos and L. de Teresa, Boundary controllability of parabolic coupled equations, *J. Funct. Anal.* **259** (2010), 1720–1758.
- [20] E. Fernández-Cara and E. Zuazua, The cost of approximate controllability for heat equations: the linear case, *Adv. Differential Equations* **5** (2000), 465–514.
- [21] M. González-Burgos and L. Ouaili, Sharp estimates of the one-dimensional boundary control cost for parabolic systems, in Identification and Control: Some New Challenges, *Contemporary Mathematics* vol. **757**, Amer. Math. Soc., Providence, RI, 2020, 109–126.
- [22] M. González-Burgos and G. R. Sousa-Neto, Boundary controllability of a one-dimensional phase-field system with one control force, *J. Differential Equations* **269** (2020), 4286–4331.
- [23] E.N. Güichal, A lower bound of the norm of the control operator for the heat equation, *J. Math. Anal. Appl.* **110** (1985), 519–527.
- [24] S.W. Hansen, Bounds on functions biorthogonal to sets of complex exponentials; control of damped elastic systems, *J. Math. Anal. Appl.* **158** (1991), 487–508.
- [25] E. Hille, *Analytic Function Theory*, volume II, Second edition, AMS Chelsea Publishing/American Mathematical Society, Providence, RI, 2002.
- [26] A. Kirsch, *An introduction to the Mathematical Theory of Inverse Problems*, Second edition, Applied Mathematical Sciences, 120, Springer, New York, 2011.
- [27] W. A. J. Luxemburg and J. Korevaar, Entire functions and Müntz-Szász type approximation, *Trans. Amer. Math. Soc.* **157** (1971), 23–37.
- [28] P. Lissy, On the cost of fast controls for some families of dispersive or parabolic equations in one space dimension, *SIAM J. Control Optim.* **52** (2014), 2651–2676.
- [29] P. Lissy, Explicit lower bounds for the cost of fast controls for some 1-D parabolic or dispersive equations, and a new lower bound concerning the uniform controllability of the 1-D transport-diffusion equation, *J. Differential Equations* **259** (2015), 5331–5352.
- [30] P. Lissy, The cost of the control in the case of a minimal time of control: the example of the one-dimensional heat equation, *J. Math. Anal. Appl.* **451** (2017), 497–507.
- [31] P. Lissy and E. Zuazua, Internal observability for coupled systems of linear partial differential equations, *SIAM J. Control Optim.* **57** (2019), 832–853.
- [32] Y. Liu, T. Takahashi and M. Tucsnak, Single input controllability of a simplified fluid-structure interaction model, *ESAIM Control Optim. Calc. Var.* **19** (2013), 20–42.
- [33] L. Miller, Geometric bounds on the growth rate of null-controllability cost for the heat equation in small time, *J. Differential Equations* **204** (2004), 202–226.
- [34] L. Ouaili, Minimal time of null controllability of two parabolic equations, *Math. Control Relat. Fields* **10** (2020), 89–112.

- [35] L. Ouaili, *Contrôlabilité de Quelques Systèmes Paraboliques*, Ph.D thesis, Aix-Marseille University, 2020.
- [36] J. Pöschel and E. Trubowitz, *Inverse Spectral Theory*, Pure and Applied Mathematics, 130, Academic Press, Inc., Boston, MA, 1987.
- [37] R. M. Redheffer, Completeness of sets of complex exponentials, *Advances in Math.* **24** (1977), 1–62.
- [38] El H. Samb, Internal null-controllability of the N-dimensional heat equation in cylindrical domains, *C. R. Math. Acad. Sci. Paris* **353** (2015), 925–930.
- [39] L. Schwartz, *Étude des Sommes d'Exponentielles Réelles*, Actualités Sci. Ind., no. 959. Hermann et Cie., Paris, 1943.
- [40] T.I. Seidman, Two results on exact boundary control of parabolic equations, *Appl. Math. Optim.* **11** (1984), 145–152.
- [41] T.I. Seidman, S.A. Avdonin and S.A. Ivanov, The “window problem” for series of complex exponentials, *J. Fourier Anal. Appl.* **6** (2000), 233–254.
- [42] G. Tenenbaum and M. Tucsnak, New blow-up rates for fast controls of Schrödinger and heat equations, *J. Differential Equations* 243 (2007), p. 70-100.
- [43] G. Tenenbaum and M. Tucsnak, On the null-controllability of diffusion equations, *ESAIM Control Optim. Calc. Var.* **17** (2011), 1088–1100.
- [44] R.M. Young, *An Introduction to Nonharmonic Fourier Series*, Pure and Applied Mathematics, 93, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980.

Received xxxx 20xx; revised xxxx 20xx; early access xxxx 20xx.