

Original

Hospital costs associated with nosocomial infections in a pediatric intensive care unit



Áurea Morillo-García^{a,*}, Josefa M. Aldana-Espinal^b, Antonio Olry de Labry-Lima^{c,d}, Raquel Valencia-Martín^a, Reyes López-Márquez^e, Mercedes Loscertales-Abril^f, Manuel Conde-Herrera^a

^a Servicio de Medicina Preventiva y Salud Pública, Hospital Universitario Virgen del Rocío, Sevilla, España

^b Planes Integrales de Salud, Dirección General de Asistencia Sanitaria y Resultados en Salud, Servicio Andaluz de Salud, Sevilla, España

^c Escuela Andaluza de Salud Pública, Granada, España

^d CIBER de Epidemiología y Salud Pública (CIBERESP), España

^e Servicio de Medicina Preventiva y Salud Pública, Hospital Infanta Elena, Huelva, España

^f Unidad de Cuidados Críticos y Urgencias Pediátricas, Hospital Universitario Virgen del Rocío, Sevilla, España

ARTICLE INFO

Article history:

Received 19 June 2014

Accepted 17 February 2015

Available online 25 March 2015

Keywords:

Intensive Care Units, Pediatric

Nosocomial infection

Healthcare costs

Hospital costs

Cohort studies

Regression analysis

ABSTRACT

Objective: To estimate the additional cost attributable to nosocomial infection (NI) in a pediatric intensive care unit (PICU) and related factors.

Methods: A prospective cohort study was conducted in all children admitted to the PICU of a tertiary-care pediatric hospital between 2008 and 2009. Descriptive and bivariate analyses were conducted of total direct costs due to PICU stay and medical procedures in patients with and without NI. A log-linear regression model was performed to determine the factors associated with higher total cost.

Results: A total of 443 patients were studied and the prevalence of NI was 11.3%. The difference in the median total cost was €30,791.4 per patient between groups with and without NI. The median cost of PICU length of stay in patients with NI was almost eight times higher than the median cost of patients without NI. In patients with NI, the highest costs related to medical procedures were associated with antibiotics, enteral and parenteral feeding, and imaging tests. In the multivariate model, the factors associated with higher cost were infection, the performance of cardiovascular surgery, urgent admission, a higher pediatric risk mortality score, and the presence of immunosuppression. By contrast, older children and those with surgical admission generated lower cost.

Conclusions: NI was associated with an increase in total cost, which implies that the prevention of these infections through specific interventions could be cost-effective and would help to increase the safety of healthcare systems.

© 2014 SESPAS. Published by Elsevier España, S.L.U. All rights reserved.

Coste asociado a la infección nosocomial en una unidad de cuidados intensivos pediátricos

RESUMEN

Objetivo: El objetivo del estudio es estimar el coste adicional atribuible a la infección nosocomial (IN) en una Unidad Pediátrica de Cuidados Intensivos (UCIP) y sus factores asociados.

Método: estudio de cohortes prospectivo de todos los pacientes ingresados en una UCIP de tercer nivel entre 2008 y 2009. Se realizó un análisis descriptivo y bivalente del coste total asociados a estancia en UCIP y procedimientos en pacientes con y sin IN. Mediante regresión lineal múltiple, se estimaron los factores asociados al incremento del coste total.

Resultados: se estudiaron 443 pacientes, la incidencia de IN fue 11,3%. La diferencia de las medianas en el coste total fue de 30.791,4€ por paciente entre los grupos con y sin IN. El coste mediano de la estancia de pacientes con IN fue casi ocho veces mayor que el coste mediano de los pacientes sin IN. En pacientes con IN, el coste asociado a procedimientos más elevado fue el de antibióticos, nutrición enteral y parenteral y pruebas de imagen. En el modelo multivariante los factores asociados con un mayor coste fueron: presencia de infección nosocomial, cirugía cardiovascular, tipo ingreso urgente, mayor índice pronóstico de mortalidad al ingreso y la presencia de inmunosupresión. Por el contrario, los de mayor edad y aquellos ingresados por cirugía presentaron un menor coste.

Palabras clave:

Unidad de Cuidados Intensivos Pediátricos

Infección nosocomial

Costes

Costes hospitalarios

Cohortes

Regresión

* Corresponding author. Servicio de Medicina Preventiva y Salud Pública, Hospital Universitario Virgen del Rocío, Avda. Manuel Siurot s/n 41013, Sevilla, España.

Tel.: +34 955013287; fax: +34 955013285.

E-mail address: aurea.morillo@gmail.com (Á. Morillo-García).

Conclusiones: La IN está asociada al incremento del coste total, lo que implica que la prevención de estas infecciones mediante intervenciones específicas podría resultar costo-efectiva, redundando en sistemas de salud más seguros.

© 2014 SESPAS. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

In Pediatric Intensive Care Units (PICU), the prevalence of nosocomial infection (NI) is variable, ranging from 10 to 33%^{1,2}, and incidence rates from 5.9 to 15 per 1,000 patient-days^{3,4}. This is explained by the greater complexity of these patients affected by a variety of medical and surgical pathologies¹.

NI is also considered one of the most costly problems for hospitals. The most important direct NI-associated cost comes from the increase in the excess of length of stay (LOS)⁵. Therefore, most reports focus on NI-related costs due to LOS, as it is quite easy to translate such costs to currency value, applying the local cost per hospitalization day⁶. Further studies also have taken into account the costs of diagnostic and medical procedures, derived from or due to the infection, as the most accurate way to calculate the NI-attributable cost^{7–10}.

The study of NI-associated cost has evolved in time with the introduction of new cost estimation methods^{11–13}. The most common comparative method used to analyse the cost of NI, is the prospective or retrospective observational study of the hospital environment¹⁴.

Few studies have assessed the impact of NI on total costs in PICU¹⁵, some of them have analysed the cost attributable to any type of NI in PICU^{7,11,12,16}, others have focused only on nosocomial primary bloodstream infection^{8,9}. The average cost attributable to NI described in literature ranges between €1,510¹⁷–53,085⁹ per patient.

The Spanish Public Healthcare System is a tax-financed system that offers universal coverage, which is different from privately-funded healthcare systems, where most of the published reports on NI-related cost have been performed^{5,7}. The lack of previous Spanish reports about NI costs in PICU and the actual economical crisis in this country, justifies the present paper. The aim of the study is to estimate the cost attributable to nosocomial infection in a pediatric intensive care unit and its related factors.

MATERIAL AND METHODS

Study Design

Prospective cohort study between September 2008 and July 2009. The study was carried out in a 16-bed multidisciplinary PICU of a tertiary-care university hospital of the Andalusian Public Health System. This Unit is a regional referent institution for highly complex procedures, and offers assistance to patients older than 28 days of age and also to neonates during the cardiovascular surgery postoperative period. All patients admitted in the PICU for less than 24 hours were excluded (a total of 426 patients were excluded). Nine patients remained hospitalized at the end of the study.

Variables

The dependent variable was total cost (only took into account the costs incurred in the PICU), which included the sum of the following cost items: PICU LOS, invasive mechanical ventilation, imaging and microbiological tests, enteral and total

parenteral nutrition, antibiotherapy, consumables (central line, urinary catheter) and structure costs.

The following independent variables were collected: a) Sociodemographic and admission variables: Age, sex, type and cause of PICU admission and cause of discharge; b) Clinical variables: Presence of NI, classified according the definitions of the Centers of Disease Control and Prevention¹⁸; c) Risk factors for NI¹: factors related to patient's intrinsic characteristics and factors related to surgery and other procedures made during PICU stay.

Sources of information

The consumption of different resources (LOS, special feeding formulas and parenteral nutrition, days of mechanical ventilation, imaging and microbiological tests, antibiotherapy and consumables) was obtained from medical records and direct observation, all of which were validated with PICU pediatrician.

The cost of imaging procedures was obtained from the Andalusian Official Bulletin Board¹⁹. Cost of consumables, those related to PICU stay, nutrition, mechanical ventilation and antibiotherapy were obtained from the hospital accounting system. Costs of microbiological tests were obtained from an external reference laboratory²⁰. All costs were updated to the present-day currency in Spain, the Euro (2009).

Costs estimation

1) The estimation of PICU LOS (it includes the costs of PICU staff, consumable goods, catering, instruments and tools, inversions, cleaning, office material, conservation and repair, professional services and supplies for the PICU) was calculated according to the method suggested by Brazzi et al.²¹ 2) The average cost per day of enteral feeding and total parenteral nutrition was calculated and attributed to each patient. 3) The average cost of invasive mechanical ventilation was calculated per patient according to the cost of acquisition, usage and lifespan. 4) For the estimation of the structure costs we used the method suggested by Mason et al.²², which means add 25% to the direct cost. All of them were validated with PICU pediatrician.

Data were collected from the clinical records by trained personnel in epidemiological surveillance and tabulated in a spreadsheet which was specifically designed for this study. The data collection model applied included consumable goods used per patient. These data were classified according to whether they were attributable or not to NI, depending on the most prevalent NI types. The criteria considered to attribute procedures to NI are shown in [Table 1](#). Finally, the project was approved by the hospital Ethics Committee.

Statistical Analysis

NI definition, and infection rates (with 95% confidence intervals) proposed by the National Nosocomial Infections Surveillance System were estimated²³. The chi-squared test was used to compare proportions (NI presence or absence).

NI costs were calculated from the hospital perspective. As a result, we only considered direct tangible costs of PICU for global NI (includes bloodstream infection, pneumonia, urinary tract infection and surgical site infection). In order to calculate average cost,

Table 1
Criteria used to attribute medical procedures-related costs to nosocomial infection (NI).

Type of medical procedure	Description
Use of central venous catheter	Catheter replacement due to suspected bloodstream infection will be attributed to NI, whenever the criteria set by the Centers for Disease Control and Prevention (CDC) are met.
Invasive Mechanical Ventilation	The onset and/or extension of invasive mechanical ventilation in case of clinical worsening of a patient with lower respiratory tract infection or as a result of worsening of other NI considered within the context of cost analysis.
Urinary catheter	The cost of the urinary catheter or its replacement will be attributed to NI whenever urinary tract infection acquired during the urinary catheterization period is confirmed.
Microbiological, Imaging and antibiotherapy tests	Any microbiological cultures, Imaging tests and antibiotic therapy required for the diagnosis and treatment of NI and which are not included in the medical protocol required by the cause of admission will be attributed to NI.

the procedures required per patient, which were attributable to NI, were considered. To check whether these procedures were justified or not, patients' clinical record and interviewed with pediatrician were reviewed.

First of all, a descriptive analysis was performed through mean, median, standard deviation and range. Subsequently, univariate analysis established differences between total cost and independent variables (Mann-Whitney U and Kruskal-Wallis tests for independent categorical variables, and linear regression analysis for independent continuous variables).

A multivariate linear regression was performed to determine the NI cost adjusted by different variables. The variable cost was log transformed, because it was not normally distributed. Clinically relevant variables, those which a priori would be expected to predict costs and those that were shown to be associated at a statistically significant level, were selected for inclusion in the model. The non-collinearity, homogeneity of variance of the variables and the homoscedasticity of the model as well as data fitting using Cook's Distance, the relationship between outliers and standard points and the normal distribution of outliers, were checked. The analysis was carried out using the statistical software package SPSS.v19.

Results

The number of patients enrolled in the study was 443 (57% were boys), with a median age of 1.5 years (percentile 25: 5.3 months-percentile 75: 6.2 years). The cause of admission was surgery in 58.5% of patients, being cardiovascular surgery the most common intervention (46.33%). Global mortality in the Unit was 6.3%. The most frequent procedures were: use of antibiotics, central venous catheterization and invasive mechanical ventilation (85%, 64% and

Table 2
Incidence of nosocomial infection sites.

Nosocomial Infection Site	Number of episodes	Frequency* (%)	Incidence Rate† (%)	Incidence Density Rate‡
Bloodstream infection	35	42.7	7.9	14.43
Pneumonia	23	28.0	5.2	15.76
Surgical site infection	11	13.4	2.5	4.25
Urinary tract infection	7	8.5	1.6	-
Other infection	6	7.3	1.4	-

* No. Nosocomial infections (NI)/total number of infections*100 †No. NI/total number of patients*100; ‡No. bloodstream infections per 1,000 days central line catheter; No. pneumonias per 1,000 days with mechanical ventilation; No. surgical site infection/total of surgical interventions *100.

Table 3
Bivariate Analysis between nosocomial infection and sociodemographic and admission variables.

	Without Nosocomial Infection N (%)	With Nosocomial Infection N (%)
Sex^b		
Boy	228 (90.5%)	24 (9.5%)
Girl	165 (86.4%)	26 (13.6%)
Procedence^b		
Other hospital	44 (88%)	6 (12%)
Home	48 (92.3%)	4 (7.7%)
Operating room	119 (93.7%)	8 (6.3%)
Other service	182 (85%)	32 (15%)
Presence of malformation^b		
No	195 (90.3%)	21 (9.7%)
Yes	198 (87.2%)	29 (12.8%)
Cause of PICU admission^b		
Medical	159 (88.3%)	21 (11.7%)
Surgical	208 (88.1%)	28 (11.9%)
Trauma	12 (100%)	0 (0%)
Transplant	10 (90.9%)	1 (9.1%)
Others	4 (100%)	0 (0%)
Age groups^a		
0-2 months	44 (74.6%)	15 (25.4%)
3 months - 1 years	109 (85.8%)	18 (14.2%)
2-5 years	126 (95.5%)	6 (4.5%)
6-12 years	74 (90.2%)	8 (9.8%)
Over 12 years	40 (93%)	3 (7%)

^a p < 0.001

^b p > 0.05N: number of patients.

55%, respectively). Fifty patients suffered, at least, one episode of NI, which means a prevalence of patients with NI of 11.3%. The total number of NIs was 82 in 443 patients (14.71%). Table 2 shows the number of infections according to site and infection rates which ranged between 7.9% (bloodstream infections incidence) to 1.4% (other infections). 60.6% of the infected patients had a single episode of infection, 18.2% had two infections, and 16.7% had three infections. Two patients had up to 4 infections during admission to the Unit.

Non significant differences were found between patients who had NI and those who didn't according to gender, patient origin, presence of malformation and reason for admission. On the contrary, those with a younger age had a higher risk of NI in comparison to those who were older (Table 3).

Table 4 shows the relationship between the total cost and independent variables for the patients admitted in the PICU. It highlights the higher median cost in children with less than 2 months of age (€6,903.30) and the difference in median cost between bloodstream infection and pneumonia (€37,356.68 and €34,912.96, respectively). Furthermore, the cost increase was proportional to the number of NI diagnosed, with a difference in median cost between having a single infection and 3 or 4 infections of €38,202.57.

Table 4
Cost analysis of variables related to admission, patient's intrinsic characteristics, and variables related or due to procedures during pediatric intensive care unit (PICU) stay.

	Total Cost € Average (Median)	P
<i>Age groups</i>		
0-2 months	12,391.54 (6,903.30)	
3 months - 1 years	10,028.14 (4,047.99)	
2-5 years	6,626.95 (3,047.07)	
6-12 years	5,936.10 (3,064.88)	
Over 12 years	6,112.18 (3,092.41)	<0.001 ^b
<i>Cause of PICU admission</i>		
Medical	10,831.99 (4,234.83)	
Surgical	6,466.42 (3,079.48)	
Trauma	2,566.42 (2,087.45)	
Transplant	8,646.17 (6,548.23)	
Others	6,819.02 (4,862.94)	<0.001 ^b
<i>Type of admission</i>		
Elective	6,175.98 (3,061.63)	
Emergency	10,291.44 (4,215.44)	<0.001 ^a
<i>PRISM index</i>		
0-10	5,823.08 (3,137.58)	
>10	14,416.39 (8,040.86)	<0.001 ^a
<i>Immunosuppression</i>		
No	7,782.94 (3,179.44)	
Yes	12,096.62 (6,174.37)	<0.001 ^a
<i>Previous respiratory disease</i>		
No	7,018.44 (3,378.61)	
Yes	27,812.31 (4,176.73)	0.026 ^a
<i>Infection at admission in PICU</i>		
No	6,866.32 (3,127.06)	
Yes	11,759.96 (4,501.40)	<0.001 ^a
<i>Cause of discharge</i>		
Healing	6,946.21 (3,203.70)	
Exitus	21,681.17 (6,626.98)	
Unknown ^c	22,420.13 (11,438.87)	<0.001 ^b
<i>Surgery during admission</i>		
No	8,091.67 (4,048.02)	
Yes	8,245.86 (3,183.97)	0.106 ^a
<i>Cardiovascular surgery</i>		
No	7,122.94 (3,059.62)	
Yes	11,069.21 (5,157.50)	<0.001 ^a
<i>Mechanical ventilation</i>		
No	3,559.82 (2,291.40)	
Yes	11,969.72 (5,100.19)	<0.001 ^a
<i>Total parenteral feeding</i>		
No	3,214.45 (3,430.33)	
Yes	13,714.52 (9,089.37)	0.03 ^a
<i>Immunosuppressive therapy</i>		
No	5,908.44 (2,724.87)	
Yes	10,506.52 (4,468.25)	<0.001 ^a
<i>Use of antibiotherapy</i>		
No	3,571.58 (2,485.69)	
Yes	9,000.77 (4,004.44)	0.012 ^a
<i>No. of nosocomial infections</i>		
None	4,657.19 (3,084.24)	
One	19,041.84 (15,085.93)	
Two	42,683.96 (35,019.31)	
Three-four	63,213.73 (53,288.50)	<0.001 ^b
<i>Bloodstream infection</i>		
No	5,830.76 (3,162.45)	
Yes	43,187.44 (28,012.01)	<0.001 ^a
<i>Pneumonia</i>		
No	6,458.08 (3,218.90)	
Yes	41,371.04 (31,827.37)	<0.001 ^a

^a Mann Whitney U test (2 categories)

^b Kruskal-Wallis test (> 2 categories).

^c Patients who remained in the PICU after the study period. PICU: Pediatric Intensive Care Unit; PRISM: Pediatric Risk of Mortality

Table 5
Cost in Euros (€) of children admitted in the Pediatric Intensive Unit (PICU) per patient with and without nosocomial infection.

	Without Infection Average (median)	With Infection Average (median)
PICU-stay cost ^a	3601.21 (2386.64)	27875.97 (18695.35)
Mechanical ventilation cost ^a	4.27 (0)	41.25 (34.09)
Imaging test cost ^b	63.67 (27.69)	143.98 (101.53)
Feeding cost ^{a,b}	15.34 (5.80)	20.85 (116.18)
Antibiotherapy cost ^a	20.85 (0.93)	140.68 (33.40)
Microbiological test cost ^a	83.25 (49.58)	240.49 (135.54)
Central venous catheter related cost	29.40 (39.16)	70.50 (58.75)
Arterial catheter related cost	2.70 (0)	5.55 (8.41)
Direct cost ^a	3773.26 (2479.76)	28406.39 (19238.12)
Total cost ^a	4716.58 (3099.70)	35507.99 (24047.64)

PICU: Pediatric Intensive Care Unit

^a p < 0.001

^b p < 0.005; [‡]Parenteral feeding or specific infant formulas provided by the Pharmacy Service; [§]Total cost= direct cost plus 25%²¹

Table 5 shows the differences between infection and non-infection regarding cost. It is worth noting that about 70% of the total cost was due to LOS expenses. In this sense, median LOS of patients with NI was 24 days versus 3 days for those without NI (p < 0.001). The difference in the median cost of LOS in NI-patients, was statistically different comparing it to those without NI (€18,695.35 and €2,386.64, respectively). Regarding the total cost of the patients with NI included, a difference in median cost of €20,947.94 (€30,791.41 average cost) was observed between patients with and without NI (p < 0.001).

The multivariate analysis shows that those with bloodstream infections (p < 0.001) and pneumonia (p = 0.008) have a greater cost than those without NI. It is worth noting that a higher cost in PICU was associated with cardiovascular surgery (p < 0.001), urgent type of admission (p = 0.004), higher Pediatric Risk of Mortality index (p < 0.001) and the presence of immunosuppression (p < 0.001). By contrast, older children and surgical type of admission (p = 0.009) presented a lower cost (Table 6).

Discussion

The analysis of NI-related costs started in countries with privately-funded health care systems, followed by others with a hybrid system of private and public insurance, like Spain^{10,24–26}. In these studies, the type of patients included and the healthcare systems where they were performed were very different, which hinders any possible comparison. However, all of them report a remarkable cost associated to NI⁷.

In the present study, estimated costs have been based upon prolonged PICU LOS and variable costs of different medical procedures, the latter being a direct attribution resulting from the NI presence, which is a better estimation of attributable costs. In this sense, Pittoni and Scatto²⁷ remark the importance of taking into account direct variable costs to understand the real costs generated from a patient admitted in the intensive care unit, including non complex medical procedures as they may affect the result of cost analysis in an accumulative way.

The authors have performed a multivariate analysis using multiple linear regression models in order to assess factors associated with the total cost, as it is difficult to choose the matching factors between cases and controls without committing a selection bias. This follows the current trend in NI-related cost analysis (from 1990 to 1999, 14.3% of the studies performed multiple linear regression and from 2000–2006, 34.5% did so²⁸).

Table 6
Multivariate model of factors associated with the increase in total costs at the pediatric intensive care unit (n=443; R² 48%).

		P	β (95% CI)
Constant		<0.001	3.554 (3.397; 3.710)
Age groups	0-2 months		
	2 months - 1 year	0.003	-0.172 (-0.287; -0.057)
	1-5 years	0.003	-0.181 (-0.298; -0.063)
	5-12 years	0.001	-0.217 (-0.343; -0.091)
	Over 12 years	0.376	-0.067 (-0.215; 0.081)
Bloodstream infection	No		
	Yes	<0.001	0.648 (0.499; 0.797)
Pneumonia	No		
	Yes	0.008	0.254 (0.067; 0.442)
Cardiovascular surgery	No		
	Yes	<0.001	0.287 (0.196; 0.378)
Type of admission	Medical		
	Surgical	0.009	-0.149 (-0.260; -0.038)
Immunosuppression	No		
	Yes	0.001	0.199 (0.085; 0.312)
Type of admission	Elective		
	Urgent	0.004	0.157 (0.050; 0.264)
PRISM index		<0.001	0.014 (0.007; 0.021)

PRISM: Pediatric Risk of Mortality

In this study, we observed a difference of 21 days in PICU LOS between patients with and without NI. Although, the authors did not adjust by other variables, there was a difference in costs between both groups of more than €30,000. In spite of being statistically significant, it may not be exclusively due to NI. This analysis, although it has been the most frequently used in studies of costs for NI, has limitations, as it results from an average measure. Also, it does not reflect the real expenditure, considering that this is not uniform throughout the stay, and that, as the NI is time-dependent, it should measure the extension of the stay only after its diagnosis, and not for the entire stay¹². Nevertheless, other studies carried out in PICU, despite of using a different methodology, and adjusted due to other factors like patients' age or cause of PICU admission, show similar results, like, Slonim et al.⁸, who found a difference of 14 days, or Elward et al.⁹, with a mean hospital LOS cost of €24,619 in patients with NI.

It also must be taken into account the association between LOS and NI, which may be contradictory, since, although it is recognized by the literature that NI increases hospital stay²⁹, some authors also consider LOS as a risk factor for the NI³⁰ (longer hospitalizations leading to colonization of patients by pathogenic microorganisms, longer duration of invasive procedures and pharmacotherapy which favours the development of NI). However, the cost study presented here was part of a prospective study which also analyzed the risk factors associated with NI, not finding a significant association between LOS and NI. Therefore, in this paper, the authors have assumed that NI has influenced the increase in LOS, considering what referred by the literature and the results of the study for risk factors of NI.

Average cost of medical procedures in the current study was higher in the case of NI patients when compared with non-infected patients, where the use of antibiotherapy, feeding and imaging tests should be highlighted. In the case of bloodstream infection, the cost of catheters and antibiotics doubled the observed in patients with pneumonia, as in Pittoni and Scatto's case²⁷, which remarks that one of the most difficult challenges is the bloodstream infection management, because of its wide array of possible treatments and elevated associated cost.

Globally, the cost of NI is within the range described in other studies performed in PICU^{5,7,9,11,17,31} (from €1,510¹⁷ to €53,085⁹). The last study showed higher costs, which may, in part, be justified by the use of a cost-estimation method based upon cost to charge ratios and the inclusion of more complex patients. The BI was the

most costly NI, showing significant difference when compared to pneumonia, which it agrees with other studies^{5,32}.

Nevertheless, the results from other studies which only report average values must be interpreted cautiously. Their costs estimations were obtained in different countries with a healthcare management which may not be equivalent to the Spanish Health System. Likewise, other costs should be considered, such as those derived from patients who die as a result of the NI; the cost of the medical procedures required by the treatment of NI beyond 48 hours after discharge from PICU, or the indirect or intangible costs referred to productivity loss, and more specifically what the patient loses as a result of the infection and/or treatment (pain, suffering, loss of leisure time), which affects their quality of life.

Some of the variables that were associated with an increase of total cost (presence of NI, emergency admission, previous cardiovascular surgery and immunosuppression), were also reported by other studies^{9,16,29}. Other factors that may also increase the cost of the Unit were the cause of non surgical admission and comorbidities such as severe lesions in primary organs^{7,9,12,33}. In the present study, only immunosuppression, as a patient's intrinsic factor, contributed in a statistically significant way to the model. In this sense, Gemke et al.³⁴ stated that non previously programmed admissions did increase the use of resources and, consequently, total costs.

Another factor included in our model as a risk factor is cardiovascular surgery, which could be due to the high number of patients undergoing such surgery in the PICU, postoperative LOS, frequent infectious complications and prolonged use of medical procedures, which means an increase of total economic costs. The inclusion of this factor is supported by a recent report published by González et al.³⁵, which states that prolonged admissions convey a greater use of medical resources, higher risk of NI and higher mortality rates among children admitted to PICU³⁵. Other studies mention hospital stay^{1,14} and non survival in the Unit^{7,9,24} as factors capable of rising costs. These variables were only statistically significant in the bivariate analysis.

In conclusion, this paper has approached to NI-related economic cost, where most of the expenses are derived from the medical procedures made to the patient during PICU stay and its complications. The higher total cost per patient requires the implementation of specific cost-effective preventive measures to control NI, in order to minimize its effects in the health of pediatric patients and in medical costs.

What is known about the topic?

Nosocomial infection in Pediatric Intensive Care Units has a significant impact on morbidity, mortality and economic cost. However, very few studies have performed an estimate of the economic cost of nosocomial infection in public health systems, since most of them are confined to privately funded healthcare systems and adult population.

What does this study adds?

The present study provides an estimate of the economic cost of nosocomial infection in critically ill children, which may be useful to both economic managers and physicians about its repercussion in economic terms.

Editor in charge

Miguel Ángel Negrín

Author's Contribution

The project's original idea was provided by R. López. A. Morillo and JM Aldana adapted the idea for the research project and its implementation in a Pediatric Intensive Care Unit. R. Valencia developed the project's methods, and performed the database and its deputation. The clinical data collection and the attribution of medical procedure-related costs to nosocomial infection were made by A. Morillo and R. López. A. Olry performed the economic analysis. M. Conde and M. Loscertales requested the costs information from the Departments of Pharmacology and Economic Management. A. Morillo wrote the first draft of the article, and was subsequently reviewed and approved by all authors.

Funding

This project has been funded by the Andalusian Regional Ministry of Health.

Conflicts of interest

None

Acknowledgements

To all the staff of the Pediatric Intensive Care Unit and the Department of Economic Management from Virgen del Rocío University Hospital, for their cooperation in this project.

References

1. de Mello MJ, de Albuquerque MF, Lacerda HR, et al. Risk factors for healthcare-associated infection in pediatric intensive care units: a systematic review. *Cad Saude Publica*. 2009;25 Suppl 3:S373–91.
2. Joram N, de Saint L, Stamm D, et al. Healthcare-associated infection prevention in pediatric intensive care units: a review. *Eur J Clin Microbiol Infect Dis*. 2012;31:2481–90.
3. Urrea M, Pons M, Serra M, et al. Prospective incidence study of nosocomial infections in a pediatric intensive care unit. *Pediatr Infect Dis J*. 2003;22:490–3.
4. Stover BH, Shulman ST, Bratcher DF, et al. Pediatric Prevention Network. Nosocomial infection rates in US children's hospitals' neonatal and pediatric intensive care units. *Am J Infect Control*. 2001;29:152–7.
5. Stone P, Larson E, Kavar L. A systematic audit of economic evidence linking nosocomial infections and infection control interventions. *Am J Infect Control*. 2002;30:145–52.
6. Vrijens F, Hulstaert F, Van de Sande S, et al. Hospital-acquired, laboratory-confirmed bloodstream infections: linking national surveillance data to clinical

- and financial hospital data to estimate increased length of stay and health care cost. *J Hosp Infect*. 2010;15:8–162.
7. Chalom R, Raphaely RC, Costarino AT. Hospital costs of pediatric intensive care. *Crit Care Med*. 1999;27:2079–85.
8. Slonim A, Kurtines H, Sprague B, et al. The costs associated with nosocomial bloodstream infection in the pediatric intensive care unit. *Pediatr Crit Care Med*. 2001;2:170–4.
9. Elward AM, Hollenbeak CS, Warren DK, et al. Attributable cost of nosocomial primary bloodstream infection in pediatric intensive care unit patients. *Pediatrics*. 2005;115:868–72.
10. Morís de la Tassa J, Fernández P, Antuña Egocheaga A, et al. Estimación de los costes asociados a la infección urinaria nosocomial. Un estudio de casos y controles. *Rev Clín Esp*. 2003;203:119–24.
11. Stone PW, Braccia D, Larson E. Systematic review of economic analyses of health care-associated infections. *Am J Infect Control*. 2005;33:501–9.
12. Roberts R, Scott RD II, Cordell R, et al. The use of economic modelling to determine the Hospital Costs associated with nosocomial infections. *CID*. 2003;36:1424–32.
13. Yalcin AN. Socioeconomic burden of nosocomial infections. *Indian J Med Sci*. 2003;57:450–6.
14. Scott RD II. The direct medical costs of Healthcare-Associated Infections in U.S. Hospitals and the benefits of prevention. Economist Division of Healthcare Quality Promotion National Center for Preparedness, detection and control of Infectious Diseases coordinating Center for Infectious Diseases. Centers for Disease Control and Prevention. March 2009. [Accessed 29 July 2012]. Available at: http://www.cdc.gov/hai/pdfs/hai/scott_costpaper.pdf
15. Sachdeva RS. Cost of nosocomial infections in the pediatric intensive care unit. *Seminars in Pediatric Infectious Diseases*. 1999;110:239–42.
16. Domínguez TE, Chalom R, Costarino AT. The impact of adverse patient occurrences on hospital costs in the pediatric intensive care unit. *Crit Care Med*. 1999;160:976–81.
17. Gurskis V, Kévalas R, Kerienė V, et al. Economic evaluation of nosocomial infections in pediatric intensive care units in Lithuania. *Medicina (Kaunas)*. 2010;46:781–9.
18. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control*. 2008;36:309–32.
19. ORDEN de 14 de octubre de 2005, por la que se fijan los precios públicos de los servicios sanitarios prestados por Centros dependientes del Sistema Sanitario Público de Andalucía. *BOJA* n° 210 de 2005:46–83.
20. Laboratorio Reference. [Accessed 2 May 2012]. Available at: <http://www.referencelaboratory.es/index.html>
21. Brazzi L, Bertolini G, Arrighi E, et al. Top-down costing: problems in determining staff costs in intensive care medicine. *Intensive Care Med*. 2002;28:1661–3.
22. Mason A, Epstein D, Smith PC, et al. International comparison of costs: An exploration of within- and between-country variations for ten healthcare services in nine EU member states. [Accessed January 9, 2011]. Available at: www.healthbasket.org
23. Emori TG, Culver DH, Horan TC, et al. National nosocomial infections surveillance system (NNIS): description of surveillance methodology. *Am J Infect Control*. 1991;19:19–35.
24. García S, Ruza F, Alvarado F, et al. Analysis of costs in a pediatric ICU. *Int Care Med*. 1997;23:218–25.
25. Díaz C, García M, Bueno A, et al. The estimation of the cost of nosocomial infection in an intensive care unit. *Med Clin (Barc)*. 1993;100:329–32.
26. Fernández M, Peinado F, Teba F, et al. Nosocomial infection in BPH: economic costs and increase in hospital stay. *Actas Urol Esp*. 1996;20:269–73.
27. Pittoni GM, Scatto A. Economics and outcome in the intensive care unit. *Curr Opin Anaesthesiol*. 2009;22:232–6.
28. Fukuda H, Lee J, Imanaka Y. Variations in analytical methodology for estimating costs of hospital-acquired infections: a systematic review. *J Hosp Infect*. 2011;77:93–105.
29. Srinivasan R, Asselin J, Gildengorin G, et al. A Prospective Study of Ventilator-Associated Pneumonia in Children. *Pediatrics*. 2009;123:1108–15.
30. Foglia EE, Fraser VJ, Elward AM. Effect of Nosocomial Infections Due to Antibiotic-Resistant Organisms on Length of Stay and Mortality in the Pediatric Intensive Care Unit. *Am J Infect Control*. 2007;28:299–306.
31. Navarrete S, Armengol G. Costos secundarios por infecciones nosocomiales en dos unidades pediátricas de cuidados intensivos. *Salud Publica Mex*. 1999;41:S51–8.
32. Chen Y, Wang F, Liu CY, et al. Incidence rate and variable cost of nosocomial infections in different types of intensive care units. *Infect Control Hosp Epidemiol*. 2009;30:39–46.
33. Chen YY, Chou YC, Chou P. Impact of nosocomial infection on cost of illness and length of stay in intensive care units. *Infect Control Hosp Epidemiol*. 2005;26:281–7.
34. Gemke R, Bonsel GJ, McDonnell J, et al. Patient characteristics and resource utilization in pediatric intensive care. *Arch Dis Child*. 1994;71:291–6.
35. González-Cortés R, López-Herce-Cid J, García A, et al. Ingreso prolongado en la unidad de cuidados intensivos pediátricos: mortalidad y consumo de recursos asistenciales. *Med Intensiva*. 2011 [Accessed 15 February 2013]. Available at: [http://www.elsevier.es/sites/default/files/elsevier/eop/S0210-5691\(11\)00115-X.pdf](http://www.elsevier.es/sites/default/files/elsevier/eop/S0210-5691(11)00115-X.pdf)