

Regulation of oxygen sensing by ion channels

José López-Barneo, Raquel del Toro, Konstantin L. Levitsky, María D. Chiara and Patricia Ortega-Sáenz

J Appl Physiol 96:1187-1195, 2004. doi:10.1152/jappphysiol.00929.2003

You might find this additional info useful...

This article cites 92 articles, 68 of which can be accessed free at:

</content/96/3/1187.full.html#ref-list-1>

This article has been cited by 19 other HighWire hosted articles, the first 5 are:

Vasodilation induced by oxygen/glucose deprivation is attenuated in cerebral arteries of SUR2 null mice

Adebowale Adebisi, Elizabeth M. McNally and Jonathan H. Jaggar

Am J Physiol Heart Circ Physiol, October, 2011; 301 (4): H1360-H1368.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

The Ventilatory Response to Hypoxia in Mammals: Mechanisms, Measurement, and Analysis

Luc J. Teppema and Albert Dahan

Physiol Rev, April, 2010; 90 (2): 675-754.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Molecular diversity and function of K⁺ channels in airway and alveolar epithelial cells

Olivier Bardou, Nguyen Thu Ngan Trinh and Emmanuelle Brochiero

Am J Physiol Lung Cell Mol Physiol 2009; 296 (2): L145-L155.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Heme oxygenase is necessary for the excitatory response of cultured neonatal rat rostral ventrolateral medulla neurons to hypoxia

Dominic D'Agostino, Emilio Mazza, Jr. and Judith A. Neubauer

Am J Physiol Regul Integr Comp Physiol 2009; 296 (1): R102-R118.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Functional mitochondria are required for O₂ but not CO₂ sensing in immortalized adrenomedullary chromaffin cells

J. Buttigieg, S. T. Brown, M. Lowe, M. Zhang and C. A. Nurse

Am J Physiol Cell Physiol 2008; 294 (4): C945-C956.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Updated information and services including high resolution figures, can be found at:

</content/96/3/1187.full.html>

Additional material and information about *Journal of Applied Physiology* can be found at:

<http://www.the-aps.org/publications/jappl>

This information is current as of September 22, 2014.

HIGHLIGHTED TOPIC | *Oxygen Sensing in Health and Disease*

Regulation of oxygen sensing by ion channels

José López-Barneo, Raquel del Toro, Konstantin L. Levitsky, María D. Chiara, and Patricia Ortega-Sáenz

Laboratorio de Investigaciones Biomédicas, Departamento de Fisiología and Hospital
Universitario Virgen del Rocío, Universidad de Sevilla, E-41013 Sevilla, Spain

López-Barneo, José, Raquel del Toro, Konstantin L. Levitsky, María D. Chiara, and Patricia Ortega-Sáenz. Regulation of oxygen sensing by ion channels. *J Appl Physiol* 96: 1187–1195, 2004; 10.1152/jappphysiol.00929.2003.—O₂ sensing is of critical importance for cell survival and adaptation of living organisms to changing environments or physiological conditions. O₂-sensitive ion channels are major effectors of the cellular responses to hypoxia. These channels are preferentially found in excitable neurosecretory cells (glomus cells of the carotid body, cells in the neuroepithelial bodies of the lung, and neonatal adrenal chromaffin cells), which mediate fast cardiorespiratory adjustments to hypoxia. O₂-sensitive channels are also expressed in the pulmonary and systemic arterial smooth muscle cells where they participate in the vasomotor responses to low O₂ tension (particularly in hypoxic pulmonary vasoconstriction). The mechanisms underlying O₂ sensing and how the O₂ sensors interact with the ion channels remain unknown. Recent advances in the field give different support to the various current hypotheses. Besides the participation of ion channels in acute O₂ sensing, they also contribute to the gene program developed under chronic hypoxia. Gene expression of T-type calcium channels is upregulated by hypoxia through the same hypoxia-inducible factor-dependent signaling pathway utilized by the classical O₂-regulated genes. Alteration of acute or chronic O₂ sensing by ion channels could participate in the pathophysiology of human diseases, such as sudden infant death syndrome or primary pulmonary hypertension.

electrophysiology; gene expression; hypoxia-inducible factors

OXYGEN SENSING IS OF PARAMOUNT importance for cell survival due to the central role of O₂ as acceptor of the electrons in the mitochondrial respiratory chain, thus making possible the synthesis of adenosine triphosphate (ATP) by oxidative phosphorylation. The provision of sufficient O₂ to the tissues in variable habitats or physiological situations is a homeostatic challenge because even transient localized O₂ deficits can produce irreversible cellular damage. The lack of O₂ participates critically in the pathogenesis of major causes of mortality, such as stroke, myocardial infarction, and chronic lung disease. In mammals, acute hypoxia triggers fast respiratory and cardiovascular counterregulatory adjustments (occurring over a time scale of seconds to minutes) to ensure sufficient O₂ supply to the most critical organs such as the brain or the heart. These acute responses to hypoxia depend on the modulation of O₂-regulated ion channels, which mediate adaptive changes in cell excitability, contractility, and secretory activity (Fig. 1). O₂-regulated ion channels are preferentially expressed in excitable cells of specific tissues such as the arterial and airway chemoreceptors (carotid and neuroepithelial bodies), smooth muscle from the pulmonary and systemic vasculature, and neonatal adrenal medulla (40). Chronic exposure to hypoxia (for hours to days) regulates the expression of numerous genes

encoding enzymes, growth factors, or transporters, which induce molecular and histological modifications to reduce the cellular need and dependence on O₂ and increase O₂ supply to the tissues (Fig. 1). Chronic adaptation to hypoxia, observed in almost every cell type, critically depends on transcriptional mechanisms that determine the level of expression of numerous genes (9, 66). Transcriptional activity induced by hypoxia has been shown to rely on O₂-dependent protein hydroxylases, which regulate the activity and nuclear translocation of hypoxia-inducible transcription factors (HIF-1 and isoforms) (42, 66).


In this review, we summarize the participation of ion channels in the cellular and systemic adaptive responses to hypoxia. We discuss the role of ion channels as effectors of the hypoxia signal transduction pathway in cells acutely responding to low P_{O₂}, emphasizing the present hypotheses on the mechanisms underlying O₂ detection. We also stress the growing importance of ion channels as part of the gene expression program triggered by chronic hypoxia. Finally, we comment on the pathophysiological implications of acute and chronic regulation of ion channel function by O₂ tension.

ACUTE RESPONSES TO HYPOXIA MEDIATED BY ION CHANNELS

Changes of local O₂ tension can induce modifications in the electrical activity of numerous cell types, including neurons; however, neurosecretory cells located in chemoreceptor organs and smooth muscle cells in pulmonary and systemic arteries are

Address for reprint requests and other correspondence: J. López-Barneo, Laboratorio de Investigaciones Biomédicas, Edificio de Laboratorios, 2ª planta, Hospital Universitario Virgen del Rocío, Avenida Manuel Siurot s/n, E-41013 Sevilla, Spain (E-mail: Jose.L.Barneo.sspa@juntadeandalucia.es).

Fig. 1. Hypoxia signaling pathway with indication of the major adaptive responses to acute and chronic hypoxia.




those directly involved in the major systemic adjustments to acute hypoxia.

O₂-Sensitive Neurosecretory Cells

The carotid and aortic bodies or the neuroepithelial bodies of the lung, which are organs capable of sensing global O₂ tension, participate in the compensatory cardiorespiratory adjustments to hypoxia. These organs contain O₂-sensitive neurosecretory cells, which release transmitters on exposure to environmental low P_{O₂} (<50 to 60 Torr) (40). The classical O₂-sensing chemoreceptors are the carotid bodies, composed of clusters of sensory (glomus) cells innervated by afferent nerve fibers. These sensory fibers convey chemosensory discharges to the brain stem respiratory center to evoke hyperventilation during hypoxemia. Similar clusters of excitable neurosecretory cells have also been described in the neuroepithelial bodies of the lung and in adrenal chromaffin cells from the neonate where they detect P_{O₂} changes in the inspired air and in blood, respectively. Excitation of chemoreceptor cells by hypoxia mainly depends on the presence of membrane channels whose activity is modulated by low P_{O₂}. The “O₂-sensitive” channels studied in more detail are K⁺ channels, initially described in glomus cells of the carotid body (7, 15, 23, 39, 53, 71) and found in all the hypoxia-responsive neurosecretory cells studied so far (61, 75, 87, 92). Nevertheless,

the kind of O₂-sensitive K⁺ channel appears to change among the various chemoreceptor cells or among cells in different animal species. Some K⁺-channel types proposed to participate in acute O₂ sensing are voltage-dependent channels of the voltage-gated K⁺ (K_v) or *Shaker* family, Ca²⁺-activated K⁺ (K_{Ca}) channels, and TASK-like background K⁺ channels (see Refs. 38 and 40 and references therein). Recordings illustrating the reversible inhibition by hypoxia of whole cell K⁺ currents in several chemoreceptor cell types are shown in Fig. 2A, and the “membrane model” of chemosensory transduction is summarized by a scheme in Fig. 2B. At least in the case of carotid body glomus cells, all of the indicated steps involved in stimulus-secretion coupling have strong experimental support. Transmitter release induced by hypoxia in isolated cells is 1) mimicked by depolarization with high extracellular K⁺ or application of K⁺ channels blockers (51, 78), 2) paralleled by an increase in cytosolic Ca²⁺ concentration ([Ca²⁺]_i) (44, 78), and 3) abolished by removal of extracellular Ca²⁺ or blockade of voltage-gated Ca²⁺ channels (44, 78). In addition, it has directly been shown in patch-clamped cells that hypoxia or K⁺-channel blockers produce glomus cell depolarization (8, 85) and an increase of action potential firing frequency (8, 41, 44). The rise of cytosolic [Ca²⁺]_i induced by hypoxia is prevented in voltage-clamped cells held at negative membrane potentials (8). Altogether, these data indicate that chemosen-

Fig. 2. A: reversible reduction of macroscopic K⁺ currents by hypoxia in 4 representative O₂-sensitive neurosecretory cells. B: scheme of the membrane model of O₂ sensing in neurosecretory cells. [Ca²⁺]_i, Ca²⁺ concentration; CNS, central nervous system; I_K, K⁺ current.



sory transduction is initiated by the closure of K^+ channels by low P_{O_2} , which leads to membrane depolarization and/or increase of action potential firing frequency, extracellular Ca^{2+} influx through voltage-gated channels, and transmitter release to the extracellular milieu (38).

Vascular Smooth Muscle Cells

Vascular smooth muscle cells (VSMCs) control blood flow and tone, and their contraction is directly influenced by blood O_2 tension. The acute vascular responses to hypoxia studied in more detail are pulmonary vasoconstriction and dilation of systemic vessels (82, 91).

Hypoxia-induced vasoconstriction (pulmonary vasculature). Hypoxic pulmonary vasoconstriction (HPV), which occurs predominantly in small resistance arteries, is essential for fetal life because it helps to maintain the high pulmonary vascular resistance that diverts blood through the ductus arteriosus. In adults, HPV reduces blood flow through poorly ventilated alveoli, thus contributing to matching perfusion to ventilation and preventing systemic hypoxemia when atelectasis is present. Although the adaptive changes of pulmonary myocytes to low P_{O_2} are complex, these cells respond, similar to the O_2 -sensitive neurosecretory cells, with reduction in amplitude of the macroscopic voltage-dependent K^+ currents (49, 56, 89, 90). Inhibition of one or several types of K^+ channels by hypoxia leads to membrane depolarization, opening of voltage-gated Ca^{2+} channels, and myocyte contraction (82, 89).

Hypoxia-induced vasodilation. Hypoxic vasodilation is another fast response to hypoxia of VSMCs, particularly well manifested in coronary and cerebral vessels, that helps to increase the perfusion of blood to the O_2 -deprived tissues. A major component of hypoxic vasodilation is mediated by ATP-sensitive K^+ (K_{ATP}) channels of vascular myocytes, which open in response to hypoxia due to decreased ATP production (13). However, there are other O_2 -sensitive ionic mechanisms causing myocyte relaxation because it occurs with P_{O_2} levels that do not compromise energy metabolism. K_{Ca} channels potentiated by low P_{O_2} have been described in isolated cerebral resistance myocytes (24), and a somewhat similar mechanism (inhibition of K^+ channels by normoxia) has been proposed to induce contraction of the ductus arteriosus at birth once the blood in the newborn is oxygenated (77). Moreover, there is considerable evidence indicating that, in arterial myocytes, transmembrane Ca^{2+} influx is also directly inhibited by low P_{O_2} . Relaxation by hypoxia is produced in arteries precontracted with K^+ (a condition that prevents repolarization by opening of K_{ATP} or K_{Ca} channels), and in isolated myocytes the elevation of cytosolic $[Ca^{2+}]$ induced by high K^+ concentration is reversibly reduced by low P_{O_2} (20, 79). Inhibition of L-type Ca^{2+} channels by hypoxia has been described in patch-clamped systemic arterial myocytes (20, 21, 68).

MECHANISMS OF O_2 SENSING BY ION CHANNELS

O_2 Sensor in Close Contact with the Ion Channels

Although the role of ion channels as effectors in the acute cellular responses to hypoxia is well established, the identity of the O_2 sensor molecules and the signaling pathways linking the sensors to the effectors remain, however, enigmatic. Because some channels retain the hypoxia responsiveness in excised

membrane patches, it has been suggested that the O_2 sensor is closely associated with the channel oligomer, either attached to the pore-forming α -subunit or as part of an auxiliary subunit (23, 32, 37, 59). Switching of the sensor between oxo- and deoxo-conformations would result in alteration of the channel gating owing to direct allosteric interactions. Some specific types of K^+ channel α - and/or β -subunits are expressed in O_2 -sensitive cells (4, 10, 14, 25, 30, 62). The levels of protein expression of $Kv\alpha 1.5$ and $Kv\alpha 3.1$ as well as $Kv\beta 1.1$ subunits are reported to be higher in pulmonary resistance arterial myocytes than in other arterial VSMCs (10). It has been shown that mice lacking the $Kv\alpha 1.5$ subunits have impaired HPV and reduced sensitivity of whole cell voltage-gated K^+ currents to hypoxia (2) and that antibodies against $Kv2.1$ diminish O_2 -sensitive currents in rat pulmonary myocytes (30). However, whether $Kv1.5$, $Kv2.1$, or any other K^+ channel α -subunit acts as an O_2 sensor or whether they are simply effectors in the hypoxia signaling cascade of pulmonary VSMCs is not known. Some recombinant subunits of K^+ and Ca^{2+} channels expressed in heterologous cells have been shown to be O_2 sensitive (18, 31, 35, 36, 50, 52). These studies lead us to expect rapid progress in the identification of the O_2 sensors; however, advances produced so far have been relatively minor, and the data available are inconclusive. The O_2 sensitivity of α or combinations of α and β K^+ and Ca^{2+} channel subunits changes, depending on the experimental conditions used in the various laboratories and on the cell types used for heterologous protein expression (see Ref. 40 for a detailed discussion). Therefore, it seems that O_2 sensitivity is not absolutely intrinsic to the ion channels but requires the interaction between the O_2 -sensing signaling molecules and the pore-forming channel subunits.

Redox Model of Acute O_2 Sensing

An alternative view to the existence of an O_2 sensor attached to the ion channels is the redox model of O_2 sensing based on the conversion of O_2 into reactive oxygen species (ROS), which would then alter the cellular redox status and the function of the ion channels (which contain numerous residues susceptible to redox modification) (1, 11). The two ROS-producing systems postulated as O_2 sensors are the NADPH oxidase and mitochondria.

NADPH oxidase as possible O_2 sensor. NADPH oxidase has been proposed to transduce O_2 levels by changing the rate of superoxide anion (O_2^-) production, which after conversion to H_2O_2 oxidizes ion channels (11). Although mice lacking the gp91 catalytic subunit of the neutrophil's oxidase have impaired O_2 sensitivity of airway chemoreceptor cells (22), the hypoxia responsiveness of carotid bodies (60), neonatal adrenal medulla (74), and pulmonary VSMCs (3) remains unaltered. Furthermore, the histological appearance of glomus cells and the modulation of the O_2 -sensitive K^+ current by P_{O_2} are also unchanged in the gp91 mutant mice (27). Surprisingly, this same group reported that genetic suppression of another component of the neutrophil's oxidase ($p47^{phox}$) results in mutant mice with increased basal activity in the carotid sinus nerve and exacerbated ventilatory response to hypoxia (63). Whether this phenotype, reflecting overexcitability of the glomus cell-afferent fiber synapse, is a nonspecific side effect of the $p47^{phox}$ deletion or whether it is due to selective alteration of the carotid body O_2 -sensing machinery is presently unknown. Although these studies may suggest that the

phagocytic NADPH oxidase is not a general O₂ sensor, other isoforms, existing in numerous tissues (see Ref. 38), could participate in O₂ sensing.

Mitochondria as possible O₂ sensor. Mitochondria have also been considered by some authors to be the site for acute O₂ sensing because, similar to hypoxia, inhibitors of the electron transport chain (ETC) and metabolic poisons stimulate the carotid body. The concept behind the “mitochondrial hypothesis” of O₂ sensing is that the lack of O₂ would reduce the activity of cytochrome *c* oxidase in complex IV, thus resulting in mitochondrial depolarization and Ca²⁺ release (72). This form of mitochondria involvement in O₂ sensing lost support after the discovery that cell responsiveness to low P_{O₂} requires membrane depolarization and Ca²⁺ entry through plasmalemmal voltage-gated channels (see Ref. 37). Nevertheless, the interest in mitochondria has resurged in the past years owing to studies testing the redox model of acute O₂ sensing (1, 34, 43, 81). Mitochondria consume almost all available O₂ and are major sources of O₂⁻ due to inefficient transfer of electrons along the respiratory chain. Although there is no general agreement on whether hypoxia decreases or increases cell ROS production, it has recently been proposed for pulmonary arterial myocytes that hypoxia is sensed by the decrease in the velocity of electron transfer from cytochrome *c* to O₂, thus leading to accumulation of ETC intermediates in the reduced state and the production of ROS. It is thought that radicals are preferentially generated at the semiubiquinone site, where an electron can leak out to produce O₂⁻ (34, 79). This view contrasts with earlier observations indicating that hypoxia shifts pulmonary VSMCs to a more reduced state (1, 84). Although participation of mitochondria in the response to hypoxia of arterial pulmonary myocytes cannot be discarded (1, 34, 43, 81), the assumption that these organelles have a general role in O₂ sensing is questioned by numerous experimental findings. O₂-sensitive maxi-K⁺ channels of rat glomus cells respond to P_{O₂} changes independently of redox modification (59), and reduction of K⁺ currents by hypoxia is maintained in airway chemoreceptor cells devoid of mitochondria or after mitochondrial inhibition (65). In whole carotid bodies, the reduced (GSH)-to-oxidized (GSSG) glutathione ratio remains unchanged during exposure to hypoxia despite the fact that this quotient increases after incubation of carotid bodies with *N*-acetylcysteine, a precursor to GSH and ROS scavenger (64). In addition, it has been shown that hypoxia responsiveness of intact glomus cells is unaffected by the complete blockade of the mitochondrial electron flow with saturating concentrations of ETC inhibitors acting at the different mitochondrial complexes. Interestingly, rotenone selectively occludes responsiveness to hypoxia, an effect not mimicked by other complex I inhibitors and unaltered by feeding electrons through complex II with succinate (48). Therefore, it seems, that although

a rotenone-inhibited molecule is essential for carotid body O₂ sensing, this phenomenon is independent of mitochondrial electron flow. Discrete rotenone binding sites outside mitochondria have not been reported, but the existence of cytosolic aggregates of preassembled complex I proteins of unknown function has been documented (see Ref. 48). Rotenone has a relatively high affinity (in the nM range) for the carotid body O₂-sensing machinery; therefore, it could be used as a probe to investigate its location and nature in glomus cells.


O₂-Dependent Hydroxylases, HIF, and Acute O₂ Sensing

Another possible form of acute O₂ sensing that has recently been explored involves O₂-dependent prolyl and asparaginyl hydroxylases, which are known to regulate the activity of hypoxia-inducible transcription factors (mainly HIF isoforms 1 α and 2 α) (42). Protein hydroxylation does not seem, however, to participate directly on acute O₂ sensing, as incubation of carotid body slices with dimethylxalylglycine, a membrane-permeant competitive inhibitor of oxoglutarate that completely inhibits hydroxylases and induces the expression of O₂-sensitive genes (16), does not alter the responsiveness of glomus cells to hypoxia (47). However, HIFs appear to be necessary for setting the appropriate level of expression of the O₂-sensing machinery in carotid body cells and pulmonary myocytes. Heterozygous HIF-1 α ^{+/-} mice, with apparently normal carotid body histology, have impaired responses to low P_{O₂} and adaptability to chronic hypoxia (33). Similarly, in HIF-1 α ^{+/-} mice, the changes of pulmonary arterial myocyte membrane potential and K⁺ channel density induced by chronic hypoxia are blunted (67).

EFFECTS OF CHRONIC HYPOXIA ON ION CHANNEL GENE EXPRESSION

Despite the progress in the understanding of the role of ion channels in the acute cellular responses to lowering O₂ tension, the participation of these proteins in the adaptive long-term changes induced by chronic hypoxia has not been studied in detail. It is known that prolonged hypoxia downregulates various Kv channel genes in pulmonary artery smooth muscle cells (55, 69, 80). Chronic hypoxia also reduces K⁺ current amplitude (26, 86) but increases the density of Na⁺ and Ca²⁺ channels in carotid body glomus cells (29, 70). Recently, detailed molecular biology and electrophysiological studies have shown that T-type Ca²⁺ channels are upregulated by hypoxia in PC12 cells and possibly in other tissues (16). A five- to eightfold increase in the level of the T-type subunit α_{1H} mRNA is found in PC12 cells when exposed to hypoxia (~20 Torr), and mRNA induction is paralleled by an increase in the density of T-type Ca²⁺ currents (Fig. 3, A–D). These

Fig. 3. Upregulation by chronic hypoxia of T-type Ca²⁺ channel gene (α_{1H}). *A*: Northern blot analysis of α_{1H} mRNAs from cells exposed to either normoxia (21% oxygen, *time 0*) or hypoxia (3% oxygen) for the indicated periods of time. The levels of the cyclophilin (CP) and tyrosine hydroxylase (TH) mRNAs were analyzed to normalize the amount of RNA in each line and to check for the ability of PC12 cells to induce the expression of an O₂-sensitive gene. *B*: average fold induction of α_{1H} mRNA levels expressed as -fold change \pm SE in the hypoxic samples compared with the normoxic sample (*time 0*). *Inset*: example of a PCR experiment (30 cycles) where β -actin was used to normalize the amount of RNA in each line. N, normoxic; H, hypoxic. *C* and *D*: electrophysiological identification of slowly deactivating T-type channels in PC12 cells. Current density due to T-type Ca²⁺ channel activity increased 2.3-fold on exposure to hypoxia (3% oxygen) for 18–24 h. *E*: alignment of the nucleotide sequences of the rat, mouse, and human α_{1H} 5'-flanking region containing 6 putative hypoxia-inducible factor (HIF) consensus DNA binding sites. The core motifs are in bold face and upperlined by arrows to indicate the plus (rightward arrow) or minus (leftward arrow) DNA strand location. Note that 2 putative HIF binding sites (indicated by letters of larger size) are conserved among the 3 species. (Modified from Ref. 16.)



observations have suggested that upregulation of T-type Ca^{2+} channels by hypoxia may contribute to cellular functions susceptible of modulation by low O_2 concentration, such as cellular excitability, differentiation, growth, and proliferation. T-type Ca^{2+} channel gene induction is also stimulated by desferroxamine, cobalt, or dimethylallylglycine. These compounds mimic hypoxia by inhibiting oxygen-, Fe^{2+} -, and oxoglutarate-dependent dioxygenases that under normoxic conditions hydroxylate specific proline and asparagine residues in HIF before its degradation (42). In addition, it has been shown that stabilization of HIF or induction of $\alpha 1\text{H}$ mRNA by hypoxia is blocked when the cells are incubated with HIF antisense oligonucleotides (16). The involvement of HIF in the hypoxic upregulation of the $\alpha 1\text{H}$ Ca^{2+} channel suggested by these experiments is further supported by the presence of hypoxia responsive elements (HIF to DNA binding sites) in the 5'-flanking region of the $\alpha 1\text{H}$ gene (Fig. 3E). This promoter region is highly conserved among mammals, with more than 71% similarity between rodents and humans and 93% similarity between rats and mice. These results indicate that T-type Ca^{2+} channels, and possibly other ion channels, are part of the gene program developed under chronic hypoxia. The data summarized in Fig. 3 represent the first example of an ion channel gene whose expression, similar to erythropoietin and other classical O_2 -sensitive genes, is regulated by the O_2 -sensitive hydroxylase-HIF pathway (9, 42, 66).

PATHOPHYSIOLOGY ASSOCIATED WITH ION CHANNEL-DEPENDENT O_2 SENSING

Primary Alterations of Acute O_2 Sensing

There are several human diseases that seem to be related to primary alterations of the acutely responding O_2 -sensitive cells. Some cases of congenital central hypoventilation syndrome (CCHS) appear without alterations in central respiratory centers but with marked decrease in the number of glomus cells and hypoplasia of carotid bodies despite a two- to threefold increase of sustentacular cells (12). In this same study, compensatory hyperplasia of the neuroepithelial bodies of the lung was also observed. In ~10–20% of patients, CCHS is associated with Hirschsprung disease, thus raising the possibility that the RET protooncogene, altered in Hirschsprung disease, participates in the mechanisms of O_2 sensing. Interestingly, RET is part of the multicomponent receptor complex of the glial cell line-derived neurotrophic factor (GDNF); in addition, both RET and GDNF are highly expressed in adult carotid bodies (76). Therefore, GDNF activation of RET is probably required for the maintenance of the O_2 sensitivity of glomus cells. Increased sustentacular cell number (28) and decreased carotid body size (45) have also been reported for some cases of sudden infant death syndrome (SIDS). Unexpected sudden death has been reported after bilateral carotid body denervation in humans and animals (17, 73), and infants prone to apnea have altered responses to mild hypoxia (6). Carotid body dysfunction in these syndromes could be the result of a primary alteration of either the O_2 sensor or the ion channels acting as effectors. Chronic exposure to hypoxia or application of chemostimulants induce glomus cell overexcitability, increased Ca^{2+} entry, and carotid body hyperplasia (29, 48, 86). Thus glomus cell hypoexcitability could underlie the hypoplasia observed in CCHS and SIDS. Perrin et al. (54) reported in

patients affected by SIDS the presence of higher carotid body dopamine content than in normal children. This could also be the cause of carotid body hypoexcitability, as it is known that dopamine inhibits Ca^{2+} currents in glomus cells (5).

Alteration of O_2 -sensitive K^+ channels could also participate in the pathophysiology of primary pulmonary hypertension, a condition characterized by increased resistance of the fine branches of the pulmonary artery. VSMCs taken from small pulmonary arteries of patients with primary pulmonary hypertension appear to be depolarized and to have higher cytosolic $[\text{Ca}^{2+}]$ levels relative to cells from patients with secondary pulmonary hypertension (88). In addition, several anorexic drugs (aminorex, fenfluramine, and others), known to produce pulmonary hypertension, have been shown to inhibit macroscopic K^+ currents in pulmonary arterial smooth muscle (83). A direct link between maxi- K^+ channels and pulmonary hypertension has been demonstrated in newborn lambs (46). Interestingly, it has recently been reported that *in vivo* transfer of Kv1.5 channels reduces pulmonary hypertension and restores HPV in chronically hypoxic rats (57).

Chronic Hypoxia and Modifications of Ion Channel Gene Expression

Different forms of chronic hypoxia (sustained or intermittent) cause alterations of various O_2 -sensitive tissues (58). Maintained reductions of O_2 tension (either in high altitude or in cages for experimental animals) induce a marked carotid body hypertrophy and blunted response to low Po_2 . It has been reported that glomus cells from chronically hypoxic carotid bodies are more excitable, due to the overexpression of Na^+ and Ca^{2+} channels, but they also have reduced voltage-dependent K^+ current amplitude (26, 29, 70, 86). Intermittent hypoxia is known to cause hypertension, secondarily to activation of arterial chemoreceptors and subsequent sympathetic stimulation (19). However, it is also possible that hypoxia causes alteration in the expression of channels that regulate smooth muscle excitability in the systemic vasculature. In the pulmonary arterial tree, chronic hypoxia reduces the amplitude of macroscopic K^+ currents (69) and downregulates various voltage-gated K^+ channels (55, 80). As described in the preceding section, the expression of T-type Ca^{2+} channel genes is increased by hypoxia in PC12 and other cell types (16). Given the broad distribution of these channels, they probably have a major role in cell adaptation to chronic hypoxia.

FUTURE RESEARCH DIRECTIONS

Over the past decade, we have witnessed a rapid development and maturation of the field of O_2 sensing due to the identification and characterization of effector molecules, particularly ion channels and transcription factors. A major recent advance has been the discovery of the O_2 -dependent hydroxylases that regulate the stabilization and transcriptional activity of HIF (42). Regarding the role of ion channels in O_2 sensing, a significant contribution has been the demonstration that, as the classical O_2 -sensitive genes, they are also regulated by the HIF signaling pathway and therefore are part of the gene program developed under chronic hypoxia (16, 66). The study of ion channel-encoding O_2 -sensitive genes will surely receive higher attention in the near future. Although the present experimental work further supports the notion that O_2 -regulated

ion channels participate in the acute cellular responses to hypoxia, there are several unresolved pivotal questions pertaining to the nature of the O₂ sensors and the mechanisms of interaction of the sensors with the ion channels. We have summarized in this review recent work that, in our view, helps to distinguish between the mechanisms that need to be explored further and those that lack experimental support. It is possible that acute O₂ sensing does not utilize a single mechanism but that different O₂ sensors are expressed in the various hypoxia responsive cells. Nevertheless, new experimental work is urgently needed to fully characterize the involvement of ion channels in the hypoxia signaling pathway. The combination of biophysical, molecular biology, and pharmacological techniques applied to in vitro preparations and animal models are experimental approaches that are already yielding important results. These techniques are helping to clarify the critical role of ion channels in hypoxic pulmonary hypertension (57, 80, 83, 88). Similarly, rotenone, a drug that selectively occludes sensitivity to hypoxia of carotid body glomus cells (48), could be a useful tool to search for the location and nature of the carotid body O₂ sensor. Finally, it can be expected that this basic knowledge will have a medical impact. Ion channels and other O₂-sensitive effectors are involved in vasomotor and cardiorespiratory control, and localized lacks of O₂ are critical in the pathogenesis of major causes of mortality. Therefore, amplification or inhibition of adaptive responses to hypoxia is a promising pharmacological strategy that may result in effective therapies for human diseases.

ACKNOWLEDGMENTS

The authors thank María García-Fernández for comments on the manuscript.

GRANTS

J. López-Barneo is a recipient of the "Ayuda a la Investigación 2000" of the Juan March Foundation.

REFERENCES

1. Archer SL, Huang J, Henry T, Peterson D, and Weir EK. A redox-based O₂ sensor in rat pulmonary vasculature. *Circ Res* 73: 1100–1112, 1993.
2. Archer SL, London B, Hampl V, Wu X, Nsair A, Puttagunta L, Hashimoto K, Waite RE, and Michelakis ED. Impairment of hypoxic pulmonary vasoconstriction in mice lacking the voltage-gated potassium channel Kv 1.5. *FASEB J* 15: 1801–1803, 2001.
3. Archer SL, Reeve HL, Michelakis E, Puttagunta L, Waite R, Nelson DP, Dinauer MC, and Weir EK. O₂ sensing is preserved in mice lacking the gp91 phox subunit of NADPH oxidase. *Proc Natl Acad Sci USA* 96: 7944–7949, 1999.
4. Archer SL, Souil E, Dinh-Xuan AT, Chremmer B, Mercier JC, Yaagoubi AE, Nguyen-Huu L, Reeve HL, and Hampl V. Molecular identification of the role of voltage-gated K⁺ channels, Kv 1.5 and Kv 2.1, in hypoxic pulmonary vasoconstriction and control of resting membrane potential in rat pulmonary arterial myocytes. *J Clin Invest* 101: 2319–2330, 1998.
5. Benot AR and López-Barneo J. Feedback inhibition of Ca²⁺ currents by dopamine in glomus cells of the carotid body. *Eur J Neurosci* 2: 809–812, 1990.
6. Brady JP, Ariagno RL, Watts JL, Goldman SL, and Dumpit FM. Apnea, hypoxemia, and aborted sudden infant death syndrome. *Pediatrics* 62: 686–691, 1978.
7. Buckler KJ. A novel oxygen-sensitive potassium current in rat carotid body type I cells. *J Physiol* 498: 649–662, 1997.
8. Buckler KJ and Vaughan-Jones RD. Effects of hypoxia on membrane potential and intracellular calcium in rat neonatal carotid body type I cells. *J Physiol* 476: 423–428, 1994.
9. Bunn HF and Poyton RO. Oxygen sensing and molecular adaptation to hypoxia. *Physiol Rev* 76: 839–885, 1996.
10. Coppock EA and Tamkun MM. Differential expression of Kv channel α - and β -subunits in the bovine arterial pulmonary circulation. *Am J Physiol Lung Cell Mol Physiol* 281: L1350–L1360, 2001.
11. Cross AR, Henderson L, Jones OT, Delpiano MA, Hentschel J, and Acker H. Involvement of an NAD(P)H oxidase as a Po₂ sensor protein in the rat carotid body. *Biochem J* 272: 743–747, 1990.
12. Cutz E, Ma TKF, Perrin DG, Moore AM, and Becker LE. Peripheral chemoreceptors in congenital central respiratory syndrome. *Am J Respir Crit Care Med* 155: 358–363, 1997.
13. Dart C and Standen NB. Activation of ATP-dependent K⁺ channels by hypoxia in smooth muscle cells isolated from the pig coronary artery. *J Physiol* 483: 29–39, 1995.
14. Davies ARL and Kozlowski RZ. Kv channel subunit expression in rat pulmonary arteries. *Lung* 179: 147–161, 2001.
15. Delpiano MA and Hescheler J. Evidence for a Po₂-sensitive K⁺ channel in the type-I cell of the rabbit carotid body. *FEBS Lett* 249: 195–198, 1989.
16. Del Toro R, Levitsky KL, López-Barneo J, and Chiara MD. Induction of T-type calcium channel gene expression by chronic hypoxia. *J Biol Chem* 278: 22316–22324, 2003.
17. Donnelly DF and Haddad GG. Prolonged apnea and impaired survival in piglets after sinus and aortic nerve section. *J Appl Physiol* 68: 1048–1052, 1990.
18. Fearon IM, Palmer AC, Balmforth AJ, Ball SG, Mikala G, Schwartz A, and Peers C. Hypoxia inhibits the recombinant α 1C subunit of the human cardiac L-type Ca²⁺ channel. *J Physiol* 500: 551–556, 1997.
19. Fletcher EC. Physiological consequences of intermittent hypoxia: systemic blood pressure. *J Appl Physiol* 90: 1600–1605, 2001.
20. Franco-Obregón A and López-Barneo J. Low Po₂ inhibits calcium channel activity in arterial smooth muscle cells. *Am J Physiol Heart Circ Physiol* 271: H2290–H2299, 1996.
21. Franco-Obregón A, Ureña J, and López-Barneo J. Oxygen-sensitive calcium channels in vascular smooth muscle and their possible role in hypoxic arterial relaxation. *Proc Natl Acad Sci USA* 92: 4715–4719, 1995.
22. Fu XW, Wang D, Nurse CA, Dinauer MC, and Cutz E. NADPH oxidase is an O₂ sensor in airway chemoreceptors: evidence from K⁺ current modulation in wild-type and oxidase-deficient mice. *Proc Natl Acad Sci USA* 97: 4374–4379, 2000.
23. Ganfornina MD and López-Barneo J. Single K⁺ channels in membrane patches of arterial chemoreceptor cells are modulated by O₂ tension. *Proc Natl Acad Sci USA* 88: 2927–2930, 1991.
24. Gebremedhin D, Bonnet P, Greene AS, England SK, Rusch NJ, Lombard JH, and Harder DR. Hypoxia increases the activity of Ca²⁺-sensitive K⁺ channels in cat cerebral arterial muscle cell membranes. *Pflügers Arch* 428: 621–630, 1994.
25. Hartness ME, Lewis A, Searle GJ, O'Kelly I, Peers C, and Kemp PJ. Combined antisense and pharmacological approaches implicate hTASK as an airway O₂ sensing K⁺ channel. *J Biol Chem* 276: 26499–26508, 2001.
26. Hatton CJ, Carpenter E, Pepper DR, Kumar P, and Peers C. Developmental changes in isolated rat type I carotid body cell K⁺ currents and their modulation by hypoxia. *J Physiol* 501: 49–58, 1997.
27. He L, Chen J, Dinger B, Sanders K, Sundar K, Hoidal J, and Fidone S. Characteristics of carotid body chemosensitivity in NADPH oxidase-deficient mice. *Am J Physiol Cell Physiol* 282: C27–C33, 2002.
28. Heath D, Khan Z, and Smith P. Histopathology of the carotid body in neonates and infants. *Histopathology* 17: 511–520, 1990.
29. Hempleman SC. Increased calcium current in carotid body glomus cells following in vivo acclimatization to chronic hypoxia. *J Neurophysiol* 76: 1880–1886, 1996.
30. Hogg DS, Davies AR, McMurray G, and Kozlowski RZ. Kv2.1 channels mediate hypoxic inhibition of IKv in native pulmonary arterial smooth muscle cells of the rat. *Cardiovasc Res* 55: 233–235, 2002.
31. Hulme JT, Coppock EA, Felipe A, Martens JR, and Tamkun MM. Oxygen sensitivity of cloned voltage-gated K⁺ channels expressed in the pulmonary vasculature. *Circ Res* 85: 489–497, 1999.
32. Jiang C and Haddad GG. A direct mechanism for sensing low oxygen levels by central neurons. *Proc Natl Acad Sci USA* 91: 7198–7201, 1994.
33. Kline DD, Peng YJ, Manalo DJ, Semenza GL, and Prabhakar NR. Defective carotid body function and impaired ventilatory responses to chronic hypoxia in mice partially deficient for hypoxia-inducible factor 1 α . *Proc Natl Acad Sci USA* 99: 821–826, 2002.

34. Leach RM, Hill HM, Snetkov VA, Robertson TP, and Ward JPT. Divergent roles of glycolysis and the mitochondrial electron transport chain in hypoxic pulmonary vasoconstriction of the rat: identity of the hypoxic sensor. *J Physiol* 536: 211–224, 2001.
35. Lewis A, Hartness ME, Chapman CG, Fearon IM, Meadows HJ, Peers C, and Kemp PJ. Recombinant hTASK1 is an O₂-sensitive K⁺ channel. *Biochem Biophys Res Commun* 285:1290–1294, 2001.
36. Lewis A, Peers C, Ashford ML, and Kemp PJ. Hypoxia inhibits human recombinant large conductance, Ca²⁺-activated K⁺ (maxi-K) channels by a mechanism which is membrane delimited and Ca²⁺ sensitive. *J Physiol* 540: 771–780, 2002.
37. López-Barneo J. Oxygen-sensitive ion channels: how ubiquitous are they? *Trends Neurosci* 17: 133–135, 1994.
38. López-Barneo J. Oxygen and glucose sensing by carotid body glomus cells. *Curr Opin Neurobiol* 13: 493–499, 2003.
39. López-Barneo J, López-López JR, Ureña J, and González C. Chemotransduction in the carotid body: K⁺ current modulated by PO₂ in type I chemoreceptor cells. *Science* 241: 580–582, 1988.
40. López-Barneo J, Pardal R, and Ortega-Sáenz P. Cellular mechanisms of oxygen sensing. *Annu Rev Physiol* 63: 259–287, 2001.
41. Lopez-Lopez J, Gonzalez C, Ureña J, and Lopez-Barneo J. Low PO₂ selectively inhibits K channel activity in chemoreceptor cells of the mammalian carotid body. *J Gen Physiol* 93: 1001–1015, 1989.
42. Masson N and Ratcliffe PJ. HIF prolyl and asparaginyl hydroxylases in the biological response to intracellular O₂ levels. *J Cell Sci* 116: 3041–3049, 2003.
43. Michelakis ED, Hampl V, Nsair A, Wu X, Harry G, Haromy A, Gurtu R, and Archer SL. Diversity in mitochondrial function explains differences in vascular oxygen sensing. *Circ Res* 90: 1307–1315, 2002.
44. Montoro R, Ureña J, Fernández-Chacón R, Álvarez de Toledo G, and López-Barneo J. Oxygen sensing by ion channels and chemotransduction in single glomus cells. *J Gen Physiol* 107: 133–143, 1996.
45. Naeye RL, Fischer R, Ryser M, and Waken P. Carotid body in the sudden infant death syndrome. *Science* 191: 567–569, 1976.
46. Olschewski A, Hong Z, Linden BC, Porter VA, Weir EK, and Cornfield DN. Contribution of the K(Ca) channel to membrane potential and O₂ sensitivity is decreased in an ovine PPHN model. *Am J Physiol Lung Cell Mol Physiol* 283: L11103–L11109, 2002.
47. Ortega-Sáenz P, García-Fernández M, Pardal R, Alvarez E, and López-Barneo J. Studies on glomus cell sensitivity to hypoxia in carotid body slices. *Adv Exp Med Biol* 536: 65–73, 2003.
48. Ortega-Sáenz P, Pardal R, García-Fernández M, and López-Barneo J. Rotenone selectively occludes sensitivity to hypoxia in rat carotid body glomus cells. *J Physiol* 548: 789–800, 2003.
49. Osipenko ON, Evans AM, and Gurney AM. Regulation of the resting potential of rabbit pulmonary artery myocytes by a low threshold, O₂-sensing potassium current. *Br J Pharmacol* 120: 1461–1470, 1997.
50. Osipenko ON, Tate RJ, and Gurney AM. Potential role for Kv3.1b channels as oxygen sensors. *Circ Res* 86: 534–540, 2000.
51. Pardal R, Ludewig U, García-Hirschfeld J, and López-Barneo J. Secretory responses of intact glomus cells in thin slices of rat carotid body to hypoxia and tetraethylammonium. *Proc Natl Acad Sci USA* 97: 2361–2366, 2000.
52. Patel AJ, Lazdunski M, and Honoré E. Kv2.1/Kv9.3, a novel ATP-dependent delayed-rectifier K⁺ channel in oxygen-sensitive pulmonary artery myocytes. *EMBO J* 16: 6615–6625, 1997.
53. Peers C. Hypoxic suppression of K⁺ currents in type I carotid body cells: selective effect on the Ca²⁺-activated K⁺ current. *Neurosci Lett* 119: 253–256, 1990.
54. Perrin DG, Cutz E, Becker LE, Bryan AC, Madapallimatum A, and Sole MJ. Sudden infant death syndrome: increased carotid body dopamine and noradrenaline content. *Lancet* 2: 535–537, 1984.
55. Platoshyn O, Yu Y, Golovina VA, McDaniel SS, Krick S, Li L, Wang JY, Rubin LJ, and Yuan JX. Chronic hypoxia decreases Kv channel expression and function in pulmonary artery myocytes. *Am J Physiol Lung Cell Mol Physiol* 280: L801–L812, 2001.
56. Post JM, Hume JR, Archer SL, and Weir EK. Direct role for potassium channel inhibition in hypoxic pulmonary vasoconstriction. *Am J Physiol Cell Physiol* 262: C882–C890, 1992.
57. Pozeg ZI, Michelakis ED, McMurtry MS, Thebaud B, Wu XC, Dyck JR, Hashimoto K, Wang S, Moudgil R, Harry G, Sultanian R, Koshal A, and Archer SL. In vivo gene transfer of the O₂-sensitive potassium channel Kv1.5 reduces pulmonary hypertension and restores hypoxic pulmonary vasoconstriction in chronically hypoxic rats. *Circulation* 107: 2037–2044, 2003.
58. Prabhakar N. Oxygen sensing during intermittent hypoxia: cellular and molecular mechanisms. *J Appl Physiol* 90: 1986–1994, 2001.
59. Riesco-Fagundo AM, Pérez-García MT, González C, and López-López JR. O₂ modulates large conductance Ca²⁺-dependent K⁺ channels of rat chemoreceptor cells by a membrane-restricted and CO-sensitive mechanism. *Circ Res* 89: 430–436, 2001.
60. Roy A, Rozanov C, Mokashi A, Daudu P, Al-Medhi AB, Shams H, and Lahiri S. Mice lacking gp91 phox subunit of NAD(P)H oxidase showed glomus cell [Ca²⁺]_i and respiratory responses to hypoxia. *Brain Res* 872: 188–193, 2000.
61. Rychkov GY, Adams MB, McMillen IC, and Roberts ML. Oxygen-sensing mechanisms are present in the chromaffin cells of the sheep adrenal medulla before birth. *J Physiol* 509: 887–893, 1998.
62. Sánchez D, López-López JR, Pérez-García MT, Sanz-Alfayate G, Obeso A, Ganfornina MD, and González C. Molecular identification of Kv α subunits that contribute to the oxygen-sensitive K⁺ current of chemoreceptor cells of the rabbit carotid body. *J Physiol* 542: 369–382, 2002.
63. Sanders K, Sundar K, He L, Dinger B, Fidone S, and Hoidal JR. Role of components of the phagocytic NADPH oxidase in oxygen sensing. *J Appl Physiol* 93: 1357–1364, 2002.
64. Sanz-Alfayate G, Obeso A, Agapito MT, and González C. Reduced to oxidized glutathione ratios and oxygen sensing in calf and rabbit carotid body chemoreceptor cells. *J Physiol* 537: 209–220, 2001.
65. Searle GJ, Hartness ME, Hoareau R, Peers C, and Kemp PJ. Lack of contribution of mitochondrial electron transport to acute O₂ sensing in model airway chemoreceptors. *Biochem Biophys Res Commun* 291: 332–337, 2002.
66. Semenza GL. HIF-1: mediator of physiological and pathophysiological responses to hypoxia. *J Appl Physiol* 88: 1474–1480, 2000.
67. Shimoda LA, Manalo DJ, Sham JSK, Semenza GL, and Silvestre JT. Partial HIF-1 α deficiency impairs pulmonary arterial myocyte electrophysiological responses to hypoxia. *Am J Physiol Lung Cell Mol Physiol* 281: L202–L208, 2001.
68. Smani T, Hernandez A, Ureña J, Castellano AG, Franco-Obregón A, Ordoñez A, and López-Barneo J. Reduction of Ca²⁺ channel activity by hypoxia in human and porcine coronary myocytes. *Cardiovasc Res* 53: 97–104, 2002.
69. Smirnov SV, Robertson TP, Ward JPT, and Aaronson PI. Chronic hypoxia is associated with reduced delayed rectifier K⁺ current in rat pulmonary artery smooth muscle. *Am J Physiol Heart Circ Physiol* 266: H365–H370, 1994.
70. Stea A, Jackson A, and Nurse CA. Hypoxia and N₆,O₂'-dibutyryladenosine 3',5'-cyclic monophosphate, but not nerve growth factor, induce Na⁺ channels and hypertrophy in chromaffin-like arterial chemoreceptors. *Proc Natl Acad Sci USA* 89: 9469–9473, 1992.
71. Stea A and Nurse CA. Whole-cell and perforated-patch recordings from O₂-sensitive rat carotid body cells grown in short- and long-term culture. *Pflügers Arch* 418: 93–101, 1991.
72. Streller T, Huckstorf C, Pfeiffer C, and Acker H. Unusual cytochrome a₅₉₂ with low PO₂ affinity correlates as putative oxygen sensor with rat carotid body chemoreceptor discharge. *FASEB J* 16: 1277–1279, 2002.
73. Sullivan CE. Bilateral carotid body resection in asthma: vulnerability to hypoxic death in sleep. *Chest* 78: 354, 1980.
74. Thompson RJ, Farragher SM, Cutz E, and Nurse CA. Developmental regulation of O₂ sensing in neonatal adrenal chromaffin cells from wild-type and NADPH-oxidase-deficient mice. *Pflügers Arch* 444: 539–548, 2002.
75. Thompson RJ, Jackson A, and Nurse CA. Developmental loss of hypoxic chemosensitivity in rat adrenomedullary chromaffin cells. *J Physiol* 498: 503–510, 1997.
76. Toledo-Aral JJ, Méndez-Ferrer S, Pardal R, Echevarría M, and López-Barneo J. Trophic restoration of the nigrostriatal dopaminergic pathway in long-term carotid body-grafted parkinsonian rats. *J Neurosci* 23: 141–148, 2003.
77. Tristani-Firouzi M, Reeve HL, Tolarova S, Weir EK, and Archer SL. Oxygen-induced constriction of rabbit ductus arteriosus occurs via inhibition of a 4-aminopyridine-, voltage-sensitive potassium channel. *J Clin Invest* 98: 1959–1965, 1996.
78. Ureña J, Fernández-Chacón R, Benot AR, Álvarez de Toledo G, and López-Barneo J. Hypoxia induces voltage-dependent Ca²⁺ entry and

- quantal dopamine secretion in carotid body glomus cells. *Proc Natl Acad Sci USA* 91: 10208–10211, 1994.
79. **Ureña J, Franco-Obregón A, and López-Barneo J.** Contrasting effects of hypoxia on cytosolic Ca^{2+} spikes in conduit and resistance myocytes of the rabbit pulmonary artery. *J Physiol* 496: 103–109, 1996.
80. **Wang J, Juhaszova M, Rubin LJ, and Yuan XJ.** Hypoxia inhibits gene expression of voltage-gated K^+ channel α subunits in pulmonary artery smooth muscle cells. *J Clin Invest* 100: 2347–2353, 1997.
81. **Waypa GB, Chandel NS, and Schumacker PT.** Model for hypoxic pulmonary vasoconstriction involving mitochondrial oxygen sensing. *Circ Res* 88: 1259–1266, 2001.
82. **Weir EK and Archer SL.** The mechanism of acute hypoxic pulmonary vasoconstriction: the tale of two channels. *FASEB J* 9: 183–189, 1995.
83. **Weir EK, Reeve HL, Huang JM, Michelakis E, Nelson DP, Hampl V, and Archer SL.** Anorexic agents aminorex, fenfluramine, and dexfenfluramine inhibit potassium current in rat pulmonary vascular smooth muscle and cause pulmonary vasoconstriction. *Circulation* 94: 2216–2220, 1996.
84. **White CW, Jackson JH, McMurtry IF, and Repine JE.** Hypoxia increases glutathione redox cycle and protects rat lungs against oxidants. *J Appl Physiol* 65: 2607–2616, 1988.
85. **Wyatt CN and Peers C.** Ca^{2+} -activated K^+ channels in isolated type I cells of the neonatal rat carotid body. *J Physiol* 483: 559–565, 1995.
86. **Wyatt CN, Wright C, Bee D, and Peers C.** O_2 -sensitive K^+ currents in carotid body chemoreceptor cells from normoxic and chronically hypoxic rats and their roles in hypoxic chemotransduction. *Proc Natl Acad Sci USA* 92: 295–299, 1995.
87. **Youngson C, Nurse C, Yeger H, and Cutz E.** Oxygen sensing in airway chemoreceptors. *Nature* 365: 153–155, 1993.
88. **Yuan JX, Aldinger AM, Juhaszova M, Wang J, Conte JV Jr, Gaine SP, Orens JB, and Rubin LJ.** Dysfunctional voltage-gated K^+ channels in pulmonary artery smooth muscle cells of patients with primary pulmonary hypertension. *Circulation* 98: 1400–1406, 1998.
89. **Yuan XJ.** Oxygen sensitive ion channel(s): where and what? *Am J Physiol Lung Cell Mol Physiol* 281: L1345–L1349, 2001.
90. **Yuan XJ, Goldman WF, Tod ML, Rubin LJ, and Blaustein MP.** Hypoxia reduces potassium currents in cultured rat pulmonary but not mesenteric arterial myocytes. *Am J Physiol Lung Cell Mol Physiol* 264: L116–L123, 1993.
91. **Yuan XJ, Tod ML, Rubin LJ, and Blaustein MP.** Contrasting effects of hypoxia on tension in rat pulmonary and mesenteric arteries. *Am J Physiol Heart Circ Physiol* 259: H281–H289, 1990.
92. **Zhu WH, Conforti L, Czyzyk-Krzeska MF, and Millhorn DE.** Membrane depolarization in PC12 cells during hypoxia is regulated by an O_2 -sensitive K^+ current. *Am J Physiol Cell Physiol* 271: C658–C665, 1996.

