

# A plant proton-pumping inorganic pyrophosphatase functionally complements the vacuolar ATPase transport activity and confers bafilomycin resistance in yeast

José R. PÉREZ-CASTIÑEIRA, Agustín HERNÁNDEZ, Rocío DRAKE and Aurelio SERRANO<sup>1</sup>

Instituto de Bioquímica Vegetal y Fotosíntesis, Universidad de Sevilla-CSIC, Avda. Americo Vespucio, 49, 41092 Sevilla, Spain

V-ATPases (vacuolar H<sup>+</sup>-ATPases) are a specific class of multi-subunit pumps that play an essential role in the generation of proton gradients across eukaryotic endomembranes. Another simpler proton pump that co-localizes with the V-ATPase occurs in plants and many protists: the single-subunit H<sup>+</sup>-PPase [H<sup>+</sup>-translocating PPase (inorganic pyrophosphatase)]. Little is known about the relative contribution of these two proteins to the acidification of intracellular compartments. In the present study, we show that the expression of a chimaeric derivative of the *Arabidopsis thaliana* H<sup>+</sup>-PPase AVP1, which is preferentially targeted to internal membranes of yeast, alleviates the phenotypes associated with V-ATPase deficiency. Phenotypic complementation was achieved both with a yeast strain with its V-ATPase specifically inhibited by bafilomycin A1 and with a *vma1*-null mutant lacking a catalytic V-ATPase subunit. Cell staining with vital fluorescent dyes showed that AVP1 recovered vacuole acidification and normalized the endocytic pathway of the

*vma* mutant. Biochemical and immunochemical studies further demonstrated that a significant fraction of heterologous H<sup>+</sup>-PPase is located at the vacuolar membrane. These results raise the question of the occurrence of distinct proton pumps in certain single-membrane organelles, such as plant vacuoles, by proving yeast V-ATPase activity dispensability and the capability of H<sup>+</sup>-PPase to generate, by itself, physiologically suitable internal pH gradients. Also, they suggest new ways of engineering macrolide drug tolerance and outline an experimental system for testing alternative roles for fungal and animal V-ATPases, other than the mere acidification of subcellular organelles.

**Key words:** chimaeric membrane protein, green fluorescent protein, heterologous expression, proton-translocating inorganic pyrophosphatase, *Saccharomyces cerevisiae*, vacuolar-type ATPase.

## INTRODUCTION

The occurrence of an electrochemical proton gradient across the internal membranes of eukaryotic cells is one of the known requirements for the correct function of the endo- and exo-cytic pathways [1]. Moreover, pH gradients are necessary not only for the correct functioning of the vesicle transport machinery, but also for many other organellar functions, such as, among others, dissociation of ligand–receptor complexes in receptor-mediated endocytosis or glycosylation of proteins in the Golgi apparatus [2]. A specific class of proton-pumping ATPases, V-ATPases (vacuolar ATPases or V-type ATPases), play an essential role in the generation of these positive-inside proton gradients [1,3]. V-ATPases were originally described in the vacuolar membrane of plants and fungi (hence their name), thereafter they were found in many single-membrane organelles of eukaryotic cells. In addition, they have also been reported to occur at the plasma membrane of certain mammalian cell types [2]. All in all, these multi-subunit proteins fulfil important roles in diverse scenarios, both physiological and in disease, including cancer [3,4]. Not surprisingly then, there is an active search for specific inhibitors of these pumps that could be used both in basic research and in therapeutic applications. Bafilomycin A1 was the first reported specific inhibitor of V-ATPases [5], it consists of a large macrocyclic lactone ring which places it in the macrolide antibiotic group. Unlike other eukaryotes,

*Saccharomyces cerevisiae* cells can grow either in the presence of V-ATPase inhibitors [6] or when a null mutation occurs in one of the genes encoding certain subunits of the V-ATPase complex [1]. However, they show a range of conditional-lethal phenotypes and thus become very sensitive to neutral/alkaline pH values [6,7], high extracellular calcium concentrations [8] and certain metal cations, such as zinc [9]; actually, the combination of sensitivity to high pH and extracellular calcium is widely accepted as the most characteristic *Vma*<sup>−</sup> phenotype [1].

V-ATPases are very complex multi-subunit pumps that contain both membrane-bound and peripheral components. This paradoxical complexity, in comparison with other transporters, has been justified by proposing moonlighting functions of some subunits or even whole domains of the V-ATPases [10,11]. However, to study non-transport functions of V-ATPases is a complicated task if organelle acidification is not taken over by some other protein(s). Strikingly, this situation may be the natural scenario in plant cells and many protists where V-ATPases co-exist with alternative proton pumps. However, to date, no work has proven that V-ATPases can be unambiguously substituted by any other proton pump *in vivo*.

H<sup>+</sup>-PPases [proton-translocating PPases (inorganic pyrophosphatases)] are a type of homo-oligomeric membrane-embedded proteins containing a single highly hydrophobic subunit type that couples the energy obtained in the hydrolysis of PPI (inorganic pyrophosphate) to the transport of protons

Abbreviations used: ACMA, 9-amino-6-chloro-2-methoxy acridine; DIC, differential interference contrast; DTT, dithiothreitol; GFP, green fluorescent protein; H<sup>+</sup>-PPase, H<sup>+</sup>-translocating inorganic pyrophosphatase; PPase, inorganic pyrophosphatase; PPI, inorganic pyrophosphate; V-ATPase, vacuolar H<sup>+</sup>-ATPase; yEGFP, yeast enhanced GFP.

<sup>1</sup>To whom correspondence should be addressed (email aurelio@ibvf.csic.es).

across biological membranes [12–14]. They are the structurally simplest class of primary proton pumps known to date, and was the last one established, in the middle of the 1980s [12,13].  $H^+$ -PPases are reportedly located in cell membrane vesicles and acidocalcisomes of prokaryotes and diverse acidic single-membrane organelles (acidocalcisomes, lysosomes, vacuoles) of certain eukaryotes: protists and plants [13,15–17].  $H^+$ -PPases seem to co-localize with V-ATPases in those eukaryotes where they occur [18], thus, one point of discussion is the physiological significance of this apparent functional redundancy and whether  $H^+$ -PPases contribute in varying degrees to the generation of proton gradients under diverse physiological scenarios.

We recently reported that translational fusions with certain N-terminal signal peptides enhance the expression levels and/or alter the subcellular distribution of diverse  $H^+$ -PPases heterologously expressed in *S. cerevisiae*. When this approach was applied to the vacuolar  $H^+$ -PPase AVP1 from *Arabidopsis thaliana*, a protein expressed at moderate levels in yeast in its natural form [19], its subcellular distribution was altered; thus, chimaeric proteins constructed by fusing the N-terminal signal sequence of the  $H^+$ -PPase from the protist *Trypanosoma cruzi* [20] to AVP1 were shown to preferentially accumulate in intracellular acidic vesicles [21]. In the present study, we show that the expression of such an internal membrane-targeted chimaeric AVP1 derivative confers yeast resistance to the antibiotic macrolide bafilomycin A1, a specific V-ATPase inhibitor. Further experiments performed with a yeast *vma* mutant defective in its V-ATPase activity demonstrated that the  $H^+$ -PPase acts by functionally complementing the transport function of the former proton pump. These results raise again the question of the presence of both proton pumps in certain organelles, such as the plant vacuole. Potential applications of this system in order to elucidate the roles of V-ATPases and/or  $H^+$ -PPases are also discussed.

## EXPERIMENTAL

### Yeast strains

The YPC3 mutant strain was generated from the *S. cerevisiae* haploid strain W303-1A (MATa, *ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1, ura3-1*) by the single-step transplacement procedure as described previously [21]. This mutant has the *IPP1* gene coding for the cytosolic soluble PPase under the control of the yeast galactokinase (*GALI*) promoter. YPC4 was generated from *S. cerevisiae* mutant RS-1144 (W303-1A *vma1::LEU2*) following the same procedure. Mutant RS-1144, provided by Professor Ramón Serrano (Universidad Politécnica de Valencia, Valencia, Spain), was derived from strain W303-1A by disrupting its *VMA1* (*TFP1*) gene with the yeast *LEU2* cassette [22]. The *VMA1* gene of *S. cerevisiae* is translated by protein splicing into two different proteins: the 69 kDa catalytic subunit of the vacuolar  $H^+$ -ATPase and a 50 kDa DNA endonuclease [23]. Vacuolar protease-deficient mutant BJ5457 was provided by Professor Andrés Aguilera (Universidad de Sevilla, Sevilla, Spain). This mutant carries mutations both in the *PEP4* gene and the *PRB1* gene, coding for the yeast vacuolar proteases PrA (Proteinase A) and PrB (Proteinase B) respectively [24]. Yeast strains used in the present study are described in Supplementary Table S1 at <http://www.BiochemJ.org/bj/437/bj4370269add.htm>.

### Transformation of YPC3 and YPC4

*S. cerevisiae* mutant strains YPC3 and YPC4 were transformed with plasmids pRS699b (negative control), pAVP1, pTcAVP1 and pTcGFP-VP1 by using the method described by Schiestl and Gietz [25]. All plasmids are 2  $\mu$ m-based derivatives of the *URA3*-containing *Escherichia coli/S. cerevisiae* shuttle plasmid pRS699

[26], as described previously [21] (Supplementary Table S2 in [21]). These plasmids bear the yeast *PMA1* promoter for constitutive expression of inserts [26]. Three different constructs were made using the sequence coding for the  $K^+$ -dependent  $H^+$ -PPase from the higher plant *A. thaliana* (*AVP1*): pAVP1, carrying an intron-less open reading frame construct of AVP1; pTcAVP1, a 5' in-frame fusion of the sequence coding for the N-terminal signal peptide of the *T. cruzi*  $H^+$ -PPase (TcVP) with that of AVP1; and pTcGFP-VP1, an in-frame fusion where the sequence coding for yEGFP [yeast enhanced GFP (green fluorescent protein)] is placed between the coding sequence of the TcVP N-terminal signal peptide and that of AVP1. In addition to these, plasmids pIPP1, bearing the whole open reading frame encoding the cytosolic soluble PPase from yeast (*IPP1*), and pVMA1, bearing the genomic sequence of *VMA1* coding for the A catalytic subunit of the yeast V-ATPase, were constructed using the above mentioned expression plasmid backbone (Supplementary Figure S1 at <http://www.BiochemJ.org/bj/437/bj4370269add.htm>). YPC3 and YPC4 cells were initially grown at 30 °C in galactose-containing synthetic medium [27] devoid of histidine (YPC3) or histidine and leucine (YPC4); transformants were selected by growing cells on 2% agar plates in culture medium without histidine and uracil (YPC3) or without histidine, leucine and uracil (YPC4). Sorbitol (0.5 M) was added to the plates where transformants of YPC4 were selected as an osmotic protectant.

### Phenotype complementation tests

Complementation studies were performed by pre-inoculating 2 ml of galactose-containing selective medium with transformed cells from the plates and growing overnight at 30 °C with agitation (200 rev./min). The following day, 2 ml of glucose-containing selective medium or YPD medium [1% (w/v) yeast extract/2% (w/v) peptone/2% (w/v) glucose] were inoculated with 20  $\mu$ l (1:100 dilution) of cells previously grown on galactose and these were allowed to grow as described above. This treatment is necessary to bring down the PPase activity associated with Ipp1p [28]. After overnight growth on glucose, 10-fold serial dilutions of the cultures were made in sterile water and 5  $\mu$ l drops of each dilution were spotted on to YPD and YPGal [1% (w/v) yeast extract/2% (w/v) peptone/2% (w/v) galactose] agar plates containing 50 mM Tris/Mes adjusted to pH 5.5, 7.5 or 8.  $CaCl_2$  (100 mM) or  $ZnCl_2$  (4 mM) was added to the plates at pH 5.5 where indicated. Plates were typically grown at 30 °C for 2–5 days.

### Fluorescence microscopy

Vacuolar lumen acidification was assessed by fluorescence staining with quinacrine as described previously [29]. Cells were visualized after 10 min of fluorophore accumulation with a fully automated Leica DM6000B microscope (Leica Microsystems) with FITC green fluorescence filters (excitation filter 480/40 nm, dichromatic mirror 505 nm, suppression filter 527/30 nm), a 100 $\times$  objective and equipped with a cooled CCD (charge-coupled device) camera (ORCA-AG, Hamamatsu Photonics). Vacuolar membrane labelling was carried out with the vital lipophilic dye FM4-64 as described previously [29]. Cells were visualized 30 min after incubation with the dye as described above, except that Texas Red filters (excitation filter 560/40 nm, dichromatic mirror 595 nm, suppression filter 645/75 nm) were used. In all cases, cells were also viewed using DIC (differential interference contrast) microscopy to observe cell morphology.

### Isolation of yeast vacuolar membranes

Colonies of transformed YPC4 cells were collected from a plate and liquid-grown up to stationary phase in galactose-containing

selective medium; then, 4 ml of stationary cultures were used to inoculate 400 ml of YPD adjusted to pH 5.5 with 50 mM Tris/Mes and grown overnight to early exponential phase ( $A_{660}$  was approximately 1). Yeast vacuoles were obtained from these cultures using the method described by Roberts et al. [30]

### Isolation of total membrane fractions from yeast

Preparation of membrane fractions from YPC4 cells transformed with the different plasmids were obtained by a modification of a method described previously [21]. Yeast colonies were collected from a plate and liquid-grown up to stationary phase in galactose-containing selective medium; then, 4 ml of stationary cultures were used to inoculate 400 ml of YPD. After overnight growth, cells were sedimented by centrifugation at 700 *g* for 10 min, washed thoroughly with water, resuspended in 5 ml of ice-cold buffer A [25 mM Tris/HCl (pH 8), 10% glycerol, 4 mM 2-mercaptoethanol, 2 mM DTT (dithiothreitol), 2 mM EDTA, 10 mM MgCl<sub>2</sub>, 1 mM benzamidine, 2 mM  $\epsilon$ -aminocaproic acid and 1 mM PMSF], and homogenized by vigorous shaking with glass beads. The homogenate was diluted up to 25 ml with buffer B [10 mM Tris/HCl (pH 7.6), 10% glycerol, 2 mM DTT and 1 mM EDTA] and centrifuged for 10 min at 700 *g* to remove beads and debris. The resulting supernatant was centrifuged for 30 min at 40 000 rev./min (Beckman 60Ti rotor) thus yielding a total membrane pellet. This pellet was homogenized in stripping buffer [60 mM Tris/HCl (pH 8), 12% glycerol, 0.72 M KCl and 1.2 mM CaCl<sub>2</sub>] [31] and centrifuged for 20 min at 40 000 rev./min (Beckman 60Ti rotor). This pellet was subsequently washed with buffer B and centrifuged [40 000 rev./min (Beckman 60Ti rotor) for 20 min]. The final pellets were resuspended and homogenized in 1 ml of buffer B.

### PPase activity and H<sup>+</sup>-translocation assays

Membrane-associated PPase activity was measured by spectrophotometric detection of released phosphate [32]. Proton-translocation activities were assayed by monitoring the fluorescence quenching of ACMA (9-amino-6-chloro-2-methoxy acridine) [33]. The statistical significance of the differences observed between sets of activity data was determined by Student's unpaired *t* tests using the application available at <http://graphpad.com/quickcalcs/ttest1.cfm>.


### Western blotting and total protein estimation

Western blot analysis was carried out after SDS/PAGE as described previously [34]. For immunodetection of H<sup>+</sup>-PPases, an affinity-purified polyclonal antibody raised in rabbit against the membrane-bound PPase (TVP) from the bacterium *Thermotoga maritima* [35] was used at a 1:200 dilution. For Vma1p, GFP and Pma1p immunodetection, commercial monoclonal antibody 8B1 (Molecular Probes), a polyclonal 'Living Colors' antibody from BD Biosciences and a polyclonal antibody provided by Professor Ramón Serrano (Universidad Politécnica de Valencia, Valencia, Spain) were used respectively, all of them at a 1:2000 dilution. The total protein concentration was estimated using the Bradford method [36] with ovalbumin as a standard.

## RESULTS

### Chimaeras of AVP1 bearing the N-terminal domain of TcVP induce bafilomycin resistance in YPC3 cells

YPC3 is a yeast mutant whose *IPP1* gene coding for the essential cytosolic PPase is under the control of the galactokinase (*GALI*)




**Figure 1** Drop tests of yeast mutant YPC3 cells grown in the presence of the macrolide inhibitor bafilomycin A1 at pH 5.5 and 8.0 (A) and in the presence of 100 mM CaCl<sub>2</sub> or 4 mM ZnCl<sub>2</sub> at pH 5.5 (B)

Cells were transformed with plasmids pRS699b (Control-), pIPP1, pAVP1, pTcAVP1 and pTcGFPVAVP1 and grown as described in the Experimental section. Serial dilutions of the cultures were made in sterile water and spotted on to YPGal and YPD plates containing 2  $\mu$ M bafilomycin A1 and buffered at the indicated pH values with 50 mM Tris/Mes. Plates containing Ca<sup>2+</sup> and Zn<sup>2+</sup> salts were adjusted to pH 5.5 with the same buffer. In (A) 1:10, 1:100 and 1:1000 serial dilutions are shown. In (B) 1:10 and 1:100 dilutions are shown. Growth was recorded after 2 days at 30°C in all cases, except for Zn<sup>2+</sup> plates which were grown for 4 days. No sensitivity to pH 8 or to Ca<sup>2+</sup> or Zn<sup>2+</sup> was observed for YPC3 cells in the absence of bafilomycin A1 (results not shown). baf., bafilomycin A1; w/o, without.

promoter, therefore these cells are non-viable when glucose is present as a carbon source. It has been shown previously that plant and bacterial membrane-bound H<sup>+</sup>-PPases can functionally complement the yeast cytosolic sPPase (Ipp1p), especially when fused to certain N-terminal signal peptides [21,28]. Addition of macrolide bafilomycin A1 to YPC3 cells transformed with different plasmids bearing constitutively expressed genes coding for soluble or membrane-bound PPases resulted in sensitivity to neutral/alkaline pH and to the divalent cations Ca<sup>2+</sup> and Zn<sup>2+</sup>, when grown on galactose (Figure 1), that is, when Ipp1p is expressed. In contrast, when cells were grown on glucose in the presence of bafilomycin, those expressing different chimaeric derivatives of AVP1, the K<sup>+</sup>-dependent H<sup>+</sup>-PPase from the higher plant *A. thaliana*, recover the capacity to grow at pH 7.5–8.0 (Figure 1A). In the presence of 100 mM calcium at pH 5.6, some growth was observed in YPC3 cells overexpressing Ipp1p, although they became significantly sensitive to this cation. Cells expressing the different AVP1 chimaeras showed a somewhat smaller sensitivity to this CaCl<sub>2</sub> (Figure 1B). In the presence of Zn<sup>2+</sup>, only those cells transformed with plasmids pTcAVP1 and pTcGFPVAVP1 expressing internal membrane-targeted chimaeras could grow after 4 days, particularly the latter (Figure 1B).

### Chimaeras of AVP1 bearing the N-terminal domain of TcVP reverse the sensitivity of YPC4 Vma1<sup>-</sup> cells to neutral/alkaline pH, Ca<sup>2+</sup> and Zn<sup>2+</sup>

In order to fully ascertain whether the observed bafilomycin A1 tolerance was actually brought about by complementation of V-ATPase proton-transport activity, similar experiments to those described above were performed with YPC4 cells. This mutant also has *IPP1* under the control of the *GALI* promoter




**Figure 2** Drop tests of yeast mutant YPC4 cells grown at pH 5.5, 7.5 and 8.0 (A) and in the presence of 100 mM CaCl<sub>2</sub> or 4 mM ZnCl<sub>2</sub> at pH 5.5 (B)

Cells were transformed with plasmids pRS699b (Control-), pIPP1, pAVP1, pTcAVP1, pTcGFPVP1 and pVMA1 and grown as described in Experimental section. Serial dilutions of the cultures were made in sterile water and spotted on to YPGal and YPD plates buffered at the indicated pH values with 50 mM Tris/Mes. Plates containing Ca<sup>2+</sup> and Zn<sup>2+</sup> salts were adjusted to pH 5.5 with the same buffer. In (A) 1:10, 1:100 and 1:1000 serial dilutions are shown. In (B) 1:10 and 1:100 dilutions are shown. Growth was recorded after 4 days in all cases, except for those at pH 8 which were grown for 7 days. baf., bafilomycin A; w/o, without.

but, additionally, it has the gene coding for the catalytic subunit VMA1 of the vacuolar H<sup>+</sup>-ATPase disrupted. YPC4, like YPC3, is unable to grow on glucose, but, as expected, when grown on galactose it exhibits the typical range of Vma<sup>-</sup> phenotypes: very high sensitivity to neutral/alkaline pH values and to Ca<sup>2+</sup> and Zn<sup>2+</sup> cations (Figures 2A and 2B). However, when YPC4 cells were transformed with AVP1 and its chimaeric derivatives, they recovered the capacity to grow on glucose at pH 5.5 and 7.5, especially in the case of the TcGFPVP1 chimaera. In the presence of Ca<sup>2+</sup>, only cells transformed with plasmids pTcAVP1 and pTcGFPVP1 were able to grow, whereas only the latter showed growth in the presence of Zn<sup>2+</sup>. Again, as in the case of YPC3 with bafilomycin A1, the construct bearing the gene coding for Ipp1p could only support growth on glucose at pH 5.6 and in the absence of divalent cations. On the other hand, transformation of YPC4 with VMA1 resulted in the reversal of the phenotypes associated with the disruption of this gene only in the presence of galactose (Figures 2A and 2B).

#### Expression of TcGFPVP1 allows acidification of vacuoles in YPC4 cells

To verify that AVP1 and its derivatives were able to promote acidification of internal compartments, we analysed the accumulation of quinacrine, a vital dye that concentrates in acidic lumina. Figure 3 shows typical patterns of quinacrine accumulation in vacuoles of YPC4 cells transformed with the different constructs tested. No internal acidification could be observed in cells expressing the soluble cytosolic PPase Ipp1p, whereas cells expressing AVP1 and TcAVP1 showed a punctate pattern of green fluorescence suggesting acidification of prevacuolar/endosomal compartments. In contrast with this, cells transformed with the plasmid pTcGFPVP1 showed a different fluorescence pattern, with a clear vacuolar accumulation




**Figure 3** Vacuolar acidification of transformed YPC4 cells studied by accumulation of quinacrine

Cells were transformed with the indicated plasmids and grown as described in the Experimental section. Vacuole acidification was assessed by accumulation of the fluorescent probe quinacrine as described previously [28]. Cells were visualized in a Leica DM6000B microscope. The left-hand column shows DIC microscopy of a typical field of transformed YPC4 cells, the middle column shows the pattern of quinacrine accumulation (observed with a FITC green filter) in the same cells, and the right-hand column shows the overlap of both.

of quinacrine just like YPC4 cells transformed with plasmid pVMA1 grown in galactose. The former cells showed a punctate green fluorescence pattern, corresponding to the TcGFPVP1 polypeptide, when visualized in the absence of quinacrine (see below). It is important to point out that some of the cells transformed with pTcGFPVP1 exhibited clusters of smaller vacuoles rather than the single large vacuole usually observed in cells expressing Vma1p (see Figure 3). These results are in agreement with the previously presented drop tests (compare with Figures 1 and 2), which show comparatively higher growth of TcGFPVP1-expressing clones at neutral/alkaline pH values and in the presence of Ca<sup>2+</sup>/Zn<sup>2+</sup>, these tolerance phenotypes being strictly dependent on physiologically competent acidic vacuoles.

#### Expression of all AVP1 derivatives allows normal internalization of the vital dye FM4-64

The vital fluorescent dye FM4-64 binds to the plasma membrane and follows the endocytic pathway to reach the vacuole [29]. FM4-64 internalization has been shown to be severely impaired in *vma* mutants [37]. Figure 4 shows labelling of vacuole membranes in YPC4 cells after incubation with FM4-64 for 30 min. It can be observed that cells expressing the AVP1



**Figure 4** Endocytic internalization of the fluorescent marker FM4-64 in transformed YPC4 cells

The rate of endocytosis was studied by incubating transformed YPC4 cells grown to early exponential phase with FM4-64 for 15 min, as described previously [28]. Cells were then transferred to YPD and visualized as described in Figure 3 after 30 min of incubation with the fluorescent dye. In (A), the left-hand column shows DIC microscopy of a typical field of transformed YPC4 cells, the middle column shows the pattern of FM4-64 labelling in the same cells, and the right-hand column shows the overlap of both. (B) YPC4 cells transformed with plasmid pTcGFPVAVP1. From left-to-right: visualization of the cells by DIC microscopy, with red (Texas Red) and green (FITC) filters, overlap of red and green filter microphotographs, and overall overlap of the three former microphotographs.


and TcGFPVAVP1 proteins have their vacuoles clearly labelled, whereas cells expressing TcAVP1 showed labelling of vacuoles and other smaller vesicles. Control cells expressing Vma1p also showed a clear pattern of vacuolar membrane staining. No internal labelling was obtained in cells transformed with pIPP1. Cells transformed with pTcGFPVAVP1 were also visualized with a green (FITC) fluorescence filter showing a punctate pattern that suggested either a preferential localization of TcGFPVAVP1 in internal membranes other than the vacuolar membrane, or a degradation of the GFP moiety by vacuolar proteases (Figure 4B), a possibility that was subsequently studied by using a vacuolar protease-deficient mutant (see below). In some of these cells, clusters of small acidic vesicles were also observed. This effect had previously been described with YPC3 cells [21].

#### Membrane-bound H<sup>+</sup>-PPases are present in YPC4 vacuoles

In order to check whether the different natural and chimaeric plant H<sup>+</sup>-PPases expressed in YPC4 cells were localized in

the vacuoles, activity assays and immunodetection experiments were carried out with purified vacuole preparations obtained from YPC4 cells transformed with the different plasmids described above. Cells transformed with pTcGFPVAVP1 showed a characteristic trace of ACMA fluorescence quenching that was dependent on the addition of PP<sub>i</sub>, this quenching was reversed by the addition of the H<sup>+</sup>/K<sup>+</sup> ionophore nigericin. Smaller quenching rates were obtained with cells transformed with pTcAVP1 and pAVP1 (Figure 5A). Vacuoles from cells transformed with pVMA1 did not show PP<sub>i</sub>-dependent ACMA quenching, just like cells transformed with plasmids pRS699b and pIPP1 (results not shown). When ATP-dependent proton transport was recorded, only vacuoles isolated from cells transformed with plasmid pVMA1 exhibited ACMA quenching (Figure 5A). This quenching was sensitive to bafilomycin A1 (results not shown). Membrane-associated PPase activity levels determined for vacuolar membranes obtained from the different YPC4 transformants were consistent with the H<sup>+</sup>-translocation results (Figure 5B), thus no PPase activity was detected in samples obtained from cells transformed with






**Figure 5** PP<sub>i</sub>- and ATP-dependent proton-pumping activity measured as quenching of ACMA fluorescence (A), membrane-associated PP<sub>i</sub> hydrolysis activity (B) and Western blot analyses (C) of vacuolar membrane preparations obtained from transformed YPC4 cells

(A) Proton-translocation activity was measured by recording the PP<sub>i</sub>-dependent (left-hand panels) and ATP-dependent (right-hand panels) ACMA fluorescence quenching. The upper panels show typical traces obtained in these experiments. The lower panels show initial rates of fluorescence quenching for the different preparations. Values are means  $\pm$  S.E.M. for three independent experiments ( $n = 3$ ) and asterisks indicate differences that are statistically significant ( $P < 0.005$ ). n.d., not detected. Approximately 30  $\mu$ g of protein was used for these assays. (B) Specific PP<sub>i</sub> hydrolysis activity associated with vacuolar membranes obtained from YPC4 cells transformed with the different plasmids tested. Activity values shown are means  $\pm$  S.E.M. ( $n = 3$ ) using preparations obtained from YPC4 cells transformed with pRS699b as a reference. Asterisks indicate differences that are statistically significant ( $P < 0.005$ ). Open and closed bars represent activities in the absence and presence of potassium chloride respectively. n.d., no detected activity. (C) Immunodetection of AVP1 (upper panel) and yeast Vma1p (middle panel) in vacuolar membranes of transformed YPC4 cells. Approximately 30  $\mu$ g of protein was loaded per lane. Asterisks indicate bands associated with chimaeric protein TcGFPAVP1 (approximately 120 kDa) and AVP1 (72 kDa).

plasmids pRS699b, pIPP1 and pVMA1. Samples from cells expressing the different AVP1 chimaeras did show PPase activity that was increased by the addition of K<sup>+</sup>, a feature of AVP1 [19]. Immunodetection in Western blots of vacuolar membranes of the diverse native and chimaeric plant H<sup>+</sup>-PPases used in the present study (Figure 5C) gave somewhat unpredicted results despite the use of protease inhibitor cocktails during the isolation procedure, thus a polypeptide of the expected size for the native AVP1 protein (approximately 72 kDa) was obtained in the case of cells transformed with pAVP1, pTcAVP1 and pTcGFPAVP1; additionally, another band of approximately 64 kDa was present in these preparations. The relative amounts of H<sup>+</sup>-PPases immunodetected in these preparations were, however, in accordance with the estimated activity values. A minor band of approximately 120 kDa, the size expected for TcGFPAVP1, could also be detected only in samples obtained from cells transformed with plasmid pTcGFPAVP1 (compare with Figure 5C).

### Vacuolar proteases are involved in the processing of native AVP1 and its chimaeric derivatives targeted to the vacuolar membrane

The possibility that AVP1 and its chimaeras might be subjected to proteolytic processing by vacuolar proteases *in vivo* was investigated by using the vacuolar-protease-deficient yeast mutant BJ5457 [24]. Immunodetection analyses carried out with total membrane preparations obtained with mutants YPC4 and BJ5457 transformed with plasmids pAVP1, pTcAVP1 and pTcGFPAVP1 using an antibody against a bacterial membrane-bound PPase (anti-TVP) showed remarkable results (Figure 6A). Thus, in the case of YPC4, a polypeptide of approximately 72 kDa was detected in all cases, although in cells transformed with pTcGFPAVP1, two more bands of approximately 120 kDa and 64 kDa were clearly recognized by the anti-TVP antibody. Peptide mass fingerprint analysis of the 120 kDa polypeptide by MALDI-TOF (matrix-assisted laser-desorption ionization-time-of-flight) MS showed that it actually corresponded to a fusion between GFP and AVP1 (results not shown). Consistently, an antibody



**Figure 6** Western blot analysis (A) and membrane-associated PP<sub>i</sub> hydrolysis activity (B) of total membrane preparations obtained from transformed YPC4 and BJ5457 cells, and fluorescence microscopy visualization of YPC4 and BJ5457 cells transformed with pTcGFP-PPase (C)

(A) Immunodetection of AVP1 and derivatives with antibodies against a membrane-bound PPase (anti-TVP, upper panel) and GFP (anti-GFP) in total membranes of YPC4 (left-hand panels) and BJ5457 (right-hand panels) cells transformed with plasmids pRS699b, pIPP1, pAVP1, pTcAVP1 and pTcGFP-PPase. Approximately 70 μg of protein was loaded per lane. Asterisks indicate bands associated with chimaeric protein TcGFP-PPase (approximately 120 kDa) and AVP1 (72 kDa). (B) Specific PP<sub>i</sub> hydrolysis activity associated with total membranes obtained from YPC4 and BJ5457 cells transformed with the different plasmids tested. Activity values shown are means ± S.E.M. (*n* = 3) using preparations obtained from cells transformed with pRS699b as a reference. Asterisks indicate differences that are statistically significant (*P* < 0.005). Open and closed bars represent activities in the absence and presence of potassium chloride respectively. n.d., not detected. (C) YPC4 and BJ5457 cells were transformed with plasmid pTcGFP-PPase and visualized in a Leica DM6000B microscope. Left-hand column shows DIC microscopy of a typical field of cells, the middle column shows the pattern of green fluorescence distribution (FITC green filter) in the same cells, and the right-hand column shows the overlap of both. White arrowheads indicate vacuolar membranes.

against GFP (anti-GFP) only recognized the 120 kDa band. In the case of mutant BJ5457, the anti-TVP antibody recognized polypeptides of approximately 72 kDa only in preparations obtained with cells transformed with plasmids pAVP1 and pTcAVP1, whereas a single polypeptide of approximately 120 kDa was detected in membranes of cells transformed with plasmid pTcGFP-PPase. The anti-GFP antibody could only detect the latter (Figure 6A).

Analysis of membrane-associated PP<sub>i</sub> hydrolytic activity showed that, in YPC4, the presence of GFP at the N-terminus of AVP1 increased the specific activity of the resulting protein 3–4-fold with respect to AVP1 and TcAVP1. In mutant BJ5457, the specific PPase activities associated with AVP1, TcAVP1 and

TcGFP-PPase were significantly increased with respect to the same proteins expressed in YPC4. The increases were much higher in the cases of AVP1 (approx. 6-fold) and TcAVP1 (approx. 4-fold) than in the case of TcGFP-PPase (approx. 1.7-fold) (Figure 6B).

Fluorescent microscopy visualization of YPC4 and BJ5457 cells transformed with plasmid pTcGFP-PPase remarkably showed different patterns of green fluorescence distribution, thus, in YPC4 cells, a punctate pattern was observed, suggesting a preferential accumulation of green fluorescence in internal membrane systems other than the vacuolar membrane. In contrast, green fluorescence mostly associated with the latter was clearly observed in BJ5457 cells (Figure 6C).

## DISCUSSION

In previous studies, we emulated in yeast the cytosolic  $\text{PP}_i$  metabolism of the plant cell, i.e. absence of a soluble cytosolic PPase, so that the  $\text{PP}_i$  generated by the anabolism was efficiently hydrolysed by a membrane-embedded  $\text{PP}_i$ -dependent proton pump [21,28]. In the present study, we have taken one step forward using a metabolic background in which the ATP-dependent proton pump of yeast internal membranes, the V-ATPase, has been either chemically inhibited or functionally impaired by mutation of a gene encoding an essential catalytic subunit. The results obtained strongly suggest that a distinct type of  $\text{PP}_i$ -dependent proton pump can efficiently substitute for the V-ATPase of *S. cerevisiae* in diverse physiological scenarios. Thus overexpression of the  $\text{K}^+$ -dependent  $\text{H}^+$ -PPase from the higher plant *A. thaliana*, a protein that occurs along with a V-ATPase in plant tonoplast [18], alleviates the range of phenotypes associated with the lack of the latter. The presence of an appropriate N-terminal domain proved to be important in order to observe the optimal phenotypic complementation in our system. The N-terminal domain with signal sequence features of a protistan  $\text{H}^+$ -PPase, when fused to AVP1, alters the subcellular distribution of the latter so that it becomes preferentially located in internal membranes of yeast [21]. Not surprisingly, as shown in the present study, chimaeric proteins of AVP1, constructed with that signal peptide (TcAVP1 and TcGFP AVP1), functionally substituted for the V-ATPase more efficiently than the native AVP1 protein with no extra N-terminal domain. The difference in favour of TcGFP AVP1 compared with TcAVP1 is probably due to the higher overall expression levels attained by the former (see below). Actually, under non-restrictive conditions towards  $\text{Vma}^-$  defects, TcGFP AVP1 supports growth of YPC4 more efficiently than TcAVP1 (Figure 2A) which means that the former is more efficient at complementing the phenotype associated with the absence of Ipp1p. Addition of GFP at the N-terminus of some membrane proteins has been previously reported either because it stabilizes proteins expressed in heterologous systems or because the presence of GFP at the C-terminus yields polypeptides with no activity [21,38,39]. In fact, both assumptions are true for the plant  $\text{H}^+$ -PPase AVP1 expressed in yeast, as the present study and our previous study [21] demonstrates.

The results with both bafilomycin-treated YPC3 and YPC4 cells also showed that no version of heterologously expressed AVP1 was able to complement  $\text{Vma}^-$  phenotypes in galactose, even though the corresponding genes are under the control of the constitutive *PMA1* promoter. Similarly, when the parental strains (bafilomycin-treated W303-1A and RS-1114) were transformed with the same plasmids no complementation of the V-ATPase by AVP1 was observed either on galactose or on glucose (results not shown). These results suggested that AVP1 cannot pump protons when the soluble PPase is present in the cytosol, probably due to the competition for the substrate, therefore it was necessary to bring down the activity of the latter to negligible levels in order to have the former efficiently working *in vivo*. YPC3 and YPC4 were designed and generated so that this could be accomplished by transferring the cells from galactose to glucose and allowing them to grow for several hours. Actually, this reflects the metabolic scenario in cells of plant green tissues, where sPPases are restricted to energy-linked organelles (mitochondria and chloroplasts) and, probably, nuclei [17], although it must be mentioned that other  $\text{PP}_i$ -utilizing enzymes involved in central carbon metabolism, such as a  $\text{PP}_i$ -dependent phosphofructokinase, are also present in the plant cell cytosol [40]. All in all, these results suggest that the presence of active soluble PPases is incompatible with  $\text{H}^+$ -PPase functions, and helps to explain why the former are not found

in the cytosol of eukaryotic cells if  $\text{H}^+$ -PPases are present in endomembranes.

Acidification of internal compartments of YPC4 cells was studied with the fluorescent probe quinacrine, which accumulates in acidic vacuoles. TcGFP AVP1 was able to acidify the vacuole in YPC4 cells, although it also seemed to induce some fragmentation of this compartment. This has already been reported for YPC3 cells and may be related to the accumulation of this heterologous chimaeric protein in the internal compartments of yeast cells [21]. It should be noted in this respect that a more or less severe vacuolar fragmentation takes place in yeast under several, and not necessarily pathological, scenarios. In contrast, cells transformed with AVP1 and TcAVP1 showed a punctate pattern of quinacrine accumulation which probably means acidification of prevacuolar/endosomal vesicles. This pattern was more significant in the case of TcAVP1, probably due to the effect of the addition of the N-terminal signal domain added to the latter that favours accumulation of this chimaeric protein in internal membranes of yeast [21]. In the presence of quinacrine, cells expressing TcGFP AVP1 only exhibited the green fluorescence associated with the dye; however, these cells were also expected to show green fluorescence due to the GFP. Consequently, they were also visualized in the absence of quinacrine showing, in this case, a punctate pattern of fluorescence (see below). This suggests that quinacrine fluorescence masks that of GFP in our system when both are monitored simultaneously.

Results with FM4-64 are consistent with those obtained with quinacrine in the case of TcGFP AVP1; however, they further suggest that full acidification of the vacuole may not be absolutely essential to have the endocytic system operational at the right pace, acidification of the prevacuolar/endosomal compartments being sufficient. The fact that the green fluorescence observed in YPC4 cells transformed with plasmid pTcGFP AVP1 does not necessarily co-localize with the red fluorescence associated with acidic vesicles further suggests that either TcGFP AVP1 polypeptide is not significantly translocated to the vacuolar membrane or the GFP moiety of this chimaera has been lost, presumably by proteolysis.

Biochemical analysis of vacuolar membrane preparations from YPC4 transformed with the different plasmids tested showed that AVP1 and all of its chimaeric derivatives were heterologously expressed and located, at least partially, in the vacuoles, as demonstrated by the data of membrane-associated  $\text{H}^+$ -translocating and PPase activities, as well as by immunodetection analysis of vacuolar membrane preparations. Activity data were consistent with the phenotypes observed, thus TcGFP AVP1 showed higher PPase and  $\text{H}^+$ -translocating activities than TcAVP1 which, in turn, produced higher activities than the native AVP1. Immunodetection with the anti-TVP antibody revealed that TcAVP1 and TcGFP AVP1 gave higher expression levels in this compartment than AVP1. This is consistent with previous reports regarding the role of the N-terminal signal domain of *T. cruzi*  $\text{H}^+$ -PPase on the subcellular distribution of AVP1 [21]. On the other hand, a significant processing of all of the proteins examined was also found; actually, only a tiny amount of a polypeptide of approximately 120 kDa, corresponding to a fusion of GFP and AVP1, was detected, and another band of approximately 64 kDa was consistently observed in all cases. This evidence suggested that vacuolar proteases may be involved in the cleavage of AVP1 and its derivatives, with a concomitant degradation of the GFP in the case of TcGFP AVP1. In order to study this situation, the vacuolar-deficient yeast mutant BJ5457 [24] was transformed with plasmids pAVP1, pTcAVP1 and pTcGFP AVP1 and total membrane preparations obtained with the resulting strains were analysed and compared with preparations obtained with YPC4



cells transformed with the same plasmids. Immunochemical analysis strongly suggested that vacuolar proteases are involved in the processing of AVP1 and chimaeric derivatives observed in mutant YPC4. These results, along with the membrane-associated PPase activity data, further support the idea that, although AVP1 and TcAVP1 can be found in YPC4 vacuoles, a high proportion of these polypeptides is subjected to proteolysis. This effect is much smaller in the case of TcGFP-VP1, thus suggesting that the combination of the *T. cruzi* H<sup>+</sup>-PPase N-terminal signal peptide and yEGFP is more efficient than the signal peptide alone at altering the subcellular distribution of the resulting chimaera. Fluorescence microscopy supported the data obtained by immunodetection and activity assays, thus, unlike in mutant BJ5457, the green fluorescence corresponding to TcGFP-VP1 in mutant YPC4 showed a pattern not clearly associated with the vacuolar membrane. There are two contributions to this effect according to our results: on the one hand, a significant fraction of TcGFP-VP1 is targeted to other internal membranes and, on the other hand, the polypeptide located in the vacuolar membrane is subjected to proteolysis so that most of the GFP moiety is degraded. Therefore the role of vacuolar proteases provides an explanation of why the fluorescence associated with GFP is not located in the vacuolar membrane and further suggests that the proton pump responsible for the acidification of the vacuole in YPC4 cells transformed with pTcGFP-VP1 is actually a native-like AVP1 protein and not the chimaeric derivative.

Taken together our results show that the PP<sub>i</sub>-dependent proton pump AVP1 can alleviate the most characteristic phenotypes associated to the lack of V-ATPase in yeast, especially when the former is preferentially targeted to internal membranes by attaching an appropriate signal peptide to its N-terminus [21]. In the opinion of the authors, this is the first direct proof that a H<sup>+</sup>-translocating PPase can sustain growth and significant acidification of intracellular lumina in the absence of any V-ATPase activity.

These results raise another question related to the occurrence of two proton pumps in the same subcellular location in certain organisms, one of them being an extremely complex molecular machine (the V-ATPase), and the other a much simpler membrane pump. There are numerous recent data supporting that V-ATPases may play more roles than the acidification of internal compartments, such as the participation of the V<sub>0</sub> domain in membrane fusion [41] and others [42]. On the other hand, the V-ATPase complex may offer many more possibilities of activity modulation and, thus, may be more finely tuned than the comparatively simpler H<sup>+</sup>-PPase. Additionally, reports of co-regulation and physical interactions between V-ATPases and H<sup>+</sup>-PPases in plant tonoplast may complicate further the task of discriminating the respective roles of both pumps [18]. A yeast expression system like the one in the present study can help ascertain the importance of different V-ATPase polypeptides or whole domains in moonlighting functions independent of lumen acidification.

In any case, the H<sup>+</sup>-PPases probably imply an adaptive advantage to the organisms where they occur because they offer an alternative pump to generate proton gradients in internal organelles by using an abundant by-product of anabolism, PP<sub>i</sub>. [43]. This is likely to be a critical issue under conditions that require the optimization of cellular energy usage. In this respect, it is worthwhile pointing out that the H<sup>+</sup>-PPase seems to be the main tonoplast pump in other plant systems, such as grape berry cells [44].

The yeast system described in the present work may have biotechnological applications, thus it is a good experimental system to accomplish studies of random mutagenesis aimed at

obtaining enhanced versions of AVP1 and other PP<sub>i</sub>-dependent ion pumps. On the other hand, YPC4 transformed with diverse H<sup>+</sup>-PPases might be a useful system to check for possible roles other than the acidification of internal organelles proposed for fungal and animal V-ATPases, and it may be a suitable system to study the effect of H<sup>+</sup>-PPases inhibitors *in vivo*, as these proton pumps occur in micro-organisms responsible for diseases such as malaria, leishmaniasis, Chagas disease and sleeping sickness [15,45]. Finally, this experimental system envisages novel ways of engineering cell tolerance to macrolides, a class of drugs of increasing relevance in therapeutics of a range of diseases in which cellular proton homeostasis may play crucial roles, such as cancer.

## AUTHOR CONTRIBUTION

José Pérez-Castiñeira carried out most of the experimental work and wrote the paper; Agustín Hernández generated the YPC4 mutant and corrected the paper; Rocío Drake performed preliminary studies; and Aurelio Serrano supervised the work and wrote the final version of the paper.

## ACKNOWLEDGEMENTS

We thank Professor Ramón Serrano for providing the anti-Pma1p antibody and yeast mutant strain RS-1144, and Professor Andrés Aguilera for providing yeast strain BJ5457. We also thank Dr Alicia Orea for excellent technical assistance with fluorescence microscopy and Ms Isabel Jiménez for yeast culturing.

## FUNDING

This work was supported by the Regional Andalusian Government and the Spanish Ministerio de Ciencia e Innovación financial support to PAIDI group BIO-261 [grant numbers P07-CVI-03082, BFU2007-61887 and BFU2010-15622], partially funded by the EU FEDER (Fondo Europeo de Desarrollo Regional) programme.

## REFERENCES

- Kane, P. M. (2006) The where, when, and how of organelle acidification by the yeast vacuolar H<sup>+</sup>-ATPase. *Microbiol. Mol. Biol. Rev.* **70**, 177–191
- Nishi, T. and Forgac, M. (2002) The vacuolar H<sup>+</sup>-ATPases: nature's most versatile proton pumps. *Nat. Rev. Mol. Cell Biol.* **3**, 94–103
- Forgac, M. (2007) Vacuolar ATPases: rotary proton pumps in physiology and pathophysiology. *Nat. Rev. Mol. Cell Biol.* **8**, 917–929
- Hernández, A., Serrano, G., Herrera-Palau, R., Pérez-Castiñeira, J. R. and Serrano, A. (2010) Intraorganellar acidification by V-ATPases: a target in cell proliferation and cancer therapy. *Recent Pat. Anticancer Drug Discovery* **5**, 88–98
- Bowman, E. J., Siebers, A. and Altendorf, K. (1988) Bafilomycins: a class of inhibitors of membrane ATPases from microorganisms, animal cells, and plant cells. *Proc. Natl. Acad. Sci. U.S.A.* **85**, 7972–7976
- Bowman, E. J., Graham, L. A., Stevens, T. H. and Bowman, B. J. (2004) The bafilomycin/concanamycin binding site in subunit c of the V-ATPases from *Neurospora crassa* and *Saccharomyces cerevisiae*. *J. Biol. Chem.* **279**, 33131–33138
- Nelson, H. and Nelson, N. (1990) Disruption of genes encoding subunits of yeast vacuolar H<sup>+</sup>-ATPase causes conditional lethality. *Proc. Natl. Acad. Sci. U.S.A.* **87**, 3503–3507
- Ohya, Y., Umemoto, N., Tanida, I., Ohta, A., Iida, H. and Anraku, Y. (1991) Calcium-sensitive *cls* mutants of *Saccharomyces cerevisiae* showing a Pet<sup>-</sup> phenotype are ascribable to defects of vacuolar membrane H<sup>+</sup>-ATPase activity. *J. Biol. Chem.* **266**, 13971–13977
- Ramsay, L. M. and Gadd, G. M. (1997) Mutants of *Saccharomyces cerevisiae* defective in vacuolar function confirm a role for the vacuole in toxic metal ion detoxification. *FEMS Microbiol. Lett.* **152**, 293–298
- Hinton, A., Bond, S. and Forgac, M. (2009) V-ATPase functions in normal and disease processes. *Pflugers Arch.* **457**, 589–598
- Morel, N. (2003) Neurotransmitter release: the dark side of the vacuolar-H<sup>+</sup> ATPase. *Biol. Cell* **95**, 453–457
- Drozdowicz, Y. M. and Rea, P. A. (2001) Vacuolar H<sup>+</sup> pyrophosphatases: from the evolutionary backwaters into the mainstream. *Trends Plant Sci.* **6**, 206–211

- 13 Maeshima, M. (2001) Tonoplast Transporters: organization and function. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **52**, 469–497
- 14 Serrano, A., Pérez-Castiñeira, J. R., Baltscheffsky, M. and Baltscheffsky, H. (2007) H<sup>+</sup>-PPases: yesterday, today and tomorrow. *IUBMB Life* **59**, 76–83
- 15 Docampo, R., de Souza, W., Miranda, K., Rohloff, P. and Moreno, S. N. (2005) Acidocalcisomes: conserved from bacteria to man. *Nat. Rev. Microbiol.* **3**, 251–261
- 16 Pérez-Castiñeira, J. R., López-Marqués, R. L., Losada, M. and Serrano, A. (2001) A thermostable K<sup>+</sup>-stimulated vacuolar-type pyrophosphatase from the hyperthermophilic bacterium *Thermotoga maritima*. *FEBS Lett.* **496**, 6–11
- 17 Pérez-Castiñeira, J. R., Gómez-García, R., López-Marqués, R. L., Losada, M. and Serrano, A. (2001) Enzymatic systems of inorganic pyrophosphate bioenergetics in photosynthetic and heterotrophic protists: remnants or metabolic cornerstones? *Int. Microbiol.* **4**, 135–142
- 18 Schumacher, K. (2006) Endomembrane proton pumps: connecting membrane and vesicle transport. *Curr. Opin. Plant Biol.* **9**, 595–600
- 19 Kim, E. J., Zhen, R. G. and Rea, P. A. (1994) Heterologous expression of plant vacuolar pyrophosphatase in yeast demonstrates sufficiency of the substrate-binding subunit for proton transport. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 6128–6132
- 20 Hill, J. E., Scott, D. A., Luo, S. and Docampo, R. (2000) Cloning and functional expression of a gene encoding a vacuolar-type proton-translocating pyrophosphatase from *Trypanosoma cruzi*. *Biochem. J.* **351**, 281–288
- 21 Drake, R., Serrano, A. and Pérez-Castiñeira, J. R. (2010) N-terminal chimaeras with signal sequences enhance the functional expression and alter the subcellular localization of heterologous membrane-bound inorganic pyrophosphatases in yeast. *Biochem. J.* **426**, 147–157
- 22 Ros, R., Montesinos, C., Rimon, A., Padan, E. and Serrano, R. (1998) Altered Na<sup>+</sup> and Li<sup>+</sup> homeostasis in *Saccharomyces cerevisiae* cells expressing the bacterial cation antiporter NhaA. *J. Bacteriol.* **180**, 3131–3136
- 23 Kane, P. M., Yamashiro, C. T., Wolczyk, D. F., Neff, N., Goebel, M. and Stevens, T. H. (1990) Protein splicing converts the yeast *TFP1* gene product to the 69-kD subunit of the vacuolar H<sup>+</sup>-adenosine triphosphatase. *Science* **250**, 651–657
- 24 Jones, E. W. (1991) Tackling the protease problem in *Saccharomyces cerevisiae*. *Methods Enzymol.* **194**, 428–453
- 25 Schiestl, R. H. and Gietz, R. D. (1989) High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. *Curr. Genet.* **16**, 339–346
- 26 Yenush, L., Mulet, J. M., Arino, J. and Serrano, R. (2002) The Ppz protein phosphatases are key regulators of K<sup>+</sup> and pH homeostasis: implications for salt tolerance, cell wall integrity and cell cycle progression. *EMBO J.* **21**, 920–929
- 27 Treco, D. A. and Lunblad, V. (1993) Preparation of yeast media. *Curr. Protoc. Mol. Biol. Unit* **13.1**,
- 28 Pérez-Castiñeira, J. R., López-Marqués, R. L., Villalba, J. M., Losada, M. and Serrano, A. (2002) Functional complementation of yeast cytosolic pyrophosphatase by bacterial and plant H<sup>+</sup>-translocating pyrophosphatases. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 15914–15919
- 29 Conibear, E. and Stevens, T. H. (2002) Studying yeast vacuoles. *Methods Enzymol.* **351**, 408–432
- 30 Roberts, C. J., Raymond, C. K., Yamashiro, C. T. and Stevens, T. H. (1991) Methods for studying the yeast vacuole. *Methods Enzymol.* **194**, 644–661
- 31 Lenoir, G., Menguy, T., Corre, F., Montigny, C., Pedersen, P. A., Thines, D., le Maire, M. and Falson, P. (2002) Overproduction in yeast and rapid and efficient purification of the rabbit SERCA1a Ca<sup>2+</sup>-ATPase. *Biochim. Biophys. Acta* **1560**, 67–83
- 32 Rathbun, W. B. and Betlach, M. V. (1969) Estimation of enzymically produced orthophosphate in the presence of cysteine and adenosine triphosphate. *Anal. Biochem.* **28**, 436–445
- 33 Pérez-Castiñeira, J. R. and Apps, D. K. (1990) Vacuolar H<sup>+</sup>-ATPase of adrenal secretory granules: rapid partial purification and reconstitution into proteoliposomes. *Biochem. J.* **271**, 127–131
- 34 Valverde, F., Losada, M. and Serrano, A. (1997) Functional complementation of an *Escherichia coli* gap mutant supports an amphibolic role for NAD(P)-dependent glyceraldehyde-3-phosphate dehydrogenase of *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **179**, 4513–4522
- 35 López-Marqués, R. L., Pérez-Castiñeira, J. R., Buch-Pedersen, M. J., Marco, S., Rigaud, J. L., Palmgren, M. G. and Serrano, A. (2005) Large-scale purification of the proton pumping pyrophosphatase from *Thermotoga maritima*: a 'Hot-Solve' method for isolation of recombinant thermophilic membrane proteins. *Biochim. Biophys. Acta* **1716**, 69–76
- 36 Bradford, M. M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248–254
- 37 Perzov, N., Padler-Karavani, V., Nelson, H. and Nelson, N. (2002) Characterization of yeast V-ATPase mutants lacking Vph1p or Stv1p and the effect on endocytosis. *J. Exp. Biol.* **205**, 1209–1219
- 38 Jasti, J., Furukawa, H., Gonzales, E. B. and Gouaux, E. (2007) Structure of acid-sensing ion channel 1 at 1.9 Å resolution and low pH. *Nature* **449**, 316–323
- 39 Sato, K., Sato, M. and Nakano, A. (2001) Rer1p, a retrieval receptor for endoplasmic reticulum membrane proteins, is dynamically localized to the Golgi apparatus by coatomer. *J. Cell Biol.* **152**, 935–944
- 40 Mertens, E. (1991) Pyrophosphate-dependent phosphofructokinase, an anaerobic glycolytic enzyme? *FEBS Lett.* **285**, 1–5
- 41 Baars, T. L., Petri, S., Peters, C. and Mayer, A. (2007) Role of the V-ATPase in regulation of the vacuolar fission-fusion equilibrium. *Mol. Biol. Cell* **18**, 3873–3882
- 42 Gaxiola, R. A., Palmgren, M. G. and Schumacher, K. (2007) Plant proton pumps. *FEBS Lett.* **581**, 2204–2214
- 43 López-Marqués, R. L., Pérez-Castiñeira, J. R., Losada, M. and Serrano, A. (2004) Differential regulation of soluble and membrane-bound inorganic pyrophosphatases in the photosynthetic bacterium *Rhodospirillum rubrum* provides insights into pyrophosphate-based stress bioenergetics. *J. Bacteriol.* **186**, 5418–5426
- 44 Fontes, N., Silva, R., Vignault, C., Lecourieux, F., Geros, H. and Delrot, S. (2010) Purification and functional characterization of protoplasts and intact vacuoles from grape cells. *BMC Res. Notes* **3**, 19
- 45 Pérez-Castiñeira, J. R., Alvar, J., Ruíz-Pérez, L. M. and Serrano, A. (2002) Evidence for a wide occurrence of proton-translocating pyrophosphatase genes in parasitic and free-living protozoa. *Biochem. Biophys. Res. Commun.* **294**, 567–573

Received 10 March 2011/3 May 2011; accepted 5 May 2011


Published as BJ Immediate Publication 26 May 2011, doi:10.1042/BJ20110447

## SUPPLEMENTARY ONLINE DATA

# A plant proton-pumping inorganic pyrophosphatase functionally complements the vacuolar ATPase transport activity and confers bafilomycin resistance in yeast

José R. PÉREZ-CASTIÑEIRA, Agustín HERNÁNDEZ, Rocío DRAKE and Aurelio SERRANO<sup>1</sup>

Instituto de Bioquímica Vegetal y Fotosíntesis, Universidad de Sevilla-CSIC, Avda. Americo Vespucio, 49, 41092 Sevilla, Spain



**Figure S1** Schematic diagram showing the constructions used for plasmidic expression in yeast of AVP1, AVP1-derived chimaeric H<sup>+</sup>-PPases, IPP1 and the vacuolar ATPase subunit VMA1

**Table S1** Yeast strains utilized in this article and their parental strains

| Strains          | Genotype  | References        |
|------------------|---|-------------------|
| Parental strains |   |                   |
| W303-1A          | <i>MATa, ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1</i>            | [1]               |
| RS-1144          | <i>MATa, ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 vma1::LEU2</i> | [2]               |
| Strains          |   |                   |
| YPC3             | <i>W303-1A ipp1::P<sub>GAL1</sub>-IPP1 HIS3</i>                             | [3]               |
| YPC4             | <i>RS-1144 ipp1::P<sub>GAL1</sub>-IPP1 HIS3</i>                             | The present study |
| Other strains    |   |                   |
| BJ5457           | <i>MATα pep4::HIS3 prb1-Δ trp1 ura3-52 leu2-Δ his3-Δ lys2-801 can1</i>      | [4]               |

## REFERENCES

- 1 Thomas, B. J. and Rothstein, R. (1989) Elevated recombination rates in transcriptionally active DNA. *Cell* **56**, 619–630
- 2 Ros, R., Montesinos, C., Rimon, A., Padan, E. and Serrano, R. (1998) Altered Na<sup>+</sup> and Li<sup>+</sup> homeostasis in *Saccharomyces cerevisiae* cells expressing the bacterial cation antiporter NhaA. *J. Bacteriol.* **180**, 3131–3136
- 3 Drake, R., Serrano, A. and Pérez-Castiñeira, J. R. (2010) N-terminal chimaeras with signal sequences enhance the functional expression and alter the subcellular localization of heterologous membrane-bound inorganic pyrophosphatases in yeast. *Biochem. J.* **426**, 147–157
- 4 Jones, E. W. (1991) Tackling the protease problem in *Saccharomyces cerevisiae*. *Methods Enzymol.* **194**, 428–453

Received 10 March 2011/3 May 2011; accepted 5 May 2011

Published as BJ Immediate Publication 26 May 2011, doi:10.1042/BJ20110447

<sup>1</sup>To whom correspondence should be addressed (email aurelio@ibvf.csic.es).