

# A stochastic pitchfork bifurcation in a reaction-diffusion equation

BY TOMÁS CARABALLO, JOSÉ A. LANGA, & JAMES C. ROBINSON<sup>†</sup>

*Departamento de Ecuaciones Diferenciales y Análisis Numérico,  
Universidad de Sevilla,  
Apdo. de Correos 1160,  
41080-Sevilla.  
Spain*

We study in some detail the structure of the random attractor for the Chafee-Infante reaction-diffusion equation perturbed by a multiplicative white noise,

$$du = (-Au + \beta u - u^3) dt + \sigma u \circ dW_t \quad x \in D \subset \mathbb{R}^m.$$

First we prove, for  $m \leq 5$ , a lower bound on the dimension of the random attractor which is of the same order in  $\beta$  as the upper bound we derived in an earlier paper, and is the same as that obtained in the deterministic case. Then we show, for  $m = 1$ , that as  $\beta$  passes through  $\lambda_1$  (the first eigenvalue of the negative Laplacian) from below, the system undergoes a stochastic bifurcation of pitchfork type. We believe that this is the first example of such a stochastic bifurcation in an infinite-dimensional setting. Central to our approach is the existence of a random unstable manifold.

**Keywords:** Random attractors, stochastic bifurcation, Hausdorff dimension

## 1. Introduction

Attractors for infinite-dimensional dynamical systems have proven very useful tool in the study of the asymptotic behaviour of many partial differential equations (see for example Hale (1988), Ladyzhenskaya (1991), Temam (1988, 2nd ed. 1997)). Much attention has been focused on the dimension of these attractors, since in many cases one can prove that this is finite, and deduce that the long-time behaviour of the system depends on only a finite number of degrees of freedom.

The definition of an attractor has been generalized to the stochastic case by Crauel & Flandoli (1994) and Schmalfuß (1992), and has once again been fruitful in furthering our understanding of the associated random and stochastic differential equations, in particular in finite dimensions (Arnold 1998).

Roughly speaking, a random attractor is a family of compact random sets, which are invariant for the stochastic flow and attract all solutions ‘from  $t = -\infty$ ’. Once again, it is possible to show that certain of these attractors are finite-dimensional, although the number of examples is much more limited than in the deterministic case. Debussche (1997) has adapted the most powerful deterministic technique to

<sup>†</sup> Permanent address: Mathematics Institute, University of Warwick, Coventry CV4 7AL. UK.

treat the stochastic case, and in a previous paper (Caraballo *et al.* 2000) we used his method to obtain an upper bound on the Hausdorff dimension of the random attractor for the Chafee-Infante reaction-diffusion equation perturbed by a multiplicative noise in the sense of Stratonovich,

$$du = (-Au + \beta u - u^3) dt + \sigma u \circ dW_t \quad x \in D \subset \mathbb{R}^m. \quad (1.1)$$

Different techniques are required, for both stochastic and deterministic systems, to obtain lower bounds on the dimension of these attractors. Such bounds are generally based on finding some invariant manifolds which must be subsets of the attractor (Babin & Vishik 1983).

Here we adopt this approach, using some ideas due to DaPrato & Debussche (1996) to show (in §4) that there is an  $n$ -dimensional unstable manifold near the zero solution when  $\lambda_n < \beta < \lambda_{n+1}$  (the  $\lambda_j$  are the eigenvalues of the negative Laplacian, see §3). This gives a lower bound on the attractor dimension which is of the same order as our previous upper bound. Remarkably these bounds do not depend on the level of the noise ( $\sigma$  in (1.1)), and are of the same order as the bounds in the deterministic case.

In the last decade there has been some research in stochastic bifurcation theory, although it still seems to be unclear how this theory can be set up in general (Arnold 1998, 2000). However, to our knowledge there are as yet no studies of bifurcations in infinite-dimensional stochastic differential equations. We adopt here the *dynamical* concept of a stochastic bifurcation, which is understood as a qualitative change in the invariant sets (or invariant measures) of the system (see §5).

For the one-dimensional case  $m = 1$  we use the manifold structure, which gave us the lower bound on the dimension, to investigate in more detail what happens to the random attractor as  $\beta$  passes through  $\lambda_1$  from below. For  $\beta < \lambda_1$  we showed previously that the attractor is just the point  $\{0\}$ , and our lower bound show that for  $\beta > \lambda_1$  the Hausdorff dimension of the attractor is at least 1. We show that for  $\lambda_1 < \beta < \lambda_2$  the unstable manifold is in fact tangent to the space spanned by the first eigenfunction of the Laplacian, and thus actually has one branch in the cone of positive solutions and the other one in the cone of negative solutions (§5). It is only a short step from here to the required pitchfork bifurcation, using the theory of monotone random dynamical systems developed by Arnold & Chueshov (1998a & b).

We end with some conclusions and open problems.

## 2. Formulation of the problem and deterministic results

Let  $D \subset \mathbb{R}^m$ ,  $m \leq 5$ , be an open bounded set with regular boundary. We consider the following Chafee-Infante reaction-diffusion equation in  $D$  perturbed by a linear multiplicative white noise

$$du = (\Delta u + \beta u - u^3) dt + \sigma u \circ dW_t, \quad (2.1)$$

with  $u(x, t) = 0$  for  $x \in \partial D$ , and where  $W_t(\omega) : \Omega \rightarrow C^0(\mathbb{R}, \mathbb{R})$  is a one-dimensional Wiener process on the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

We rewrite (2.1) as the following differential equation on  $L^2(D)$ ,

$$du = (-Au + \beta u - u^3) dt + \sigma u \circ dW_t \quad (2.2)$$

where  $A = -\Delta$  on  $D$  with the appropriate (Dirichlet) boundary conditions. The operator  $A$  is positive, linear, self-adjoint and with compact inverse. Thus there exists a sequence  $\{\lambda_j\}$  of positive eigenvalues, whose associated eigenfunctions  $w_j$  (with  $Aw_j = \lambda_j w_j$ ) form an orthonormal basis for  $H$  (e.g. Renardy & Rogers 1992). We order these so that  $\lambda_{n+1} \geq \lambda_n$ .

(a) *The deterministic case ( $\sigma = 0$ )*

When  $\sigma = 0$ , existence and uniqueness results are proved for the equation in Marion (1987), Temam (1988), and Robinson (2001). There exists a unique weak solution

$$u(t; u_0) \in L^2(0, T; H_0^1(D)) \cap L^4((0, T) \times D) \cap C([0, T]; L^2(D)),$$

so that in particular we can use the solutions to define a semigroup  $S(t)$  on  $L^2(D)$ , via

$$S(t)u_0 = u(t; u_0).$$

$S(t)$  satisfies the usual semigroup properties,

$$S(0) = \text{id}, \quad S(t)S(s) = S(t+s), \quad \text{and} \quad S(t)u_0 \text{ continuous in } t \text{ and } u_0. \quad (2.3)$$

It is shown in all three of the above references that the equation also enjoys the existence of a global attractor  $\mathcal{A}$ , that is, a compact invariant set which attracts the orbits of all bounded sets, i.e.

$$S(t)\mathcal{A} = \mathcal{A} \quad \text{for all} \quad t \in \mathbb{R} \quad (2.4)$$

and

$$\text{dist}(S(t)B, \mathcal{A}) \rightarrow 0 \quad \text{as} \quad t \rightarrow \infty, \quad (2.5)$$

where  $B$  is any bounded subset of  $L^2(D)$  and  $\text{dist}(A, B)$  is the Hausdorff semidistance between  $A$  and  $B$ ,

$$\text{dist}(A, B) = \sup_{a \in A} \inf_{b \in B} |a - b|. \quad (2.6)$$

Estimates of the Hausdorff dimension of this attractor in terms of  $\beta$  can also be obtained, namely

$$d_H(\mathcal{A}) \leq C\beta^{m/2}.$$

(Recall that  $D \subset \mathbb{R}^m$ .) By showing the existence of an unstable manifold near the origin, Babin & Vishik (1983) show that a similar lower bound holds, so that in fact there exists a constant  $c$  such that

$$c\beta^{m/2} \leq d_H(\mathcal{A}) \leq C\beta^{m/2}$$

(see also Temam 1988). We write this more compactly as

$$d_H(\mathcal{A}) = O(\beta^{m/2}). \quad (2.7)$$

We will obtain, below, the same behaviour in the stochastic case.

In the particular case  $m = 1$  the structure of this attractor is extremely well understood: the attractor consists of a collection of stationary points, which bifurcate from the origin as  $\beta$  passes through each eigenvalue  $\lambda_j$ , and the unstable manifolds joining them (Hale 1988; Henry 1984). In particular, as  $\beta$  passes through  $\lambda_1$  the attractor changes from a single stable fixed point at  $u \equiv 0$  to a set, homeomorphic to an interval, which consists of the one-dimensional unstable manifold of the origin. This has two distinct components, one of which lies in the cone of positive solutions and on which all solutions approach a new positive fixed point, and one which lies in the cone of negative solutions and on which all solutions approach a new negative fixed point. It is this pitchfork bifurcation that we will seek to mirror in the stochastic case.

### 3. Random attractors

We now discuss briefly the definition of a random dynamical system and a random attractor, unsurprisingly using our equation (2.2) as an illustrative example.

#### (a) Random dynamical systems

Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space and  $\{\theta_t : \Omega \rightarrow \Omega, t \in \mathbb{R}\}$  a family of measure preserving transformations such that  $(t, \omega) \mapsto \theta_t \omega$  is measurable,  $\theta_0 = \text{id}$ , and  $\theta_{t+s} = \theta_t \theta_s$  for all  $s, t \in \mathbb{R}$ . The flow  $\theta_t$  together with the corresponding probability space,

$$(\Omega, \mathcal{F}, \mathbb{P}, (\theta_t)_{t \in \mathbb{R}})$$

is called a (*measurable*) *dynamical system*.

In our case we take  $(\Omega, \mathcal{F}, \mathbb{P})$  to be the probability space which generates the one-dimensional Wiener process  $dW_t$ . The shift  $\theta_t$  acts on  $\Omega$  so that

$$W_t(\theta_s \omega) = W_{t+s}(\omega) - W_s(\omega), \quad (3.1)$$

the additional subtracted term ensuring that  $W_t(\theta_s \omega)$  is still a Brownian motion. For this example it also follows that the shift  $\theta_t$  is ergodic (Arnold 1998).

A continuous *random dynamical system* (RDS) on a Polish space  $(X, d)$  with Borel  $\sigma$ -algebra  $\mathcal{B}$  over  $\theta$  on  $(\Omega, \mathcal{F}, \mathbb{P})$  is a measurable map

$$\begin{aligned} \varphi : \mathbb{R}^+ \times \Omega \times X &\rightarrow X \\ (t, \omega, x) &\mapsto \varphi(t, \omega)x \end{aligned}$$

such that  $\mathbb{P}$  – a.s.

- i)  $\varphi(0, \omega) = \text{id}$  on  $X$
- ii)  $\varphi(t + s, \omega) = \varphi(t, \theta_s \omega) \circ \varphi(s, \omega)$  for all  $t, s \in \mathbb{R}^+$  (cocycle property)
- iii)  $\varphi(t, \omega) : X \rightarrow X$  is continuous.

When  $\sigma \neq 0$  it is known (Pardoux 1975) that for each  $u_0 \in L^2(D)$  and  $T > 0$ , there exists a unique solution  $u(t; x_0)$  of (2.2), with

$$u(t; x_0) \in L^2(\Omega \times (0, T); H_0^1(D)) \cap L^4(\Omega \times (0, T) \times D) \cap L^2(\Omega; C(0, T; L^2(D))).$$

In particular, it follows that the solutions of (2.2) can be used to generate a random dynamical system if we define

$$\varphi(t, \omega)u_0 = u(t; \omega, u_0),$$

where  $u(t; \omega, u_0)$  is the solution of (2.2) with noise  $\omega$  and initial condition  $u(0) = u_0$ .

(b) *Random attractors*

A random set  $\mathcal{A}(\omega)$  is said to be a *random attractor* for the RDS  $\varphi$  if

- i)  $\mathcal{A}(\omega)$  is a random compact set, that is,  $\mathbb{P}$  – a.s.,  $\mathcal{A}(\omega)$  is compact and for all  $x \in X$ , and the map  $\omega \mapsto \text{dist}(x, \mathcal{A}(\omega))$  is measurable with respect to  $\mathcal{F}$ .
- ii)  $\mathbb{P}$  – a.s.  $\varphi(t, \omega)\mathcal{A}(\omega) = \mathcal{A}(\theta_t\omega)$  for all  $t \geq 0$  (invariance) and
- iii) for all  $B \subset X$  bounded (and non-random),  $\mathbb{P}$  – a.s.,

$$\lim_{t \rightarrow \infty} \text{dist}(\varphi(t, \theta_{-t}\omega)B, \mathcal{A}(\omega)) = 0,$$

where  $\text{dist}(\cdot, \cdot)$  denotes the Hausdorff semidistance in  $X$  (cf. (2.6)).

Since  $\varphi(t, \theta_{-t}\omega)u_0$  can be interpreted as the position at  $t = 0$  of the trajectory which was at  $u_0$  at time  $-t$ , this *pullback convergence* property is essentially attraction ‘from  $t = -\infty$ ’.

In Caraballo *et al.* (2000) we proved the existence of a random attractor for our equation, using a theorem due to Crauel & Flandoli (1994). We also showed that if  $\beta < \lambda_1$  then  $\mathcal{A}(\omega) = \{0\}$ , and more generally (using a result of Debussche 1998) that if

$$\beta < \frac{1}{d} \sum_{j=1}^d \lambda_j$$

then  $d_H(\mathcal{A}(\omega)) < d$ . Since one can bound  $\sum_{j=1}^d \lambda_j \leq Cd^{(m+2)/m}$ , this implies the upper bound

$$d_H(\mathcal{A}(\omega)) \leq c_1\beta^{m/2}. \tag{3.2}$$

Note that this is of the same order as the deterministic bound in (2.7), and of course is extremely suggestive of the fact that the attractor becomes more complicated as  $\beta$  increases through  $\lambda_1$ .

#### 4. A lower bound on the attractor dimension

We now turn to proving the promised lower bound on the dimension of the random attractor  $\mathcal{A}(\omega)$ . We make use of an idea from the deterministic theory, which is to show that the attractor must contain an unstable manifold of a certain dimension. However, this approach usually requires various differentiability properties of the flow, and such technicalities are by no means straightforward in the stochastic case. Furthermore, there is currently no well-developed theory of unstable manifolds for general stochastic PDEs.

So we proceed in a manner which we believe is slightly unusual, and could prove useful not only for other stochastic examples but also in the deterministic case, by proving not the existence of a  $C^1$  unstable manifold, but of a Lipschitz manifold in a neighbourhood of the origin. The theory of inertial manifolds, introduced by Foias *et al.* (1988) in the deterministic case, and developed by various authors for stochastic equations (Bensoussan & Flandoli 1994; Chueshov & Girya 1994), is well suited to this, and we follow some ideas from a paper of DaPrato & Debussche (1996) in our proof.

In this way, we prove the following theorem.

**Theorem 4.1.** *If  $\lambda_n < \beta < \lambda_{n+1}$  then there exists a set  $\mathcal{M}_\delta(\omega)$ , a (locally) invariant  $n$ -dimensional Lipschitz manifold, which is part of the unstable set of the origin, i.e.*

$$\lim_{t \rightarrow \infty} \text{dist}(\varphi(-t, \omega)u, 0) = 0 \quad (4.1)$$

for all  $u \in \mathcal{M}(\omega)$ .

In (4.1),  $\varphi(-t, \omega)u$  is the point  $v \in \mathcal{M}(\theta_{-t}\omega)$  such that  $\varphi(t, \theta_{-t}\omega)v = u$  (the fact that  $\mathcal{M}(\omega)$  is a finite-dimensional invariant manifold ensures that such a point exists – see proposition 4.6 for more details).

We now appeal to a result of Crauel (1999), which guarantees that the unstable set of the origin (defined precisely as in (4.1)) must be a subset of the random attractor, and so obtain the lower bound on the attractor dimension given in the following theorem.

**Theorem 4.2.** *Provided that  $m \leq 5$ , if  $\lambda_n < \beta < \lambda_{n+1}$  then  $d_H(\mathcal{A}(\omega)) \geq n$ . In particular, the dimension of the attractor satisfies  $d_H(\mathcal{A}(\omega)) = O(\beta^{m/2})$ .*

(a) *Truncating the equation*

We prove the existence of  $\mathcal{M}_\delta(\omega)$  in a series of propositions. We apply a version of the theory in DaPrato & Debussche (1996) developed to prove the existence of inertial manifolds for stochastic PDEs with a general multiplicative noise term. However, here we do not look for an inertial manifold, but in fact an unstable manifold in a neighbourhood of the origin.

The first task is to truncate the equation in a neighbourhood of the origin, to ensure that the nonlinear term is globally Lipschitz, with a small Lipschitz constant. [Our approach is, in fact, similar to that of Boxler (1991) for centre manifolds, although here we can truncate our equation in a manner which is independent of  $\omega$ .] Although this is a standard approach in the deterministic theory, in which one has a compact absorbing ball, it is perhaps the main weakness of the theory of inertial manifolds for stochastic equations, where in general the radius of the absorbing ball will depend on  $\omega$ , and as yet there is no proof which allows the Lipschitz constant of the nonlinearity to vary in this way. However, since we require only a local unstable manifold, we will be able to truncate the equation in a consistent fashion.

We restrict to the case  $m \leq 5$ , since then we can use the embedding  $H^{2\alpha} \subset L^6$  with  $3/4 < \alpha < 1$  to show that  $F(u) = u^3$  is Lipschitz from  $H^{2\alpha}$  into  $L^2$  (we suppress the  $D$  in all our function spaces from now on),

$$\|u^3 - v^3\|_{L^2} \leq C(\|u\|_{H^{2\alpha}}^2 + \|v\|_{H^{2\alpha}}^2)\|u - v\|_{H^{2\alpha}}. \quad (4.2)$$

Indeed, we have

$$\begin{aligned}
 |u^3 - v^3|_{L^2}^2 &= \int_D (u^3(x) - v^3(x))^2 dx \\
 &= \int_D \left( \int_{v(x)}^{u(x)} 3s^2 ds \right)^2 dx \\
 &\leq 9 \int_D |u(x) - v(x)|^2 (u(x)^2 + v(x)^2)^2 dx \\
 &\leq 9 \left( \int_D |u(x) - v(x)|^{2p} dx \right)^{1/p} \left( \int_D |u(x)^2 + v(x)^2|^{2q} dx \right)^{1/q} \\
 &\leq 9 |u - v|_{L^{2p}}^2 (|u|_{L^{4q}} + |v|_{L^{4q}})^4 \\
 &\leq 9 |u - v|_{L^6}^2 (|u|_{L^6} + |v|_{L^6})^4,
 \end{aligned}$$

taking  $(p, q) = (3, 3/2)$ .

For a  $C^1$  cut-off function  $\theta : \mathbb{R}^+ \rightarrow [0, 1]$ , such that

$$\theta(r) = \begin{cases} 1 & r \leq 1 \\ 0 & r \geq 2 \end{cases}$$

with  $|\theta'(r)| \leq 2$ , it is straightforward to verify (cf. Temam, 1988) that

$$F(u) = -\theta\left(\frac{\|u\|_{H^{2\alpha}}}{R}\right) u^3 \tag{4.3}$$

is globally bounded,

$$|F(u)| \leq M_0 \quad \text{for all } u \in H^{2\alpha},$$

and globally Lipschitz,

$$|F(u) - F(v)| \leq \tilde{L}_f \|u - v\|_{H^{2\alpha}} \quad \text{for all } u, v \in H^{2\alpha}. \tag{4.4}$$

with  $\tilde{L}_f \leq c_0 R^2$  for some constant  $c_0$ . Note in particular that  $F(u) = -u^3$  in  $B_{H^{2\alpha}}(0, R)$ , and so the two equations agree in a small  $H^{2\alpha}$ -neighbourhood of the origin.

We will find it more convenient to use the norm equivalence between the  $H^{2\alpha}$  norm and the norm in  $D(A^\alpha)$ . Writing  $|A^\alpha u| = |u|_\alpha$ , where  $A$  is the negative Laplacian on  $D$  with Dirichlet boundary conditions, we have

$$\|u\|_{H^{2\alpha}} \leq c |u|_\alpha. \tag{4.5}$$

We can re-write (4.4) as

$$|F(u) - F(v)| \leq L_f |u - v|_\alpha \quad \text{for all } u, v \in D(A^\alpha), \tag{4.6}$$

where  $L_f = c\tilde{L}_f$  (with  $c$  as in (4.5)).

Note that we can make  $L_f$  as small as we wish provided that  $R$  in (4.3) is chosen small enough.

(b) *An invariant manifold for the truncated equation*

We now work with the truncated equation

$$du = (-Au + \beta u + F(u)) dt + \sigma u \circ dW_t, \quad (4.7)$$

and prove the existence of an invariant manifold of dimension  $n$  when  $\lambda_n < \beta < \lambda_{n+1}$ . In the statement of the theorem we write  $P_n$  for the orthogonal projection onto the first  $n$  eigenfunctions of  $A$ , ordered so that the corresponding eigenvalues are non-decreasing in  $n$ , and  $Q_n$  for its orthogonal complement,  $Q_n = I - P_n$ .

**Theorem 4.3.** *If  $\lambda_n < \beta < \lambda_{n+1}$ , and  $R$  in (4.3) is chosen small enough that  $L_f$  satisfies*

$$\beta - \lambda_n > 4L_f\lambda_n^\alpha \quad (4.8)$$

and

$$\lambda_{n+1} - \beta > (2KL_f)^{1/(1-\alpha)}, \quad (4.9)$$

then (4.7) has an  $n$ -dimensional invariant manifold  $\mathcal{M}(\omega)$ ,

$$\varphi(\lambda, \omega)\mathcal{M}(\omega) = \mathcal{M}(\theta_\lambda\omega), \quad \lambda \geq 0.$$

Furthermore,  $\mathcal{M}(\omega)$  is given as the graph of a Lipschitz function  $\Phi_\omega : P_n H \rightarrow Q_n H \cap D(A^\alpha)$ ,

$$|\Phi_\omega(p_1) - \Phi_\omega(p_2)|_\alpha \leq 2|p_1 - p_2|_\alpha. \quad (4.10)$$

We analyse the equation  $\omega$ -by- $\omega$ . To this end we consider not equation (2.2) for  $u$ , but, setting  $v = e^{-\sigma W_t}u$  and observing that

$$dv = e^{-\sigma W_t}du - e^{-\sigma W_t}\sigma u \circ dW_t,$$

we can treat the non-autonomous equation for  $v$ ,

$$\frac{dv}{dt} = -Av + \beta v + e^{-\sigma W_t}F(e^{\sigma W_t}v). \quad (4.11)$$

We write this as

$$\frac{dv}{dt} = -Av + \beta v + F_\sigma(v),$$

where

$$F_\sigma(v) = e^{-\sigma W_t}F(e^{\sigma W_t}v).$$

Note that

$$\begin{aligned} |F_\sigma(u) - F_\sigma(v)| &\leq e^{-\sigma W_t}|F(e^{\sigma W_t}u) - F(e^{\sigma W_t}v)| \\ &\leq e^{-\sigma W_t}L_f|e^{\sigma W_t}u - e^{\sigma W_t}v|_\alpha \\ &= L_f|u - v|_\alpha, \end{aligned} \quad (4.12)$$



so that the Lipschitz property of  $F$  in (4.6) is transferred to  $F_\sigma$ . Furthermore it follows, since the support of  $F$  is contained in a bounded set in  $D(A^\alpha)$ , and  $F(0) = 0$ , that

$$|F_\sigma(u)| \leq M_1$$

for some constant  $M_1$ .

The solution operator corresponding to the transformed equation we write as  $\psi(t, \omega)$ . Clearly,

$$\psi(t, \omega) = e^{-\sigma W_t(\omega)} \varphi(t, \omega). \quad (4.13)$$

For a function

$$y : (-\infty, 0] \rightarrow P_n H \quad (4.14)$$

we define a norm

$$\|y\|_E = \sup_{t \in (-\infty, 0]} |y(t)|_\alpha, \quad (4.15)$$

and similarly for

$$z : (-\infty, 0] \rightarrow Q_n D(A^\alpha). \quad (4.16)$$

For a pair  $(y, z)$ , we define

$$\|(y, z)\|_E = \max(\|y\|_E, \|z\|_E). \quad (4.17)$$

The space of all functions  $(y, z)$  as in (4.14) and (4.16) with finite  $E$  norm (the norm in (4.17)) we denote by  $E$ .

We will need the following standard bound on the operator norm of  $A^\alpha e^{-At} Q_n$  from  $L^2$  into  $L^2$  (see Temam 1988, for example),

$$\|A^\alpha e^{-At} Q_n\|_{op} \leq \tilde{K}(t^{-\alpha} + \lambda_{n+1}^\alpha) e^{-\lambda_{n+1} t} \quad \text{for all } t \geq 0. \quad (4.18)$$

We also set

$$K = \tilde{K} \int_0^\infty (t^{-\alpha} + 1) e^{-t} dt. \quad (4.19)$$

From now on we write  $P = P_n$  and  $Q = Q_n$ .

Roughly speaking, we follow the proof in DaPrato & Debussche (1996). However, here the argument is somewhat different, since our particular choice of multiplicative noise allows us to consider the transformed equation (4.11)  $\omega$ -by- $\omega$ , and we do not have to deal with the problems which arise when trying to solve a more general stochastic equation backwards in time. We also need to tailor the analysis to make sure that the dependence on  $\beta$  remains explicit.

**Proposition 4.4.** *For each  $\omega \in \Omega$ , there exists a unique solution  $(y, z)$  of the coupled equations*

$$\begin{aligned} \frac{dy}{dt} &= -Ay + \beta y + PF_\sigma(y + z) & y(0) &= y_0 \\ \frac{dz}{dt} &= -Az + \beta z + QF_\sigma(y + z) & z(t) &\rightarrow 0 \text{ as } t \rightarrow -\infty, \end{aligned} \quad (4.20)$$

on the interval  $(-\infty, 0]$ .

*Proof.* We use a contraction mapping argument. For a given pair  $(\eta, \xi) \in E$ , we define

$$\mathcal{T}_{y_0}(\eta, \xi) = (y, z),$$

where  $(y, z)$  solve the coupled system

$$\begin{aligned} \frac{dy}{dt} &= -Ay + \beta y + PF_\sigma(\eta + \xi) & y(0) &= y_0 \\ \frac{dz}{dt} &= -Az + \beta z + QF_\sigma(\eta + \xi) & z(t) &\rightarrow 0 \text{ as } t \rightarrow -\infty. \end{aligned} \quad (4.21)$$

Note that such a solution is well-defined, since, using the variation of constants formula on the second equation we have

$$z(t) = e^{-(A-\beta I)(t-\tau)}z(\tau) + \int_\tau^t e^{-A(t-s)}QF_\sigma(\xi(s) + \eta(s)) ds.$$

Since  $(A - \beta I)$  is a positive operator, letting  $\tau \rightarrow -\infty$  and using the condition that  $z(\tau) \rightarrow 0$  we obtain the unique solution

$$z(t) = \int_{-\infty}^t e^{-A(t-s)}QF_\sigma(\xi(s) + \eta(s)) ds. \quad (4.22)$$

To show that  $\mathcal{T}_{y_0}$  is a contraction on  $E$ , first we estimate the difference of two solutions of the equation

$$\frac{dy_i}{dt} = -Ay_i + \beta y_i + PF_\sigma(\eta_i + \xi_i) \quad y_i(0) = y_0, \quad (4.23)$$

corresponding to two different choices of  $(\eta, \xi) \in E$  in (4.21). Writing  $y(t) = y_1(t) - y_2(t)$ , it follows that  $y$  satisfies

$$\frac{dy}{dt} = -Ay + \beta y + PF_\sigma(\eta_1 + \xi_1) - PF_\sigma(\eta_2 + \xi_2) \quad y(0) = 0. \quad (4.24)$$

Using the variation of constants formula, we can write, for  $t < 0$ ,

$$y(0) = e^{(A-\beta I)t}y(t) + \int_t^0 e^{(A-\beta I)(t-s)}(PF_\sigma(\eta_1(s) + \xi_1(s)) - PF_\sigma(\eta_2(s) + \xi_2(s))) ds,$$

and so, since  $y(0) = 0$ ,

$$y(t) = - \int_t^0 e^{-(A-\beta I)s}(PF_\sigma(\eta_1(s) + \xi_1(s)) - PF_\sigma(\eta_2(s) + \xi_2(s))) ds. \quad (4.25)$$

Taking the norm in  $D(A^\alpha)$  throughout (4.25) we obtain, since  $-(A - \beta I)$  is a positive operator on  $PH$ , with eigenvalues bounded below by  $\beta - \lambda_n$ ,

$$|y(t)|_\alpha \leq L_f \lambda_n^\alpha \int_t^0 e^{(\beta - \lambda_n)s}(|\eta(s)|_\alpha + |\xi(s)|_\alpha) ds,$$

where  $\eta = \eta_1 - \eta_2$ , and similarly for  $\xi$ .

This becomes

$$\begin{aligned} |y(t)|_\alpha &\leq 2L_f \lambda_n^\alpha \int_t^0 e^{(\beta-\lambda_n)s} \|(\eta, \xi)\|_E ds \\ &\leq \frac{2L_f \lambda_n^\alpha}{\beta - \lambda_n} \|(\eta, \xi)\|_E [1 - e^{(\beta-\lambda_n)t}]. \end{aligned}$$

Since  $\beta > \lambda_n$ ,

$$\|y\|_E \leq \frac{2L_f \lambda_n^\alpha}{\beta - \lambda_n} \|(\eta, \xi)\|_E. \quad (4.26)$$

(4.8) now guarantees that  $\|y\|_E \leq \frac{1}{2} \|(\eta, \xi)\|_E$ .

For a similar choice of  $(\eta_i, \xi_i)$ , we now estimate the difference of solutions of the two equations

$$\frac{dz_i}{dt} = -Az_i + \beta z_i + QF_\sigma(\eta_i + \xi_i) \quad z_i(t) \rightarrow 0 \quad \text{as } t \rightarrow -\infty.$$

Setting  $z = z_1 - z_2$ , we have

$$\frac{dz}{dt} = -Az + \beta z + QF_\sigma(\eta_1 + \xi_1) - QF_\sigma(\eta_2 + \xi_2).$$

As before we use the variation of constants formula, and since  $z(t) \rightarrow 0$  as  $t \rightarrow -\infty$  we have

$$z(t) = \int_{-\infty}^t e^{-(A-\beta I)(t-s)} Q[F_\sigma(\eta_1 + \xi_1) - F_\sigma(\eta_2 + \xi_2)] ds.$$

Once more, taking the norm in  $D(A^\alpha)$ , using the Lipschitz property of  $F_\sigma$  in (4.12), and the bound on  $A^\alpha e^{-A(t-s)} Q$  in (4.18) we have

$$|z(t)|_\alpha \leq \tilde{K} L_f \int_{-\infty}^t ((t-s)^{-\alpha} + (\lambda_{n+1} - \beta)^\alpha) e^{-(\lambda_{n+1} - \beta)(t-s)} (|\eta(s)|_\alpha + |\xi(s)|_\alpha) ds.$$

Using the definition of the norm in  $E$  (4.17) now gives

$$\begin{aligned} |z(t)|_\alpha &\leq \tilde{K} L_f \int_{-\infty}^t ((t-s)^{-\alpha} + (\lambda_{n+1} - \beta)^\alpha) e^{-(\lambda_{n+1} - \beta)(t-s)} \|(\eta, \xi)\|_E ds \\ &\leq \tilde{K} L_f \|(\eta, \xi)\|_E \int_{-\infty}^0 (|\tau|^{-\alpha} + (\lambda_{n+1} - \beta)^\alpha) e^{(\lambda_{n+1} - \beta)\tau} d\tau \\ &\leq \tilde{K} L_f \|(\eta, \xi)\|_E \int_{-\infty}^0 (|\tau|^{-\alpha} + (\lambda_{n+1} - \beta)^\alpha) e^{(\lambda_{n+1} - \beta)\tau} d\tau \\ &\leq \frac{\tilde{K} L_f}{\lambda_{n+1} - \beta} \|(\eta, \xi)\|_E (\lambda_{n+1} - \beta)^\alpha \int_{-\infty}^0 [|u|^{-\alpha} + 1] e^u du. \end{aligned}$$

It follows that

$$\|z\|_E \leq \frac{K L_f}{(\lambda_{n+1} - \beta)^{1-\alpha}} \|(\eta, \xi)\|_E,$$

where the constant  $K$  is defined in (4.19).

The condition on  $L_f$  in (4.9) now ensures that  $\|z\|_E \leq \frac{1}{2} \|(\eta, \xi)\|_E$ . It follows that  $\|(y, z)\|_E \leq \frac{1}{2} \|(\eta, \xi)\|_E$ , and so  $\mathcal{T}_{y_0}$  is a contraction, and we obtain the unique solution of our coupled pair of equations.  $\square$

We now define

$$\Phi_\omega(y_0) = z(0),$$

where  $z$  is the solution of (4.20) with  $y(0) = y_0$  and noise  $W_t(\omega)$ . Observe in particular that  $\Phi_\omega(0) = 0$  for all  $\omega \in \Omega$ .

**Proposition 4.5.** *For almost every  $\omega \in \Omega$ ,*

$$|\Phi_\omega(y_1^0) - \Phi_\omega(y_2^0)|_\alpha \leq 2|y_1^0 - y_2^0|_\alpha,$$

and the graph  $\mathcal{M}(\omega)$  of  $\Phi_\omega$  is invariant for  $\varphi$ ,

$$\varphi(\lambda, \omega)\mathcal{M}(\omega) = \mathcal{M}(\theta_\lambda \omega) \quad \lambda \geq 0,$$

i.e for any  $y_0 \in PH$  and  $\lambda \geq 0$  we have

$$\varphi(\lambda, \omega)(y_0 + \Phi_\omega(y_0)) = y_\lambda + \Phi_{\theta_\lambda \omega}(y_\lambda), \quad (4.27)$$

$\mathbb{P}$  - a.s., where

$$y_\lambda = P\varphi(\lambda, \omega)(y_0 + \Phi_\omega(y_0)).$$

*Proof.* First we show the Lipschitz property of  $\Phi_\omega$ , by estimating

$$\|\mathcal{T}_{y_0^1}(\eta, \xi) - \mathcal{T}_{y_0^2}(\eta, \xi)\|_E.$$

We will write

$$(y^i, z^i) = \mathcal{T}_{y_0^i}(\eta, \xi).$$

Since  $z^1 = z^2$ , we need only estimate the difference  $y = y_1 - y_2$  between two solutions of

$$\frac{dy_i}{dt} = -(A - \beta I)y_i + PF_\sigma(\eta + \xi) \quad y_i(0) = y_0^i,$$

but with the same pair  $(\eta, \xi) \in E$ . So,

$$\frac{dy}{dt} = -(A - \beta I)y.$$

Thus

$$y(0) = e^{(A - \beta I)t}y(t),$$

and so, for  $t < 0$ ,

$$|y(t)|_\alpha \leq e^{(\beta - \lambda_n)t}|y_0^1 - y_0^2|_\alpha.$$

It follows, since  $\beta > \lambda_n$ , that

$$\|\mathcal{T}_{y_0^1}(\eta, \xi) - \mathcal{T}_{y_0^2}(\eta, \xi)\|_E \leq |y_0^1 - y_0^2|_\alpha.$$

So we have

$$\begin{aligned} \|(y^1, z^1) - (y^2, z^2)\|_E &= \|\mathcal{T}_{y_0^1}(y^1, z^1) - \mathcal{T}_{y_0^2}(y^2, z^2)\|_E \\ &\leq \|\mathcal{T}_{y_0^1}(y^1, z^1) - \mathcal{T}_{y_0^2}(y^1, z^1)\|_E \\ &\quad + \|\mathcal{T}_{y_0^2}(y^1, z^1) - \mathcal{T}_{y_0^2}(y^2, z^2)\|_E \end{aligned}$$

$$\leq |y_0^1 - y_0^2|_\alpha + \frac{1}{2} \|(y^1, z^1) - (y^2, z^2)\|_E,$$

and so

$$\|(y^1, z^1) - (y^2, z^2)\|_E \leq 2|y_0^1 - y_0^2|_\alpha.$$

Since  $\Phi_\omega(y_0^i) = z^i(0)$ , the Lipschitz bound follows.

To show the invariance, observe that  $\Phi_\omega(y_0)$  is defined as the solution at time 0 of the  $z$  component of (4.20) when  $y(0) = y_0$ . Solving (4.11) with the initial condition  $y_0 + \Phi_\omega(y_0)$  produces a solution of the coupled equations

$$\begin{aligned} \frac{dy}{dt} &= -Ay + \beta y + PF_\sigma(y + z) & y(\lambda) &= P\psi(\lambda, \omega)[y_0 + \Phi_\omega(y_0)] \\ \frac{dz}{dt} &= -Az + \beta z + QF_\sigma(y + z) & z(t) &\rightarrow 0 \quad \text{as } t \rightarrow -\infty, \end{aligned}$$

which is unique, by proposition 4.4. Now, observe that  $z(\lambda)$  is the same as the  $z(0)$  produced as the solution of (4.11) with  $W_t(\omega)$  replaced by

$$W_{t+\lambda}(\omega) = W_t(\theta_\lambda \omega) + W_\lambda(\omega).$$

With this change we obtain the solution of (4.11) with  $\omega$  replaced by  $\theta_\lambda \omega$ , but multiplied by an additional factor of  $e^{\sigma W_\lambda(\omega)}$ .

We therefore obtain not invariance of  $\mathcal{M}(\omega)$  for the transformed equation, but rather

$$\psi(\lambda, \omega)\mathcal{M}(\omega) = e^{\sigma W_\lambda(\omega)}\mathcal{M}(\theta_\lambda \omega).$$

The required invariance property for the untransformed equation is now immediate using (4.13).  $\square$

(c) *A local unstable manifold for the original equation*

Finally, we show that the inertial manifold found in the previous section is in fact part of the unstable set of the origin, within the small neighbourhood on which the truncated and original equations agree.

**Proposition 4.6.** *There exists a  $\delta(\omega)$  such that  $\mathcal{M}_\delta(\omega)$ , the intersection of  $\mathcal{M}(\omega)$  with the ball  $B_{H^{2\alpha}}(0, \delta(\omega))$ , is part of the unstable set of the origin for the original equation.*

*Proof.* We consider the stochastic ‘‘inertial form’’, i.e. the equation on the inertial manifold for the  $P$  components of  $u$ , and show that there exists a  $\delta(\omega) < R$  such that, for all  $p_0$  with  $|p_0|_\alpha \leq \delta$

- (i)  $|p(t)|_\alpha \leq R$  for all  $t \leq 0$ , and
- (ii)  $p(t) \rightarrow 0$  as  $t \rightarrow -\infty$ .

Point (i) shows that the trajectory ‘‘backwards’’ from  $p_0 + \Phi_\omega(p_0)$  remains within  $B_{H^{2\alpha}}(0, R)$ , so that it is fact a solution of the original (untruncated) equation, while (ii) shows that  $p_0 + \Phi_\omega(p_0)$  is an element of the unstable set of the origin.

Note that we can consider trajectories ‘‘backwards in time’’ since  $\mathcal{M}(\omega)$  is invariant:  $\varphi(t, \theta_{-t}\omega)\mathcal{M}(\theta_{-t}\omega) = \mathcal{M}(\omega)$ , and so for  $u \in \mathcal{M}(\omega)$  there exists at least one point  $v$  in  $\mathcal{M}(\theta_{-t}\omega)$  such that  $\varphi(t, \theta_{-t}\omega)v = u$ . Since the governing equation

reduces to a finite-dimensional random ordinary differential equation on  $\mathcal{M}(\omega)$  (see below) this point is unique.

The inertial form is

$$dp = \{-(A - \beta I)p + PF(p + \Phi_{\theta_t\omega}(p))\} dt + \sigma p \circ dW_t.$$

Writing  $x(t) = e^{-\sigma W_t} p(t)$  this becomes

$$\frac{dx}{dt} = -(A - \beta I)x + e^{-\sigma W_t} PF(e^{\sigma W_t} x + \Phi_{\theta_t\omega}(e^{\sigma W_t} x)).$$

Now, we have

$$|A^\alpha PF(u)| \leq \lambda_n^\alpha |F(u)| \leq L_f \lambda_n^\alpha |u|_\alpha,$$

since  $F$  is Lipschitz and  $F(0) = 0$ . Therefore we have, noting that the Lipschitz constant of  $\Phi_\omega$  in (4.10) is independent of  $\omega$ ,

$$\frac{1}{2} \frac{d}{dt} |x(-t)|_\alpha^2 \leq -(\beta - \lambda_n) |x(-t)|_\alpha^2 + e^{-\sigma W_{-t}} \lambda_n^\alpha 3L_f e^{\sigma W_{-t}} |x(-t)|_\alpha^2,$$

or

$$\frac{d}{dt} |x(-t)|_\alpha \leq -(\beta - \lambda_n - 3\lambda_n^\alpha L_f) |x(-t)|_\alpha.$$

Note that it follows from (4.8) that

$$\gamma \equiv \beta - \lambda_n - 3\lambda_n^\alpha L_f > \lambda_n^\alpha L_f,$$

and in particular is positive. We therefore have, for  $t < 0$ ,

$$|x(t)|_\alpha \leq e^{\gamma t} |x_0|_\alpha, \tag{4.28}$$

which gives, for  $p(t)$ ,

$$|p(t)|_\alpha \leq e^{\gamma t} e^{\sigma W_t} |p_0|_\alpha.$$

Now, for each fixed  $\omega$ , given  $\epsilon > 0$  there exists a constant  $C_\epsilon(\omega)$  such that

$$e^{\sigma W_t} \leq C_\epsilon e^{-\epsilon t} \quad \text{for all } t \leq 0. \tag{4.29}$$

Therefore we have

$$|p(t)|_\alpha \leq e^{\gamma t} e^{-\epsilon t} C_\epsilon(\omega) |p_0|_\alpha,$$

and so if we choose  $\delta(\omega) = \min(R, R/C_\epsilon(\omega))$  the result follows.  $\square$

## 5. A bifurcation of pitchfork type

We now make further use of the unstable manifold to help us prove the existence of a stochastic bifurcation as  $\beta$  increases through  $\lambda_1$ .

(a) *A definition of a stochastic pitchfork bifurcation*

To motivate our definition we briefly recall a simple example in which such a bifurcation has been studied by Arnold & Boxler (1992).

The stochastic ordinary differential equation

$$dx = (\beta x - x^3) dt + \sigma x \circ dW_t,$$

can be solved explicitly. As  $\beta$  passes through zero from below, behaviour similar to that in the deterministic case can be observed. A *random fixed point* is a random set which, for almost all  $\omega$ , consists of just one point,  $a(\omega)$ , and is invariant,

$$\varphi(t, \omega)a(\omega) = a(\theta_t \omega).$$

Arnold & Boxler (1992) show that for  $\beta < 0$  the random attractor is just the fixed point  $\{0\}$ , whereas for  $\beta > 0$  two new random fixed points appear,  $\pm a(\omega)$ . The random attractor now consists of the interval  $[-a(\omega), +a(\omega)]$ .

Crauel *et al.* (1999) discuss such bifurcations in one-dimensional systems in more detail, and give a definition of a stochastic pitchfork bifurcation in terms of random invariant measures. They define a stochastic pitchfork bifurcation as the appearance of two new invariant measures, associated with new random fixed points, which are stable in an appropriate sense.

Below, we adopt a definition of a pitchfork bifurcation based on the appearance of new random fixed points rather than in terms of invariant measures.

**Definition 5.1.** *Let  $(\varphi_\beta)_{\beta \in \mathbb{R}}$  a family of RDS parametrized by  $\beta$ . We say that  $(\varphi_\beta)$  undergoes a stochastic pitchfork bifurcation at  $\beta = \beta_0$  if*

- i) For  $\beta \leq \beta_0$ , the random attractor consists solely of one (random) fixed point,  $\{0\}$ , and*
- ii) for  $\beta > \beta_0$ , the random attractor contains  $\{0\}$  and two new random fixed points,  $a^+(\omega)$  and  $a^-(\omega)$ , and  $\mathbb{P}$ -a.s.  $a^\pm(\omega) \rightarrow 0$  as  $\beta \downarrow \beta_0$ .*

Although, unlike Crauel *et al.* (1999), we do not include a criterion of stability for the new fixed points, we believe that this type of bifurcation still merits the title ‘‘pitchfork’’, and we now prove that, in the case of a one-dimensional domain, such a bifurcation occurs as  $\beta$  passes through  $\lambda_1$ . Some discussion of the stability of the new fixed points is given at the end of §5.

We showed in a previous paper (Caraballo *et al.* 2000) that for  $\beta < \lambda_1$  the random attractor consists solely of the fixed point  $u \equiv 0$ . To study the structure of the attractor for  $\beta > \lambda_1$ , we argue as follows. First we show that the manifold  $\mathcal{M}(\omega)$  is tangent to the linear space spanned by the first eigenfunction of the Laplacian, and with a little care it follows that this manifold intersects the cone  $\mathcal{K}^+$  of positive functions. Since  $\mathcal{K}^+$  is in fact invariant for the cocycle (e.g. Kotelenetz 1992) the restriction of the cocycle to  $\mathcal{K}^+$  has a non-trivial attractor  $\mathcal{A}^+(\omega)$ . Now, since the cocycle is order-preserving, the theory developed by Arnold & Chueshov (1998) shows that functions in  $\mathcal{A}^+(\omega)$  are bounded above and below by random fixed points: below by zero, and above by a new random fixed point  $a^+(\omega)$ . The same argument applies in  $\mathcal{K}^-$  and proves the presence of a pitchfork bifurcation.

(b) *Differentiability of  $\mathcal{M}(\omega)$  at 0*

First we prove the differentiability property of the manifold. In particular, the cubic bound (5.1) will be used to show that a portion of  $\mathcal{M}(\omega)$  lies in  $\mathcal{K}^+$ .

**Proposition 5.2.** *The manifold  $\mathcal{M}(\omega)$  is tangent to  $P_n H$  at the origin, and moreover*

$$|\Phi_\omega(p)|_\alpha \leq C_\omega |p|_\alpha^3. \quad (5.1)$$

*Proof.* First, observe that

$$F_\sigma(u) = -e^{-\sigma W_t} \theta \left( \frac{|ue^{\sigma W_t}|_\alpha}{R} \right) (ue^{\sigma W_t})^3$$

is essentially cubic, so that

$$\begin{aligned} |F_\sigma(u)| &\leq e^{2\sigma W_t} \left( \int u^6 dx \right)^{1/2} \\ &\leq e^{2\sigma W_t} \|u\|_{L^6}^3 \\ &\leq C e^{2\sigma W_t} |u|_\alpha^3. \end{aligned}$$

Now,  $\Phi_\omega(y_0)$  is  $z(0)$  from the coupled equations in proposition 4.4, so we know (cf. (4.22)) that

$$z(0) = \int_{-\infty}^0 e^{(A-\beta I)s} Q F_\sigma(y(s) + z(s)) ds,$$

and hence

$$|z(0)|_\alpha \leq \tilde{K} \int_{-\infty}^t ((-s)^{-\alpha} + (\lambda_{n+1} - \beta)^\alpha) e^{(\lambda_{n+1} - \beta)s} C e^{2\sigma W_s} |u|_\alpha^3 ds.$$

Since

$$|u|_\alpha^3 \leq (|y|_\alpha + |z|_\alpha)^3 \leq 8(|y|_\alpha^3 + |z|_\alpha^3),$$

we have, using (4.29),

$$|z(0)|_\alpha \leq K_\omega [\|y\|_E^3 + \|z\|_E^3]. \quad (5.2)$$

Now, we know that

$$z(t) = \int_{-\infty}^t e^{-(A-\beta I)(t-s)} Q F_\sigma(y + z) ds,$$

and we can estimate  $|z(t)|_\alpha$  as in the proof of proposition 4.4, using the fact that  $F_\sigma(0) = 0$ ,

$$\begin{aligned} |z(t)|_\alpha &\leq \tilde{K} L_f \int_{-\infty}^t ((t-s)^\alpha + (\lambda_{n+1} - \beta)^\alpha) e^{-(\lambda_{n+1} - \beta)(t-s)} |y + z|_\alpha ds \\ &\leq \left( \tilde{K} L_f \int_{-\infty}^t ((t-s)^\alpha + (\lambda_{n+1} - \beta)^\alpha) e^{-(\lambda_{n+1} - \beta)(t-s)} ds \right) \end{aligned}$$



$$\times (\|y\|_E + \|z\|_E).$$

Now, we saw in proposition 4.4 that the integral expression here is bounded by  $\frac{1}{2}$ , and so we have  $\|z\|_E \leq \frac{1}{2}(\|y\|_E + \|z\|_E)$ , i.e.  $\|z\|_E \leq \|y\|_E$ .

Equation (5.2) now becomes

$$|z(0)|_\alpha \leq 2K_\omega \|y\|_E^3, \tag{5.3}$$

so it only remains to estimate  $\|y\|_E$ . However, this is the solution of the governing equation which has  $y(0) = y_0$  and  $z(0) = \Phi_\omega(y_0)$ , and so which lies on  $\mathcal{M}(\omega)$ . The inequality (4.28) from proposition 4.6 shows that in fact

$$|y(t)|_\alpha \leq |y_0|_\alpha \quad \text{for all } t \leq 0,$$

and so  $\|y\|_E \leq |y_0|_\alpha$ . Combining this with (5.3) gives the required bound.  $\square$

(c) *The bifurcation theorem*

We now prove our second main result.

**Theorem 5.3.** *Let  $m = 1$ . Then (2.2) undergoes a stochastic pitchfork bifurcation at  $\beta = \lambda_1$ . In particular, for  $\beta > \lambda_1$  there exist positive and negative random fixed points  $\pm a(\omega)$ , and  $a(\omega) \rightarrow 0$  as  $\beta \downarrow \lambda_1$ .*

Note that one could easily recast this result in terms of the appearance of two new invariant measures, namely the random Dirac measures concentrated at  $\pm a(\omega)$ .

*Proof.* Choose and fix  $\beta > \lambda_1$ . Since the first eigenfunction of the Laplacian is positive, we use the cubic estimate to show that  $\mathcal{M}_\delta(\omega)$  must contain a positive function. More precisely, we let

$$\mathcal{K}^+ = \{u \in L^2(D) : u(x) \geq 0 \text{ almost everywhere}\},$$

and

$$\mathcal{K}^- = \{u \in L^2(D) : u(x) \leq 0 \text{ almost everywhere}\}.$$

The main idea is to show that one portion of the unstable manifold must be a subset of  $\mathcal{K}^+$ , and another a subset of  $\mathcal{K}^-$ .

On a one-dimensional domain  $[0, L]$  we know that the eigenfunctions of the Laplacian with Dirichlet boundary conditions are proportional to

$$w_n(x) = \sin(n\pi x/L).$$

Since the manifold  $\mathcal{M}_\delta(\omega)$  is given as a graph over the space spanned by the first eigenfunction, for each fixed  $\omega$  the manifold consists of a family of functions (here parametrised by  $\epsilon$ ) given in the form

$$u(x, \epsilon) = \epsilon \sin(\pi x/L) + \sum_{j=2}^{\infty} c_j \sin(j\pi x/L),$$

where  $c_j = c_j(\epsilon, \omega)$ .

Since

$$\left| \frac{\sin(j\pi x/L)}{\sin(\pi x/L)} \right| \leq j$$

(simply rewrite the sine terms using complex exponentials) we have

$$u(x, \epsilon) \geq \sin(\pi x/L) \left[ \epsilon - \sum_{j=2}^{\infty} |c_j| j \right].$$

Using the cubic bound on  $\Phi_\omega$ , we know that

$$\left| \sum_{j=2}^{\infty} c_j \sin(j\pi x/L) \right|_\alpha = C \left( \sum_{j=2}^{\infty} |c_j|^2 j^{4\alpha} \right)^{1/2} \leq C_\omega |\epsilon \sin(\pi x/L)|_\alpha^3 = K_\omega \epsilon^3.$$

Since  $\alpha > 3/4$  we can now write

$$\sum_{j=2}^{\infty} |c_j| j \leq \left( \sum_{j=2}^{\infty} j^{4\alpha} |c_j|^2 \right)^{1/2} \left( \sum_{j=2}^{\infty} j^{2(1-2\alpha)} \right)^{1/2} \leq C \epsilon^3,$$

and so

$$u(x, \epsilon) \geq \sin(\pi x/L) [\epsilon - C \epsilon^3].$$

Thus for  $\epsilon > 0$  small enough,  $u(x, \epsilon) \geq 0$  on  $[0, L]$ ; similarly for  $\epsilon < 0$  and small enough,  $u(x, \epsilon) \leq 0$  on  $[0, L]$ . In other words, there are portions of  $\mathcal{M}_\delta(\omega)$  near the origin which intersect non-trivially with  $\mathcal{K}^+$  and  $\mathcal{K}^-$ .

Now, Kotelenetz (1992) shows that the cocycle generated by the equation is order-preserving on  $L^2(D)$ , so that if  $u_0 \geq v_0$  almost everywhere then

$$\varphi(t, \omega) u_0 \geq \varphi(t, \omega) v_0.$$

In particular, since zero is a fixed point of the equation,  $\mathcal{K}^\pm$  are invariant subspaces for  $\varphi$ . (These two facts are easy to check if  $u_0$  and  $v_0$  are continuous functions by applying the standard deterministic theory (essentially the maximum principle, see Smith 1995) to the equation for  $v(t) = e^{-\sigma W_t} u(t)$ .)

By following the analysis in Caraballo *et al.* (2000) one can show that the flow in each of these subspaces has a positively invariant compact absorbing set, namely a bounded set in  $H^1(D)$  with radius  $r(\omega)$ , where  $r(\omega)$  is a tempered random variable. In particular therefore, using the theory in Flandoli & Schmalfuss (1996) the flow in each of these subspaces has its own random attractor,  $\mathcal{A}^+(\omega)$  and  $\mathcal{A}^-(\omega)$ , and these attract all tempered random sets. Since  $\mathcal{M}_\delta(\omega) \cap \mathcal{K}^\pm$  forms part of the unstable set of the origin, it must be a subset of  $\mathcal{A}^\pm(\omega)$ , and so in particular  $\mathcal{A}^\pm(\omega)$  is non-trivial.

In their 1998 paper Arnold & Chueshov develop an extensive theory for order-preserving random dynamical systems, and in particular one of their results (theorem 2) provides the existence of new stochastic fixed points in this example (essentially we follow Chueshov, 2000): if  $\varphi$  is an order-preserving random dynamical system for which there exists a random interval

$$\mathcal{J}(\omega) = \{u : b(\omega) \leq u \leq c(\omega)\}$$

which is attracted to and contains  $\mathcal{A}(\omega)$ , then there exist random fixed points  $u_-(\omega) \leq u_+(\omega)$  such that  $u_-(\omega) \leq u \leq u_+(\omega)$  for all  $u \in \mathcal{A}(\omega)$ .

Since  $\mathcal{A}(\omega)$  lies within a random ball in  $H^1(D)$  of radius  $r(\omega)$ , and since  $H^1(D) \subset C^0(D)$  with  $\|u\|_\infty \leq C(D)\|u\|_H^{\frac{1}{2}}$ , it follows in particular that

$$0 \leq u \leq c(D)r(\omega) \quad \text{for all} \quad u \in \mathcal{A}^+(\omega).$$

Using the above result we can deduce that there exist two random fixed points  $u_-(\omega)$  and  $u_+(\omega)$  such that

$$u_-(\omega) \leq u \leq u_+(\omega) \quad \text{for all} \quad u \in \mathcal{A}^+(\omega).$$

Clearly  $u_-(\omega) = 0$ , but since  $\mathcal{A}^+(\omega)$  is non-trivial this proves the existence of a new random (positive) fixed point  $a(\omega) = u_+(\omega)$ . By symmetry there is also a new random (negative) fixed point  $-a(\omega)$ . (For a similar argument see Chueshov (2000).)

That  $a(\omega) \rightarrow 0$  as  $\beta \downarrow \lambda_1$  follows from a slight variant of the general result on the upper-semicontinuity of random attractors to be found in Caraballo & Langa (2001, theorem 3), which guarantees under conditions which are easily verified in our case (convergence of the corresponding cocycles; convergence of the compact absorbing sets; and positive invariance of the compact absorbing set for the limit equation ( $\beta = \lambda_1$ )) that  $\mathbb{P}$ -a.s.

$$\lim_{\beta \downarrow \lambda_1} \text{dist}(\mathcal{A}_\beta^+(\omega), \mathcal{A}_{\lambda_1}(\omega)) = 0.$$

Since  $a(\omega) \subset \mathcal{A}^+(\omega)$  and  $\mathcal{A}_{\lambda_1}(\omega) = \{0\}$  we have  $a(\omega) \rightarrow 0$ . □

In fact, we know a little more than is stated in the theorem. For  $\beta > \lambda_1$  the attractor contains the two distinct, non-trivial, invariant subsets  $\mathcal{A}^+(\omega)$  and  $\mathcal{A}^-(\omega)$ , which attract all initial conditions in  $\mathcal{K}^+$  and  $\mathcal{K}^-$  respectively. One can think of this, loosely, as a transfer of stability of the origin to  $\mathcal{A}^+(\omega)$  and  $\mathcal{A}^-(\omega)$ .

For a more concrete stability result, theorem 2 in Arnold & Chueshov (1998) guarantees a limited type of stability for the new fixed points. In particular,  $a(\omega)$  is stable from above, in that if  $u(\omega) \geq a(\omega)$  (note that this “initial condition” cannot in general be taken to be deterministic) then

$$\lim_{t \rightarrow \infty} \varphi(t, \theta_{-t}\omega)u(\theta_{-t}\omega) = a(\omega)$$

(stability “from below” holds for  $-a(\omega)$ ). Since trajectories near zero in  $\mathcal{K}^+$  move away from the origin, it seems reasonable to expect that  $a(\omega)$  is attracting in  $\mathcal{K}^+$ . If one could show that solutions eventually enter  $\mathcal{K}^+$  or  $\mathcal{K}^-$  this would give the stability of the fixed points. (A general result guaranteeing the existence of stable fixed points is given by Schmalfuß (1996), but we have not been able to apply this in our case.)

## 6. Conclusion

We have shown that the structure of the random attractor for our example, the Chafee-Infante equation with a multiplicative noise term, is surprisingly close to that of the deterministic equation. In particular, we have shown that the dimension of the attractor, asymptotically in  $\beta$ , exhibits the same behaviour as in the deterministic case, and is independent of the level of noise ( $\sigma$ ).

Analysing further the structure of this attractor we have shown that as  $\beta$  passes through  $\lambda_1$  the attractor grows and two new random fixed points appear. Noting that  $a^\pm(\omega) \in \mathcal{A}^\pm(\omega)$ , what we expect is that in fact, for  $\lambda_1 < \beta < \lambda_2$ ,

$$\mathcal{A}(\omega) = \mathcal{A}^+(\omega) \cup \mathcal{A}^-(\omega),$$

with  $\mathcal{A}^\pm(\omega)$  consisting of a one-dimensional manifold (the image of  $\mathcal{M}_\delta(\omega) \cap \mathcal{K}^\pm$  under the flow) joining the origin to  $\pm a(\omega)$ . This would ensure that the attractor's structure is exactly that in the deterministic case.

Some support for this picture, at least for small  $\sigma$ , is given by the upper semi-continuity result in Caraballo *et al.* (1998). We proved there that as  $\sigma \rightarrow 0$ , for each  $\omega$ ,

$$\text{dist}(\mathcal{A}_\sigma(\omega), \mathcal{A}) \rightarrow 0.$$

Thus the random attractor, a set which is at least one-dimensional, must lie within a small neighbourhood of  $\mathcal{A}$ , which is itself homeomorphic to a line when  $\lambda_1 < \beta < \lambda_2$ .

Finally, we comment that we would not expect similar ideas to work in the case of an additive noise term (e.g.  $+\epsilon\phi dW_t$ , for some  $\phi \in H$ ). Without the fixed point  $\{0\}$  there is no clear way to show that an invariant manifold for a truncated version of the equation is a subset of the random attractor. Indeed, the one-dimensional example in Crauel & Flandoli (1998) suggests that with an additive noise the attractor will be very much simpler than in the deterministic case. We hope to investigate this further in a future paper.

Many thanks to Igor Chueshov, Hans Crauel, Franco Flandoli, and Antonio Suárez for some interesting and helpful conversations. James Robinson is currently a Royal Society University Research Fellow, and would like to thank the Society for their support. He would also like to thank EDAN for their hospitality, and Iberdrola for their financial assistance during his visit. Tomás Caraballo and José Langa have been partially supported by DGICYT Project PB98-1134.

## References

- Arnold, L. 1998 *Random dynamical systems*. New York: Springer.
- Arnold, L. 1998 Recent progress in stochastic bifurcation theory. Report no. 439, University of Bremen.
- Arnold, L. & Boxler, P. 1992 Stochastic bifurcation: instructive examples in dimension one. In *Diffusion processes and related problems in analysis, Vol II* (eds. Pinsky & Wihstutz), pp. 241–256. Stuttgart: Birkhäuser.
- Arnold, L. & Chueshov, I. 1998 Order-preserving random dynamical systems: Equilibria, attractors, applications. *Dyn. Stab. Sys.* **13**, 265–280.
- Babin, A. V. & Vishik, M. I. 1983 Attractors of partial differential evolution equations and estimates of their dimension. *Russian Math. Surveys* **38:4**, 151–213.
- Bensoussan, A. & Flandoli, F. 1995 Stochastic inertial manifolds. *Stoch. and Stoch. Reports* **53**, 13–39.
- Boxler, P. 1991 How to construct stochastic center manifolds on the level of vector fields. In *Lyapunov exponents*. Springer Lectures Notes in Math., no. 1486, pp. 141–158. Berlin: Springer.
- Caraballo, T. & Langa, J. A. 2001 On the upper semicontinuity of cocycle attractors for non-autonomous and random dynamical systems. *Dynamics of Continuous, Discrete and Impulsive Systems*, to appear.

- Caraballo, T., Langa, J. A. & Robinson, J. C. 1998 Upper semicontinuity of attractors for small random perturbations of dynamical systems. *Commun. Part. Diff. Eq.* **23**, 1557–1581.
- Caraballo, T., Langa, J. A. & Robinson, J. C. 2000 Stability and random attractors for a reaction-diffusion equation with multiplicative noise. *Discrete Cont. Dyn. Sys.* **6**, 875–892.
- Chueshov, I. D. & Girya, T. V. 1994 Inertial manifolds for stochastic dissipative dynamical systems. *Doklady of Acad. Sci. Ukraine* **7**, 42–45.
- Chueshov, I. D. 2000 Gevrey regularity of random attractors for stochastic reaction-diffusion equations. *Random Op. Stoch. Equ.* **8**, 143–162.
- Crauel, H. 1995 Random probability measures on polish spaces. Habilitationsschrift, University of Bremen.
- Crauel, H. 1999 Random point attractors versus random set attractors. *Proc. London Math. Soc.*, to appear.
- Crauel, H. & Flandoli, F. 1994 Attractors for random dynamical systems. *Prob. Th. Rel. Fields* **100**, 365–393.
- Crauel, H. & Flandoli, F. 1998 Additive noise destroys a pitchfork bifurcation. *J. Dyn. Diff. Eqn.* **10**, 259–274.
- Crauel, H., Debussche, A. & Flandoli, F. 1997 Random attractors. *J. Dyn. Diff. Equ* **9**, 307–341.
- Crauel, H., Imkeller, P. & Steinkamp, M. 1999, Bifurcations of one-dimensional stochastic differential equations. In *Stochastic dynamics* (ed. H. Crauel & V. M. Gundlach). New York: Springer, pp. 27–47.
- Da Prato, G. & Debussche, A. 1996 Construction of stochastic inertial manifolds using backward integration. *Stoch. and Stoch. Reports* **59**, 305–324.
- Debussche, A. 1998 Hausdorff dimension of a random invariant set. *J. Math. Pures Appl.* **77**, 967–988.
- Flandoli, F. & Schmalfuss, B. 1996 Random attractors for the stochastic 3-D Navier-Stokes equation with multiplicative white noise. *Stochastics and Stoch. Reports* **59**, 21–45.
- Hale, J. K. 1988 *Asymptotic behavior of dissipative dynamical systems*. Mathematical Surveys and Monographs, vol. 25. Providence, R.I.: American Mathematical Society.
- Kotelenez, P. 1992 Comparison methods for a class of function valued stochastic partial differential equations. *Prob. Th. Rel. Fields* **93**, 1–19.
- Marion, M. 1987 Attractors for reaction-diffusion equations; existence and estimate of their dimension. *Appl. Anal.* **25**, 101–147.
- Pardoux, E. 1975 *Équations aux Dérivées Partielles Stochastiques non Linéaires Monotones*. Thesis, Univ. Paris XI.
- Renardy, M. R. & Rogers, R. C. 1992 *An introduction to partial differential equations*. New York: Springer.
- Robinson, J. C. 2001 *Infinite-dimensional dynamical systems*. Cambridge University Press.
- Schmalfuß, B. 1992 Backward cocycles and attractors of stochastic differential equations. In *International Seminar on Applied Mathematics - Nonlinear Dynamics: Attractor Approximation and Global Behaviour* (ed. V. Reitmann, T. Riedrich & N. Koks), pp. 185–192. Technische Universität, Dresden.
- Schmalfuß, B. 1996 A random fixed point theory based on Lyapunov exponents. *Random Comput. Dynam.* **4**, 257–268.
- Temam, R. 1988 (second edition 1997) *Infinite dimensional dynamical systems in mechanics and physics*. New York: Springer.