

Agricultural Water Management
Manuscript Draft

Manuscript Number: AGWAT3330

Title: Seasonal changes of maximum daily shrinkage reference equations for irrigation scheduling in olive trees: influence of fruit load

Article Type: Research Paper

Keywords: LVDT; RDI; trunk diameter fluctuations.

Abstract: Maximum daily shrinkage (MDS) is the parameter of daily cycle of trunk diameter most widely suggest in irrigation scheduling of several fruit trees. However, as in other plant-measured approach, the irrigation decision may be difficult due to the influence of the environment in the values obtained. Reference equations of MDS have been established in order to avoid the effects of environmental conditions. Such equations are usually related with simple meteorological data, in order to estimate easily MDS values in full-irrigated conditions. This work studies the influence of the fruit load and the inter-annual variations in the reference equation of MDS in olive trees. These reference equations were calculated during 4 seasons in a full-irrigated orchard and the equations were validated with the data of a different season. The values of MDS were related with vapour pressure deficit (VPD) and temperature obtained near the experimental orchard. In addition, meteorological data were considered as mean daily or as midday values. The validation of the equations were made using the fits with all the meteorological data considered (midday and mean daily of VPD and temperature). In each meteorological data, in addition, two different fit, one according fruit load and other with the complete pool data were used. The equations fit were significantly different each season in all the meteorological data considered. Although, seasons with similar fruit load were more similar. In both meteorological data considered (VPD and temperature) the midday values improve the fit respect to mean daily values. The reference equations in which temperature was used obtained best fit that the ones calculated with VPD. No significant differences were found in the validation when equations according with fruit load or using the complete pool data were compared. The limitations and usefulness of these reference equations is also discussed.

1 1 **Seasonal changes of maximum daily shrinkage reference equations for irrigation**
2
3
4 2 **scheduling in olive trees: influence of fruit load**

5 3
6
7
8 4 **A. Moriana^a, F. Moreno^{b,*}, I.F. Girón^b, W. Conejero^c, M.F. Ortuño^c, M. Corell^a, A.**
9
10 5 **Torrecillas^{c,d}**

11
12
13
14 6
15
16 7 *^aEscuela Universitaria de Ingeniería Técnica Agrícola. University of Seville, Sevilla,*
17
18 8 *Spain*

19
20
21 9 *^bInstituto de Recursos Naturales y Agrobiología (CSIC), P.O. Box 1052, E-41080*
22
23 10 *Sevilla, Spain*

24
25
26 11 *^cDpto. Riego. Centro de Edafología y Biología Aplicada del Segura (CSIC). P.O. Box*
27
28 12 *164, E-30100 Espinardo (Murcia), Spain*

29
30
31 13 *^dUnidad Asociada al CSIC de Horticultura Sostenible en Zonas Áridas (UPCT-*
32
33 14 *CEBAS), Paseo Alfonso XIII s/n. E-30203 Cartagena (Murcia), Spain*

34
35
36 15 * Corresponding author: fmoreno@irnase.csic.es Phone: (+34) 954624711; Fax: (+34)
37
38 16 954624002


1 25 **Abstract**


2
3 26 Maximum daily shrinkage (MDS) is the parameter of daily cycle of trunk diameter most
4
5 27 widely suggest in irrigation scheduling of several fruit trees. However, as in other plant-
6
7 28 measured approach, the irrigation decision may be difficult due to the influence of the
8
9 29 environment in the values obtained. Reference equations of MDS have been established
10
11 30 in order to avoid the effects of environmental conditions. Such equations are usually
12
13 31 related with simple meteorological data, in order to estimate easily MDS values in full-
14
15 32 irrigated conditions. This work studies the influence of the fruit load and the inter-
16
17 33 annual variations in the reference equation of MDS in olive trees. These reference
18
19 34 equations were calculated during 4 seasons in a full-irrigated orchard and the equations
20
21 35 were validated with the data of a different season. The values of MDS were related with
22
23 36 vapour pressure deficit (VPD) and temperature obtained near the experimental orchard.
24
25 37 In addition, meteorological data were considered as mean daily or as midday values.
26
27 38 The validation of the equations were made using the fits with all the meteorological data
28
29 39 considered (midday and mean daily of VPD and temperature). In each meteorological
30
31 40 data, in addition, two different fit, one according fruit load and other with the complete
32
33 41 pool data were used. The equations fit were significantly different each season in all the
34
35 42 meteorological data considered. Although, seasons with similar fruit load were more
36
37 43 similar. In both meteorological data considered (VPD and temperature) the midday
38
39 44 values improve the fit respect to mean daily values. The reference equations in which
40
41 45 temperature was used obtained best fit that the ones calculated with VPD. No significant
42
43 46 differences were found in the validation when equations according with fruit load or
44
45 47 using the complete pool data were compared. The limitations and usefulness of these
46
47 48 reference equations is also discussed.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 49 **Keyword:** LVDT, RDI, trunk diameter fluctuations.
2
3

4 50
5


6 51 **1. Introduction**

7
8 52 Irrigated agriculture is actually the largest fresh water consumer in the world.  In the last
9
10 53 decades, olive production in the Mediterranean region has intensified, and the
11
12 54 traditional rainfed crop is now frequently irrigated (Eris and Barut, 1995). The scarcity
13
14 55 of water supplies and the increasing demand of other water-user sectors impose to the
15
16 56 Mediterranean agriculture an increasing pressure to limit its water consumption, and so
17
18 57 there is a constant need to improve the water use by the crops using better irrigation
19
20 58 management (Fereres and Evans, 2006). Among the tools that olive growers can use to
21
22 59 achieve this goal are more precise irrigation scheduling methods which involve the
23
24 60 determination of water requirements by crop and/or the application of regulated deficit
25
26 61 irrigation.
27
28
29
30
31

32
33 62 Measurement of the plant water condition may be useful for irrigation
34
35 63 scheduling because of its dynamic nature, which is directly related with climatic and
36
37 64 soil conditions, as well as crop productivity (Fereres and Goldhamer, 2003; Goldhamer
38
39 65 et al., 2003). 

40
41
42 66 The trunk or stem of all plants presents daily cycles of swelling and shrinking
43
44 67 that is known as trunk diameter variations (Kozlowski, 1967). Continuous records of
45
46 68 stem diameter have been proposed as a management tool for irrigation scheduling
47
48 69 (Huguet et al., 1992; Cabibel and Isberie, 1997; Cohen et al., 2001; Goldhamer and
49
50 70 Fereres, 2001). In a recent paper Ortuño et al. (2010) have reviewed the state of the art
51
52 71 regarding the use of trunk diameter variations derived parameters for irrigation
53
54 72 scheduling in woody crops. As so far as we know, Goldhamer and Fereres (2004) were
55
56
57
58
59
60
61
62
63
64
65

1 73 the first to demonstrate that is possible to develop a deficit irrigation schedule based
2
3 74 only on maximum daily trunk shrinkage (MDS) in almond trees. García Orellana et al.
4
5 75 (2007), Velez et al. (2007) and Ortuño et al. (2009c) confirmed that in citrus MDS is a
6
7 76 good indicator for scheduling deficit irrigation. Other useful parameter derived from the
8
9 77 trunk daily cycles of swelling and shrinking is the trunk growth rate (TGR) as defined
10
11 78 by Goldhamer and Fereres (2001) that can be used for irrigation scheduling of fruit
12
13 79 trees.
14
15
16
17

18 80 The use of the absolute values of the plant-based water status indicators could be
19
20 81 meaningless and thus we need to obtain reference values for these indicators. Reference
21
22 82 values can be obtained by maintaining trees under conditions of non-limiting soil water
23
24 83 supply. At the same time is necessary to develop reference equations to help us to
25
26 84 interpret the values of a plant-based water status indicator. These reference equations
27
28 85 can be obtained by relating their values in trees under non-limiting soil water conditions
29
30 86 with evaporative demand of the atmosphere (Moreno et al., 2006; Conejero et al.,
31
32 87 2007b; Ortuño et al., 2009b and 2010).
33
34
35
36
37

38 88 MDS values can be affected by several factors, such as tree age (Morianana and
39
40 89 Fereres, 2004), phenological period (Marsal et al., 2002; Intrigliolo and Castel, 2004;
41
42 90 Moriana and Fereres, 2004; Conejero et al., 2007b) and fruit load (Conejero et al., 2010;
43
44 91 Marsal et al., 2002; Intrigliolo and Castel, 2006). In olive trees the alternate bearing can
45
46 92 be a factor that can affect MDS values. In a recent paper by Moriana et al. (2010) have
47
48 93 shown that MDS is no the best indicator for optimal irrigation scheduling in olive trees
49
50 94 but can be a good tool to be used in deficit irrigation scheduling. In this case, the stress
51
52 95 level will be indicated by MDS values lower than the one obtained in the base lines or
53
54 96 reference equations.
55
56
57
58
59
60
61
62
63
64
65

1 97 The objectives of this paper were: (1) to obtain reference equations of MDS for
2
3 98 olive trees based on its relation with the evaporative demand of the atmosphere; (2) to
4
5 99 study the interannual variation of the reference equations, and (3) to evaluate the
6
7
8 100 influence of fruit load on the MDS vs evaporative demand parameters relationships.
9

10
11 101

12 102 **2. Material and Methods**


13 103 2.1. Description and design of the experiment


14
15 104 Experiments were conducted at La Hampa, the experimental farm of the Instituto de
16
17
18 105 Recursos Naturales y Agrobiología (CSIC), which is located at Coria del Río near
19
20
21 106 Seville (Spain) (37°17'N, 6°3'W, 30m altitude) during 5 consecutive seasons (from
22
23
24 107 2005 to 2009). The sandy loam soil (about 2 m deep) of the experimental site was
25
26
27 108 characterized by a volumetric water content of 0.33 m³ m⁻³ at saturation, 0.21 m³ m⁻³ at
28
29
30 109 field capacity and 0.1 m³ m⁻³ at permanent wilting point, and 1.30 (0-10cm) and 1.50
31
32
33 110 (10-120 cm) g cm⁻³ bulk density.

34
35 111 The experiment was performed on 37-year-old olive trees (*Olea europaea* L cv
36
37 112 Manzanillo). Tree spacing followed a 7m x 5m square pattern. Pest control and
38
39
40 113 fertilization practices were those commonly used by the growers and no weeds were
41
42
43 114 allowed to develop within the orchard.

44
45 115 Irrigation was carried out during the night by drip using one lateral pipe per tree
46
47 116 row and five emitters per plant, delivering 3 L h⁻¹ each. Plants irrigation requirements
48
49
50 117 were determined according to daily reference evapotranspiration (ET_o) and a crop factor
51
52
53 118 based on the time of the year and the percent of ground area shaded by the tree canopy
54
55 119 (Fernández et al., 1998). During the experimental period (from end of april until
56
57
58
59
60
61
62
63
64
65


1 120 beginning of October), total crop evapotranspiration (ET_c) was 430 mm (2005), 413
2
3 121 mm (2006), 414 mm (2007), 430 mm (2008), 392 mm (2009).
4

5
6 122 During the experimental period, olive trees were irrigated daily above their
7
8 123 water requirements in order to obtain non-limiting soil water conditions. A total amount
9
10 124 of water (rainfall not included) of 476 mm (2005), 442 mm (2006), 410 mm (2007),
11
12 125 644mm (2008), 605mm (2009), measured with in-line water meters, was applied during
13
14 126 the experiment. 

15
16
17 127 The design of the experiment was completely randomized  with four replications,
18
19 128 each replication consisting of the three adjacent rows of five trees. Measurements were
20
21 129 made in the inner tree of the central row of each replicate, the other trees served as
22
23 130 borders.
24

25 131 2.2 Measurements

26
27
28 132 Micrometeorological 30 min data, namely air temperature, solar radiation, air relative
29
30 133 humidity and wind speed at 2 m above the soil surface were collected by an automatic
31
32 134 weather station located some 40 m from the experimental site. Daily reference
33
34 135 evapotranspiration (ET_o) was calculated using the Penman-Monteith equation (Allen et
35
36 136 al., 1998). Mean daily vapour pressure deficit (VPD_m) was calculated from the mean
37
38 137 daily vapour pressure and relative humidity (Goldhamer and Fereres, 2001).
39

40
41
42 138 Trunk diameter fluctuations were measured throughout the experimental periods
43
44 139 in four trees,  using a set of linear variable displacement transducer (LVDT) (model
45
46 140 DF±2.5 mm, accuracy ±10 µm, Solartron Metrology, Bognor Reiss, UK) attached to the
47
48 141 trunk, with a special bracket made of Invar, an alloy of Ni and Fe with a thermal
49
50 142 expansion coefficient close to zero (Katerji et al., 1994). Measurements were taken
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 143 every 10 s and the datalogger (model CR10X with AM 416 multiplexer, Campbell Sci.
2
3 144 Ltd., Logan, USA) was programmed to report 30 min means. Maximum

4
5
6 145 The data obtained during the five seasons were analyzed taking into account the
7
8 146 years with low fruit load (2005, 2007 and 2009) and years with full fruit load (2006 and
9
10 147 2008). Data from 2009 were used to validate the relationships obtained in previous
11
12 148 years. Linear regression analysis was carried out to explore relationships between
13
14 149 variables (MDS and climatic variables). Differences between regression lines were
15
16 150 determined with a T-test of the slope and y-intercept.
17
18
19
20
21 151

22 152 **3. Results**

23
24
25 153 The MDS vs mean daily temperature relationship during the four years of the
26
27 154 experiment showed the best fit in a lineal form (Table 1 and Fig. 1a). The increase in
28
29 155 temperature produces an increase in the MDS in a rate around $0.04 \text{ mm } ^\circ\text{C}^{-1}$. The range
30
31 156 of variations in mean daily temperature was wide enough for the Seville conditions of
32
33 157 olive growth and varied from around 10 to 30°C . The equations for each year for mean
34
35 158 daily temperature are shown in Table 1. All the equations were significantly different in
36
37 159 the slope and the intercept. The coefficient of determination was significant in all the
38
39 160 years but low, except in the 2008 season when it was clearly higher ($r^2=0.82$). When the
40
41 161 data were grouped in full fruit load (FFL) and low fruit load (LFL) years there were no
42
43 162 significant differences in the slope but it was in the intercept. The LFL equations tended
44
45 163 to lower values of MDS than the FFL equations when the same mean daily temperature
46
47 164 is considered.

48
49
50
51
52 165 When the temperature considered is the ones that occurred at midday the scatter
53
54 166 of the points is reduced (Table 1 and Fig. 1b) in comparison with that of mean daily



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

167 temperature (Fig. 1a). The range of variations in temperature (Fig. 1b) is similar to that
168 in Fig. 1a, and changes from around 20 to 40°C. The equations in each year were
169 significantly different between them, as in the case of mean daily temperature. The
170 coefficients of determination were slightly higher than the ones of mean daily
171 temperature (Table 1). The equations of LFL and FFL years were significantly different
172 for the intercept but not for the slope. As in the data of Fig. 1a, the values during LFL
173 year tended to be lower than the ones of the FFL year when the same range of
174 temperature is considered.

175 The relationship between MDS and VPD was also lineal. The increase in VPD
176 produces an increase in the MDS in full irrigated conditions (Fig. 2). When the mean
177 daily VPD is considered the range of data were from near 0 to 4 KPa (Fig. 2a). The
178 equations of each year were significantly different in the intercept and the scatter was
179 slightly higher than in the midday temperature relationship (Table 1 and Figs. 1b and
180 2a). There were also significant differences between the equations when they were
181 grouped in FFL and LFL. The MDS in FFL year tended to higher values than in LFL
182 year when the same mean daily VPD is considered.

183 The scatter in the MDS vs VPD relationship is slightly reduced when the values
184 at midday (Figure 2b) are considered instead of the daily average (Figure 2a). Although
185 the coefficient of determination was slightly higher than the mean daily VPD, they were
186 lower than the ones obtained in midday temperature relationships (Table 1). The range
187 of variations of midday VPD was also higher than mean daily VPD and it extended until
188 6 KPa (Fig. 2). There were significantly differences in the intercept but not in the slope
189 of the equations between years. There were also significant differences between the
190 equations when they were grouped in full fruit load and low fruit load years (Table 1).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


191 The MDS in FFL year tended to higher values than in LFL year when the same midday
192 VPD is considered.


193 The equations obtained with the data of 2005 to 2008 seasons were validated
194 with the data of 2009 season (Figs. 3 and 4; Table 2). Although all the seasons were
195 statistically different in both meteorological parameters (temperature and VPD)
196 considered (Table 1), from the point of view of irrigation scheduling in a commercial
197 orchard the variation between seasons was considered similar. Only the influence of fruit
198 load was evaluated. However, even though, alternative bearing may be common in field
199 conditions, in commercial orchards is difficult to identify most of the seasons as low
200 fruit load or as full fruit load year. Therefore, the validation was made with two
201 equations, one of them related to the fruit load and the other with the one that
202 considered all the seasons, which so called from here “total” equation (Table 1). In 2009
203 season, the orchard had very low yield (around 4 kg per tree), therefore for each
204 variable (midday and mean daily temperature and midday and mean daily VPD) the
205 validations were made with the low fruit load year equations (Table 1). The fit of the
206 observed and estimated MDS when the temperature is considered (Fig. 3) was
207 significantly different from line 1:1 in all the cases (Table 2). The midday temperature,
208 however, tended to nearer values to the 1:1 line than the mean daily temperature (slope
209 0.80 and 0.73 respectively, Table 2). The data of the fits with mean daily temperature
210 showed higher scatter (higher MSE, lower r^2) than the midday temperature (Fig. 3 and
211 Table 2). However, there were no significant differences between the equations of Table
212 2. When the same kind of temperature is considered the low fruit load equations were
213 nearer to 1:1 line than the “total” equations. Nevertheless, in all the cases the fit

1 214 obtained with LFL or “total” data were not significantly different in slope but it was in
2
3 215 intercept (always lower in LFL equations).
4

5
6 216 The validation of the VPD equations (Fig. 4) showed that the prediction were
7
8 217 poorer than the ones obtained with any of the temperatures (Fig. 3). The parameters of
9
10 218 the relationship MDS observed vs measured were significantly different from the line
11
12 219 1:1 and significantly lower than the ones obtained with temperature, especially the slope
13
14 220 that were around 0.5 while in temperature were around 0.8 (Table 2). There were no
15
16 221 significant differences between the slope of the LFL and “total” equations but it was in
17
18 222 the intercepts. The LFL equations tended to intercept nearer to zero than the “total”
19
20 223 although in all the cases were higher than the ones obtained with the temperature.
21
22
23
24
25
26
27

28 225 **4. Discussion**

29
30 226 MDS is considered a good indicator of the transpiration stream (Herzog et al 1995) but
31
32 227 the relationship with VPD was poorer than the ones obtained with temperature (Tables 1
33
34 228 and 2). Similar results have been reported in several works in different fruit trees
35
36
37 229 mond, Fereres and Goldhamer (2003); plum (Intrigliolo and Castel, 2006); olive
38
39 230 (Moreno et al, 2006); lemon (Ortuño et al, 2009)). In addition, the relationship along the
40
41 231 season was steady and lineal and apparently, there was no influence of the phenological
42
43 232 stage of trees as in other fruit trees (plum, Intrigliolo and Castel, 2007). The midday
44
45 233 parameters presented a better fit than the daily average (Tables 1 and 2). MDS is a
46
47 234 parameter that is calculated during the most active transpiration phase and the “mean
48
49 235 VPD or mean temperature” included values for the complete day where there are
50
51
52 236 periods in which transpiration even is null. “Midday parameters”, however, are likely
53
54
55 237 more related with the phase of shrinkage because the higher rate of shrinkage occurred
56
57
58
59
60
61
62
63
64
65

1 238 around this moment of the day. All the equations were significantly different each
2
3 239 season, though the ones with the similar crop load tended to be nearer. Such differences
4
5 240 between seasons may indicate that the MDS is an accurate measurement that is likely
6
7 241 affected in several ways for the physiolo of the plant. Genard et al (2001) suggested
8
9 242 that the trunk diameter varied according to several factors such as xylem, osmotic and
10
11 243 turgor water potential and for the elasticity of the wall. Therefore, in theory, is difficult
12
13 244 that the same relationship between MDS and temperature may be obtained each year
14
15 245 even in the same orchard.

16
17
18 246 MDS has been traditionally considered the best indicator of trunk diameter
19
20 247 variations for irrigation scheduling in most of the fruit trees (Huguet et al, 1992;
21
22 248 Goldhamer and Fereres, 2001; Ortuño et al 2010). However, in olive trees, this indicator
23
24 249 presented several limitations for using in full irrigated conditions. There are several
25
26 250 works in olive trees that presented no variations in MDS in conditions of mild water
27
28 251 stress (Moriana et al 2003; Moriana and Fereres, 2002), only in conditions of very
29
30 252 severe water stress MDS is reduced (Moriana et al 2000; Moriana et al 2003). Such
31
32 253 response has been suggested that is related with the physiology of the specie (Moriana
33
34 254 et al, 2010). On one hand, MDS increase in full irrigated conditions quickly due to the
35
36 255 evaporative demand, while the ones of the stressed trees increase slower. Therefore,
37
38 256 conditions of mild water stress produced clear differences in water potential meanwhile
39
40 257 similar values in MDS (Moriana et al. 2010). Nevertheless, the deficit irrigation
41
42 258 strategies in olive trees suggest a moderate or even severe water stress conditions during
43
44 259 the pit hardening (Goldhamer, 1999; Moriana et al 2003; Tognetti et al 2007). In these
45
46 260 conditions reference values of MDS may be probably very useful for controlling the
47
48 261 level of water stress but using in the opposite way that in the rest of fruit trees. During
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 262 the pit hardening the reduction of MDS from reference values will indicate moderate or
2
3 263 severe water stress conditions. Several questions arise then. The first, how much MDS
4
5 264 may be reduced should be answered in further experimental works. The others are about
6
7
8 265 which reference equation may be used. According to the results of this work (Tables 1
9
10 266 and 2) in commercial orchard the differences between the crop load and the equation
11
12
13 267 that included all the data (“total equation”) is small. The validation of both VPD and
14
15 268 temperature equations (Table 2) suggest that the estimation is very close, even though
16
17
18 269 the 2009 season was a clear low fruit load year (the yield was almost null). Therefore, in
19
20 270 commercial conditions when commonly low fruit load and full fruit load years are
21
22 271 difficult to identify the “total” midday temperature will be the best selection. On the
23
24
25 272 other hand, there is no data about the feasibility of this equation out of the experimental
26
27
28 273 farm even though the same cultivar would be used. Moriana and Fereres (2004)
29
30 274 suggested different baselines in cv Picual using mean VPD, with different age and
31
32 275 density but similar conditions to the present work (this experimental farm is around 150
33
34
35 276 Km far from the plot of this work and with very similar climatic conditions). The one-
36
37 277 year equations presented by these authors (Moriana and Fereres, 2004), were similar in
38
39
40 278 slope to the ones obtained in the present work (Table 1) in full fruit load and low fruit
41
42 279 load years in mature trees. According with the results of the present work, VPD
43
44 280 estimation would be worse than temperature estimation. The baselines of midday
45
46
47 281 temperature obtained in the present work (Table 1) may be a good tool for irrigation
48
49 282 scheduling of olive trees, at least from the point of view of commercial management, if
50
51
52 283 the orchard is under similar climatic conditions to that of our experimental farm.
53
54
55 284

56
57 285 **5. Conclusions**
58
59
60
61
62
63
64
65

1 286 MDS was related with the VDP and temperature, although the fits calculated with
2
3 287 temperature were better than the ones obtained with VPD. The best fits were obtained
4
5 288 with values measured at midday instead of the mean daily. This better agreement is
6
7
8 289 likely related with the period when the shrinkage is produced. The equations obtained
9
10 290 were different each season, though the season with similar fruit load presented similar
11
12 291 equations. The MDS values of full fruit load (FFL) seasons tended to be higher than the
13
14 292 low fruit load (LFL) seasons. However, when the equations were validated with an
15
16 293 additional low fruit load season, there were no significant differences between equations
17
18 294 that considered LFL data or the one that considered the completed pool of data. The
19
20 295 parameters of other MDS reference equations found in the literature were similar in
21
22 296 mature trees when the same fruit load was considered. Therefore, though cultivar or
23
24 297 density may be factors that affect the reference equations, fruit load and age of the tree
25
26 298 are probably the most important. The reference equations of midday temperature
27
28 299 obtained in the present work (Table 1) may be a good tool for irrigation scheduling of
29
30 300 olive trees, at least from the point of view of commercial management, if the orchard is
31
32 301 under similar climatic conditions to that of our experimental farm.
33
34
35
36
37
38
39
40
41

42 302

43 303 **Acknowledgements**

44 304 This research was supported by the Spanish Ministerio de Ciencia e Innovación
45
46 305 (MICINN), (CICYT/FEDER AGL2004-0794-C03-02 and AGL2007-66279-C03-
47
48 306 02/AGR). Thanks are due to J. Rodriguez for help with field measurements.
49
50
51

52 307

53 308 **References**

54

55

56

57

58

59

1 309 Cabibel, B. and Isberie, C. 1997. Flux de sieve et alimentation hydrique de cerisiers
2
3 310 irrigués ou non en localisation. *Agronomie* 17, 97-112.
4
5 311 Cohen, M., Goldhamer, D., Fereres, E., Girona, J., Mata, M., 2001. Assessment of
6
7 312 peach tree responses to irrigation water deficits by continuous monitoring of
8
9 313 trunk diameter changes. *J. Hort. Sci. Biotech.* 76, 55–60.
10
11 314 Conejero, W., Alarcón, J.J., García-Orellana, Y., Nicolás, E., Torrecilla, A. 2007.
12
13 315 Evaluation of sap flow and trunk diameter sensors using for irrigation
14
15 316 scheduling in early maturing peach trees. *Tree Physiol.* 27, 1753-1759.
16
17 317 Eris, A. and Barut, E. 1994. Olive growing in Turkey. *Chron. Hortic.* 35, 14-16.
18
19 318 Fereres, E., Goldhamer, D.A., 2003. Suitability of stem diameter variations and water
20
21 319 potential as indicators for irrigation scheduling of almond trees. *J. Hort. Sci.*
22
23 320 *Biotech.* 78, 139–144.
24
25 321 Fereres, E. and Evans, R.G. 2006. Irrigation of fruit trees and vines: an introduction.
26
27 322 *Irrig. Sci.* 24, 55-57.
28
29 323 Fernández, J.E., Díaz, A., Martín-Palomo, M.J., Moreno, F., Girñon, I. 1998. Riego y
30
31 324 fertilización del olivar en la comarca del Aljarafe. Instituto de Recursos Naturales
32
33 325 y Agrobiología de Sevilla (CSIC) y Comisión Interministerial de Ciencia y
34
35 326 Tecnología (CICYT), Sevilla, 32pp.
36
37 327 García-Orellana, Y., Ruiz-Sánchez, M.C., Alarcón, J.J., Conejero, W., Ortuño, M.F.,
38
39 328 Nicolás, E., Torrecillas, A., 2007. Preliminary assessment of the feasibility of
40
41 329 using maximum daily trunk shrinkage for irrigation scheduling in lemon trees.
42
43 330 *Agric. Water Manage.* 89, 167–171.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


1 331 Genard, M., Fishman, S., Vercambre, G., Huguet, J.G., Bussi, C., Besset, J., Habib, R.,
2
3 332 2001. A biophysical analysis of stem and root diameter variations in woody
4
5
6 333 plants. *Plant Physiol.* 126, 188-202.
7
8 334 Goldhamer, D.A., Fereres, E. 2001. Irrigation scheduling protocols using continuously
9
10 335 recorded trunk diameter measurements. *Irrig. Sci.* 20, 115-125.
11
12 336 Goldhamer, D.A., Fereres, E. 2004. Irrigation scheduling of almond trees with trunk
13
14 337 diameter sensors. *Irrig. Sci.* 23, 11-19.
15
16 338 Goldhamer, D.A., Fereres, E., Salinas, M. 2003. Can almond trees directly dictate their
17
18 339 irrigation needs? *Calif. Agric.* 57, 138-144.
19
20
21 340 Herzog, K.M., Häsler, R., Thum, R., 1995. Diurnal changes in the radius of a subalpine
22
23 341 Norway spruce stem: their relation to the sap flow and their use to estimate
24
25 342 transpiration. *Trees* 10, 94-101.
26
27
28 343 Huguet, J.G., Li, Sh., Lorendeau, J. Y., Pelloux, G., 1992. Specific micromorphometric
29
30 344 reactions of fruit trees to water stress and irrigation scheduling automation. *J.*
31
32 345 *Hort. Sci. Bio.* 67, 631-640
33
34
35 346 Intrigliolo, D.S. and Castel, J.R., 2004. Continuous measurement of plant and soil water
36
37 347 status for irrigation scheduling in plum. *Irrig. Sci.* 23, 93–102.
38
39
40 348 Intrigliolo, D.S. and Castel, J.R., 2006. Usefulness of diurnal trunk shrinkage as a water
41
42 349 stress indicator in plum trees. *Tree Physiol.* 26, 303–311.
43
44
45 350 Intrigliolo, D.S. and Castel, J.R., 2007. Crop load affects maximum daily trunk
46
47 351 shrinkage of plum trees. *Tree Physiol.* 27, 89–96.
48
49
50 352 Katerji, N., Tardieu, F., Bethenod, O., Quetin, P., 1994. Behavior of Maize stem
51
52 353 diameter during drying cycles: comparison of two methods for detecting water
53
54 354 stress. *Crop Sci.* 34, 165-169
55
56
57
58
59
60
61
62
63
64
65


1 355 Kozłowski, T.T. 67. Diurnal variations in stem diameters of small trees. Bot. Gaz.
2
3 356 123, 60-68.
4
5 357 Marsal, J., Gelly, M., Mata, M., Arbone´s, J., Rufat, J., Girona, J., 2002. Phenology and
6
7 358 drought affects the relationship between daily trunk shrinkage and midday stem
8
9 359 water potential of peach trees. J. Hort. Sci. Biotechnol. 77, 411–417.
10
11 360 Moreno, F., Conejero, W., Martín-Palomo, M.J., Girón, I.F., Torrecillas, A., 2006.
12
13 361 Maximum daily trunk shrinkage reference values for irrigation scheduling in
14
15 362 olive trees. Agric. Water Manage. 84, 290–294.
16
17 363 Moriana, A. and Fereres, E., 2002. Plant indicators for scheduling irrigation for young
18
19 364 olive trees. Irrig. Sci. 21:83-90.
20
21 365 Moriana, A. and Fereres, E., 2004. Establishing references values of trunk diameter
22
23 366 fluctuations and stem water potential for irrigation scheduling of olive trees.
24
25 367 Acta Hort. 664:407-412.
26
27 368 Moriana, A., Orgaz, F., Fereres, E., Pastor, M., 2003. Yield responses of a mature olive
28
29 369 orchard to water deficits. J. Amer. Soc. Hort. Sci. 128:425-431
30
31 370 Moriana, A., Girón, I.J., Martín-Palomo, M.J., Conejero, W., Ortuño, M.F., Torrecilla,
32
33 371 A., Moreno, I. 2010. New approach for olive trees irrigation schedulings using
34
35 372 trunk diameter sensors. Agric. Water Manage. 97, 1822-1828.
36
37 373 Ortuño, M.F., Brito, J.J., Conejero, W., García-Orellana, Y, Torrecillas, A. 2009a.
38
39 374 Using continuously recorded trunk diameter fluctuations for estimating water
40
41 375 requirements of lemon trees. Irrig. Sci. 27, 271-276.
42
43 376 Ortuño, M.F., Brito, J.J., Conejero, W., García-Orellana, Y, Torrecillas, A. 2009b.
44
45 377 Maximum daily trunk shrinkage and stem water potencial baselines for irrigation
46
47 378 scheduling of lemon trees. Irrig. Sci. 27, 121-127.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



379 Ortuño, M.F., García-Orellana, Y., Conejero, W., Pérez-Sarmiento, F., Torrecillas, A.
380 2009c. Assessment of maximum daily trunk shrinkage signal intensity threshold
381 values for deficit irrigation in lemon trees. *Agric. Water Manage.* 96, 80-86.

382 Ortuño, M.F., Conejero, W., Moreno, F., Moriana, A., Intrigliolo, D.S., Biel, C.,
383 Mellisho, C.D., Pérez-Pastor, A., Domingo, R., Ruíz-Sánchez, M.C., Casadesus,
384 J., Bonany, J., Torrecillas, A. 10. Could trunk diameter sensors be used in
385 woody crops for irrigation scheduling? A review of current knowledge and
386 future perspective. *Agric. Water Manage.* 97, 1-11.

387 Velez, J.E., Intrigliolo, D.S., Castel, J.R. 2007. Scheduling deficit irrigation of citrus
388 trees with maximum daily trunk shrinkage. *Agric. Water Manage.* 90, 197-204.

389
390
391
392
393
394
395
396
397
398
399
400
401
402

1 403 **Figure Captions**

2
3 404 Figure 1. Relationship of MDS with the mean daily temperature (a) and the midday
4 temperature (b) during four consecutive seasons (2005 to 2008). ■ 2005; ● 2006; □
5 405 2007; ○ 2008. The regression equations obtained with each season, the “FFL (full fruit
6 406 load)” and “LFL (low fruit load)” season and the total pool of data is presented in Table
7
8 407 1.
9
10
11
12
13 408 1.

14
15 409
16
17
18 410 Figure 2. Relationship of MDS with the mean daily VPD (a) and the midday VPD (b)
19 during four consecutive seasons (■ 2005; ● 2006; □ 2007; ○ 2008). Line represent the
20 411 fit of all the data. The regression equations obtained with each season, the “FFL (Full
21 412 fruit load)” and “LFL (Low fruit load)” season and the total pool of data is presented in
22
23 413 Table 1.
24
25
26
27
28 414 Table 1.

29
30 415
31
32
33 416 Figure 3. Validation of the reference equations with the measured data of MDS in full
34 irrigated trees during 2009 season. The equations used are the ones obtained with the
35 417 mean daily temperature (a) and the midday temperature (b). White circle are the
36
37 418 equation obtained with the LFL (low fruit load) years and black circle are the equation
38
39 419 using the pool data, total equation (see Table 1 for equations). In all the cases the
40 420 relationship between MDS measured and estimated are significantly different from the
41
42 421 line 1:1.
43
44
45
46
47
48
49
50
51

52 424 Figure 4. Validation of the references equations with the measured data of MDS in full
53 irrigated trees during 2009 season. The equations used are the ones obtained with the
54 425 mean daily VPD (a) and the midday VPD (b). White circle are the equations obtained
55
56
57 426
58
59
60
61
62
63
64
65

1 427 with the LFL (Low fruit load) years and black circle are the equations with the pool
2
3 428 data, “total equation (Table 1). In all the cases the relationship between MDS measured
4
5
6 429 and estimated are significantly different from the line 1:1.
7
8 430
9
10 431
11
12 432
13
14 433
15
16 434
17
18 435
19
20 436
21
22 437
23
24 438
25
26 439
27
28 440
29
30 441
31
32 442
33
34 443
35
36 444
37
38 445
39
40 446
41
42 447
43
44 448
45
46 449
47
48 450
49
50 451
51
52 452
53
54 453
55
56 454
57
58 455
59
60 456
61
62 457
63
64 458
65

451 Table 1. Equations, coefficient of determinations (r^2) obtained in the relationships of
 452 Figs. 1 and 2. Each season is presented and, in addition, the results when they are
 453 grouped according to the crop load (LFL, low fruit load; FFL, full fruit load). “Total” is
 454 the equation considering all data (four seasons). RMSE: residual mean squared error. N:
 455 number of data. Statistic Dif: statistical differences between equations
 456

Season	Equations	r^2	RMSE	N	Statistic Dif.
MDS vs Mean Temperature					
2005	-0.79+0.053X	0.68***	0.09	105	All of them
2006	-0.26+0.035X	0.48***	0.12	111	Statistical
2007	-0.52+0.038X	0.65***	0.09	160	Different
2008	-0.59+0.046X	0.82***	0.08	175	
LFL (05&07)	-0.67+0.046X	0.68**	0.10	265	Intercept different
FFL (06&08)	-0.55+0.045X	0.73***	0.10	286	
Total	-0.59+0.045X	0.67***	0.11	551	
MDS vs Midday Temperature					
2005	-0.78+0.042X	0.79***	0.07	105	All of them
2006	-0.37+0.033X	0.67***	0.09	111	Statistical
2007	-0.57+0.034X	0.73***	0.08	160	Different
2008	-0.61+0.038X	0.85***	0.07	175	
LFL (05&07)	-0.65+0.037X	0.80***	0.08	265	Intercept different
FFL (06&08)	-0.58+0.038X	0.78***	0.09	286	
Total	-0.58+0.037X	0.73***	0.10	551	
MDS vs Mean VPD					
2005	0.03+0.16X	0.69***	0.08	105	
2006	0.34+0.16X	0.57***	0.10	111	Intercept
2007	0.09+0.20X	0.63***	0.10	163	different
2008	0.08+0.24X	0.78***	0.09	167	
LFL (05&07)	0.17+0.13X	0.67***	0.10	268	All of them
FFL (06&08)	0.16+0.22X	0.64***	0.12	278	Different
Total	0.23+0.14X	0.48***	0.14	546	
MDS vs Midday VPD					
2005	-0.07+0.13X	0.82***	0.06	105	Intercept
2006	0.30+0.12X	0.68***	0.09	111	different
2007	0.06+0.13X	0.67***	0.09	160	
2008	0.11+0.14X	0.51***	0.13	167	
LFL (05&07)	0.13+0.10X	0.66***	0.10	265	All of them
FFL (06&08)	0.16+0.14X	0.53***	0.13	278	different
Total	0.20+0.10X	0.45***	0.14	543	

457

1 458
 2 459
 3 460 Table 2. Best fits of the relationship between MDS observed and estimated using
 4
 5
 6 461 different meteorological variables. The adjusted validated were obtained from 2005 to
 7
 8 462 2008 (Table 1), while the data used to compared such validations were measured during
 9
 10 463 2009 season (n=148). LFL, low fruit load equation. “Total” is the equation considering
 11
 12
 13 464 all data (four seasons). RMSE: residual mean squared error
 14
 15
 16 465
 17 466

Eq. validated	Equations	RMSE	r²
LFL Mean Temperature	Y= 0.11+0.74X	0.10	0.66***
“Total” Mean Temperature	Y=0.17+0.73X	0.09	0.66***
LFL Midday Temperature	Y=0.09+0.81X	0.06	0.85***
“Total” Midday Temperature	Y=0.15+0.80X	0.06	0.85***
LFL Mean VPD	Y=0.20+0.44X	0.05	0.75***
“Total” Mean VPD	Y=0.26+0.48X	0.05	0.75***
LFL Midday VPD	Y=0.17+0.54X	0.05	0.81***
“Total” Midday VPD	Y=0.24+0.54X	0.05	0.81***

35 467
 36 468
 37 469
 38 470
 39 471
 40 472
 41 473
 42 474
 43 475
 44 476
 45 477
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

Figure 1





Figure 2





Figure 3





Figure 4

