

Intermolecular Displacement of S-Bound L-Methionine on Platinum(II) by Guanosine 5'-Monophosphate: Implications for the Mechanism of Action of Anticancer Drugs

Kevin J. Barnham, Milos I. Djuran, Piedad del Socorro Murdoch and Peter J. Sadler*

Department of Chemistry, Birkbeck College, University of London, Gordon House and Christopher Ingold Laboratories, 29 Gordon Square, London, UK WC1H 0PP

NMR investigations of the kinetics and thermodynamics of the competitive binding of L-methionine (Met), L-histidine (His), and 5'-monophosphates of guanosine (5'-GMP), adenosine (5'-AMP), thymidine (5'-TMP) and cytidine (5'-CMP) to $[\text{Pt}(\text{dien})\text{Cl}]^+$ (dien = 1,5-diamino-3-azapentane) in aqueous solution show that 5'-GMP selectively displaces S-bound Met, a finding which has implications for DNA platination by anticancer drugs *in vivo*.

DNA platination is thought to be a key event in the mechanism of action of platinum anticancer drugs, and there is much current interest in the mechanism of this reaction, especially in the formation of Pt-G (guanine) adducts.¹ The amino acid L-methionine (Met) is a thioether which plays an important role in the metabolism of all cells. Platinum(II) has a very high affinity for sulfur ligands and the bis-chelate- $[\text{Pt}(\text{Met} - \text{H-S}, \text{N})_2]$ has been isolated from the urine of patients treated with the anticancer drug cisplatin (*cis*- $[\text{PtCl}_2(\text{NH}_3)_2]$).² This complex, which is a mixture of diastereoisomers of the *cis* and *trans* isomers,³ appears to be a stable end-product of cisplatin metabolism and is unreactive towards nucleobases at neutral pH.⁴ However we found⁵ that cisplatin reacts with guanosine 5'-monophosphate (5'-GMP) even in the presence of Met, and in order to understand the course of the latter reaction, we investigated similar reactions

Table 1 Kinetic data for reactions of $[\text{Pt}(\text{dien})\text{Cl}]^+$ and $[\text{Pt}(\text{dien})(\text{Met-S})]^{2+}$ with 5'-GMP, where k_2 is the second-order rate constant

Complex	Reactant	pH*, T/K	$10^4 k_2/\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$
$[\text{Pt}(\text{dien})\text{Cl}]^+$ 1	Met ^a	3.9, 300	140
	GSMc ^b	5.0, 295	330
	5'-GMP ^b	5.0, 295	0.62 ^c
$[\text{Pt}(\text{dien})(\text{Met-S})]^{2+}$ 2	5'-GMP ^a	7.0, 298	0.51
		7.0, 310	1.66
		7.0, 318	3.57

^a This work. ^b Ref. 7. ^c First-order rate constant $10^4 k_1/\text{s}^{-1}$.





Fig. 1 Time-course of the reaction between $[\text{Pt}(\text{dien})\text{Cl}]^+$ 1, Met and 5'-GMP (1:2:2 mol ratio). Initially there is rapid formation of $[\text{Pt}(\text{dien})(\text{Met-S})]^{2+}$ 2 followed by displacement of Met by 5'-GMP. In the later stages the formation of $[\text{Pt}(\text{dien})(5'\text{-GMP-N}^7)]$ 3 is almost complete

of $[\text{Pt}(\text{dien})\text{Cl}]^+$ 1. Complex 1 has the advantage that the chelated dien ligand does not readily behave as a leaving group in the presence of sulfur ligands, unlike the amines on cisplatin. We show here that intermolecular displacement of S-bound L-methionine by N⁷-bound 5'-GMP readily occurs, and is selective in comparison with the N ligands of other DNA bases or L-histidine (His). These findings, together with the recent report of intramolecular displacement of a Pt-bound thioether by a guanine nucleobase,⁶ suggest that novel routes to DNA platination by platinum anticancer drugs may exist *in vivo*.

First we studied the competitive reaction of 17⁺ (10 mmol dm⁻³) with Met (20 mmol dm⁻³) and 5'-GMP (20 mol dm⁻³) in D₂O pH* 7.2.‡ In the initial stages of the reaction (<40 h), ¹H NMR (JEOL GX270) peaks for free Met [δ 2.136 (C ^{α} H₃) and 3.869 (C ^{α} H)] decreased in intensity, new peaks characteristic of $[\text{Pt}(\text{dien})(\text{Met-S})]^{2+}$ 2 appeared in the spectrum [δ 2.547 (C ^{α} H₃) and 3.940 (C ^{α} H)],§ whereas little of the 5'-GMP reacted, Fig. 1. In the later stages, the peaks for bound Met and free 5'-GMP (δ 8.202 for H⁸) decreased in intensity, whereas those for free Met increased in intensity, as did those assignable to bound 5'-GMP in $[\text{Pt}(\text{dien})(5'\text{-GMP-N}^7)]$ 3 (δ 8.856 for H⁸), Fig. 1. In a similar competition reaction between Met and adenosine 5'-monophosphate (5'-AMP) (pH* 6.3), nearly all the Met reacted with 1 within 6 h, but only ca. 2.5% of the 5'-AMP had reacted after 30 d at ambient temperature.

In separate experiments we confirmed that the reaction of 1 with Met alone is relatively fast, and that complex 3 can be formed from complex 2 by direct displacement of coordinated Met by 5'-GMP. The rate of reaction of Met with 1 (half-life 2.0 h at 300 K; second-order rate constant k_2 , Table 1) is similar to that reported previously for S-methylglutathione (GSMc).⁷ The direct reaction of the Met adduct 2 with 5'-GMP was studied at 310 K in 50 mmol dm⁻³ phosphate buffer, pH* 7.0. The appropriate plot for second-order kinetics⁸ was



linear giving a rate constant of $1.66 \times 10^{-4} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ (half-life of 167 h). The reaction was also followed at 298 and 318 K, and an Eyring plot yielded values of ΔH^\ddagger of 73.8 kJ mol⁻¹ and ΔS^\ddagger of $-79.7 \text{ J K}^{-1} \text{ mol}^{-1}$. These values are typical of those reported previously for substitution reactions of square-planar platinum(II) complexes which occur *via* an associative mechanism (five-coordinate transition state).⁹

Reactions of **2** with nucleobases were remarkably selective for guanine. In a competitive experiment between **2** (10 mmol dm⁻³, 50 mmol dm⁻³ phosphate buffer, pH* 7.0) 5'-GMP, 5'-AMP, 5'-TMP (thymine 5'-monophosphate) and 5'-CMP (cytosine 5'-monophosphate) (10 mmol dm⁻³ each), 7% of the 5'-GMP had reacted by displacing Met after 12.6 h, but none of the other bases had reacted.

Notable also was the inability of the imidazole N of His to displace S-bound Met: no reaction between **2** (10 mol dm⁻³, 50 mmol dm⁻³ phosphate buffer, pH* 7) and 1 mole equiv. of His was observed even after 3 d.

Comparisons between complexes of Pt^{II} and Pd^{II} are of interest since Pd^{II} analogues of Pt^{II} antitumour agents are usually much less active. For equimolar mixtures of [Pd(dien)Cl]⁺ **1**, Met and 5'-GMP, only GMP adducts were detected in the NMR spectrum, whereas with Met and 5'-AMP, both S-bound Met together with peaks assignable to both N¹- and N⁷-bound 5'-AMP were seen. These data are consistent with the more rapid substitution reactions of Pd^{II} compared to Pt^{II} (often > 10⁴ times faster), and again with the stronger nucleophilicity of guanine compared to adenine. In the case of Pd^{II}, initial [Pd(dien)(Met)]²⁺ may convert rapidly to [Pd(dien)(5'-GMP-N⁷)], although the reaction with GMP may also occur rapidly *via* aqua species. Palladium(II) binding to N⁷ has been established by X-ray crystallography of [Pd(dien)(guanosine)](ClO₄)₂,¹⁰ and Pd^{II} is known to bind strongly to both N⁷ and N¹ of AMP.¹¹

These findings have implications for the mechanism of action of platinum anticancer drugs. Sulfur ligands are generally thought to have a much higher affinity for Pt^{II} than nitrogen ligands and to diminish the antitumour activity of platinum complexes.¹² Indeed sulfur nucleophiles have been used as rescue agents to remove excess Pt from the body.¹³ The present work and that on intramolecular displacement,⁶ which was reported whilst this work was being written up, suggest that the binding of thioether sulfur to Pt^{II} is reversible and could provide a novel mechanism for DNA platination. The displacement of S-bound Met by N⁷-bound GMP appears to be about an order of magnitude slower than the intramolecular isomerization of [Pt(dien)(guanosylhomocysteine-S)] reported by van Boom and Reedijk,⁶ although detailed kinetics were not determined for the latter reaction and so a proper comparison of rates is not possible.

Platinum transfer reactions to DNA bases *via* Met intermediates could have biological significance in cells with high concentrations of Cl⁻ ions in their nuclei (e.g. 150 mmol dm⁻³ in liver cells¹⁴) for which the accepted pathway of DNA platination *via* aqua intermediates is likely to be quenched. The GMP-Met displacement reactions are slow, but it is notable that a very slowly excreted pool of Pt exists *in vivo* after administration of cisplatin (which has a half-life of several days).¹⁵ In small cellular compartments such as the nucleus, the concentrations of the reactants may be effectively raised, so increasing the rates of the second-order substitution reactions. Also displacement reactions may be facilitated if the methionine adduct is formed not simply by Met itself, but by an accessible Met residue on a DNA-binding protein. Our finding of selective transfer to G as opposed to other DNA bases, or to His which is a common residue in proteins, is notable since G bases are known to be major targets for Pt attack on DNA.¹⁶ It will be interesting to investigate the effects which Pt-bound Met and methionine-containing peptides and proteins have on the DNA sequence specificity for G attack. Since monodentate S-bound Met has free amino and

carboxylate groups, rapid transport mechanisms could exist for such Pt complexes through cellular compartments *via* amino acid receptors in membranes. Although thiols such as glutathione are also abundant S-containing ligands in cells, it is notable that thioethers react faster with [Pt(dien)Cl]⁺ than thiols.^{7,17} Reactions between cisplatin and 5'-GMP in the presence of Met are complicated by the loss of ammine ligands, but it appears¹⁸ that displacement of S-bound Met can occur in this case also and therefore may be a general reaction available to platinum anticancer complexes.

We thank the Medical Research Council, Sybil Evans Trust, International Association for Cancer Research, the Royal Society, SERC and University of London Intercollegiate Research Service for their support for this work. We acknowledge EC support under the HCM programme (grant CHRX-CT92-0016) allowing regular scientific exchange with colleagues in Europe.

Received, 22nd November 1993; Com. 3/06964H

Footnotes

† The compounds [Pt(dien)Cl]Cl **1** and [Pt(dien)(NO₃)]NO₃ were prepared as previously described.⁷ [Pt(dien)(Met-S)]²⁺ **2** was prepared by mixing pH 2 solutions (adjusted with 1 mol dm⁻³ HNO₃) of [Pt(dien)(NO₃)]NO₃ (2 cm³, 50 mmol dm⁻³) and Met (2 cm³, 50 mmol dm⁻³). The reaction was carried out at low pH to avoid coordination of the amino group. Complete formation of the product was checked by ¹H NMR spectroscopy and any slight excess of one component was corrected if necessary by addition of the other component. The solution was lyophilized and the solid complex was stored at -20 °C and used as required.

‡ pH* is the pH meter reading in D₂O solution.

§ The single set of peaks is indicative of rapid inversion at the chiral S centre on the ¹H NMR time-scale consistent with monodentate S-bound Met (ref. 3).

References

- N. Farrell, *Transition Metal Complexes as Drugs and Chemotherapeutic Agents*, Kluwer Academic Press, Dordrecht, 1989.
- C. M. Riley, L. A. Sternson, A. J. Repta and S. A. Slyter, *Anal. Biochem.*, 1983, **130**, 203.
- P. d. S. Murdoch, J. D. Ranford, P. J. Sadler and S. J. Berners-Price, *Inorg. Chem.*, 1993, **32**, 2249.
- A. Lepre, P. d. S. Murdoch and P. J. Sadler, unpublished work.
- M. D. Rhodes, J. D. Ranford and P. J. Sadler, in *Metallothioneins: Synthesis Structure and Properties of Metallothioneins, Phytochelatins and Metal-thiolate Complexes*, ed. M. J. Stillman, C. F. Shaw III and K. T. Suzuki, VCH, 1992, ch. 16, p. 423; J. D. Ranford, P. J. Sadler and R. E. Norman, *Int. Chem. Congr. Pacific Basin Soc.*, ACS, Washington, 1989, Abstr. BIOS88.
- S. S. G. E. van Boom and J. Reedijk, *J. Chem. Soc., Chem. Commun.*, 1993, 1397.
- M. I. Djuran, E. L. M. Lempers and J. Reedijk, *Inorg. Chem.*, 1991, **30**, 2648.
- W. J. Moore, *Physical Chemistry*, Longman, London, 1978, ch. 9.
- R. G. Wilkins, *Kinetics and Mechanism of Reactions of Transition Metal Complexes*, VCH, Weinheim, 1991, p. 235.
- F. D. Rochon, P. C. Kong, B. Coulombe and R. Melanson, *Can. J. Chem.*, 1980, **58**, 381.
- R. B. Martin, *ACS Symp. Ser.*, 1983, **208**, 231.
- N. Farrell, *Transition Metal Complexes as Drugs and Chemotherapeutic Agents*, Kluwer Academic Press, Dordrecht, 1989, p. 51.
- P. C. Dedon, R. Qazi and R. F. Borch, in *Biochemical Mechanisms of Platinum Antitumour Drugs*, ed. D. C. H. McBrien and T. F. Slater, IRL Press, Oxford, 1986, p. 199.
- G. Siebert, *Sub.-Cell. Biochem.*, 1972, **1**, 277.
- A. W. Prestayko, in *Cisplatin, Current status and New Developments*, eds. A. W. Prestayko, S. T. Crooke and S. K. Carter, Academic Press, London, 1980, p. 2.
- S. E. Sherman and S. J. Lippard, *Chem. Rev.*, 1987, **87**, 1153.
- E. L. M. Lempers, K. Inagaki and J. Reedijk, *Inorg. Chim. Acta*, 1988, **152**, 201.
- K. J. Barnham, M. I. Djuran, P. d. S. Murdoch, J. D. Ranford and P. J. Sadler, unpublished work.