

The module $\mathcal{D}f^s$ for locally quasi-homogeneous free divisors

Francisco Calderón-Moreno and Luis Narváez-Macarro*

Abstract

We find explicit free resolutions for the \mathcal{D} -modules $\mathcal{D}f^s$ and $\mathcal{D}[s]f^s/\mathcal{D}[s]f^{s+1}$, where f is a reduced equation of a locally quasi-homogeneous free divisor. These results are based on the fact that every locally quasi-homogeneous free divisor is Koszul free, which is also proved in this paper.

Introduction

In this paper we study the module $\mathcal{D}f^s$, where \mathcal{D} is the ring of germs at $0 \in \mathbb{C}^n$ of linear holomorphic differential operators and f is a reduced local equation of a locally quasi-homogeneous free divisor $D \subset (\mathbb{C}^n, 0)$.

The module $\mathcal{D}f^s$ encodes an enormous amount of geometric information of the singularity $f = 0$, but usually it is hard to work with in an explicit way. We prove the following results (see corollary 5.8 and theorem 5.9):

(A) Let $f = 0$ be a reduced local equation of a locally quasi-homogeneous free divisor of \mathbb{C}^n , and let $\{\delta_1, \dots, \delta_{n-1}\}$ be a basis of the module of vector fields vanishing on f . Then

1. The δ_i generate the ideal $\text{Ann}_{\mathcal{D}} f^s$.

*The authors are supported by PB97-0723.

⁰Keywords: Free divisor, de Rham complex, D-module, locally quasi-homogeneous, Koszul complex, Spencer complex, ideal of linear type.

⁰Mathematical Subject Classification: 14F40, 32S20, 32S40.

2. There exist explicit Koszul-Spencer type free resolutions for the modules $\mathcal{D}f^s$ and $\mathcal{D}[s]f^s/\mathcal{D}[s]f^{s+1}$ built on $\delta_1, \dots, \delta_{n-1}$ and $f, \delta_1, \dots, \delta_{n-1}$, respectively.

Locally quasi-homogeneous free divisors form an important class of divisors with non isolated singularities: normal crossing divisors, the union of reflecting hyperplanes of a complex reflection group, free hyperplane arrangements or the discriminant of stable mappings in Mather’s “nice dimensions” are examples of such divisors.

Let X be a complex analytic manifold. Given a divisor $D \subset X$, let us write $j : U = X \setminus D \hookrightarrow X$ for the corresponding open inclusion and $\Omega^\bullet(*D)$ for the meromorphic de Rham complex with poles along D . In [11], Grothendieck proved that the canonical morphism $\Omega^\bullet(*D) \rightarrow \mathbf{R}j_*(\mathbb{C}_U)$ is an isomorphism (in the derived category). This result is usually known as (a version of) *Grothendieck’s Comparison Theorem*.

In [17], K. Saito introduced the *logarithmic de Rham complex* associated with D , $\Omega_X^\bullet(\log D)$, generalizing the well known case of normal crossing divisors (cf. [8]). In the same paper, K. Saito also introduced the important notion of *free divisor*.

In [7], it is proved that the logarithmic de Rham complex $\Omega_X^\bullet(\log D)$ computes the cohomology of the complement U if D is a locally quasi-homogeneous free divisor (we say that D satisfies the *logarithmic comparison theorem*). In other words, the canonical morphism $\Omega_X^\bullet(\log D) \rightarrow \mathbf{R}j_*(\mathbb{C}_U)$ is an isomorphism, or using Grothendieck’s result, the inclusion $\Omega_X^\bullet(\log D) \hookrightarrow \Omega^\bullet(*D)$ is a quasi-isomorphism. In fact, in [5] it is proved that, in the case of $\dim X = 2$, D is locally quasi-homogeneous if and only if it satisfies the logarithmic comparison theorem.

Since the derived direct image $\mathbf{R}j_*(\mathbb{C}_U)$ is a perverse sheaf (it is the de Rham complex of the holonomic module of meromorphic functions with poles along D [15], II, th. 2.2.4), we deduce that $\Omega_X^\bullet(\log D)$ is perverse for every locally quasi-homogeneous free divisor.

On the other hand, the first author proved the following results [4]:

Let $D \subset X$ be a Koszul free divisor (see definition 1.6) and \mathcal{J} the left ideal of the ring \mathcal{D}_X of differential operators on X generated by the logarithmic vector fields with respect to D . Then

- 1) The left \mathcal{D}_X -module $\mathcal{D}_X/\mathcal{J}$ is holonomic.

2) There is a canonical isomorphism in the derived category

$$\Omega_X^\bullet(\log D) \simeq \mathbf{R} \operatorname{Hom}_{\mathcal{D}_X}(\mathcal{D}_X/\mathcal{J}, \mathcal{O}_X).$$

As a consequence, the logarithmic de Rham complex associated with a Koszul free divisor is a perverse sheaf.

The proof of **(A)** depends strongly on the following result, which has been suggested by the above results (see theorem 4.3):

(B) Every locally quasi-homogeneous free divisor is Koszul free.

In the first three sections we introduce some material concerning locally quasi-homogeneous free divisors, Koszul free divisors, the notion of linear type ideal and the module $\mathcal{D}f^s$.

In the fourth section we include the proof of **(B)** in our previous paper [6].

The fifth section is the main part of this paper and contains the proof of **(A)** and some related results.

In the sixth section we study some examples and we state some problems and conjectures.

The first part of **(A)** has been proposed (without proof) in [1, page 240] in the particular case of discriminants of versal deformations of simple hypersurface singularities. The normal crossing divisors case has been treated in [10].

1 Locally quasi-homogeneous and Koszul free divisors

1.1 Let X be a n -dimensional complex analytic manifold. We denote by $\pi : T^*X \rightarrow X$ the cotangent bundle, \mathcal{O}_X the sheaf of holomorphic functions on X , \mathcal{D}_X the sheaf of linear differential operators on X (with holomorphic coefficients), $Gr_{F^\bullet}(\mathcal{D}_X)$ the graded ring associated with the filtration F^\bullet by the order, $\sigma(P)$ the principal symbol of a differential operator P and $\{-, -\}$ the Poisson bracket on \mathcal{O}_{T^*X} or $Gr_{F^\bullet}(\mathcal{D}_X)$. We will note $\mathcal{O} = \mathcal{O}_{X,p}$, $\mathcal{D} = \mathcal{D}_{X,p}$ and $Gr_{F^\bullet}(\mathcal{D}) = Gr_{F^\bullet}(\mathcal{D}_X)_p$ the respective stalks at p , with p a point of X . If $J \subset \mathcal{D}$ is a left ideal, we denote by $\sigma(J)$ the corresponding graded ideal of $Gr_{F^\bullet}(\mathcal{D})$. Given a divisor $D \subset X$, we denote by $Der(\log D)$ the \mathcal{O}_X -module of the logarithmic vector fields with respect to D [17]. If f is a local

equation of D at p , we denote by $\text{Der}(\log f)$ the stalk at p of $\text{Der}(\log D)$, whose elements are germs at p of vector fields δ such that $\delta(f) \in (f)$.

Definition 1.2.— A divisor D is Euler-homogeneous at $p \in D$ if there is a local equation h for D around p , and a germ of (logarithmic) vector field δ such that $\delta(h) = h$. A such δ is called a local Euler vector field for h .

The set of points where a divisor is Euler-homogeneous is open.

Definition 1.3.— (cf. [7]) A germ of divisor $(D, p) \subset (X, p)$ is quasi-homogeneous if there are local coordinates $x_1, \dots, x_n \in \mathcal{O}_{X,p}$ with respect to which (D, p) has a weighted homogeneous defining equation (with strictly positive weights). A divisor D in a n -dimensional complex manifold X is locally quasi-homogeneous if the germ (D, p) is quasi-homogeneous for each point $p \in D$. A germ of divisor $(D, p) \subset (X, p)$ is locally quasi-homogeneous if the divisor D is locally quasi-homogeneous in a neighborhood of p .

Obviously a locally quasi-homogeneous divisor is Euler-homogeneous at every point.

Definition 1.4.— We say that a reduced germ $f \in \mathcal{O}_{X,p}$ is locally quasi-homogeneous if the germ of divisor $(\{f = 0\}, p)$ is.

Remark 1.5.— A reduced germ $f \in \mathcal{O}_{X,p}$ is locally quasi-homogeneous if and only if for every $q \in \{f = 0\}$ near p there are local coordinates $z_1, \dots, z_n \in \mathcal{O}_{X,q}$ and a quasi-homogeneous polynomial $P(t_1, \dots, t_n)$ (with strictly positive weights) such that $f_q = P(z_1, \dots, z_n)$.

Definition 1.6.— ([17], [4], def. 4.1.1) Let $D \subset X$ be a divisor. We say that D is free at $p \in X$ if $\text{Der}(\log D)_p$ is a free \mathcal{O} -module (of rank n). We say that D is a Koszul free divisor at $p \in X$ if it is free at p and there exists a basis $\{\delta_1, \dots, \delta_n\}$ of $\text{Der}(\log D)_p$ such that the sequence of symbols $\{\sigma(\delta_1), \dots, \sigma(\delta_n)\}$ is regular in $\text{Gr}_{F^\bullet}(\mathcal{D}) = \text{Gr}_{F^\bullet}(\mathcal{D}_X)_p$. If D is a free (resp. Koszul free) divisor at each point of X , we simply say that it is free (resp. Koszul free). We say that a reduced germ $f \in \mathcal{O}_{X,p}$ is free if the divisor $f^{-1}(0)$ is free at p .

Let's remark that a divisor D is automatically Koszul free at every $p \in X \setminus D$.

Remark 1.7.— The ideal $I_{D,p} = \text{Gr}_{F^\bullet}(\mathcal{D}) \text{Der}(\log D)_p$ is generated by the elements of any basis of $\text{Der}(\log D)_p$. As D is Koszul free at p if and only if $\text{depth}(I_{D,p}, \text{Gr}_{F^\bullet}(\mathcal{D})) = n$ (cf. [14], cor. 16.8), it is clear that the definition

of Koszul free divisor does not depend on the election of a particular basis. By the coherence of $Gr_{F^\bullet}(\mathcal{D}_X)$, if a divisor is Koszul free at a point, then it is Koszul free near that point.

We have not found a reference for the following well known proposition (see [14], th. 17.4 for the local case).

Proposition 1.8.— Let $\mathbb{C}\{x\}$ be the ring of convergent power series in the variables $x = x_1, \dots, x_n$ and let G be the graded ring of polynomials in the variables ξ_1, \dots, ξ_t with coefficients in $\mathbb{C}\{x\}$. A sequence $\sigma_1, \dots, \sigma_s$ of homogeneous polynomials in G is regular if and only if the set of zeros $V(I)$ of the ideal I generated by $\sigma_1, \dots, \sigma_s$ has dimension $n + t - s$ in $U \times \mathbb{C}^t$, for some open neighborhood U of 0 (then each irreducible component has dimension $n + t - s$).

Proof: Let $\mathbb{C}\{x, \xi\}$ be the ring of convergent power series in the variables $x_1, \dots, x_n, \xi_1, \dots, \xi_t$. As the σ_i are homogeneous in G and the ring $\mathbb{C}\{x, \xi\}$ is a flat extension of G , the σ_i are a regular sequence in G if and only if they are a regular sequence in $\mathbb{C}\{x, \xi\}$. But the last condition is equivalent to the equality (*loc. cit.*):

$$\dim_{(0,0)}(V(I)) = \dim(\mathbb{C}\{x, \xi\}/I) = n + t - s.$$

Finally, using the fact that all the σ_i are homogeneous in the variables ξ , the local dimension of $V(I)$ at $(0, 0)$ coincides with its dimension in $U \times \mathbb{C}^t$ for some neighborhood U of 0. C.Q.D.

Corollary 1.9.— Let $D \subset X$ be a free divisor. Let J be the ideal in \mathcal{O}_{T^*X} generated by $\pi^{-1} Der(\log D)$. Then, D is Koszul free if and only if the set $V(J)$ of zeros of J has dimension n (in this case, each irreducible component of $V(J)$ has dimension n).

Proposition 1.10.— Let X be a complex manifold of dimension n and let $D \subset X$ be a divisor. Then:

1. Let $X' = X \times \mathbb{C}$ and $D' = D \times \mathbb{C}$. The divisor $D \subset X$ is Koszul free if and only if $D' \subset X'$ is Koszul free.
2. Let Y be another complex manifold of dimension r and let $E \subset Y$ be a divisor. Then: a) The divisor $(D \times Y) \cup (X \times E)$ is free if $D \subset X$ and $E \subset Y$ are free.

b) The divisor $(D \times Y) \cup (X \times E)$ is Koszul free if $D \subset X$ and $E \subset Y$ are Koszul free.

Proof:

1. It is a consequence of [7], lemma 2.2, (iv) and the fact that $\sigma_1, \dots, \sigma_n$ is a regular sequence in $\mathcal{O}_{X,p}[\xi_1, \dots, \xi_n]$ if and only if $\xi_{n+1}, \sigma_1, \dots, \sigma_n$ is a regular sequence in $\mathcal{O}_{X',(p,t)}[\xi_1, \dots, \xi_n, \xi_{n+1}]$.
2. a) It is an immediate consequence of Saito's criterion (cf. [7], lemma 2.2, (v)).
b) It is a consequence of a) and Corollary 1.9.

C.Q.D.

Example 1.11.— Examples of Koszul free divisors are:

- 1) Nonsingular divisors.
- 2) Normal crossing divisors.
- 3) Plane curves: If $\dim_{\mathbb{C}} X = 2$, we know that every divisor $D \subset X$ is free [17], cor. 1.7. Let $\{\delta_1, \delta_2\}$ be a basis of $Der(\log D)_x$. Their symbols $\{\sigma_1, \sigma_2\}$ are obviously linearly independent over \mathcal{O} , and by Saito's criterion [17], 1.8, they are relatively primes in $\text{Gr}_{F^\bullet}(\mathcal{D}) = \mathcal{O}[\xi_1, \xi_2]$. So they form a regular sequence in $\text{Gr}_{F^\bullet}(\mathcal{D})$, and D is Koszul free (see [4], cor. 4.2.2).
- 4) Proposition 1.10 gives a way to obtain Koszul free divisors in any dimension.
- 5) There are irreducible Koszul free divisors in dimensions greater than 2, which are not constructed from divisors in lower dimension [16]: $X = \mathbb{C}^3$ and $D \equiv \{f = 0\}$, with

$$f = 2^8 z^3 - 2^7 x^2 z^2 + 2^4 x^4 z + 2^4 3^2 x y^2 z - 2^2 x^3 y^2 - 3^3 y^4.$$

A basis of $Der(\log f)$ is $\{\delta_1, \delta_2, \delta_3\}$, with

$$\begin{aligned} \delta_1 &= 6y \partial_x + (8z - 2x^2) \partial_y - xy \partial_z, \\ \delta_2 &= (4x^2 - 48z) \partial_x + 12xy \partial_y + (9y^2 - 16xz) \partial_z, \\ \delta_3 &= 2x \partial_x + 3y \partial_y + 4z \partial_z, \end{aligned}$$

and the sequence $\{\sigma(\delta_1), \sigma(\delta_2), \sigma(\delta_3)\}$ is $\text{Gr}_{F^\bullet}(\mathcal{D})$ -regular.

2 Ideals of linear type

Definition 2.1.— (cf. [18], §7.2) Let A be a commutative ring, $I \subset A$ an ideal, $\mathcal{R}(I) = \bigoplus_{i=0}^{\infty} I^d t^d \subset A[t]$ the Rees algebra of I and $\text{Sim}(I)$ the symmetric algebra of the A -module I . We say that I is of *linear type* if the canonical (surjective) morphism of graded A -algebras

$$\text{Sim}(I) \rightarrow \mathcal{R}(I)$$

is an isomorphism.

Lemma 2.2.— Given a commutative ring A and an ideal $I \subset A$ generated by a family of elements $\{a_i\}_{i \in \Lambda}$, the following properties are equivalent:

- a) I is of linear type.
- b) If $\varphi : A[\{X_i\}_{i \in \Lambda}] \rightarrow \mathcal{R}(I)$ is the morphism of graded algebras defined by $\varphi(X_i) = a_i t$, then the kernel of φ is generated by homogeneous elements of degree 1.

Proof: We consider the kernel of the surjective morphism of graded A -algebras $\Phi : A[\{X_i\}_{i \in \Lambda}] \rightarrow \text{Sim}(I)$, defined by $\Phi(X_{i_1} \cdots X_{i_d}) = a_{i_1} \cdots a_{i_d}$. Then $\ker(\Phi) = \ker(\varphi)$ if and only if I is of linear type, $\ker(\Phi)$ is an ideal generated by its homogeneous elements of degree 1, $\ker(\Phi)_1$, and $\ker(\Phi)_1 = \ker(\varphi)_1$. C.Q.D.

The definition and the lemma above sheafify in the obvious way.

The following results concern the case where the ideal I is generated by a regular sequence.

Lemma 2.3.— Let $\{a_1, \dots, a_m\}$ be an A -sequence. For $p \leq m$, if $\alpha a_1^{s_1} \cdots a_m^{s_m} \in (a_1^{s_1+k_1}, \dots, a_p^{s_p+k_p})$, then $\alpha \in (a_1^{k_1}, \dots, a_p^{k_p})$.

Proof: For $j = p+1, \dots, m$, $\{a_1^{s_1+k_1}, \dots, a_p^{s_p+k_p}, a_{p+1}^{s_{p+1}}, \dots, a_j^{s_j}\}$ is a regular A -sequence, and we can prove inductively that

$$\alpha a_1^{s_1} \cdots a_{j-1}^{s_{j-1}} \in (a_1^{s_1+k_1}, \dots, a_p^{s_p+k_p}).$$

For $i = p-1, \dots, 0$, $\{a_1^{s_1+k_1}, \dots, a_i^{s_i+k_i}, a_{i+1}^{k_{i+1}}, \dots, a_p^{k_p}\}$ is a regular A -sequence, and we inductively prove that

$$\alpha a_1^{s_1} \cdots a_i^{s_i} \in (a_1^{s_1+k_1}, \dots, a_i^{s_i+k_i}, a_{i+1}^{k_{i+1}}, \dots, a_p^{k_p}).$$

C.Q.D.

Proposition 2.4.— Let A be a commutative ring and let $I \subset A$ be an ideal generated by a regular sequence a_1, \dots, a_n . Then, the kernel of the morphism of graded algebras

$$A[X_1, \dots, X_n] \rightarrow A[t], \quad X_i \mapsto a_i t,$$

is generated by $a_i X_j - a_j X_i$, $1 \leq i < j \leq n$. In particular, I is of linear type.

Proof: Let g be an homogeneous polynomial of degree m in $A[X_1, \dots, X_n]$ such that $g(a_1, \dots, a_n) = 0$. Let $\exp_g = cX^{e_g}$ be the greatest monomial of g in the inverse lexicographic order, with $e_g = (s_1, \dots, s_t, 0, \dots, 0)$, $s_t \neq 0$. Then

$$g(X_1, \dots, X_n) - \exp_g \in (X_1^{s_1+1}, \dots, X_t^{s_t+1}).$$

By lemma 2.3, $c = \sum_{i=1}^{t-1} \alpha_i a_i \in (a_1, \dots, a_{t-1})$. Then

$$f(X_1, \dots, X_n) = g(X_1, \dots, X_n) - \sum_{i=1}^{t-1} \alpha_i (a_i X_t - a_t X_i) X_1^{s_1} \cdots X_n^{s_n}$$

is an homogeneous polynomial of degree m such that $e_f < e_g$ and

$$f(X_1, \dots, X_n) - g(X_1, \dots, X_n) \in J = (a_i X_j - a_j X_i, 0 < i < j \leq n).$$

In particular, $f(a_1, \dots, a_n) = 0$. Consequently, after a finite number of steps, we will obtain $h(X_1) = c_m X_1^m$, such that $h(X_1) - g(X_1, \dots, X_n) \in J$. So $h(a_1) = c_m a_1^m = 0$, $c_m = 0$ and $g(X_1, \dots, X_n) \in J$.

C.Q.D.

3 The module $\mathcal{D}f^s$

Let X be a n -dimensional complex analytic manifold, p a point in X and $f \in \mathcal{O} = \mathcal{O}_{X,p}$ a non zero germ of holomorphic function with $f(p) = 0$. Let D be the (germ of) divisor defined by $f = 0$. The free module of rank one over the ring $\mathcal{O}[f^{-1}, s]$ generated by the symbol f^s has a natural left module structure over the ring $\mathcal{D}[s]$ [2]: the action of a derivation $\delta \in \text{Der}_{\mathbb{C}}(\mathcal{O})$ is given by $\delta(f^s) = \delta(f)sf^{-1}f^s$.

The following lemma is well-known and the proof is straightforward.

Lemma 3.1.– For every linear differential operator $P \in \mathcal{D}$ of order d , we have:

$$P(f^s) = C_{P,0}f^s + C_{P,1}\binom{s}{1}f^{s-1} + \cdots + C_{P,d}\binom{s}{d}f^{s-d}$$

where

$$C_{P,d} = d!\sigma(P)(df) = \{\cdots\{\{\sigma(P), f\}, f\} \cdots, f\}.$$

Denote by $J_f \subset \mathcal{O}$ the jacobian ideal associated with f . The surjection

$$\delta \in \text{Der}_{\mathbb{C}}(\mathcal{O}) \mapsto \delta(f) \in J_f$$

and the canonical isomorphism of graded \mathcal{O} -algebras

$$\text{Sim}_{\mathcal{O}}(\text{Der}_{\mathbb{C}}(\mathcal{O})) \simeq \text{Gr}_{F^{\bullet}}(\mathcal{D}) \tag{1}$$

induce a surjective graded morphism of \mathcal{O} -algebras

$$\varphi_f : \text{Gr}_{F^{\bullet}}(\mathcal{D}) \longrightarrow \mathcal{R}(J_f). \tag{2}$$

In coordinates, $\text{Gr}_{F^{\bullet}}(\mathcal{D}) = \mathcal{O}[\xi_1, \dots, \xi_n]$, $\xi_i = \sigma(\partial_i)$ and

$$\varphi_f(\sigma(P)) = \sigma(P)(\partial_1(f)t, \dots, \partial_n(f)t) = \sigma(P)(df)t^d$$

for every differential operator $P \in \mathcal{D}$ of order d .

The homogeneous part of degree 1 of $\ker \varphi_f$ is naturally identified with the \mathcal{O} -module

$$\Theta_f = \{\delta \in \text{Der}_{\mathbb{C}}(\mathcal{O}) \mid \delta(f) = 0\}$$

by means of the canonical isomorphism (1).

Lemma 3.1 implies that $\sigma(\text{Ann}_{\mathcal{D}} f^s) \subset \ker \varphi_f$.

Proposition 3.2.– With the above notations, if J_f is of linear type, then

$$\sigma(\text{Ann}_{\mathcal{D}} f^s) = \ker \varphi_f$$

and the left ideal $\text{Ann}_{\mathcal{D}} f^s$ of \mathcal{D} is generated by Θ_f .

Proof: By lemma 2.2, $\ker \varphi_f = \text{Gr}_{F^{\bullet}}(\mathcal{D})\Theta_f \subset \sigma(\text{Ann}_{\mathcal{D}} f^s)$.

The inclusion $\mathcal{D}\Theta_f \subset \text{Ann}_{\mathcal{D}} f^s$ is obvious. Let's prove that $\text{Ann}_{\mathcal{D}} f^s \subset \mathcal{D}\Theta_f$. Clearly, $F^1 \text{Ann}_{\mathcal{D}} f^s = \Theta_f$. Suppose $F^{d-1} \text{Ann}_{\mathcal{D}} f^s \subset \mathcal{D}\Theta_f$ and take a

differential operator $P \in F^d \text{Ann}_{\mathcal{D}} f^s \setminus F^{d-1} \text{Ann}_{\mathcal{D}} f^s$. Then, $\sigma(P) \in \ker \varphi_f = \text{Gr}_{F^\bullet}(\mathcal{D})\sigma(\Theta_f)$, and $\sigma(P) = \sum A_i \sigma(\delta_i)$, where $\delta_i \in \Theta_f$ and the A_i are homogeneous of degree $d-1$. Let Q_i be differential operators such that $\sigma(Q_i) = A_i$. We apply the induction hypothesis to $P - \sum_i Q_i \delta_i \in F^{d-1} \text{Ann}_{\mathcal{D}} f^s$ and we conclude the result. C.Q.D.

Proposition 3.3.– (*Isolated singularities case, cf. [13], 2.7*) If f has isolated singularity, then $\ker \varphi_f$ is generated by $\partial_i(f)\xi_j - \partial_j(f)\xi_i$, $1 \leq i < j \leq n$. In particular, the left ideal $\text{Ann}_{\mathcal{D}} f^s$ of \mathcal{D} is generated by $\partial_i(f)\partial_j - \partial_j(f)\partial_i$, $1 \leq i < j \leq n$.

Proof: It is a consequence of lemma 2.4 and proposition 3.2. C.Q.D.

4 Locally quasi-homogeneous free divisors are Koszul free

Proposition 4.1.– Let U be a connected open set of a complex n -dimensional analytic manifold X and let $\Sigma \subset U$ be a closed analytic set of dimension s . If a sequence $C = \{\sigma_1, \dots, \sigma_{n-s}\}$ of homogeneous polynomials in $\mathcal{O}_X(U)[\xi_1, \dots, \xi_n]$ is regular at every point $q \in U \setminus \Sigma$ (i.e. it is regular in $\mathcal{O}_{X,q}[\xi_1, \dots, \xi_n]$), then it is regular at every point of U .

Proof: Let $p \in \Sigma$ and let $\pi : U \times \mathbb{C}^n \rightarrow U$ be the projection. By proposition 1.8, we have to prove that the ideal $I = (\sigma_1, \dots, \sigma_{n-s})$ defines an analytic set $V = V(I) \subset U \times \mathbb{C}^n$ of dimension $n + s$. By hypothesis, we know that C is regular on $U \setminus \Sigma$, and so (*loc. cit.*) the dimension of (every irreducible component of) $V \cap \pi^{-1}(U \setminus \Sigma)$ is $n + s$. Now, let W be an irreducible component of V . It has, at least, dimension $n + s$. If W is contained in $\pi^{-1}(\Sigma) = \Sigma \times \mathbb{C}^n$, then it must be equal to $\pi^{-1}(\Sigma)$. If not, $\dim W = \dim(W \cap \pi^{-1}(U \setminus \Sigma)) \leq \dim(V \cap \pi^{-1}(U \setminus \Sigma)) = n + s$. So, we conclude that W has dimension $n + s$. C.Q.D.

Corollary 4.2.– Let D be a free divisor in some analytic manifold X and let $\Sigma \subset D$ a discrete set of points. If D is Koszul free at every point $x \in D \setminus \Sigma$, then D is Koszul free (at every point of X).

Theorem 4.3.– Every locally quasi-homogeneous free divisor is Koszul free.

Proof: We proceed by induction on the dimension t of the ambient manifold X . For $t = 1$, the theorem is trivial and for $t = 2$, the theorem is directly proved in example 1.11, 3). Now, we suppose that the result is true for $t < n$, and let D be a locally quasi-homogeneous free divisor of a complex analytic manifold X of dimension n . Let $p \in D$ and let $\{\delta_1, \dots, \delta_n\}$ be a basis of the logarithmic derivations of D at p .

Thanks to [7], prop. 2.4 and lemma 2.2, (iv), there is an open neighborhood U of p such that for each $q \in U \cap D$, with $q \neq p$, the germ of pair (X, D, q) is isomorphic to a product $(\mathbb{C}^{n-1} \times \mathbb{C}, D' \times \mathbb{C}, (0, 0))$, where D' is a locally quasi-homogeneous free divisor. Induction hypothesis implies that D' is a Koszul free divisor at 0. Then, by proposition 1.10.1., D is a Koszul free divisor at q too. We have then proved that D is a Koszul free divisor in $U \setminus \{p\}$. We conclude by using corollary 4.2. C.Q.D.

Corollary 4.4.– Every free divisor that is locally quasi-homogeneous at the complement of a discrete set is Koszul free.

In particular, the last corollary gives rise a new proof of the fact that every divisor in dimension 2 is Koszul free (cf. 1.11, 3)).

5 The module $\mathcal{D}f^s$ for locally quasi-homogeneous free divisors

5.1 In this section, $f \in \mathcal{O} = \mathcal{O}_{X,p}$ will be a reduced locally quasi-homogeneous free germ 1.4, 1.6. That means that $D = \{f = 0\}$ is a locally quasi-homogeneous free divisor near p .

We will also assume that

-) The equation f and its Euler vector field E are globally defined on X .
-) $E(q) \neq 0$ for every $q \in X \setminus \{p\}$.
-) $Der(\log D)$ is \mathcal{O}_X -free (of rank $n = \dim X$).

In order to proceed inductively on the dimension of the ambient variety when working with such f 's, we quote the following direct consequence of [9], lemmas 1.3, 1.5 (see also [7], prop. 2.4)

Proposition 5.2.— Let $f \in \mathcal{O}_{X,p}$ a reduced locally quasi-homogeneous free germ and let D be the divisor $f = 0$. For $q \in D \setminus \{p\}$ close to p , there are local coordinates $z_1, \dots, z_n \in \mathcal{O}_{X,q}$ centered at q and a quasi-homogeneous polynomial $G'(t_1, \dots, t_{n-1})$ in $n - 1$ variables which is also a locally quasi-homogeneous free germ in $\mathcal{O}_{\mathbb{C}^{n-1},0}$ and such that $f_q = G'(z_1, \dots, z_{n-1})$.

We call $\tilde{\Theta}_f$ the \mathcal{O}_X -sub-module (and Lie algebra) of $Der(\log D)$ whose sections are vector fields annihilating f . Denote by $\mathcal{J}_f \subset \mathcal{O}_X$ the jacobian ideal sheaf associated with f . The stalk of $\tilde{\Theta}_f$ (resp. of \mathcal{J}_f) at p is then Θ_f (resp. J_f).

As in (2), we have a surjective graded morphism of \mathcal{O}_X -algebras

$$\Phi_f : Gr_{F^\bullet}(\mathcal{D}_X) \longrightarrow \mathcal{R}(\mathcal{J}_f),$$

whose stalk at p is φ_f .

We have:

$$Der(\log D) = \tilde{\Theta}_f \oplus (\mathcal{O}_X E), \quad Der(\log f) = \Theta_f \oplus (\mathcal{O}E), \quad (3)$$

and $\tilde{\Theta}_f, \Theta_f$ are free of rank $n - 1$.

Proposition 5.3.— The Koszul complex associated with $\tilde{\Theta}_f \subset Der_{\mathbb{C}}(\mathcal{O}_{\mathbb{C}^n}) = Gr_{F^\bullet}^1(\mathcal{D}_X) \subset Gr_{F^\bullet}(\mathcal{D}_X)$:

$$0 \rightarrow Gr_{F^\bullet}(\mathcal{D}_X) \otimes_{\mathcal{O}_X} \bigwedge^{n-1} \tilde{\Theta}_f \xrightarrow{d_{-n+1}} \dots \xrightarrow{d_{-2}} Gr_{F^\bullet}(\mathcal{D}_X) \otimes_{\mathcal{O}_X} \bigwedge^1 \tilde{\Theta}_f \xrightarrow{d_{-1}} Gr_{F^\bullet}(\mathcal{D}_X)$$

$$d_{-k}(F \otimes (\sigma_1 \wedge \dots \wedge \sigma_k)) = \sum_{i=1}^k (-1)^{i-1} P\sigma_i \otimes (\sigma_1 \wedge \dots \wedge \hat{\sigma}_i \wedge \dots \wedge \sigma_k), \quad 0 < k < n,$$

is exact.

Proof: We need to prove that some (or any) basis $\{\delta_1, \dots, \delta_{n-1}\}$ of $\tilde{\Theta}_f$ form a regular sequence in $Gr_{F^\bullet}(\mathcal{D}_X)$, but such a basis can be augmented to a basis $\{\delta_1, \dots, \delta_{n-1}, E\}$ of $Der(\log D)$, that we know by theorem 4.3 to form a regular sequence in $Gr_{F^\bullet}(\mathcal{D}_X)$. C.Q.D.

Proposition 5.4.— With the hypothesis of 5.1, if the augmented graded complex of $Gr_{F^\bullet}(\mathcal{D}_X)$ -modules

$$0 \rightarrow Gr_{F^\bullet}(\mathcal{D}_X) \otimes_{\mathcal{O}_X} \bigwedge^{n-1} \tilde{\Theta}_f \xrightarrow{d_{-n+1}} \dots \xrightarrow{d_{-1}} Gr_{F^\bullet}(\mathcal{D}_X) \xrightarrow{\Phi_f} \mathcal{R}(\mathcal{J}_f) \rightarrow 0 \quad (4)$$

is exact on $X - \{p\}$, then it is exact everywhere.

Proof: We know that Φ_f is surjective. By proposition 5.3, the only thing to prove is $\ker \Phi_f = \text{Im } d_{-1}$. We can proceed separately on each homogeneous component:

$$0 \rightarrow \text{Gr}_{F^\bullet}^{m-n+1}(\mathcal{D}_X) \otimes_{\mathcal{O}_X} \bigwedge^{n-1} \tilde{\Theta}_f \xrightarrow{d_{-n+1}^m} \cdots \xrightarrow{d_{-1}^m} \text{Gr}_{F^\bullet}^m(\mathcal{D}_X) \xrightarrow{\Phi_f^m} \mathcal{J}_f^m \rightarrow 0.$$

Let's consider the coherent \mathcal{O}_X -module $\mathcal{F} = \text{Im } d_{-1}^m$ and the short sequence

$$0 \rightarrow \mathcal{F} \rightarrow \text{Gr}_{F^\bullet}^m(\mathcal{D}_X) \xrightarrow{\Phi_f^m} \mathcal{J}_f^m \rightarrow 0. \quad (5)$$

By proposition 5.3 and the fact that the cohomology with support $H_p^i(\mathcal{O}_X)$ vanishes for $i \neq n$, we deduce that $H_p^i(\mathcal{F}) = 0$ for $i = 0, 1$ and $H_p^0(\mathcal{J}_f^m) = 0$. These properties and the exactness of (5) on $X - \{p\}$ imply the proposition (cf. [12], (8.14)). C.Q.D.

The following lemma is clear.

Lemma 5.5.— Let $g \in \mathcal{O}_{n-1} = \mathbb{C}\{y_1, \dots, y_{n-1}\}$ and call $f = g$, but as an element in $\mathcal{O}_n = \mathbb{C}\{y_1, \dots, y_n\}$. Then:

1. $\ker \varphi_f$ is generated by $\ker \varphi_g$ and $\sigma(\partial_{y_n})$.
2. Θ_f is generated by Θ_g and ∂_{y_n} .

Theorem 5.6.— Let $f \in \mathcal{O} = \mathcal{O}_{X,p}$ be a reduced locally quasi-homogeneous free germ. Then the graded complex of $\text{Gr}_{F^\bullet}(\mathcal{D})$ -modules

$$0 \rightarrow \text{Gr}_{F^\bullet}(\mathcal{D}) \otimes_{\mathcal{O}} \bigwedge^{n-1} \Theta_f \xrightarrow{\varepsilon_{-n+1}} \cdots \xrightarrow{\varepsilon_{-1}} \text{Gr}_{F^\bullet}(\mathcal{D}) \xrightarrow{\varphi_f} \mathcal{R}(J_f) \rightarrow 0$$

is exact. In particular, the kernel of the morphism

$$\text{Gr}_{F^\bullet}(\mathcal{D}) \xrightarrow{\varphi_f} \mathcal{R}(J_f)$$

is the ideal generated by Θ_f and then the jacobian ideal J_f is of linear type.

Proof: By the exactness of (5.3), the only thing to prove is that $\ker \varphi_f$ is generated by $\sigma(\Theta_f)$. We will use induction on $n = \dim X$. If $n = 2$, we apply proposition 3.3. We suppose that the result is true if the ambient variety has

dimension $n - 1$. By Proposition 5.4, we need to prove the exactness of the complex (4) on $U \setminus \{x\}$, for some open neighborhood U of x , or equivalently, that $\ker \Phi_f$ is generated by $\sigma(\Theta_f)$ at every $q \in U \setminus \{x\}$. The result is then a consequence of proposition 5.2, lemma 5.5 and the induction hypothesis. C.Q.D.

Definition 5.7.– The Spencer complex¹ for $\tilde{\Theta}_f$ is the complex of free left \mathcal{D}_X -modules given by:

$$0 \rightarrow \mathcal{D}_X \otimes_{\mathcal{O}_X} \bigwedge^{n-1} \tilde{\Theta}_f \xrightarrow{\varepsilon_{-n+1}} \cdots \xrightarrow{\varepsilon_{-2}} \mathcal{D}_X \otimes_{\mathcal{O}_X} \bigwedge^1 \tilde{\Theta}_f \xrightarrow{\varepsilon_{-1}} \mathcal{D}_X,$$

$$\varepsilon_{-1}(P \otimes \delta) = P\delta; \quad \varepsilon_{-k}(P \otimes (\delta_1 \wedge \cdots \wedge \delta_k)) = \sum_{i=1}^k (-1)^{i-1} P\delta_i \otimes (\delta_1 \wedge \cdots \wedge \widehat{\delta}_i \cdots \wedge \delta_k)$$

$$+ \sum_{1 \leq i < j \leq k} (-1)^{i+j} P \otimes ([\delta_i, \delta_j] \wedge \delta_1 \wedge \cdots \wedge \widehat{\delta}_i \cdots \widehat{\delta}_j \cdots \wedge \delta_k), \quad 2 \leq k \leq n - 1.$$

In a similar way we define the Spencer complex for Θ_f , which is the stalk at p of the Spencer complex for $\tilde{\Theta}_f$.

Both Spencer complexes can be augmented by considering the obvious maps $\mathcal{D}_X \rightarrow \mathcal{D}_X f^s, \mathcal{D} \rightarrow \mathcal{D} f^s$.

Corollary 5.8.– With the hypothesis of 5.1, we have:

- (a) The Spencer complex for Θ_f is a resolution of $\mathcal{D} f^s$. In particular, the left ideal $\text{Ann}_{\mathcal{D}} f^s$ is generated by Θ_f .
- (b) The left ideal $\text{Ann}_{\mathcal{D}[s]} f^s$ is generated by Θ_f and $E - s$.
- (c) The left ideal $\text{Ann}_{\mathcal{D}} \eta$, where η is the class of f^s in the quotient $\mathcal{D}[s] f^s / \mathcal{D}[s] f^{s+1}$, is generated by Θ_f and f .

Proof: For (a) we proceed as in [4], prop. 4.1.3 by using proposition 3.2 and theorem 5.6. Property (b) follows easily from (a), and property (c) follows from (a) and (b). C.Q.D.

¹It should be noticed that such complex was originally used by Chevalley and Eilenberg in the setting of the cohomology of Lie algebras (cf. [19], 7.7).

Let's call $\Xi_f = \Theta_f \oplus (\mathcal{O}f)$ (resp. $\tilde{\Xi}_f = \tilde{\Theta}_f \oplus (\mathcal{O}_X f)$), which is a free sub- \mathcal{O} -module (respectively, sub- \mathcal{O}_X -module) and a Lie subalgebra of \mathcal{D} (resp. of \mathcal{D}_X). It can be also canonically embedded in $\text{Gr}_{F^\bullet}(\mathcal{D})$ (resp. $\text{Gr}_{F^\bullet}(\mathcal{D}_X)$) equipped with the Poisson bracket $\{-, -\}$. As in 5.3 and 5.7, we define the Koszul complex associated with $\Xi_f \subset \text{Gr}_{F^\bullet}(\mathcal{D})$ (resp. $\tilde{\Xi}_f \subset \text{Gr}_{F^\bullet}(\mathcal{D}_X)$) and the Spencer complex associated with $\Xi_f \subset \mathcal{D}$ (resp. $\tilde{\Xi}_f \subset \mathcal{D}_X$). The Koszul (resp. Spencer) complex associated with $\Xi_f \subset \text{Gr}_{F^\bullet}(\mathcal{D})$ (resp. with $\Xi_f \subset \mathcal{D}$) is obviously the stalk at p of the Koszul (resp. of the Spencer) complex associated with $\tilde{\Xi}_f \subset \text{Gr}_{F^\bullet}(\mathcal{D}_X)$ (resp. with $\tilde{\Xi}_f \subset \mathcal{D}_X$)

Theorem 5.9.— With the hypothesis of 5.1, the following properties hold:

1. The Koszul complex associated with $\Xi_f \subset \text{Gr}_{F^\bullet}(\mathcal{D})$ is exact.
2. The Spencer complex associated with $\Xi_f \subset \mathcal{D}$ is a free resolution of $\mathcal{D}[s]f^s/\mathcal{D}[s]f^{s+1}$.

Proof: For the first property, call \mathbf{K} the Koszul complex associated with $\tilde{\Xi}_f \subset \text{Gr}_{F^\bullet}(\mathcal{D}_X)$. The Koszul complex associated with $\Xi_f \subset \text{Gr}_{F^\bullet}(\mathcal{D})$ is the stalk at p of \mathbf{K} .

We proceed by induction on the dimension of the ambient variety. If that dimension is 1, $\Xi_f = \mathcal{O}f$, and the Koszul complex associated with f is clearly exact. Suppose the result true if the dimension of the ambient variety is $< n$.

Now, suppose $\dim X = n$.

Let $\delta_1, \dots, \delta_{n-1}$ be a basis of $\tilde{\Theta}_f$ in some small enough neighborhood U of p . According to proposition 4.1, we need to prove that \mathbf{K} is exact on $U \setminus \{p\}$.

For every $q \in U$ with $f(q) \neq 0$, the germ of f at q is a unit and by proposition 5.3, the complex \mathbf{K} is exact at q .

Let q be a point in $D = \{f = 0\}$, $q \neq p$. By proposition 5.2, there are local coordinates $z_1, \dots, z_n \in \mathcal{O}_{X,q}$ and a quasi-homogeneous polynomial $G'(t_1, \dots, t_{n-1}) \in \mathcal{O}_{\mathbb{C}^{n-1},0}$ in $n-1$ variables which is also a locally quasi-homogeneous free germ in $\mathcal{O}_{\mathbb{C}^{n-1},0}$ and such that $f_q = G'(z_1, \dots, z_n)$.

Let $G(t_1, \dots, t_n) \in \mathcal{O}_{\mathbb{C}^n,0}$ be the same polynomial as $G'(t_1, \dots, t_{n-1})$ but considered in n variables. The exactness of \mathbf{K}_q is then equivalent to the exactness of the Koszul complex associated with $\Xi_G \subset \text{Gr}_{F^\bullet} \mathcal{D}_{\mathbb{C}^n,0}$.

Let's write $\mathcal{O}_m = \mathbb{C}\{t_1, \dots, t_m\}$ and call ξ'_i the principal symbol of $\frac{\partial}{\partial t_i}$.

Let $\{\delta'_1, \dots, \delta'_{n-2}\} \subset \bigoplus_{i=1}^{n-1} \mathcal{O}_{n-1} \frac{\partial}{\partial t_i}$ be a basis of $\Theta_{G'}$. A basis of Θ_G is then $\{\delta'_1, \dots, \delta'_{n-2}, \frac{\partial}{\partial t_n}\} \subset \bigoplus_{i=1}^n \mathcal{O}_n \frac{\partial}{\partial t_i}$. Call σ'_i the principal symbol of δ'_i , $i = 1, \dots, n-2$.

By induction hypothesis we know that the Koszul complex associated with $\Xi_{G'} \subset \text{Gr}_{F^\bullet} \mathcal{D}_{\mathbb{C}^{n-1},0} = \mathcal{O}_{n-1}[\xi'_1, \dots, \xi'_{n-1}]$ is exact or, equivalently, that $\sigma'_1, \dots, \sigma'_{n-2}, G'$ is a regular sequence in $\mathcal{O}_{n-1}[\xi'_1, \dots, \xi'_{n-1}]$. That implies that $\sigma'_1, \dots, \sigma'_{n-2}, \xi'_n, G = G'$ is a regular sequence in $\mathcal{O}_n[\xi'_1, \dots, \xi'_n]$, i.e. that the Koszul complex associated with $\Xi_G \subset \text{Gr}_{F^\bullet} \mathcal{D}_{\mathbb{C}^n,0}$ is exact, and the result is proved.

For the second property, we filter the Spencer complex associated with $\Xi_f \subset \mathcal{D}$ as in [10], prop. 2.3.4:

$$\deg(\Theta_f) = 1, \deg(f) = 0.$$

Its graded complex coincides with the Koszul complex associated with $\Xi_f \subset \text{Gr}_{F^\bullet}(\mathcal{D})$, and then the Spencer complex is exact. To conclude, we use corollary 5.8, (c). C.Q.D.

6 Examples and questions

We know several (related) kind of free divisors:

[LQH] Locally quasi-homogeneous (definition 1.3).

[EH] Euler homogeneous (definition 1.2).

[LCT] Free divisors satisfying the logarithmic comparison theorem.

[KF] Koszul free (definition 1.6).

[P] Free divisors such that the complex $\Omega_X^\bullet(\log D)$ is a perverse sheaf.

We have then the following implications:

$$\begin{aligned} [\text{LQH}] &\Rightarrow [\text{EH}] \text{ (obvious),} & [\text{LQH}] &\Rightarrow [\text{LCT}] \text{ by [7], th. 1.1,} \\ [\text{LCT}] &\Rightarrow [\text{P}], \text{ by [15], II, th. 2.2.4)} & [\text{KF}] &\Rightarrow [\text{P}] \text{ by [4], th. 4.2.1,} \\ & & & [\text{LQH}] \Rightarrow [\text{KF}] \text{ by theorem 4.3.} \end{aligned}$$

Example 6.1.– (Free divisors in dimension 2) We recall theorem 3.9 from [5]: Let X be a complex analytic manifold of dimension 2 and $D \subset X$ a divisor. The following conditions are equivalent:

1. D is Euler homogeneous.

2. D is locally quasi-homogeneous.
 3. The logarithmic comparison theorem holds for D .
- Consequently, in dimension 2 we have:

$$[\text{LQH}] \Leftrightarrow [\text{EH}] \Leftrightarrow [\text{LCT}]$$

and [KF] (cf. 1.11, 3) and [P] ([4]) always hold. In particular,

$$[\text{KF}] \not\Rightarrow [\text{LQH}], [\text{EH}], [\text{LCT}].$$

Examples of plane curves not satisfying logarithmic comparison theorem are, for instance, the curves of the family (cf. [5]):

$$x^q + y^q + xy^{p-1} = 0, \quad p \geq q + 1 \geq 5.$$

Example 6.2.– (An example in dimension 3) Let's consider $X = \mathbb{C}^3$ and $D = \{f = 0\}$, with $f = xy(x + y)(x + yz)$ [4]. A basis of $\text{Der}(\log D)$ is $\{\delta_1, \delta_2, \delta_3\}$, with

$$\begin{aligned} \delta_1 &= xy\partial_x + y^2\partial_y - 4(x + yz)\partial_z, \\ \delta_2 &= x(x + 3y)\partial_x - y(3x + y)\partial_y + 4x(z - 1)\partial_z, \\ \delta_3 &= x\partial_x + y\partial_y \end{aligned}$$

the determinant of the coefficients matrix being $-16f$ and

$$\delta_1(f) = 0, \quad \delta_2(f) = 0, \quad \delta_3(f) = 4f.$$

In particular, D is Euler homogeneous ($E = (1/4)\delta_3$) and we know [5] that it satisfies the logarithmic comparison theorem. Let $I \subset \mathcal{O}_{T^*X}$ be the ideal generated by the symbols $\{\sigma_1, \sigma_2, \sigma_3\}$ of the basis of $\text{Der}(\log D)$. By corollary 1.9, D is not Koszul free, because the dimension of $V(I)$ at $((0, 0, \lambda), 0) \in T^*X$ is 4, and neither is D locally quasi homogeneous.

So:

$$[\text{LCT}] \not\Rightarrow [\text{KF}], [\text{LQH}], \quad [\text{EH}] \not\Rightarrow [\text{KF}], [\text{LQH}].$$

Finally, for the only missing relation, we quote the following conjecture from [5]:

Conjecture 6.3.– If the logarithmic comparison theorem holds for D , then D is Euler homogeneous.

Example 6.4.— Let's see in the example 6.2 that the left ideal $\text{Ann}_{\mathcal{D}}(f^s)$ is not generated by Θ_f and then, J_f is not an ideal of linear type.

Here, we set $X = \mathbb{C}^3$, $p = (0, 0, 0)$ and $E = (1/4)\delta_3$. The \mathcal{O} -modules Θ_f and $\text{Der}(\log f) = \Theta_f \oplus \mathcal{O} \cdot E$ are generated by $\{\delta_1, \delta_2\}$ and $\{\delta_1, \delta_2, E\}$, respectively. The symbols $\sigma_1 = \sigma(\delta_1)$, $\sigma_2 = \sigma(\delta_2)$ form a $\text{Gr}_{F^\bullet}(\mathcal{D})$ -regular sequence (the proof is analogous to example 1.11, 3)). Then, as in the proof of [4], prop. 4.1.2, we have

$$\sigma(\mathcal{D}\Theta_f) = \text{Gr}_{F^\bullet}(\mathcal{D})\sigma(\Theta_f).$$

For

$$P = 2y^2\partial_x\partial_y - 2y^2\partial_y^2 - (2xz + 6yz)\partial_x\partial_z + 10yz\partial_y\partial_z + 8z(1-z)\partial_z^2 + \\ +(2x - 4y)\partial_y\partial_z - x\partial_x - y\partial_y - 8z\partial_z + 4\partial_z$$

and $R = \mathbb{C}[x, y, z]$, $S = R[\xi_1, \xi_2, \xi_3]$, $\mathfrak{m} = R(x, y, z)$ we check that

1. $P \in \text{Ann}_{\mathcal{D}_X}(f^s)$,
2. $(S(\sigma_1, \sigma_2) : \sigma(P)) = S(x, y)$, and then $(S(\sigma_1, \sigma_2) : \sigma(P)) \cap R = R(x, y)$.

So, $\sigma(P) \notin R_{\mathfrak{m}}[\xi_1, \xi_2, \xi_3]\sigma(\Theta_f)$ and, by faithful flatness,

$$\sigma(P) \notin \mathcal{O}[\xi_1, \xi_2, \xi_3]\sigma(\Theta_f) = \text{Gr}_{F^\bullet}(\mathcal{D})\sigma(\Theta_f).$$

We conclude that $P \notin \mathcal{D}\Theta_f$.

Problem 6.5.— We do not know whether a free divisor defined by a quasi-homogeneous polynomial (with strictly positive weights) is locally quasi-homogeneous.

Problem 6.6.— We do not know any example of a free divisor $D \subset X$ whose logarithmic de Rham complex $\Omega_X^\bullet(\log D)$ is not perverse.

References

- [1] A.G. Aleksandrov. Nonisolated hypersurface singularities. In *Theory of singularities and its applications*, volume 1 of *Adv. Soviet Math.*, pages 211–246. A.M.S., Providence, R.I., 1990.
- [2] J. Bernstein. The analytic continuation of generalized functions with respect to a parameter. *Funk. Anal. Appl.*, 6 (1972), 26–40.

- [3] F.J. Calderón Moreno. Operadores diferenciales logarítmicos con respecto a un divisor libre. Univ. Sevilla, July 1997. Ph.D.
- [4] F.J. Calderón-Moreno. Logarithmic Differential Operators and Logarithmic De Rham Complexes Relative to a Free Divisor. *Ann. Sci. E.N.S.*, 32 (1999), 577-595.
- [5] F.J. Calderón-Moreno, D.Q. Mond, L. Narváez-Macarro and F.J. Castro-Jiménez. Logarithmic Cohomology of the Complement of a Plane Curve. *Comment. Math. Helv.*, 77 (2002), 24-38.
- [6] F.J. Calderón-Moreno and L. Narváez-Macarro. Locally quasi-homogeneous free divisors are Koszul free. *Proc. Steklov Inst. Math.*, 238 (2002), to appear.
- [7] F.J. Castro-Jiménez, D. Mond and L. Narváez-Macarro. Cohomology of the complement of a free divisor. *Transactions of the A.M.S.*, 348 (1996), 3037–3049.
- [8] P. Deligne. Équations Différentielles à Points Singuliers Réguliers, *Lect. notes in Math.*, 163, Springer-Verlag, Berlin-Heidelberg, 1970.
- [9] R. Ephraïm. Isosingular loci and the Cartesian product structure of complex analytic singularities. *Trans. Amer. Math. Soc.*, 241 (1978), 357–371.
- [10] M. Gros and L. Narváez-Macarro. Cohomologie évanescence p-adique: calculs locaux. *Rendiconti Sem. Mat. Univ. Padova*, 104 (2000), 71–90.
- [11] A. Grothendieck. On the de Rham cohomology of algebraic varieties. *Publ. Math. de l'I.H.E.S.*, 29 (1966), 95-103.
- [12] E.J.N. Looijenga. *Isolated singular points on complete intersections*, London Mathem. Soc. Lect. Notes Series, 77. Cambridge Univ. Press, Cambridge, 1984.
- [13] Ph. Maisonobe. \mathcal{D} -modules: an overview towards effectivity. In *Computer algebra and differential equations*, volume 193 of *London Math. Soc. Lecture Note Ser.*, pages 21–55. Cambridge Univ. Press, 1994.
- [14] H. Matsumura. Commutative ring theory, Cambridge University Press, Cambridge, 1992.
- [15] Z. Mebkhout. Le formalisme des six opérations de Grothendieck pour les \mathcal{D}_X -modules cohérents, *Travaux en cours*, 35, Hermann, Paris, 1989.

- [16] K. Saito. On the uniformization of complements of discriminant loci. Preprint, Williams College, 1975.
- [17] K. Saito. Theory of logarithmic differential forms and logarithmic vector fields. *J. Fac. Sci. Univ. Tokyo*, 27 (1980), 265–291.
- [18] W.V. Vasconcelos. *Computational methods in commutative algebra and algebraic geometry, Algorithms and Computation in Mathematics, 2*. Springer Verlag, New York, 1998.
- [19] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge studies in advanced mathematics*. Cambridge University Press, Cambridge, 1994.

Departamento de Álgebra, Facultad de Matemáticas, Universidad de Sevilla,
P.O. 1160, 41080 Sevilla, Spain.

E-mail:

calderon@algebra.us.es

narvaez@algebra.us.es