

Homogenization of the Poisson equation with Dirichlet conditions in random perforated domains

Carmen Calvo-Jurado¹, Juan Casado-Díaz² and Manuel Luna-Laynez²

¹ *Department of Mathematics, University of Extremadura*

² *Department of Differential Equations and Numerical Analysis, University of Sevilla*

emails: ccalvo@unex.es, jcasadod@us.es, mllaynez@us.es

Abstract

Key words: homogenization, random perforated domains

MSC 2000: 35R60, 35B27

1 Extended abstract

For a bounded open set $O \subset \mathbb{R}^N$, and a sequence of open sets $O_\varepsilon \subset O$, we consider the Poisson equation with Dirichlet conditions on ∂O_ε . The problem is to study the asymptotic behavior of the solutions, when ε tends to zero, by finding the equation satisfied by its limit. This allows us to describe the macroscopic behavior of the material corresponding to the mixture of the solid part O_ε and the holes $K_\varepsilon = O \setminus O_\varepsilon$. The obtention of this limit equation is a classical problem which has been considered since the early 80's. Usually, the set K_ε is a union of very small connected components distributed in O . It is well known that the homogenized equation is an elliptic problem in the whole of O which contains in general a new term of order zero depending on the distribution and size of the holes, the Cioranescu-Murat “strange term” ([6]). The most classical result corresponds to the homogenization of the Poisson equation with holes of size $\varepsilon^{\frac{N}{N-2}}$ if $N \geq 3$, or $\varepsilon^{-\frac{1}{\varepsilon^2}}$ if $N = 2$, which are periodically distributed with period ε . In this case the coefficient corresponding to the new term of zero order is a positive constant. The case of arbitrary holes and nonlinear equations has been considered in several papers such as [4], [5], [6], [7], [9], [10], [16].

Our purpose in the present work is to consider the case where the holes are randomly distributed. To simplify, we assume $N \geq 3$, and similarly to the classical periodic setting, we consider holes of size $\varepsilon^{\frac{N}{N-2}}$ such that the distance between two holes is of order ε . Namely:

We consider a probability space (Ω, \mathcal{F}, P) , a subset $\tilde{\Omega} \subset \Omega$, and a function $T(x) : \Omega \rightarrow \Omega$, for every $x \in \mathbb{R}^N$, which defines an ergodic measure preserving dynamical system in \mathbb{R}^N . Then, for a compact set $K \subset \mathbb{R}^N$, we define the sequence of random holes as

$$K_\varepsilon(\omega) = \bigcup_{z \in \mathbb{R}^N, T(z)\omega \in \tilde{\Omega}} \left(\varepsilon z + \varepsilon^{\frac{N}{N-2}} K \right), \quad P\text{-a.e. } \omega \in \Omega.$$

Denoting by $O_\varepsilon(\omega)$ the open set obtained from O by removing the random holes, $K_\varepsilon(\omega)$, we want to study the asymptotic behavior of the solutions u_ε of

$$\begin{cases} -\Delta_x u_\varepsilon(\omega, x) = f_\varepsilon(\omega, x) & \text{in } O_\varepsilon(\omega) \\ u_\varepsilon(\omega, x) = 0 & \text{on } \partial O_\varepsilon(\omega) \end{cases} \quad P\text{-a.e. } \omega \in \Omega, \quad (0.1)$$

where f_ε converges strongly in $L^2_P(\Omega; H^{-1}(O))$ to a function f .

The homogenization of random problems has been considered in several papers, see e.g. [2], [3], [8], [11], [12], [14], [15]. In particular, the G. Nguetseng and G. Allaire two-scale convergence method ([1], [13]), which is a very useful tool in periodic homogenization, has been extended in [2] to the setting of random homogenization problems. In general, the heterogeneities considered in those papers are given by functions of the form $F(T(\frac{x}{\varepsilon})\omega)$, with ω taking values on the probability space, T a dynamical system as above and F a random variable. However, in our problem, the homogenization process contains two different sizes ($\varepsilon^{\frac{N}{N-2}}$ and ε) to describe it, and then, it can not be analyzed by the results of [2]. To solve this difficulty, we introduce in the present paper a new extension of the two-scale method, which is based on some ideas used in [4] for the homogenization of Dirichlet elliptic problems in (deterministic) periodic domains. We show that the solution u_ε of (0.1) converges weakly in $L^2_P(\Omega; H^1_0(O))$ to the unique solution u of the problem

$$\begin{cases} -\Delta_x u(\omega, x) + \gamma \kappa u(\omega, x) = f(\omega, x) & \text{in } O \\ u(\omega, x) = 0 & \text{on } \partial O \end{cases} \quad P\text{-a.e. } \omega \in \Omega,$$

where the new term $\gamma \kappa u$ is the equivalent for our random problem of the Cioranescu-Murat strange term in the deterministic case ([6]). Similarly to the classical result, it is given by the capacity κ of the closed set K in \mathbb{R}^N , multiplied by γ , the mean density of holes in O . Thanks to the ergodic theory we prove that γ does not depend on $\omega \in \Omega$ or $x \in O$. From the physical point of view this means that the limit behavior of the material corresponding to the mixture of the solid part $O_\varepsilon(\omega)$ and the holes $K_\varepsilon(\omega)$ is deterministic. Similar results but assuming different assumptions about the random distribution of the holes and using different techniques have been obtained in [3] and [14].

Acknowledgements

This work has been partially supported by the project MTM 2011-24457 of the “Ministerio de Ciencia e Innovación” of Spain and the research group FQM-309 of the “Junta de Andalucía”.

References

- [1] G. ALLAIRE, *Homogenization and two-scale convergence*, SIAM J. Math. Anal. **23** (1992) 1482–151.
- [2] A. BOURGEAT, A. MIKELIC, S. WRIGHT, *Stochastic two-scale convergence in the mean and applications* J. Reine Angew. Math. **456** (1994) 19-51.
- [3] L.A. CAFFARELLI, A. MELLET, *Random homogenization of an obstacle problem*, Ann. Inst. H. Poincaré Anal. Non Linéaire **26** (2009) 2, 375-395.
- [4] J. CASADO-DÍAZ, *Two-Scale convergence for nonlinear Dirichlet problems in perforated domains*, Proceedings of the Royal Society of Edinburgh A, **130** (2000) 249–276.
- [5] J. CASADO-DÍAZ, A. GARRONI, *Asymptotic behavior of nonlinear elliptic systems on varying domains*, SIAM J. Math. Anal. **31** (2000) 581–624.
- [6] D. CIONARESCU, F. MURAT, *Un terme étrange venu d'ailleurs*, in Nonlinear Partial Differential Equations and Their Applications, Collège de France Seminar, Vols. **II** and **III**, Research Notes in Math. 60 and 70 Pitman, London, 1982, 98–138 and 154–178.
- [7] G. DAL MASO, A. DEFRANCESCHI, *Limits of nonlinear Dirichlet problems in varying domains*, Manuscr. Math. **61** (1988) 251–278.
- [8] G. DAL MASO, L. MODICA, *Nonlinear stochastic homogenization and ergodic theory*, J. Reine Angew. Math. **368** (1986) 28–42.
- [9] G. DAL MASO, U. MOSCO, *Wiener-criterion and Γ -convergence*, Appl. Math. Optim. **15** (1987) 15–63.
- [10] G. DAL MASO, F. MURAT, *Asymptotic behaviour and correctors for the Dirichlet problems in perforated domains with homogeneous monotone operators*, Ann. Sc. Norm. Sup. Pisa **4** (1997) 239–290.
- [11] V.V. JIKOV, S.M. KOZLOV, O.A. OLEINIK *Homogenization of differential operators and integral functionals*, Springer-Verlag, Berlin, 1994.

- [12] S.M. KOZLOV, *Homogenization of random operators*, Math. U.S.S.R. Sb. **37** (1980) 167–180.
- [13] G. NGUETSENG, *A general convergence result for a functional related to the theory of homogenization*, SIAM J. Math. Anal. **20** (1989) 608–623.
- [14] G.C. PAPANICOLAOU, S.R.S. VARADHAN, *Diffusion in regions with many small holes*, in *Stochastic differential systems*, Lecture Notes in Control and Information Sci. Springer, Berlin-New York, **25** (1980).
- [15] G.C. PAPANICOLAOU, S.R.S. VARADHAN, *Boundary value problems with rapidly oscillating random coefficients*, Colloq. Math. Soc. J. Bolyai, North-Holland, Amsterdam-New York, **27** (1981) 835–873.
- [16] I.V. SKRYPNIK, *Averaging of nonlinear Dirichlet problems in punctured domains of general structure*, Mat. Sb. 187, **8** (1996) 125–157.