

1  
2  
3 **Thermoluminescence as a complementary technique for the toxicological**  
4 **evaluation of chemicals in photosynthetic organisms**  
5  
6

7  
8 Guillermo Repetto<sup>a</sup>, Jorge L. Zurita<sup>a,b</sup>, Mercedes Roncel<sup>c</sup>, José M. Ortega<sup>c</sup>  
9

10  
11  
12 <sup>a</sup>Departamento de Biología Molecular e Ingeniería Bioquímica, Universidad Pablo de  
13 Olavide, Carretera de Utrera km. 1, 41013 Seville, Spain  
14  
15

16  
17 <sup>b</sup>Present address: Fundación Progreso y Salud, Junta de Andalucía, Américo Vespucio 5,  
18 41092 Seville, Spain  
19  
20

21  
22 <sup>c</sup>Instituto de Bioquímica Vegetal y Fotosíntesis, Universidad de Sevilla-CSIC, Américo  
23 Vespucio 49, 41092 Seville, Spain  
24  
25

26  
27  
28  
29 Corresponding author: G. Repetto, Departamento de Biología Molecular e Ingeniería  
30 Bioquímica, Área de Toxicología, Universidad Pablo de Olavide, Carretera de Utrera km. 1,  
31 41013 Seville, Spain. Fax: 34-954349813; Email: grepkuh@upo.es  
32  
33  
34  
35  
36

37  
38 **Keywords:** *Chlorella vulgaris*; chemical toxicity; lipid peroxidation, photosystem II;  
39 thermoluminescence  
40  
41  
42

43 **Abbreviations:** B-band: TL band due to  $S_{2/3}Q_B^-$  recombination; BrB: bromobenzene; C-band:  
44 TL band due to  $D^+Q_A^-$  recombination; CQ: chloroquine; D: tyrosine D of photosystem II;  
45 DEA: diethanolamine;  $EC_{50}$ : concentration of test chemical that modified the biomarker by  
46 50% in comparison with appropriate untreated control; HTL: high temperature  
47 thermoluminescence;  $InNO_3$ : indium nitrate; PpG: propyl gallate; PSII: photosystem II; Q-  
48 band: TL band due to  $S_2Q_A^-$  recombination;  $Q_A$  and  $Q_B$ : primary and secondary quinonic  
49 electron acceptors of PSII;  $S_2$  and  $S_3$ : oxidized states of the manganese oxygen evolving  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

complex of PSII; SMFA, sodium monofluoroacetate; STL: standard thermoluminescence; TL: thermoluminescence;  $t_{\max}$ : temperature of the maximum intensity of a TL band.

## Abstract

Thermoluminescence is a simple non-invasive technique very useful for study electron transfer reactions on photosystem II (standard thermoluminescence) or the level of lipid peroxidation in membranes (high temperature thermoluminescence) in photosynthetic organisms. Both techniques were used to investigate the effects produced on *Chlorella vulgaris* cells by six compounds: the chemical intermediates bromobenzene and diethanolamine, the antioxidant propyl gallate, the semiconductor indium nitrate, the pesticide sodium monofluoroacetate and the antimalarian drug chloroquine. Electron transfer activity of the photosystem II significantly decreased after the exposure of *Chlorella* cells to all the six chemicals used. Lipid peroxidation was slightly decreased by the antioxidant propyl gallate, not changed by indium nitrate and very potently stimulated by diethanolamine, chloroquine, sodium monofluoroacetate and bromobenzene. For five of the chemicals studied (not bromobenzene) there is a very good correlation between the cytotoxic effects in *Chlorella* cells measured by the algae growth inhibition test, and the inhibition of photosystem II activity. The results suggest that one very important effect of these chemicals in *Chlorella* cells is the inhibition of photosynthetic metabolism by the blocking of photosystem II functionality. In the case of sodium monofluoroacetate, diethanolamine and chloroquine this inhibition seems to be related with the induction of high level of lipid peroxidation in cells that may alter the stability of photosystem II. The results obtained by both techniques supply information that can be used as a supplement to the growth inhibition test and allows a more complete assessment of the effects of a chemical in photosynthetic organisms of aquatic ecosystems.

## 1. Introduction

Model systems and bioassays are used in environmental toxicology to provide information for risk assessment of chemicals. To improve the efficiency in evaluating the hazard of a large number of chemicals, providing mechanistic information, it would be helpful to carry out batteries of rapid and mode of action based screening tests (Katsumata et al., 2009; Repetto, 2013).

Standard (STL) and high temperature thermoluminescence (HTL) measurements are useful techniques for the study of the effects of stress conditions in photosynthetic organisms (Ducruet, 2003; Havaux, 2003). Its use in aquatic toxicology has been however very limited. Luminescence is a general term which indicates a light emitting process when the molecules in an electronically excited state decay to the stable ground state. Luminescence from organic or inorganic material can be also thermally induced and enhanced by heating the sample in the dark. This process is called thermoluminescence and describes the emission of light at characteristic temperatures from samples containing chemiluminescence active species, radical pair states or electron hole pairs (Ducruet, 2003; Havaux, 2003).

STL is normally used to study the photochemical activity of photosystem II (PSII). PSII is a large pigment-protein complex which uses light energy to water splitting producing molecular oxygen as by-product and providing an unlimited supply of reducing power needed to convert CO<sub>2</sub> into the organic molecules of life (for a review see Diner and Babcock, 1996). The reaction centre (RC) of PSII is composed of an heterodimer formed by the D1 and D2 membrane proteins that bears all the cofactors needed for charges separation and stabilization: six chlorophylls (Chl) (including four of the Chl centre, P680), two pheophytins, two plastoquinones Q<sub>A</sub> and Q<sub>B</sub>, one non-heme iron atom, one tetranuclear manganese cluster (Mn<sub>4</sub>CaO<sub>5</sub>) and two redox active tyrosines. Photons are captured by the light-harvesting pigments and driven as excitation energy to P680 forming an excited state called P680\*. The

1 P680\* state is highly reducing, and an electron is rapidly transferred to the terminal  
2 plastoquinone (PQ) electron acceptor bound to the Q<sub>B</sub> site within the D1 protein. On receiving  
3 two electrons, the PQ is protonated to plastoquinol (PQH<sub>2</sub>) and leaving the Q<sub>B</sub> site in the PSII  
4 complex. The P680 radical cation (P680<sup>+</sup>) formed in primary charge separation has a very  
5 high redox potential, estimated to be approx. +1.3 V. This high-oxidation state is required to  
6 drive the water-splitting reaction that occurs at a catalytic site, the manganese cluster (for a  
7 review see Rappaport et al., 2002; Diner and Rappaport, 2002).  
8  
9

10  
11  
12  
13  
14  
15  
16  
17 The photochemical activity of PSII complex can be measured by using the STL  
18 technique. Photosynthetic luminescence is a weak light emission originating from  
19 recombination of positive and negative charge pairs generated by a prior illumination in PSII,  
20 some of these leading to the recreation at a low yield of an exciton in the chlorophyll antenna,  
21 with a probability to deactivate as fluorescence. Luminescence decay phases can be better  
22 resolved by TL emission technique, which consists in illuminating the sample at a  
23 temperature sufficiently low to make negligibly small the recombination rate of the charge  
24 pairs under investigation, then to reveal them successively as TL bands by a progressive  
25 warming (for a review see Vass and Inoue, 1992; Ducruet, 2003; Sane, 2004).  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 After a sequence of short flashes, a so-called B-band of TL, located between 15 and 40  
40 °C depending on species and conditions, is observed in healthy dark-adapted photosynthetic  
41 material, such as leaves and algae. This emission is due to the recombination of S<sub>2</sub>/S<sub>3</sub>Q<sub>B</sub><sup>-</sup>  
42 pairs, Q<sub>B</sub> being the secondary quinonic electron acceptor and S<sub>2</sub>/S<sub>3</sub>, the states of the oxygen-  
43 evolving complex storing positive charges (Rutherford et al., 1982; Demeter and Vass, 1984).  
44  
45  
46  
47  
48  
49  
50  
51 The changes on the intensity and temperature maximum of this band reflect alterations in the  
52 functionality of PSII complex. Herbicide treatment (diuron, atrazine), which blocks the Q<sub>A</sub> to  
53 Q<sub>B</sub> electron transfer, induces the appearance of a Q-band peaking at about 5 °C, due to the  
54 S<sub>2</sub>Q<sub>A</sub><sup>-</sup> recombination. A C-band at about 55 °C can also be detected in particular conditions.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 This TL band is due to  $D^+Q_A^-$  recombination (Demeter et al., 1993, Johnson et al., 1994),  $D^+$   
2 being the oxidized form of tyrosine D on the inactive branch of PSII.  
3

4 STL measurements are useful for the study of the effects of stress conditions on PSII  
5 electron transport in intact leaves, in algal and cyanobacterial cells, as well as in thylakoids  
6 and isolated membrane complexes (Rutherford and Inoue, 1984; Rutherford et al., 1984a;  
7 Zurita et al., 2005c; Ducruet and Vass, 2009). The technique is simple, inexpensive and  
8 powerful in monitoring several aspects of the PSII electron transport. The fact that alterations  
9 in the peaks of the TL curves may be interpreted as changes in the redox potentials of donors  
10 and acceptors on PSII makes this technique a useful tool for understanding the process of  
11 adaptation, responses to abiotic and biotic stresses, and structural modifications in the PSII  
12 polypeptides.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

25 HTL is a useful technique for the study of lipid peroxidation processes in  
26 photosynthetic organisms (Havaux, 2003). Lipid peroxidation is a complex process known to  
27 occur in both plants and animals. This process involves a chain reaction initiated by the  
28 hydrogen abstraction or addition of an oxygen radical, a rearrangement of the double bonds in  
29 unsaturated lipids and the formation and propagation of lipid radicals. As a result of these  
30 reactions takes place an eventual destruction of membrane lipids with the production of a  
31 variety of breakdown products, including alcohols, ketones, alkanes, aldehydes and ethers  
32 (Dianzani and Barrera, 2008). In stress situations the reactive oxygen and nitrogen species are  
33 generated at higher than normal rates, and as a consequence, lipid peroxidation occurs. In  
34 photosynthetic organisms a large amount of oxygen is formed in thylakoids in the vicinity of  
35 powerful oxidation-reduction systems that readily reduce oxygen to dangerous superoxide and  
36 produce harmful singlet oxygen which could initiate lipid peroxidation processes (Dalton,  
37 1995; Repetto et al., 2012).  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 The lipid peroxidation is commonly detected and quantified by measuring by-products  
2 of the reaction, such as ethane, conjugated diens, malondialdehyde, and other carbonyl  
3 compounds (Heath and Packer, 1968; Pryor and Castle, 1984; Chatterjee and Agarwal, 1988;  
4 Havaux and Niyogi, 1999). The most commonly used test is called TBARS Assay  
5 (thiobarbituric acid reactive substances assay) where the concentration of malondialdehyde is  
6 determined by reaction with thiobarbituric acid to yield a fluorescent product. However, there  
7 are other sources of malondialdehyde, so this test is not completely specific for lipid  
8 peroxidation. In recent years, the development of immunochemical detection of 4-  
9 hydroxynonenal histidine adducts has opened more advanced methodological possibilities for  
10 qualitative and quantitative detection of lipid peroxidation in various human and animal  
11 tissues (Weber et al., 2013). These chemical methods cannot be however routinely used for  
12 the assessment of lipid oxidation in phytoplankton because the analysis requires high amounts  
13 of biological material, and they are often time-consuming and, therefore, not adequate for  
14 high throughput screening. The second problem is the difficulty of separating the peroxidative  
15 reactions that take place in phytoplankton from the reactions occurring in no phytoplankton  
16 materials.

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39 Recently, a new TL-based technique has been proposed as a useful method to detect  
40 lipid peroxidation in photosynthetic organisms that may solve some of the aforementioned  
41 problems (Vavilin and Ducruet, 1998; Havaux, 2003; Ducruet and Vass, 2009). When  
42 chlorophyll-containing material is gradually heated to temperatures above 70 °C, a rise of  
43 chlorophyll luminescence emission is often observed. This increase in the luminescence  
44 intensity at high temperature was named HTL (Matorin et al., 1989; Venediktov et al., 1989).  
45 A broad HTL band peaking around 135 °C (HTL2 band) is generated in the reaction of  
46 thermal decomposition of peroxidized lipids that, in turn, leads to the formation of carbonyls  
47 in triplet state followed by migration of excitation energy toward chlorophylls (Venediktov et  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

al., 1989; Vavilin et al., 1991; Vavilin and Ducruet, 1998). Its amplitude has been correlated with the accumulation of malodialdehyde, an indicator of lipid peroxidation in standard chemical tests (Vavilin and Ducruet, 1998). Consequently, HTL2 band measurements have been used to study *in vivo* lipid peroxidation in plants subjected to various oxidative stress conditions (Merzlyak et al., 1992; Havaux and Niyogi, 1999; Ducruet, 2003; Baroli et al., 2004; Havaux, 2003). In a future perspective HTL might be applied as a fast screening tool before more sophisticated methods like mass spectrometry are applied to identify the most sensitive molecular target in a given tissue.

STL and HTL techniques have been applied in several *in vitro* toxicological studies, mainly using thylakoids and cell cultures (Ducruet et al., 2007). Using thylakoids isolated from leaves of greenhouse grown peas (*Pisum sativum L.*), Mohanty et al. (1989) suggested that Cu (II) does not block electron flow between the primary and secondary quinonic electron acceptor but modifies the secondary quinonic acceptor site in such a way that it becomes unsuitable for further PSII photochemistry. Horváth et al. (1998) reported that the amplitude of the B-band was gradually decreased, which corresponds to the Cu (II) induced inactivation of Tyrosine Z. The simultaneous stepwise shift in the peak position of the B-band indicated, however, that  $S_3Q_B^-$  charge recombination is more resistant to Cu (II). Nitrite, but not formate, fluoride or nitrate anions, inhibited PSII in spinach thylakoids, probably at the donor side (Sahay et al., 2006).

Toxicological applications of HTL have been reported by several authors. A rise of thermoluminescence at 120-130 °C was detected when cultures of the unicellular green alga *Chlorella (C.) pyrenoidosa* were exposed for 5 h to 10 µM of the oxidative stress promoter methyl viologen (Vavilin and Ducruet, 1998). Hörcsik et al. (2007) found photodestruction of D1 protein in *C. pyrenoidosa* caused by Cr (VI) due to enhanced oxidative stress and lipid peroxidation, as indicated by the appearance of a HTL2 band. Schumann et al. (2009) have

1 developed a rapid and easy to handle TL based technique for evaluation of carbon  
2 tetrachloride-induced oxidative stress on rat hepatocytes.  
3

4 The aim of this work has been the application of two simple, non-invasive and  
5 inexpensive TL techniques (STL and HTL) to study the alterations produced on the fresh-  
6 water green alga *C. vulgaris* by six chemicals. This green alga was selected as a representative  
7 of primary producers. It is a genus of single-cell green algae, widely distributed in freshwater  
8 and seawater. It has a short cycle of growth, which makes it ideal for aquatic ecotoxicity  
9 studies (Wong et al., 1997). The chemical intermediates bromobenzene and diethanolamine,  
10 the antioxidant propyl gallate, the semiconductor indium nitrate, the pesticide sodium  
11 monofluoroacetate and the antimalarian drug chloroquine were selected as representative of  
12 several chemical groups according to their use. The results have been compared with those  
13 obtained by our group by using the algae growth inhibition test (Zurita et al., 2005a,b; Zurita  
14 et al., 2007a,b,c,d).  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

## 31 **2 Material and Methods**

### 32 *2.1 Cultures*

33 The unicellular green algae *Chlorella vulgaris* var. *viridis*, kindly provided by Dr. Muñoz-  
34 Reoyo (CISA, Spain), were grown as previously described (Ramos et al., 1996) in Bold's  
35 Basal Medium (BB medium) (Culture Collection of Algae and Protozoa, Cambridge, UK;  
36 <http://www.ccap.ac.uk/media/recipes.htm>) under a water-saturated sterile atmosphere  
37 containing 5% CO<sub>2</sub> at 22 °C, constant agitation and a light dark cycle of 16:8 h (white light of  
38 40 μE m<sup>-2</sup> s<sup>-1</sup>).  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



## 2.2 Chemicals

Stock solutions of bromobenzene (CAS number 108-86-1), indium nitrate (CAS number 13770-61-1), sodium monofluoroacetate (CAS number 62-74-8), diethanolamine (CAS number 111-42-2) and chloroquine (CAS number 54-05-7) were prepared daily in deionised water. Propyl gallate (CAS number 121-79-9) was solubilised in dimethyl sulfoxide. All the chemicals were from Sigma-Aldrich (USA), unless sodium monofluoroacetate (Honeywell's Riedel-de Haën). They were sonicated for 30 min and sterilized by filtration through a 0.22  $\mu\text{m}$  Millipore filter. The concentrations of the chemicals used in this work were chosen taking into account the  $\text{EC}_{50}$  values obtained for growth inhibition (biomarker) of *C. vulgaris* cells after 24 h exposure to the different chemicals, as previously published (Zurita et al., 2005a,b; Zurita et al., 2007a,b,c,d). The  $\text{EC}_{50}$  value is the concentration of the chemical that modified the biomarker (growth inhibition, for *C. vulgaris*) by 50% in comparison with appropriate untreated control. Two different concentrations of chemicals were used, one below  $\text{EC}_{50}$  values and the second one above or close to the  $\text{EC}_{50}$  values. The two concentration used and the  $\text{EC}_{50}$  values for the six chemicals are shown in Table 1. STL and HTL experiments were carried out using cultures of *C. vulgaris* in exponential phase of growth (with about  $10^6$  cell  $\text{ml}^{-1}$ ). The cultures were exposed in BB medium to the two different concentrations of chemicals selected in 80 ml flasks for 24 h at 20 °C using constant agitation at 40 r.p.m., under a continuous cold light source of  $110 \mu\text{E m}^{-2} \text{s}^{-1}$ . The suspensions were then centrifuged at 2000 r.p.m for 10 min, quantified at 665 nm and adjusted to 0.1 mg chlorophyll  $\text{ml}^{-1}$  by using BB medium. Control experiments in the absent of the chemicals were also treated as above described.

## 2.3 Thermoluminescence measurements

1 TL glow curves of *Chlorella* cell suspensions were obtained using two similar home-built  
2 devices designed by Dr. Ducruet (France) for luminescence detection from 0 °C to 80 °C  
3 (STL) and from 10 °C to 160 °C (HTL). A detailed description of these systems can be  
4 obtained elsewhere (Ducruet, 2003; Zurita et al., 2005c; Guerrero et al., 2014). Briefly,  
5 temperature regulation, signal recording and flash sequences were driven by a computer  
6 through a National Instrument DAQ-Pad1200 interface, using dedicated software.  
7 Temperature regulation of the sample cuvette was provided by a double-stage Marlow  
8 thermoelectric Peltier plate (model DT 1089-14; Marlow Industries, USA) cooled by a  
9 temperature-controlled bath. Luminescence emission was detected by a H5701-50  
10 Hamamatsu photomultiplier module. Single turn-over flashes were provided by a xenon white  
11 light (Walz XST-103). Typically, for STL measurements 50 µl *Chlorella* cell suspensions  
12 were dark-incubated for 2 min at 20 °C, then cooled to 1 °C for 1 min and illuminated at the  
13 end of this period with different numbers of saturating single turn-over flashes (separated by 1  
14 s). Luminescence emission was recorded while warming samples from 0 °C to 80 °C at a  
15 heating rate of 0.5 °C per s. For HTL measurements 50 µl *Chlorella* cell suspensions were  
16 adsorbed by filtration on a piece of filter paper (0.45 µm, Whatman) that was pressed against  
17 the copper film, dark-incubated for 10 min at 20 °C, and cooled to 10 °C for 1 min.  
18 Luminescence emission was then recorded while warming samples from 10 °C to 160 °C at a  
19 heating rate of 0.1 °C per s. N<sub>2</sub> gas was flushed on the sample during HTL experiments in  
20 order to desiccate samples and prevents any oxidation induced by high temperatures. The  
21 instruments were driven by a PC computer, with a specially developed acquisition program  
22 (Ducruet 2003 and references therein). Data acquisition, signal analysis and graphical  
23 simulation were performed as previously described (Zurita et al., 2005c; Ducruet and  
24 Miranda, 1992; Ducruet et al., 2011).

## 3 Results

### 3.1 Characterization of light-induced thermoluminescence bands in non-treated *Chlorella vulgaris* cells

Before the TL analysis of the effects of chemicals on *Chlorella* cells, the TL bands “*in vivo*” using healthy cells from cultures of this green alga were characterized by using STL technique. Excitation of dark-adapted non-treated *Chlorella* cell suspensions with a series of saturating single turn-over flashes at 1 °C induced the appearance of TL glow curves with important differences in the total signal intensity. TL curves induced by 1, 2 and 3 flashes are shown in Figure 1 as examples. The light emission curve obtained after illumination with two flashes was the best of the series and showed a  $t_{\max}$  at about 15 °C and a small band around 58-60 °C (Figure 1, 2F). The signal peaking at about 15 °C could be assigned to the TL B-band, due to the recombination of positive charges stored on  $S_2$  and  $S_3$  states of the oxygen evolving complex with an electron stabilized as  $Q_B^-$  on the secondary quinonic acceptor of PSII (Rutherford et al., 1982). The intensity of B-band exhibited a typical four-oscillation period with maximum after the 2<sup>nd</sup> and 6<sup>th</sup> flashes (data not shown). This pattern suggests that in dark-adapted *Chlorella* cells the ratio  $S_0:S_1$  and  $Q_B:Q_B^-$  are about 25:75% and 50:50%, respectively (Inoue, 1996). Thus, after one single flash, the  $S_1Q_B$  centers will go to the luminescence-emitting state  $S_2Q_B^-$ , generating a B-band peaking at about 25 °C (Figure 1, 1F), generally identified as the B<sub>2</sub>-band component. Two flashes induced the best glow curve because generate an important amount of PSII centers in both  $S_2Q_B^-$  and  $S_3Q_B^-$  luminescence states (Rutherford and Inoue, 1984). Besides, the yield from the latter recombination is higher than that from the former by a factor of 1.7-2.0 (Rutherford et al., 1984b). The much higher contribution of the second component of B-band (B<sub>1</sub>-band) after two flashes, induces the appearance of a TL glow curve significantly shifted to lower temperatures and broadened.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

A second small component of the signal peaking around 58-60 °C appears in all the TL glow curves obtained (Figure 1). We assigned this component to the C-band, originated from the recombination reaction of the positive charges stored on tyrosine D residue and electron stabilized as  $Q_A^-$  on the primary quinonic acceptor of PSII (Rutherford et al., 1982). The appearance of this band suggests that in healthy *Chlorella* cells there was a small PSII population with the electron transfer blocked both on donor and acceptor side.

### 3.2 Effects of chemicals on light-induced thermoluminescence bands in *Chlorella vulgaris* cells

STL technique was used to determine the effects of the six environmental chemicals on PSII electron transfer activity of *Chlorella* cells. In order to compare the effects at doses non-cytotoxic and cytotoxic of the chemicals compound studied in this work, TL experiments were carried out using one concentration below and other close to the  $EC_{50}$  values, respectively (Table 1). Algae cultures in exponential phase of growth were exposed for 24 h to two concentrations of each chemical. TL emissions were then recorded between 0 °C and 80 °C after the illumination of cells suspensions at 1 °C with two flashes (Figure 2).

The exposure of *Chlorella* cells for 24 h to bromobenzene did not significantly change the properties of the light-induced electron transfer on PSII (Figure 2, BrB). The incubation with 1 mM bromobenzene did not alter the TL signal assigned to B-band regarding control sample, while 2 mM bromobenzene decreased the signal intensity only by about 13%. Propyl gallate and indium nitrate caused however a moderate decrease on TL B-band of *Chlorella* cells (Figure 2, PpG and InNO<sub>3</sub>, respectively). The intensity of the TL band was decreased by about 10% and 40% after incubation with the lower and higher concentrations of both chemicals, respectively.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

The total intensity of TL emission of the B-band was not significantly modified by 5 mM of sodium monofluoroacetate (Figure 2, SMFA), but the exposure of the cells to 20 mM concentration induced a decrease of about 30% of the signal. Both concentrations of sodium monofluoroacetate induced the appearance of a more broadened TL glow curve with a significant up-shift of the temperature of the  $t_{\max}$ . TL emissions were also particularly sensitive to the exposure to chloroquine (Figure 2, CQ), showing 50% and 80% reduction of the intensity of B-band with 0.1 and 0.25 mM concentration, respectively.

The incubation of *Chlorella* cells with diethanolamine markedly reduced the electron transfer activity of PSII (Figure 2, DEA). No signal was detected after the treatment with 30 mM of diethanolamine. However, 15 mM of this chemical lowered about 15% the intensity of the B-band and induced the appearance of a more broadened TL glow curve with a shoulder peaking at about 46 °C (Figure 2, DEA). This component appearing at higher temperatures was tentatively assigned to the afterglow (AG) emission band usually induced by far red illumination (Miranda and Ducruet, 1995).

To characterize this band, TL glow curves of *Chlorella* cells suspension were obtained after illumination with only one flash (Figure 3). Under these conditions, the AG-band may be better studied as the B-band has a lower intensity. Thus, in dark-adapted *Chlorella* cell suspensions incubated with 15 mM diethanolamine, the excitation with one saturating single turn-over flash at 1 °C induced the appearance of a more broadened TL glow curve compared with the control in the absence of diethanolamine (Figure 3, DEA 1). A very clear shoulder peaking at about 46 °C was observed in these treated cells which does not appear in the control cells where only one band appears at 25 °C corresponding to B<sub>2</sub>-band. This TL curve could be well simulated by two components, one with  $t_{\max}$  of 25 °C originating from the B<sub>2</sub>-band and a clear second component with  $t_{\max}$  of 45 °C that was assigned to AG-band. The mathematical analysis of the two components allowed estimate a contribution of 48% and

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

52% to the total signal intensity for the B and AG bands, respectively. No TL emission signal due to B-band was obtained in cells incubated with the higher concentration of diethanolamine (30 mM). However, a small band of unknown origin peaking at about 70 °C could be observed. This band also appeared in TL curves induced by two flashes in cells incubated by 30 mM diethanolamine (Figure 2, DEA).

### 3.3 Effects of chemicals on the high temperature thermoluminescence emissions in *Chlorella vulgaris* cells

The effects of the six chemicals studied on the lipids of the photosynthetic membrane of *Chlorella* have been studied by using HTL technique. *Chlorella* cells in exponential phase of growth were exposed for 24 h to two different concentrations of each of the six chemicals. HTL emissions were then recorded in a large temperature range (10 °C-160 °C) at a heating rate of 0.1 °C/s as described in Materials and Methods section. Figure 4 shows that a clear HTL2 band with a  $t_{\max}$  at about 130-150 °C was present in most control and chemical treated samples. Analysis of HTL2 bands obtained, clearly suggested the existence of a high level of lipid peroxidation in photosynthetic membranes of *Chlorella* after the incubation with four of the chemicals investigated: bromobenzene, sodium monofluoroacetate, diethanolamine and chloroquine. *Chlorella* cells treated with 30 mM diethanolamine showed the most important increase in the intensity of the HTL2 band. SMFA at the higher concentration and CQ at the two concentration tested induced also important increase of the HTL2 band. The level of lipid peroxidation increased slightly with 2 mM bromobenzene, in comparison with control sample, and was not modified by either indium nitrate or propyl gallate. Interestingly, this last chemical seemed to slightly reduce the basal level of lipid peroxidation detected in non-treated cells.

## 4 Discussion

1  
2 Thermoluminescence is a technique very simple, inexpensive and powerful to study  
3  
4 some of the processes occurring in photosynthetic organisms. The effects produced by six  
5  
6 chemical compounds on the photosynthetic activity of PSII and lipid peroxidation in  
7  
8 *Chlorella vulgaris* cells were studied by using STL and HTL techniques, respectively. The  
9  
10 results have been compared with those obtained previously by our group (Zurita et al.,  
11  
12 2005a,b; Zurita et al., 2007a,b,c,d) in algae growth inhibition test studies. The additional data  
13  
14 obtained allow a more complete assessment of the toxicological effects of the chemicals  
15  
16 evaluated.  
17  
18  
19  
20

21  
22 Bromobenzene is a colourless liquid used as solvent for crystallization, intermediate in  
23  
24 organic synthesis and additive to motor oils. It can be present in atmospheric, terrestrial and  
25  
26 aquatic environments, being biologically non-degradable. However, there is limited available  
27  
28 information about the toxic hazard of bromobenzene in the environment. It is known that  
29  
30 causes centrilobular cytoplasmic alteration and hepatocyte hypertrophy in mammals (Dodd et  
31  
32 al., 2012). Bromobenzene is not a particularly potent phytotoxic compound showing low  
33  
34 inhibition activity to *Chlorella* cells growth ( $EC_{50} = 2.35$  mM), which is less sensitive than  
35  
36 other environmental receptors, including *Vibrio fischeri* bacteria, the cladocera *Daphnia*  
37  
38 *magna* or fish cell lines (Zurita et al., 2007a). A very slight decrease in PSII functionality was  
39  
40 detected after the treatment of cells with a chemical concentration close to the  $EC_{50}$  (Table 1  
41  
42 and Figure 2, BrB).  
43  
44  
45  
46  
47

48  
49 The 1.4 fold increase of lipid peroxidation by 2 mM bromobenzene observed in  
50  
51 *Chlorella* cells, according to the intensity of HTL2 band (Figure 4, BrB), is in agreement with  
52  
53 the reduction in glutathione levels in several fish cell lines exposed to concentrations higher  
54  
55 than 1 mM bromobenzene and with the increase of reactive oxygen species from 3.2 mM  
56  
57 reported by Schoonen et al. (2005). In addition, Zurita et al. (2007a) observed a maximum  
58  
59  
60  
61  
62  
63  
64  
65

1 stimulation on the antioxidant enzyme glucose-6-phosphate dehydrogenase in the PLHC-1 fish  
2 cell line after a treatment with 1.8 mM bromobenzene, confirming the situation of oxidative  
3 stress. This significant increase in lipid peroxidation induced by bromobenzene in *Chlorella*  
4 cells can be also related to the accelerated membrane phospholipid degradation described by  
5 Lamb et al. (1984). The decrease in the B-band observed after the treatment with the highest  
6 concentration of bromobenzene (Figure 2, BrB) can also be explained taking into account the  
7 effects on the PSII activity of the phospholipid degradation processes. Thus, it seems that the  
8 main cytotoxic effects of BrB seem not to be related with the inhibition of PSII activity. Other  
9 effects on the *Chlorella* metabolism induced by BrB (as lipid peroxidation) must be  
10 responsible for the growth inhibition measured in previous work (Zurita et al., 2007a).  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 Propyl gallate is an antioxidant widely used to preserve and stabilize the freshness,  
25 nutritional value, flavour and colour of foods, fats and oils, cosmetics and medicinal  
26 preparations (Van der Heijden et al., 1986). Propyl gallate is not very toxic to *Chlorella*,  
27 showing much more toxicity in other systems (Zurita et al., 2007c). A consistent decrease in  
28 PSII functionality (B-band) was detected in parallel with cytotoxicity (Table 1 and Figure 2,  
29 PpG). Propyl gallate uncouple oxidative phosphorylation by increasing the permeability of  
30 mitochondrial inner membranes to protons (Fusi et al., 1991). A similar effect on chloroplast  
31 may explain the inhibition of TL B-band of PSII.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 Although there are many studies that demonstrate the antioxidant capacity of propyl  
45 gallate (Karthikeyan et al., 2005), other authors have reported oxidant properties (Kobayashi  
46 et al., 2004; Kawanishi et al., 2005). The mitochondria alternative oxidase, that catalyzes a  
47 cyanide-resistant reduction of oxygen to water without translocation of protons, is specifically  
48 inhibited by propyl gallate, stimulating H<sub>2</sub>O<sub>2</sub> production in plant mitochondria (Josse et al.,  
49 2003). High concentrations of propyl gallate have been reported to produce loss of  
50 mitochondrial membrane potential and to be oxidant in non-vegetal systems due to depletion  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

in glutathione, stimulating glucose-6-phosphate dehydrogenase activity (Zurita et al., 2007c; Han and Park, 2010).

Therefore, in order to clarify this discrepancy, the protective and toxic effects of propyl gallate, we have investigated the possible lipid peroxidation induced by this chemical by using HTL technique (Figure 4, PpG). Our results clearly support the antioxidant properties of propyl gallate. The observed slight reduction on lipid peroxidation (Figure 4, PpG), in comparison to the control, is in agreement with its proposed antioxidant capacity (Kawanishi et al., 2005). In fact it has been recently demonstrated in PSII membranes isolated from spinach leaves, that the scavenging capacity of radicals by propyl gallate significantly suppressed  $^1\text{O}_2$  formed by oxidation of organic molecules (lipids and proteins) (Yadav and Pospíšil, 2012).

Indium nitrate is used in the industry of batteries, to prevent build-up of hydrogen gas within sealed battery casings, plating and as semiconductor in other chemical applications. It is also used for the treatment of Hodgkin and non-Hodgkin's lymphoma and for the management of hypercalcemia (Lipcamon and Sahl, 2002). However, there are no reports on its phytotoxicity. The proliferation of *Chlorella* cells is very sensitive to indium nitrate exposure, showing a parallel inhibition of PSII functionality as shown by the significant decrease of B-band (Figure 2,  $\text{InNO}_3$ ). It has been reported that treatment of rats with indium trichloride produced changes in heme biosynthesis (Woods et al., 1979). A similar effect on *Chlorella* could explain the decrease on PSII activity.

The two concentrations of indium nitrate tested in this work did not affect the lipid peroxidation level in *Chlorella* cells in comparison with control conditions (Figure 4). These results are not in agreement with the reported induction by indium of lipid peroxidation in mice liver (Verstraeten et al., 1997; Repetto and del Peso, 2012). Indium promotes membrane rigidification and lateral phase separation; both parameters are well correlated with the ability

1 of the ions to stimulate Fe<sup>2+</sup>-initiated lipid peroxidation. According to Zurita et al. (2007d),  
2 toxicity to fish cells at higher concentrations (5-7 mM) than those investigated with HTL in  
3 alga seems to be mediated by oxidative stress, and particularly by NADPH-dependent lipid  
4 peroxidation. Thus, taking into account our results, in *Chlorella* cells the toxic effect of  
5 indium nitrate seems not to be related with the lipid peroxidation process.  
6  
7  
8  
9  
10

11 Sodium monofluoroacetate is one of the most potent pesticides. It is mainly used as a  
12 rodenticide and as a vertebrate pest control agent. At least since the 1940s, it is known as one  
13 of the most toxic substances around the world, retaining the restricted use classification  
14 imposed by the USEPA in 1978. Leaf chlorosis, necrosis and suppression of seedling growth  
15 were reported in the plant *Helianthus annuus* exposed to 6.5 mM sodium monofluoroacetate  
16 (Cooke, 1976). From the soil organisms evaluated by O'Halloran et al. (2005), including the  
17 earthworm *Eisenia fetida* and microorganisms, lettuce (*Lactuca sativa*) was more sensitive  
18 than oats (*Avena sativa*). Lettuce seedling emergence and seedling shoot growth were  
19 adversely affected at a soil concentration of 7 mg per Kg of sodium monofluoroacetate. Zurita  
20 et al. (2007b) reported the lowest non observed adverse effect level in *C. vulgaris* for sodium  
21 monofluoroacetate (0.1 mM) in comparison to *D. magna*, *V. fischeri* and fish cell lines.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 Sodium monofluoroacetate triggers its main toxic action by blocking the Krebs cycle.  
40 The compound is incorporated into fluoroacetyl coenzyme A, which is condensed with  
41 oxalacetate to form fluorocitrate, a process dubbed as "lethal synthesis" (Buffa and Peters,  
42 1950). The intensity of HTL2 band was increased in *Chlorella* more than four-fold by sodium  
43 monofluoroacetate, reflecting a status of oxidative stress probably generated by the metabolic  
44 blockage. In fact, Dojcinovic et al. (2005) demonstrated that inhibition of the mitochondrial  
45 electron transport chain of the tricarboxylic acid cycle by sodium monofluoroacetate causes  
46 increased expression of nucleus-encoded alternative oxidase genes in plants. In addition,  
47 Zurita et al. (2007b) reported that glucose-6-phosphate dehydrogenase activity was the only  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 biomarker stimulated in RTG-2 fish cells exposed to sodium monofluoroacetate. Taking into  
2 account the protection of fluoroacetate toxicity observed in PLHC-1 cells pre-treated with  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

biomarker stimulated in RTG-2 fish cells exposed to sodium monofluoroacetate. Taking into account the protection of fluoroacetate toxicity observed in PLHC-1 cells pre-treated with sulphhydryl reagents, another mechanism of action of sodium monofluoroacetate seems to be the binding to sulphhydryl groups, including those of glutathione. The 30% reduction on the intensity of B-band of TL with the highest concentration of the chemical may be also related to the oxidative stress processes induced by sodium monofluoroacetate. However, this chemical induced another major change in the characteristics of the TL emission signal. Whereas a single TL band is generally observed in untreated *Chlorella* cells, sodium monofluoroacetate produced a more broadened TL curve with a significant upshift of the  $t_{max}$ . These results suggest that sodium monofluoroacetate could also induce major changes in the ratio among the two components responsible for the emission of B-band:  $S_2/S_3$  states of the manganese cluster and/or the reduced state of  $Q_B$  (Vass and Inoue, 1992).

Diethanolamine (DEA) is a viscous liquid widely used as chemical intermediate, anticorrosion agent in metalworking fluids and surface-active agent in cosmetic formulations, pharmaceuticals and agricultural products. It presents low cytotoxicity to *Chlorella* cell proliferation (Zurita et al., 2005b), while the algae *Skeletonema costatum* and *Phaeodactylum tricorutum* have been reported to be more sensitive in a recent ecotoxicological evaluation (Libralato et al., 2010). However, the TL emission bands in *Chlorella* were very sensitive and dose dependent. At concentrations around the  $EC_{50}$  for cell proliferation, the intensity of B-band showed a reduction of 15%, and disappeared when the concentration was doubled (30 mM). In addition, at this last concentration, another major change was a potent stimulation in lipid peroxidation, with an approximately 100-fold increase.

Diethanolamine probably inhibits the mitochondrial fatty acid beta-oxidation. Zurita et al. (2005b) have reported a marked steatosis in PLHC-1 cells exposed to diethanolamine. Toxicity studies on vertebrates have shown that diethanolamine inhibits cellular choline

1 uptake, causing a wide range of indirect effects, including increased susceptibility to oxidative  
2 damage, negative effects on phospholipid metabolism, altered mitochondrial function and  
3 structure, DNA methylation, altered transcription of genes involved in cell growth regulation  
4 and tumour development (Hansen et al., 2010). In the marine copepod *Calanus finmarchicus*,  
5 diethanolamine caused the up-regulation of the transcription of genes involved in protein  
6 catabolism, metal ion homeostasis, defence against oxidative stress, accompanied by altered  
7 expression of fatty acid derivates, amino acids and cholines (Hansen et al., 2010). All these  
8 metabolic alterations, including the observed important lost in the PSII activity, can be related  
9 to the interference of phospholipids accumulation in the metabolism. It has been reported that  
10 phospholipid accumulation produces a loss of mitochondrial integrity (Barbee and Hartung,  
11 1979), a process that can also occur in chloroplasts. The HTL experiments of this work  
12 showed enormous levels of lipid peroxidation in *Chlorella* cells treated with diethanolamine  
13 (Figure 4, DEA). This suggests that this chemical besides inducing an accumulation of lipids  
14 in the chloroplasts, also promotes its degradation by peroxidation processes.

15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34 Interestingly, the lower concentration (15 mM) of diethanolamine induced the  
35 appearance of a more broadened TL glow curve with a shoulder peaking at about 46 °C  
36 (Figure 2, DEA and Figure 3). We tentatively assigned this component appearing at higher  
37 temperatures to the AG emission band usually induced by far red illumination (Miranda and  
38 Ducruet, 1995). A similar AG-band has been observed in *Chlamydomonas* cells excited by  
39 white light (Ducruet et al., 2011). The presence of an AG-band in TL curves recorded in cells  
40 excited by white light flashes reflects the amount of PSII centres in the  $S_{2/3}Q_B$  non-radiative  
41 state immediately after preillumination, in which the arrival of an electron transferred from  
42 stroma along cyclic/chlororespiratory pathway(s) produces the radiative state  $S_{2/3}Q_B^-$  that  
43 emits luminescence (Ducruet et al., 2005). This back electron transfer can be induced by  
44 warming that activates the cyclic pathway(s) leading to an AG emission as  $S_{2/3}Q_B$  centers are  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 converted to  $S_{2/3}Q_B^-$  by electrons arriving from an unknown stroma reductant (Ducruet et al.,  
2 2005). In *Chlorella* cells grown in the presence of diethanolamine, phospholipids can provide  
3 additional NADPH in the stroma through the respiratory pathway (Alric, 2010), thus  
4 stimulating the cyclic versus linear electron transfer pathway to maintain the ATP/NADPH  
5 ratio required by the Benson-Calvin cycle. This stimulation of the cyclic/chlororespiratory  
6 pathway can explain the appearance of the AG-band in TL emissions induced by white light.  
7  
8 In *Chlorella* cells incubated in the presence of 30 mM diethanolamine there is not  
9 photosynthetic electron transfer on PSII (see Figure 2, DEA). However, the growth inhibition  
10 test has shown that at this concentration the percentage of *Chlorella* living cells was still  
11 about 40% (Zurira et al., 2005b). So, diethanolamine should activate alternative  
12 photosynthetic metabolism, without the participation of PSII (chlororespiration, cyclic  
13 electron transport around PSI, etc.) or not photosynthetic metabolism processes that maintain  
14 the cells viability.  
15  
16

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31 According to Foster (1994), the lysosomotropic drug chloroquine is probably the  
32 second or third most widely consumed drug in the world. It is still the main pharmaceutical  
33 used for the prophylaxis and treatment of malaria, and is also used for treatment of  
34 autoimmune diseases, collagen vascular and dermatological disorders and extraintestinal  
35 amebiasis. In addition, CQ has been proposed as a promising inexpensive drug in the  
36 management of HIV disease (Romanelli et al., 2004). It is the most potent compound  
37 evaluated according to the inhibition of the proliferation of *Chlorella* cells, with an eight-fold  
38 increase in toxicity when the exposure time period was increased from 24 to 72 h (Zurita et  
39 al., 2005a). Jjemba (2002) demonstrated that the growth of soybean plants was decreased by  
40 increasing concentrations of chloroquine, while the number of bacteria and protozoa in soil  
41 was either unchanged or increased. Although the inhibition of bioluminescence in *V. fischeri*  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 is less sensitive, the crustacean *D. magna* is even more sensitive than *Chlorella* (Zurita et al.,  
2 2005a).  
3

4 Thermoluminescence measurements in *Chlorella* cells were particularly sensitive to  
5 chloroquine, showing important reductions on B-band intensity (Figure 2, CQ), while HTL2  
6 intensity was increased 5.5-fold by 0.25 mM chloroquine, in comparison to control (Figure 4,  
7 CQ). These results agree with the previously reported oxidative stress in rats induced by  
8 chloroquine, with the consequence of lipid peroxidation as mechanism of the nephrotoxicity  
9 (Murugavel and Pari, 2004). According to Sugioka et al. (1987), the ferriprotoporphyrin IX-  
10 chloroquine complex remarkably promotes peroxidative cleavage of unsaturated  
11 phospholipids. The significant decrease of the activity of PSII can be related to lipid  
12 peroxidation on chloroplast and also with the possible binding of chloroquine to porphyrin  
13 groups of the PSII cofactors.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 In conclusion, for five of the chemicals studied (not BrB) there is a very good  
29 correlation between the cytotoxic effects in *Chlorella* cells measured by the algae growth  
30 inhibition test (Zurita et al., 2005a,b; Zurita et al., 2007a,b,c,d), and the inhibition of PSII  
31 activity measured by STL. Thus, we may propose that one very important effect of these  
32 chemicals in *Chlorella* cells is the inhibition of photosynthetic metabolism by the blocking of  
33 PSII functionality. In the case of sodium monofluoroacetate, diethanolamine and chloroquine  
34 this inhibition seems to be related with the induction of high level of lipid peroxidation in  
35 cells that may alter the stability of PSII. Other toxic mechanisms different from lipid  
36 peroxidation must be however responsible for the inhibition of PSII induced by PpG and  
37 InNO<sub>3</sub>. The inhibition of growth in *Chlorella* cells induce by BrB seems not to be related by  
38 the blocking of photosynthetic electron transfer activity on PSII.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 Consequently, although no single biomarker is viable in gaining a comprehensive  
56 understanding of xenobiotic stress (Brain and Cedergreen, 2009), the quantification of TL  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 emissions is a useful technique to study specific effects of environmental chemicals on  
2 photosynthetic organisms. STL and HTL techniques provide specific information on  
3 photosynthetic electron transport and on lipid peroxidation induced by oxidative stress,  
4 respectively. This information is complementary to that obtained by the growth inhibition test  
5 and allows a more complete assessment of the effects of a chemical in organisms of aquatic  
6 ecosystems, and about the mechanisms of the toxicity.  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16

### 17 **Conflict of interest**

18 None of the authors had any conflicts to declare.  
19  
20  
21  
22  
23

### 24 **Acknowledgements**

25 This work was supported by grants BFU2007-68107-C02-01/BMC from the Spanish  
26 Ministry of Science and Innovation and BIO2012-35271 and CTM2012-31344 from the  
27 Spanish Ministry of Economy and Competitiveness, and the PADI CVI-261, BIO-022 and  
28 CTS-996 from the Andalusia Government. All of these grants were partially found by the EU  
29 FEDER program. J.L. Zurita was the recipient of a grant for the training of research personal  
30 from the Ministry of Science and Innovation. The authors thank S. Jiménez and A. Martos for  
31 technical assistance.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

- 1  
2 Alric, J., 2010. Cyclic electron flow around photosystem I in unicellular green algae.  
3 Photosynth. Res. 106, 47–56.  
4
- 5  
6 Barbee, S.J., Hartung, R., 1979. The effect of diethanolamine on hepatic and renal  
7 phospholipid metabolism in the rat. Toxicol. Appl. Pharmacol. 47, 421–430.  
8
- 9  
10 Baroli, I., Gutman, B.J., Ledford, H.K., Shin, J.W., Chin, B.L., Havaux, M., Noyogi, K.K.,  
11 2004. Photo-oxidative stress in xanthophylls-deficient mutant of *Chlamydomonas*. J. Biol.  
12 Chem. 279, 6337-6344.  
13
- 14 Brain R.A., Cedergreen N., 2009. Biomarkers in aquatic plants: selection and utility. Rev.  
15 Environ. Contam. Toxicol. 198, 49-109.  
16
- 17  
18 Buffa, P., Peters, R.A., 1950. The in vivo formation of citrate induced by fluoroacetate and its  
19 significance. J. Phys. 110, 488 500.  
20
- 21  
22 Chatterjee, S.N., Agarwal, S., 1988. Liposomes as membrane model for study of lipid  
23 peroxidation. Free Rad. Biol. Med. 4, 51-72.  
24
- 25  
26 Cooke, J.A., 1976. The uptake of sodium fluoroacetate by plants and its physiological effects.  
27 Fluoride 9, 204–212.  
28
- 29 Dalton, D.A., 1995. Antioxidant Defenses of Plants and Fungi. In: Ahmad, S. (Ed.), Oxidative  
30 Stress and Antioxidant Defences in Biology. Chapman & Hall, N.Y., pp. 298-355.  
31
- 32  
33 Demeter S., Goussias C., Bernat G., Kovacs L., Petrouleas V., 1993. Participation of the  $g =$   
34  $1.9$  and  $g = 1.82$  EPR forms of the semiquinone-iron complex,  $Q_A^- Fe^{2+}$  of photosystem II in  
35 the generation of the Q and C thermoluminescence bands, respectively. FEBS Lett. 336, 352-  
36 356.  
37
- 38  
39 Demeter S., Vass I., 1984. Charge accumulation and recombination in Photosystem II studied  
40 by thermoluminescence. I. Participation of the primary acceptor Q and secondary acceptor B  
41 in the generation of thermoluminescence of chloroplasts. Biochim. Biophys. Acta 764, 24–32.  
42
- 43  
44 Dianzani, M., Barrera, G., 2008. Pathology and physiology of lipid peroxidation and its  
45 carbonyl products. In: Álvarez, S., Evelson, P. (Ed.), Free Radical Pathophysiology.  
46 Transworld Research Network, Kerala, India, pp. 19- 38.  
47
- 48  
49 Diner B.A., Babcock G.T., 1996. Structure, dynamics and energy conversion efficiency in  
50 photosystem II. In: Ort, D.R., Yocum, C.F. (Ed.), Oxygenic Photosynthesis: The Light  
51 Reactions. Kluwer Academic Publishers, The Netherlands, pp. 213–247.  
52
- 53  
54 Diner, B.A., Rappaport, F., 2002. Structure, dynamics, and energetics of the primary  
55 photochemistry of photosystem II of oxygenic photosynthesis. Annu. Rev. Plant. Biol. 53,  
56 551-580.  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 Dodd, D.E., Pluta, L.J., Sochaski, M.A., Banas, D.A., Thomas, R.S., 2012. Subchronic  
2 hepatotoxicity evaluation of bromobenzene in Fischer 344 rats. J. App. Toxicol. 33, 370–377.

3 Dojcinovic D., Krosting J., Harris A.J., Wagner D.J., Rhoads D.M., 2005. Identification of a  
4 region of the Arabidopsis AtAOX1a promoter necessary for mitochondrial retrograde  
5 regulation of expression. Plant Mol. Biol. 58,159-75.  
6

7  
8 Ducruet, J.M., 2003. Chlorophyll thermoluminescence of lead disc: simple instruments and  
9 progress in signal interpretation open the way to new ecophysiological indicators. J. Exp. Bot.  
10 54, 2419-2430.  
11

12  
13 Ducruet, J.M., Miranda, T., 1992. Graphical and numerical analysis of thermoluminescence  
14 and fluorescence F0 emission in photosynthetic material. Photosynth. Res. 33, 15–27.  
15

16  
17 Ducruet, J.M., Roman, M., Ortega, J.M., Janda, T., 2005. Role of the oxidized secondary  
18 acceptor QB of photosystem II in the delayed 'afterglow' chlorophyll luminescence.  
19 Photosynth. Res. 84, 161-166.  
20

21  
22 Ducruet, J.M., Peeva, V., Havaux M., 2007. Chlorophyll thermofluorescence and  
23 thermoluminescence as complementary tools for the study of temperature stress in plants.  
24 Photosynth. Res. 93, 159-171.  
25

26  
27 Ducruet, J.M., Vass, I., 2009. Thermoluminescence: experimental. Photosynth. Res. 101,  
28 195–204.  
29

30  
31 Ducruet, J.M., Serrano, A., Roncel, M., Ortega, J.M., 2011. Peculiar properties of chlorophyll  
32 thermoluminescence emission of autotrophically or mixotrophically grown *Chlamydomonas*  
33 *reinhardtii*. J. Photochem. Photobiol. B. 104, 301-307.  
34

35  
36 Foster, S., 1994. Economic prospects for a new antimalarial drug. Trans. R. Soc. Trop. Med.  
37 Hyg. 88, 55–56.

38  
39 Fusi, F., Valoti, M., Sgaragli, G., Murphy, M.P., 1991. The interaction on antioxidants and  
40 structurally related compounds with mitochondrial oxidative phosphorylation. Meth. Find.  
41 Exp. Clin. Pharmacol. 13, 599–604.  
42

43  
44 Guerrero, F., Zurita, J.L., Roncel, M., Kirilovsky, D., Ortega, J.M., 2014. The role of the  
45 high potential form of the cytochrome *b559*: Study of *Thermosynechococcus*  
46 *elongatus* mutants. Biochim. Biophys. Acta 1837, 908-919.  
47

48  
49 Han, Y.H., Park, W.H., 2010. Pyrogallol-induced calf pulmonary arterial endothelial cell  
50 death via caspase-dependent apoptosis and GSH depletion. Food Chem. Toxicol. 48, 558-563.  
51

52  
53 Havaux, M., 2003. Spontaneous and thermoinduced photon emission: new methods to detect  
54 and quantify oxidative stress in plants. Trends Plant Sci 8, 409-413.  
55

56  
57 Havaux, M., Niyogi, K.K., 1999. The violaxanthin cycle protects plants from photooxidative  
58 damage by more than one mechanism. Proc. Nat. Acad. Sci. USA 96, 8762-8767.  
59  
60  
61  
62  
63  
64  
65

1 Hansen B.H., Altin, D., Bootha, A., Vanga, S., Frenzela, M., Sørheima, K.R., Brakstada,  
2 O.G., Størseth, T.R., 2010. Molecular effects of diethanolamine exposure on *Calanus*  
3 *finmarchicus* (Crustacea: Copepoda). *Aquat. Toxicol.* 99, 212–222.

4 Heath, R.L., Parker, L., 1968. Photoperoxidation in isolated chloroplasts. I. Kinetics and  
5 stoichiometry of fatty acid peroxidation. *Arch. Biochem. Biophys.* 125, 189-198.

6  
7  
8 Horváth, G., Arellano, J.B., Droppa, M., Barón, M., 1998. Alterations in Photosystem II  
9 electron transport as revealed by thermoluminescence of Cu-poisoned chloroplasts.  
10 *Photosynth. Res.* 57, 175–182,  
11

12  
13 Hörsik, Z.T., Kovács, L., Láposi, R., Mészáros, I., Lakatos, G., Garab, G., 2007. Effect of  
14 chromium on photosystem 2 in the unicellular green alga *Chlorella pyrenoidosa*.  
15 *Photosynthetica* 45, 65-69.  
16

17  
18 Inoue, Y., 1996. Photosynthetic thermoluminescence as a simple probe of Photosystem II  
19 electron transport. In: Amesz, J. and Hoff, A.J. (Ed), *Biophysical Techniques in*  
20 *Photosynthesis*. Kluwer Academic Publishers, Dordrech, pp 93-107.  
21

22  
23 Jjemba, P.K., 2002. The effect of chloroquine, quinacrine, and metronidazole on both soybean  
24 plants and soil microbiota. *Chemosphere* 46, 1019–1025.  
25

26  
27 Johnson G.N., Boussac, A., Rutherford, A.W., 1994. The origin of 40-50 °C  
28 thermoluminescence bands in Photosystem II. *Biochim. Biophys. Acta* 1184, 85-92.  
29

30  
31 Josse, E.M., Simkin, A.J., Gaffe, J., Laboure, A.M., Kuntz, M., 2003. In vitro characterization  
32 of plastid terminal oxidase (PTOX). *Eur. J. Biochem.* 270, 3787-3794.  
33

34  
35 Karthikeyan, K., Sarala Bai, B.R., Gauthaman, K., Niranjali Deveraj, S., 2005. Protective  
36 effects of propyl gallate myocardial oxidative stress-induced injury in rat. *J. Pharm.*  
37 *Pharmacol.* 57, 67–73.  
38

39  
40 Katsumata, M., Koike, T., Kazumura, K., Takeuchi, A., Sugaya, Y., 2009. Utility of Delayed  
41 Fluorescence as Endpoint for Rapid Estimation of Effect Concentration on the Green Alga  
42 *Pseudokirchneriella subcapitata*. *Bull. Environ. Contam. Toxicol.* 83, 484–487,  
43

44  
45 Kawanishi, S., Oikawa, S., Murata, M., 2005. Evaluation for safety of antioxidant  
46 chemopreventive agents. *Antiox. Red. Sign.* 7, 1728–1739.  
47

48  
49 Kobayashi, H., Oikawa, S., Hirakawa, K., Kawanishi, S., 2004. Metal-induced oxidative  
50 damage to cellular and isolated DNA by galic acid, a metabolite of antioxidant propyl gallate.  
51 *Mutat. Res.* 558, 111–120.  
52

53  
54 Lamb, R.G., McCue, S.B., Taylor, D.R., McGuffin, M.A., 1984. The role of phospholipid  
55 metabolism in bromobenzene and carbon tetrachloride dependent hepatocyte injury. *Toxicol.*  
56 *Appl. Pharmacol.* 30, 510–520.  
57

58  
59 Libralato, G., Ghirardini, A., Avezzi, F., 2010. Seawater ecotoxicity of monoethanolamine,  
60 diethanolamine and triethanolamine. *J. Hazard. Mat.* 176, 535-539.  
61  
62  
63  
64  
65

1 Lipcamon, J.D., Sahl, B., 2002. A new weapon in the treatment of non- Hodgkin's lymphoma.  
2 Radiol. Manag. 24, 13–5.

3 Matorin, D.N., Vavilin, D.V., Kafarov, R.S., Venediktov, P.S., 1989. High temperature  
4 thermoluminescence of chlorophyll as a method for study of lipid peroxidation in plants. Dokl.  
5 Akad. Nauk. SSSR. 309, 764-768.

6  
7  
8 Merzlyak, M.N., Pavlov, V.K., Zhigalova, T.V., 1992. Effect of desiccation on chlorophyll  
9 high temperature chemiluminescence in *Acer platanoides* L. and *Aesculus hippocastanum* L.  
10 leaves. J. Plant. Physiol. 139, 629-631.

11  
12  
13 Miranda, T., Ducruet, J.M., 1995. Characterization of the chlorophyll thermoluminescence  
14 afterglow in dark-adapted or far-red-illuminated plant leaves. Plant Physiol. Biochem. 33,  
15 689-699.

16  
17  
18 Mohanty, N., Vass, I., Demeter, S., 1989. Copper toxicity affects photosystem II electron  
19 transport at the secondary quinone acceptor, Q(B). Plant Physiol. 90, 175-179.

20  
21  
22 Murugavel, P., Pari, L., 2004. Attenuation of chloroquine-induced renal damage by alpha-  
23 lipoic acid: possible antioxidant mechanism. Renal Fail. 26, 517–524.

24  
25  
26 O'Halloran, K., Jones, D., Booth, L., Fisher, P., 2005. Ecotoxicity of sodium fluoroacetate  
27 (compound 1080) to soil organisms. Environ. Toxicol. Chem. 24, 1211–1218.

28  
29 Pryor, W.A., Castle, L., 1984. Chemical methods for the detection of lipid hydroperoxides.  
30 Methods Enzymol. 105, 293-305.

31  
32  
33 Ramos, C., de la Torre, A.I., Tarazona, J.V., Muñoz, M.J., 1996. Desarrollo de un ensayo de  
34 inhibición de *Chlorella vulgaris* utilizando un test en microplacas. Rev. Toxicol. 13, 97-100.

35  
36 Rappaport, F., Guergova-Kuras, M., Nixon, P.J., Diner, B.A., Lavergne, J., 2002. Kinetics  
37 and pathways of charge recombination in photosystem II. Biochemistry 41, 8518-8527.

38  
39  
40 Repetto, G., del Peso, A., 2012. Gallium, indium, and thallium. In: Bingham, E., Cohrssen, B.  
41 (Ed.), Patty's Toxicology, Sixth Edition. vol. 1. John Wiley & Sons, New York, pp. 257-353.

42  
43  
44 Repetto, M., Semprine, J., Boveris, A., 2012. Lipid Peroxidation: Chemical Mechanism,  
45 Biological Implications and Analytical Determination. In: Catalá, A. (Ed.), Lipid  
46 Peroxidation. InTech, pp. 3-30.

47  
48  
49 Repetto, G., 2013. Test batteries in ecotoxicology. In: Férard, J., Blaise, C. (Ed.),  
50 Encyclopedia of Aquatic Ecotoxicology. Springer-Verlag, Berlin Heidelberg, pp. 1105-1128.

51  
52  
53 Romanelli, F., Smith, K.M., Hoven, A.D., 2004. Chloroquine and hydroxychloroquine as  
54 inhibitors of human immunodeficiency virus (HIV-1) activity. Curr. Pharm. Des. 10, 2643–  
55 2648.

56  
57  
58 Rutherford A.W., Crofts A.R., Inoue Y., 1982. Thermoluminescence as a probe of  
59 Photosystem II photochemistry: the origin of the flash-induced glow peaks. Biochim.  
60 Biophys. Acta 682, 457-465.

1 Rutherford, A.W., Inoue, Y., 1984. Oscillations of delayed luminescence from PSII:  
2 recombination of  $S_2Q_B^-$  and  $S_3Q_B^-$ . FEBS Lett. 165, 163-170.  
3

4 Rutherford, A.W., Renger, G., Koike, H., Inoue, Y., 1984a. Thermoluminescence as a probe  
5 of photosystem II. The redox and protonation states of the secondary acceptor quinone and  
6  $O_2$ -evolving enzyme. Biochim. Biophys. Acta 767, 548-556.  
7  
8

9 Rutherford, A.W., Govindje, Inoue, Y., 1984b. Charge accumulation and photochemistry in  
10 leaves studied by thermoluminescence and delayed light emission. Proc. Natl. Acad. Sci. USA  
11 81, 1107-1111.  
12

13 Sahay, A., Jajoo, A., Bharti, S., 2006. A thermoluminescence study of the effects of nitrite on  
14 photosystem II in spinach thylakoids. Luminescence 21, 143-147.  
15  
16

17 Sane, P.V., 2004. Thermoluminescence. A technique for probing Photosystem II. In:  
18 Carpentier, R. (Ed.), Methods in Molecular Biology, vol 274. Photosynthesis Research  
19 Protocols, Humana Press Inc, Totowa, NJ, pp 229-248.  
20  
21

22 Schumann, A., Bauer, A., Hermes, M., Gilbert, M., Hengstler, J.G., Wilhelm, C., 2009. A  
23 rapid and easy to handle thermoluminescence based technique for evaluation of carbon  
24 tetrachloride induced oxidative stress on rat hepatocytes. Arch. Toxicol. 83, 709-20.  
25  
26

27 Schoonen, W., Westerink, W., de Roos, J., Debiton, E., 2005. Cytotoxic effects of 100  
28 reference compounds on HepG2 and HeLa cells and of 60 compounds on ECC-1 and CHO  
29 cells. I Mechanistic assays on ROS, glutathione depletion and calcein uptake. Toxicol. In  
30 Vitro 19, 505-516.  
31  
32

33 Sugioka, Y., Suzuki, M., Sugioka, K., Nakano, M., 1987. A ferriprotoporphyrin IX-  
34 chloroquine complex promotes membrane phospholipid peroxidation. A possible mechanism  
35 for antimalarial action. FEBS Lett. 223, 251-254.  
36  
37

38 Van der Heijden, C.A., Janssen, P.J., Strip, J.J., 1986. Toxicology of gallates: a review and  
39 evaluation. Food Chem. Toxicol. 24, 1067-1070.  
40  
41

42 Vass, I., Inoue, Y., 1992. Thermoluminescence in the study of photosystem II. In: Barber, J.  
43 (Ed.), The photosystems: Structure, Function and Molecular Biology. Elsevier Science  
44 Publishers, Amsterdam, pp. 259-294.  
45  
46

47 Vavilin, D.V., Ducruet, J.M., 1998. The origin of 120-130 °C thermoluminescence bands in  
48 chlorophyll-containing material. Photochem. Photobiol. 68, 191-198.  
49  
50

51 Vavilin, D.V., Matorin, D.N., Kafarov, R.S., Bautina, A.L., Venediktov, P.S., 1991. High-  
52 temperature thermoluminescence of chlorophyll in lipid peroxidation. Biol. Memb. 8, 89-98.  
53  
54

55 Venediktov, P.S., Matorin, D.N., Kafarov, R.S., 1989. Chemiluminescence of chlorophyll  
56 upon lipid photoperoxidation in thylakoid membranes. Biofizika (Moscow) 34, 241-245.  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Verstraeten, S.V., Nogueira, L.V., Schreier, S., Oteiza, P.I., 1997. Effect of trivalent metal  
2 ions on phase separation and membrane lipid packing: role in lipid peroxidation. Arch.  
3 Biochem. Biophys. 338, 121–127.

4 Weber, D., Milkovic, L., Bennett, S.J., Griffiths, H.R., Zarkovic, N., Grune, T., 2013.  
5 Measurement of HNE-protein adducts in human plasma and serum by ELISA. Comparison of  
6 two primary antibodies. Redox Biol. 1, 226–233.

7  
8  
9  
10 Wong, S.L., Nakamoto, L., Wainwright, J.F., 1997. Detection of toxic organometallic  
11 complexes in waste- waters using algal assays. Arch. Environ. Cont. Toxicol 32, 358-366.

12  
13 Woods, J.S., Carver, G.T., Fowler, B.A., 1979. Altered regulation of rat hepatic heme  
14 metabolism by indium chloride. Toxicol. Appl. Pharmacol.49, 455–462.

15  
16  
17 Yadav, D.K., Pospíšil, P., 2012. Evidence on the Formation of Singlet Oxygen in the Donor  
18 Side Photoinhibition of Photosystem II: EPR Spin-Trapping Study. PLoS ONE 7, e459883

19  
20  
21 Zurita, J.L., Jos, A., del Peso, A., Salguero, M., López-Artíguez, M., Repetto, G., 2005a.  
22 Ecotoxicological evaluation of the antimalarial drug chloroquine. Aquat. Toxicol. 75, 97-107.

23  
24 Zurita, J.L., Repetto, G., Jos, A., del Peso, A., Salguero, M., López-Artíguez, M., Olano, D.,  
25 Cameán, A., 2005b. Ecotoxicological evaluation of diethanolamine using a battery of  
26 microbiotests. Toxicol. In Vitro 19, 879-886.

27  
28  
29 Zurita, J.L., Roncel, M., Aguilar, M., Ortega, J.M., 2005c. A thermoluminescence study of  
30 photosystem II back electron transfer reactions in rice leaves. Effects of salt stress.  
31 Photosynth. Res. 84, 131-137.

32  
33  
34 Zurita, J.L., Jos, A., del Peso, A., Salguero, M., López-Artíguez, M., Repetto, G., 2007a.  
35 Ecotoxicological assessment of bromobenzene using a test battery with five model systems.  
36 Food Chem. Toxicol. 45, 575-584.


37  
38  
39 Zurita, J.L., Jos, A., Cameán, A.M., Salguero, M., López-Artíguez, M., Repetto, G., 2007b.  
40 Ecotoxicological evaluation of sodium fluoroacetate using a test battery with aquatic systems.  
41 Chemosphere 67, 1-12.

42  
43  
44 Zurita, J.L., Jos, A., del Peso, A., Salguero, M., López-Artíguez, M., Repetto, G., 2007c  
45 Ecotoxicological effects of the antioxidant additive propyl gallate in five aquatic systems.  
46 Water Res. 41, 2599 – 261.

47  
48  
49 Zurita, J.L., Jos, A., del Peso, A., Salguero, M., López-Artíguez, M., Repetto, G., 2007d.  
50 Toxicological evaluation of indium nitrate on aquatic organisms and investigation of the  
51 effects in the PLHC-1 fish cell line. Sci. Tot. Environ. 387, 155-165.


1 **Figure legends**

2  
3 **Figure 1.** Thermoluminescence glow curves from non-treated *Chlorella vulgaris* cells  
4 recorded after a series of single turn-over flashes (indicated on each curve). 50  $\mu$ l of cell  
5 suspensions from *Chlorella* culture at exponential growth phase were dark-adapted for 2 min  
6 at 20 °C and subsequently cooled to 1 °C during 1 min. Then, 1, 2 or 3 saturating single turn-  
7 over flashes (separated by 1 s) were given at the end of this period. Luminescence emission  
8 was then recorded while warming samples from -5 °C to 80 °C at a heating rate of 0.5 °C per  
9 s. For other technical details see Material and Methods section.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26



1  
2  
3  
4  
5 **Figure 2.** Thermoluminescence glow curves from *Chlorella vulgaris* cells exposed for 24 h to  
6  
7 six different chemicals. Bromobenzene (BrB), propyl gallate (PpG), indium nitrate (InNO<sub>3</sub>),  
8  
9 sodium monofluoroacetate (SMFA), diethanolamine (DEA) or chloroquine (CQ). *Chlorella*  
10  
11 cells suspensions were exposed to two different concentrations of each chemical as described  
12  
13 in Material and Methods section. After the treatment, 50 µl of cells suspension were dark-  
14  
15 adapted for 2 min at 20 °C and subsequently cooled to 1 °C during 1 min. Two saturating  
16  
17 single turn-over flashes (separated by 1 s) were given at the end of this period. Luminescence  
18  
19 emission was then recorded while warming samples from 0 °C to 80 °C at a heating rate of 0.5  
20  
21 °C per s. For other technical details see Material and Methods section. Symbols: (●) Control  
22  
23 non-treated samples; (○) *Chlorella* cells exposed to the lower concentration of the chemical;  
24  
25 (Δ) *Chlorella* cells exposed to the higher concentration of the chemical. Lower and higher  
26  
27 concentrations of the chemicals, respectively: BrB, 1 and 2 mM; PpG, 0.5 and 1 mM; InNO<sub>3</sub>,  
28  
29 0.3 and 0.7 mM; SMFA, 5 and 20 mM; DEA, 15 and 30 mM; CQ, 0.1 and 0.25 mM.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65





1  
2  
3  
4  
5 **Figure 3.** Thermoluminescence glow curves from *Chlorella vulgaris* cells exposed for 24 h to  
6 diethanolamine (DEA). *Chlorella* cells suspensions were exposed to two different  
7 concentrations of diethanolamine (15 and 30 mM) as described in Material and Methods  
8 section. After the treatment, 50 µl of cells suspension were dark-adapted for 2 min at 20 °C  
9 and subsequently cooled to 1 °C during 1 min. One saturating single turn-over flash was given  
10 at the end of this period. Luminescence emission was then recorded while warming samples  
11 from 0 °C to 80 °C at a heating rate of 0.5 °C per s. For other technical details see Material  
12 and Methods section. Symbols: Control, non-treated samples; DEA 1, *Chlorella* cells exposed  
13 to 15 mM diethanolamine; DEA 2, *Chlorella* cells exposed to 30 mM diethanolamine. The B  
14 and the AG-bands resulting from the decomposition of the TL curve induced in *Chlorella*  
15 cells exposed to the lower concentration of the chemical are shown in dots.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 **Figure 4.** High temperature thermoluminescence glow curves from *Chlorella vulgaris* cells  
2  
3 exposed for 24 h to six different chemicals. Bromobenzene (BrB), propyl gallate (PpG),  
4  
5 indium nitrate (InNO<sub>3</sub>), sodium monofluoroacetate (SMFA), diethanolamine (DEA) or  
6  
7 chloroquine (CQ). *Chlorella* cells were exposed to two different concentrations of the  
8  
9 chemicals as described in Material and Methods section. After the treatment, 50 µl of cells  
10  
11 suspension were adsorbed by filtration on a piece of filter paper, dark adapted for 10 min at  
12  
13 20 °C and cooled to 1 °C for 1 min. Luminescence emission was then recorded without flash  
14  
15 excitation while warming samples from 10 °C to 160 °C at a heating rate of 0.1 °C per s. In  
16  
17 order to desiccate the samples and prevents any oxidation induced by high temperature, N<sub>2</sub>  
18  
19 was flushed on the samples during HTL experiments. For other technical details see Material  
20  
21 and Methods section. Symbols: (●) Control non-treated samples; (○) *Chlorella* cells exposed  
22  
23 to the lower concentration of the chemical; (Δ) *Chlorella* cells exposed to the higher  
24  
25 concentration of the chemical. Lower and higher concentrations of the chemicals,  
26  
27 respectively: BrB, 1 and 2 mM; PpG, 0.5 and 1 mM; InNO<sub>3</sub>, 0.3 and 0.7 mM; SMFA, 5 and  
28  
29 20 mM; DEA, 15 and 30 mM; CQ, 0.1 and 0.25 mM.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

