

MOS subject classification: 17D10; 12E20; 13F20; 16D70

A computational algebraic geometry approach to enumerate Malcev magma algebras over finite fields^{*}

Óscar J. Falcón^a, Raúl M. Falcón^{b*} and Juan Núñez^a

The set $\mathcal{M}_n(\mathbb{K})$ of n -dimensional Malcev magma algebras over a finite field \mathbb{K} can be identified with algebraic sets defined by zero-dimensional radical ideals for which the computation of their reduced Gröbner bases makes feasible their enumeration and distribution into isomorphism and isotopism classes. Based on this computation and the classification of Lie algebras over finite fields given by De Graaf and Strade, we determine the mentioned distribution for Malcev magma algebras of dimension $n \leq 4$. We also prove that every 3-dimensional Malcev algebra is isotopic to a Lie magma algebra. For $n = 4$, this assertion only holds when the characteristic of the base field \mathbb{K} is distinct of two. Copyright © 2016 John Wiley & Sons, Ltd.

Keywords: Malcev algebra; magma algebra; finite field; polynomial ring

1. Introduction

Non-associative algebras have a special importance in terms of Mathematics, as well as for their multiple applications in Natural Sciences and Engineering [1, 2]. This paper deals in particular with the open problem of enumerating and classifying *Malcev magma algebras*, with possible application in quantum mechanics and string theory. A *magma* is a finite set endowed with a binary operation. Throughout the paper we suppose this set to be $[n] = \{1, \dots, n\}$ and we denote the inner law as \cdot . An n -dimensional algebra A over a field \mathbb{K} is said to be a *magma algebra* if there exists a basis $\{e_1, \dots, e_n\}$ of the algebra and a magma $([n], \cdot)$ such that $e_i e_j = c_{ij} e_{i,j}$ for each pair of elements $i, j \leq n$ and some structure constant $c_{ij} \in \mathbb{K}$. The algebra A is then said to be *based on* the magma $([n], \cdot)$. This is said to be a *Malcev magma algebra* if it also constitutes a *Malcev algebra* [3, 4], that is, $u^2 = 0$, for all $u \in A$, and

$$((uv)w)u + ((vw)u)u + ((wu)u)v = (uv)(uw), \text{ for all } u, v, w \in A. \quad (1)$$

Every Malcev algebra is, therefore, antisymmetric, that is, $uv = -vu$, for all $u, v \in A$. Besides, (1) is equivalent to the so-called *Malcev identity*

$$M(u, v, w) = J(u, v, w)u - J(u, v, uw) = 0, \text{ for all } u, v, w \in A, \quad (2)$$

^a Faculty of Mathematics, Department of Geometry and Topology, University of Seville, c/ Tarfia s/n. 41012-Sevilla.

^b University of Seville, Department of Applied Mathematics I.

* Correspondence to: School of Building Engineering, University of Seville. Avda. Reina Mercedes 4 A, 41012, Seville, Spain. E-mail: rafalgan@us.es

* This is the accepted version of the following article: Falcón, Ó. J., Falcón, R. M., and Núñez, J. (2016) A computational algebraic geometry approach to enumerate Malcev magma algebras over finite fields. *Math. Meth. Appl. Sci.*, 39: 4901–4913. doi: 10.1002/mma.4054, which has been published in final form at [http://onlinelibrary.wiley.com/doi/10.1002/mma.4054/full]. This article may be used for non-commercial purposes in accordance with the Wiley Self-Archiving Policy [http://olabout.wiley.com/WileyCDA/Section/id-828039.html].

where J is the *Jacobian* defined as $J(u, v, w) = (uv)w + (vw)u + (wu)v$, for all $u, v, w \in A$. If the characteristic of the base field \mathbb{K} , which is denoted by $\text{char}(\mathbb{K})$, is distinct of two, then both identities (1) and (2) are equivalent to the *Sagle identity*

$$S(u, v, w, y) = ((uv)w)y + ((vw)y)u + ((wy)u)v + ((yu)v)w - (uw)(vy) = 0, \text{ for all } u, v, w, y \in A. \quad (3)$$

Unlike Malcev identity, which is not linear in its first argument $u \in A$, Sagle identity is linear in the four arguments $u, v, w, y \in A$ and invariant under cyclic permutations of the variables.

If the *Jacobi identity*, $J(u, v, w) = 0$, is satisfied for any three vectors u, v and w of an algebra, then this is called a *Lie algebra* (a *Lie magma algebra* if this is also a magma algebra). Otherwise, the algebra is said to have a *Jacobi anomaly*. In quantum mechanics, the existence of Jacobi anomalies in the underlying non-associative algebraic structure related to the coordinates and momenta of a quantum non-Hamiltonian dissipative system was already claimed by Dirac [5] in the process of taking Poisson brackets. In string theory, for instance, one such an anomaly is involved by the non-associative algebraic structure that is defined by coordinates (x) and velocities or momenta (v) of an electron moving in the field of a constant magnetic charge distribution, at the position of the location of the magnetic monopole [6]. In particular, $J(v_1, v_2, v_3) = -\vec{\nabla} \cdot \vec{B}(x)$, where $\vec{\nabla} \cdot \vec{B}(x)$ denotes the divergence of the magnetic field $\vec{B}(x)$. The underlying algebraic structure constitutes a non-Lie Malcev algebra [7], with the commutation relations $[x_a, x_b] = 0$, $[x_a, v_b] = i\delta_{ab}$ and $[v_a, v_b] = i\epsilon_{abc}B_c(x)$, where $a, b, c \in \{1, 2, 3\}$, δ_{ab} denotes the Kronecker delta and ϵ_{abc} denotes the Levi-Civita symbol. If the magnetic field is proportional to the coordinates, the latter can be normalized and $B_c(x)$ can then be supposed to coincide with x_c . The resulting algebra is then called *magnetic* [8]. A generalization to electric charges has recently been considered [9] by defining the products $[x_a, x_b] = -i\epsilon_{abc}E_c(x, v)$, where the electric field E as well as the magnetic field B must depend not only on coordinates but also on velocities. Remark that both magnetic and electric algebras constitute magma algebras.

A main open problem in the theory of Malcev algebras is their enumeration and distribution into isomorphism classes [10, 11, 12]. Over finite fields, this problem has already been dealt with for Lie algebras of dimension up to six. Particularly, De Graaf [13] made use of Gröbner bases and computational algebraic geometry in order to determine the distribution of solvable Lie algebras of dimension up to four over any field, whereas Strade [14] obtained that of nonsolvable Lie algebras of dimension up to six over a finite field. The classification of nilpotent Lie algebras of dimension up to six over any field is also known [15]. The authors in this last reference indicated explicitly that some of their results were inspired by Gröbner basis computations.

In this paper, similarly to the methodology exposed by De Graaf [13], we make use of computational algebraic geometry as an approach to deal with the classification of the sets $\mathcal{L}_n(\mathbb{K})$ and $\mathcal{M}_n(\mathbb{K})$ of n -dimensional Lie magma algebras and Malcev magma algebras, respectively, over a finite field \mathbb{K} . We deal not only with the distribution into isomorphism classes, which is the usual criterion, but also with the more general distribution into isotopism classes, which constitutes a first attempt in this regard to the best of the authors knowledge. The concept of isotopism of algebras was introduced by Albert [16], who indicated that two algebras (A, \cdot) and (A', \circ) are *isotopic* if there exist three nonsingular linear maps f, g and h between A and A' such that

$$f(u) \circ g(v) = h(u \cdot v), \text{ for all } u, v \in A. \quad (4)$$

The triple (f, g, h) is said to be an *isotopism* between both algebras A and A' , which is denoted as $A \simeq A'$. If $f = g = h$, then this constitutes an *isomorphism*, which is denoted as $A \cong A'$. To be isotopic is an equivalence relation among algebras of the same dimension and constitutes, therefore, a generalization of the concept of isomorphism. In his original paper, Albert dealt with isotopisms of division algebras, alternative algebras and Lie algebras. Shortly after, Bruck [17] dealt with isotopisms of quasigroup algebras, which constitute magma algebras in which division is possible. Since the manuscripts of Albert and Bruck, a wide amount of papers have dealt with isotopisms of distinct types of algebras as division algebras [18, 19, 20, 21], alternative algebras [22, 23], Lie algebras [24, 25], Jordan algebras [26, 27, 28, 29, 30, 31, 32], genetic algebras [33, 34], absolute valued algebras [35, 36], structural algebras [37] and real 2-dimensional commutative algebras [38]. Nevertheless, there does not exist any study on the distribution of Malcev algebras into isotopism classes.

The structure of the paper is the following. In Section 2 we expose some preliminary concepts and results on isotopisms of algebras, Malcev magma algebras and computational algebraic geometry. Section 3 deals with those ideals of polynomials

whose algebraic sets are identified with the sets of n -dimensional Lie and Malcev magma algebras. We describe in particular two algorithms whose implementation enable us to determine the enumeration of Malcev magma algebras and their distribution into isomorphism and isotopism classes. This distribution is explicitly determined in Section 4 for dimension $n \in \{3, 4\}$.

2. Preliminary results

In this section we expose some basic concepts and results on isotopisms of algebras, Malcev algebras and computational algebraic geometry that are used throughout the paper. For more details about these topics we refer to the original articles of Albert [16], Malcev [3] and Sagle [4] and to the monographs of Cox, Little and O'Shea [39, 40].

2.1. Isotopisms of algebras

Let A be an algebra over a field \mathbb{K} . This algebra is called *abelian* if all its structure constants are zero. This constitutes a trivial magma algebra for which all the products of the corresponding magma are zeros.

Lemma 2.1 *The n -dimensional abelian algebra is not isotopic to any other n -dimensional algebra.*

Proof. Let (f, g, h) be an isotopism between an n -dimensional non-abelian algebra of basis $\{e_1, \dots, e_n\}$ and the n -dimensional abelian algebra. Let $i, j \leq n$ be such that $e_i e_j \neq 0$. Then, $0 = f(e_i)g(e_j) = h(e_i e_j) \neq 0$, which is a contradiction. \square

The *derived algebra* of the algebra A is the subalgebra A^2 formed by the set of elements uv such that $(u, v) \in A \times A$. The *derived series* of the algebra A is defined as

$$\mathcal{C}_1(A) = A \supseteq \mathcal{C}_2(A) = A^2 \supseteq \dots \supseteq \mathcal{C}_k(A) = (\mathcal{C}_{k-1}(A))^2 \supseteq \dots \quad (5)$$

The algebra A is said to be *solvable* if there exists a positive integer m such that $\mathcal{C}_{m+1}(A) \equiv 0$. The smallest such an integer is called the *solvability index* of the algebra. This index and the dimension of each vector subspace $\mathcal{C}_k(A)$ are preserved by isomorphisms. Nevertheless, this is not true in general for isotopisms, for which we can only assure the next result.

Lemma 2.2 *Let A and A' be two isotopic n -dimensional algebras. Then, $\dim(A^2) = \dim(A'^2)$.*

Proof. Let (f, g, h) be an isotopism between both algebras A and A' . The regularity of f and g involves that $f(A) = g(A) = A'$. Hence, $A'^2 = f(A)g(A) = h(A^2)$ and the result follows then from the regularity of h . \square

Let us suppose now that A is a Lie algebra. The *centralizer* of a subset $S \subseteq A$ is the vector subspace $C_A(S) = \{u \in A \mid uv = 0, \text{ for all } v \in S\} \subseteq A$. The *center* of A is the set $Z(A) = C_A(A)$. The next results show the way in which isotopisms preserve centralizers and centers of Lie algebras.

Lemma 2.3 *Let (f, g, h) be an isotopism between two Lie algebras A and A' . Let S be a subset of L . Then,*

- a) $f(C_A(S)) = C_{A'}(g(S))$.
- b) $g(C_A(S)) = C_{A'}(f(S))$.
- c) $\dim(C_A(S)) = \dim(C_{A'}(f(S))) = \dim(C_{A'}(g(S)))$.

Proof. Let $u \in C_A(S)$. Then, $f(u)v = h(ug^{-1}(v)) = h(0) = 0$, for all $v \in g(S)$ and hence, $f(C_A(S)) \subseteq C_{A'}(g(S))$. Reciprocally, if $u \in C_{A'}(g(S))$, then $f^{-1}(u)v = h^{-1}(ug(v)) = h^{-1}(0) = 0$ and hence, $C_{A'}(g(S)) \subseteq f(C_A(S))$. Thus, $f(C_A(S)) = C_{A'}(g(S))$. The regularity of f and g involves the same dimension of both vector subspaces. Item (b) and the fact of being $\dim(C_A(S)) = \dim(C_{A'}(f(S)))$ in item (c) hold similarly from the antisymmetry of L . \square

Proposition 2.4 Let A and A' be two isotopic Lie algebras. Then, $f(Z(A)) = Z(A')$ and $\dim(Z(A)) = \dim(Z(A'))$.

Proof. The result is an immediate consequence of Lemma 2.3 from the definition of center of a Lie algebra. \square

Let n be the dimension of the Lie algebra A . For each positive integer $m \leq n$, we define

$$d_m(A) := \min\{\dim C_A(S) : S \text{ is an } m\text{-dimensional vector subspace of } A\}. \quad (6)$$

$$D_m(A) := \max\{\dim C_A(S) : S \text{ is an } m\text{-dimensional vector subspace of } A\}. \quad (7)$$

We prove now that both values are preserved by isotopisms.

Proposition 2.5 Let A and A' be two isotopic n -dimensional Lie algebras and let $m \leq n$ be a positive integer. Then, $d_m(A) = d_m(A')$ and $D_m(A) = D_m(A')$.

Proof. Let (f, g, h) be an isotopism between A and A' and let S be an m -dimensional vector subspace of A such that $d_m(A) = \dim C_A(S)$. The regularity of g involves the set $g(S)$ to be m -dimensional. Besides, from Lemma 2.3, $C_{A'}(g(S)) = f(C_A(S))$. Hence, $d_m(A) \geq d_m(A')$. The equality follows similarly from the isotopism (f^{-1}, g^{-1}, h^{-1}) between A' and A . The invariance of D_m holds analogously. \square

We finish with a remark on isotopisms of magma algebras. Let A and A' be two isotopic n -dimensional magma algebras over a field \mathbb{K} and let $([n], \cdot)$ and $([n], *)$ be the respective magmas on which both algebras are based. If $\{e_1, \dots, e_n\}$ and $\{e'_1, \dots, e'_n\}$ are the respective bases of A and A' , then every isotopism (f, g, h) between both algebras is characterized by three nonsingular matrices $F = (f_{ij})$, $G = (g_{ij})$ and $H = (h_{ij})$, with entries in the field \mathbb{K} , so that $\alpha(e_i) = \sum_{j=1}^n \alpha_{ij} e'_j$, for all $\alpha \in \{f, g, h\}$ and $i \in \{1, \dots, n\}$. Then,

$$c_{ij} h_{(i \cdot j)k} = \sum_{s, t \leq n | s * t = k} f_{is} g_{jt} c'_{st}, \text{ for all } i, j, k \leq n, \quad (8)$$

where c_{ij} and c'_{st} are structure constants of A and A' , respectively.

2.2. Malcev magma algebras

Every Malcev magma algebra in the current paper is described by means of a basis of vectors satisfying the conditions exposed in the next result.

Lemma 2.6 Let $n \geq 3$. Every n -dimensional non-abelian Malcev magma algebra based on a magma $([n], \cdot)$ holds, up to isomorphism, one of the next two non-isomorphic possibilities

a) $e_1 e_2 = e_2$, or

b) $e_1 e_2 = e_3$ and there does not exist a non-zero structure constant c_{ij} such that $i \cdot j \in \{i, j\}$.

Proof. Since the algebra is not abelian, there exists at least one non-zero structure constant. If there exist two distinct positive integers $i, j \leq n$ such that $c_{ij} \neq 0$ and $i \cdot j = j$, then we consider the isomorphism that maps e_i and e_j to $c_{ij} e_1$ and e_2 , respectively, and preserves the rest of basis vectors. We get in this way the product $e_1 e_2 = e_2$. Otherwise, we take a non-zero structure constant c_{uv} and the isomorphism that maps e_u , e_v and $e_{u \cdot v}$ to $c_{uv} e_1$, e_2 and e_3 , respectively, in order to get the product $e_1 e_2 = e_3$. \square

Every Malcev algebra is *binary-Lie*, that is, any two of its elements generate a Lie subalgebra. As a consequence, every Malcev algebra of dimension $n \leq 3$ is a Lie algebra. Particularly, the only 1-dimensional Malcev algebra is the abelian and the only 2-dimensional non-abelian Malcev algebra is, up to isomorphism, the magma algebra determined by the product of basis vectors $e_1 e_2 = e_2$. The next theorem deals with the distribution of 3-dimensional Malcev magma algebras over a finite field into isomorphism classes. In the statement of the result, each isomorphism class is labeled according to the notation given by De Graaf [13] and Strade [14] in their respective classifications of solvable and non-solvable Lie algebras, but we have chosen for its description a basis that follows the conditions exposed in Lemma 2.6.

Theorem 2.7 Every 3-dimensional Malcev magma algebra over a finite field \mathbb{K} is isomorphic to exactly one of the next algebras

a) The 3-dimensional abelian Lie algebra L^1 .

b) The solvable Lie algebras

- $L^2 \cong e_1 e_2 = e_2$ and $e_1 e_3 = e_3$.
- $L_0^3 \cong e_1 e_2 = e_2$.
- $L_a^4 \cong e_1 e_2 = e_3$ and $e_1 e_3 = a e_2$, for all $a \in \mathbb{K}$.
Here, $L_a^4 \cong L_b^4$ if and only if there exists $\alpha \in \mathbb{K} \setminus \{0\}$ such that $a = \alpha^2 b$.

c) The non-solvable Lie algebras

- $W(1; \underline{2})^{(1)} \cong e_1 e_2 = e_2$, $e_1 e_3 = e_3$ and $e_2 e_3 = e_1$, whenever $\text{char}(\mathbb{K}) = 2$.
- $\mathfrak{sl}(2, \mathbb{K}) \cong e_1 e_2 = e_2$, $e_1 e_3 = -e_3$ and $e_2 e_3 = e_1$, whenever $\text{char}(\mathbb{K}) \neq 2$.

Proof. The mentioned classifications of Lie algebras obtained by De Graaf and Strade enable us to assure that each Lie algebra of the list constitutes an isomorphism class in $\mathcal{M}_3(\mathbb{K})$. In order to assure that there does not exist any other isomorphism class, it is required to prove that none of the next 3-dimensional solvable Lie algebras is a magma algebra

$$L_a^3 \cong e_1 e_2 = e_2 + a e_3 \text{ and } e_1 e_3 = e_2, \text{ with } a \in \mathbb{K} \setminus \{0\}. \quad (9)$$

We prove this fact in Proposition 4.1. □

For dimension $n = 4$, there exists, up to isomorphism, a unique non-Lie Malcev algebra [41] when $\text{char}(\mathbb{K}) \notin \{2, 3\}$. This coincides with the next solvable magma algebra

$$M^0 \cong e_1 e_2 = e_2, e_1 e_3 = -e_3, e_1 e_4 = -e_4 \text{ and } e_3 e_4 = -e_2.$$

The distribution of $\mathcal{M}_4(\mathbb{K})$ into isomorphism classes is exposed in the next result, which is again based on the classifications of De Graaf and Strade.

Theorem 2.8 Let \mathbb{K} be a finite field. Every Malcev magma algebra $A \in \mathcal{M}_4(\mathbb{K})$ is isomorphic to exactly one of the next algebras

a) The non-Lie Malcev algebra M^0 .

b) The 4-dimensional abelian Lie algebra M^1 .

c) The solvable Lie algebras

- $M^2 \cong e_1 e_2 = e_2$, $e_1 e_3 = e_3$ and $e_1 e_4 = e_4$.
- $M_{-1}^3 \cong e_1 e_2 = e_2$, $e_1 e_3 = e_4$ and $e_1 e_4 = e_3$.
- $M_0^3 \cong e_1 e_2 = e_2$, $e_1 e_3 = e_4$ and $e_1 e_4 = e_4$.
- $M^4 \cong e_1 e_2 = e_2$ and $e_1 e_3 = e_2$.
- $M^5 \cong e_1 e_2 = e_3$.
- $M_{0,0}^6 \cong e_1 e_2 = e_2$, $e_1 e_3 = e_4$ and $e_1 e_4 = e_2$.
- $M_{0,0}^7 \cong e_1 e_2 = e_3$ and $e_1 e_3 = e_4$.
- $M_{a,0}^7 \cong e_1 e_2 = e_3$, $e_1 e_3 = e_4$ and $e_1 e_4 = a e_2$, for all $a \in \mathbb{K} \setminus \{0\}$.
Here, $M_{a,0}^7 \cong M_{b,0}^7$ if and only if there exists $\alpha \in \mathbb{K} \setminus \{0\}$ such that $a = \alpha^3 b$.
- $M_{0,a}^7 \cong e_1 e_2 = e_3$, $e_1 e_3 = e_4$ and $e_1 e_4 = a e_3$, for all $a \in \mathbb{K} \setminus \{0\}$.
Here, $M_{0,a}^7 \cong M_{0,b}^7$ if and only if there exists $\alpha \in \mathbb{K} \setminus \{0\}$ such that $a = \alpha^2 b$.
- $M^8 \cong e_1 e_2 = e_2$ and $e_3 e_4 = e_4$.
- $M_{1,0}^{11} \cong e_1 e_2 = e_2$, $e_1 e_4 = 1 e_4$, $e_2 e_4 = e_3$ and $e_3 e_4 = e_2$, whenever $\text{char}(\mathbb{K}) = 2$.
- $M^{12} \cong e_1 e_2 = e_2$, $e_1 e_3 = 2 e_3$, $e_1 e_4 = e_4$ and $e_2 e_4 = -e_3$, whenever $\text{char}(\mathbb{K}) \neq 2$.
- $M_0^{13} \cong e_1 e_2 = e_2$, $e_1 e_3 = e_3$, $e_1 e_4 = e_2$ and $e_2 e_4 = -e_3$.

- $M_a^{14} \cong e_1 e_2 = e_3, e_1 e_3 = a e_2$ and $e_2 e_3 = e_4$, for all $a \in \mathbb{K} \setminus \{0\}$.
Here, $M_a^{14} \cong M_b^{14}$ if and only if there exists $\alpha \in \mathbb{K} \setminus \{0\}$ such that $a = \alpha^2 b$.

d) The non-solvable Lie algebras over a field of characteristic two

- $W(1; 2) \cong e_1 e_2 = e_2, e_1 e_3 = e_3, e_2 e_3 = e_1$ and $e_3 e_4 = e_2$.
- $W(1; 2)^{(1)} \oplus Z(A) \cong e_1 e_2 = e_2, e_1 e_3 = e_3$ and $e_2 e_3 = e_1$.

e) The non-solvable Lie algebra over a field of characteristic distinct of two

- $\mathfrak{gl}(2, \mathbb{K}) \cong e_1 e_2 = e_2, e_1 e_3 = -e_3, e_2 e_3 = e_1, e_2 e_4 = e_2$ and $e_3 e_4 = -e_3$.

Proof. The classification given by De Graaf and Strade for solvable and non-solvable 4-dimensional Lie algebras over finite fields implies that all these algebras constitute distinct isomorphism classes. In order to assure that the distribution is exhaustive, it is required to prove that none of the next 4-dimensional solvable Lie algebras is isomorphic to a magma algebra

$$M_a^3 : e_1 e_2 = e_2, e_1 e_3 = e_4 \text{ and } e_1 e_4 = -a e_3 + (a + 1) e_4, \text{ with } a \notin \{0, -1\}. \quad (10)$$

$$M_{a,b}^6 : e_1 e_2 = e_2 + a e_3 + b e_4, e_1 e_3 = e_4 \text{ and } e_1 e_4 = e_2, \text{ for all } (a, b) \in \mathbb{K}^2 \setminus (0, 0). \quad (11)$$

$$M_{a,a}^7 : e_1 e_2 = e_3, e_1 e_3 = e_4 \text{ and } e_1 e_4 = a e_2 + a e_3, \text{ for all } a \in \mathbb{K} \setminus \{0\}. \quad (12)$$

$$M_{a_0}^9 : e_1 e_2 = e_2 + a_0 e_3, e_1 e_3 = e_2, e_2 e_4 = e_2 \text{ and } e_3 e_4 = e_3, \text{ whenever } \text{char}(\mathbb{K}) = 2 \text{ and } a_0 \in \mathbb{K} \text{ is such that } T^2 - T - a_0 \text{ has not root in } \mathbb{K}. \quad (13)$$

$$M_a^{13} : e_1 e_2 = e_2 + a e_4, e_1 e_3 = e_3, e_1 e_4 = e_2 \text{ and } e_2 e_4 = -e_3, \text{ for all } a \in \mathbb{K} \setminus \{0\}. \quad (14)$$

We prove this fact in Proposition 4.4. □

2.3. Computational algebraic geometry

Let X and $\mathbb{K}[X]$ be, respectively, the set of n variables $\{x_1, \dots, x_n\}$ and the related multivariate polynomial ring over a field \mathbb{K} . The algebraic set defined by an ideal I of $\mathbb{K}[X]$ is the set $\mathcal{V}(I)$ of common zeros of all the polynomials in I . The ideal I is zero-dimensional if $\mathcal{V}(I)$ is finite. It is radical if every polynomial $f \in \mathbb{K}[X]$ belongs to I whenever there exists a natural number m such that $f^m \in I$. Let $<$ be a monomial term ordering on the set of monomials in $\mathbb{K}[X]$. The largest monomial of a polynomial f in I with respect to $<$ is its leading monomial $\text{LM}(f)$, whose coefficient in f is the leading coefficient $\text{LC}(f)$. The leading term of f is the product $\text{LT}(f) = \text{LC}(f) \cdot \text{LM}(f)$. The ideal generated by all the leading monomials of I is the initial ideal $I_{<}$. Those monomials that are not in $I_{<}$ are called standard monomials of I . Regardless of the monomial term ordering, if the ideal I is zero-dimensional and radical, then the number of standard monomials in I coincides with the Krull dimension of the quotient ring $\mathbb{K}[X]/I$ and with the number of points of the algebraic set $\mathcal{V}(I)$.

A Gröbner basis of I with respect to $<$ is any subset G of polynomials in I whose leading monomials generate the initial ideal $I_{<}$. This is reduced if all its polynomials are monic and no monomial of a polynomial in G is generated by the leading monomials of the rest of polynomials in the basis. There exists only one reduced Gröbner basis of the ideal I , which becomes an optimal way to count their number of standard monomials. Its decomposition into finitely many disjoint subsets, each of them being formed by the polynomials of a triangular system of polynomial equations makes also possible to enumerate the elements of the algebraic set $\mathcal{V}(I)$ [42, 43, 44]. The reduced Gröbner basis of an ideal I can always be computed from Buchberger's algorithm [45]. Similar to the Gaussian elimination on linear systems of equations, this consists of a sequential multivariate division of polynomials, also called reduction or normal form computation, which is based on the construction of the so-called S -polynomials $S(f, g) = \text{lcm}(\text{LM}(f), \text{LM}(g))(f/\text{LT}(f) - g/\text{LT}(g))$, for all $f, g \in I$, where lcm denotes the least common multiple. In particular, a subset G of polynomials in I is a Gröbner basis if $S(f, g)$ reduces to zero after division by the polynomials in G , for all $f, g \in G$. Buchberger's algorithm can be implemented on ideals defined over any base field, but its involved exact arithmetic becomes faster on finite fields. It is due to the large numbers that appear in general as intermediate coefficients

during the computation of the reduced Gröbner basis and which constitute a major factor in the computational cost of the algorithm. Derived from Buchberger’s algorithm, a pair of more efficient direct methods are the algorithms F_4 and F_5 [46, 47] and the algorithm *slimgb* [48]. The latter is based on F_4 and reduces the computation time and the memory usage by keeping small coefficients and short polynomials during the sequential division of polynomials. In any case, the computation of a reduced Gröbner basis is always extremely sensitive to the number of variables [49, 50].

3. Algebraic sets related to Malcev magma algebras

In this section we identify the sets $\mathcal{L}_n(\mathbb{K})$ and $\mathcal{M}_n(\mathbb{K})$ of n -dimensional Lie and Malcev magma algebras over a finite field \mathbb{K} with the algebraic set of an ideal of polynomials. Let X and $\mathbb{K}[X]$ respectively be the set of n^3 variables $\{c_{ij}^k : i, j, k \leq n\}$ and its related multivariate polynomial ring over \mathbb{K} . Let \mathfrak{A} be the n -dimensional algebra over $\mathbb{K}[X]$ with basis $\{e_1, \dots, e_n\}$ so that

$$e_i e_j = \sum_{k=1}^n c_{ij}^k e_k, \tag{15}$$

for all $i, j \leq n$. Let us also consider

- i. The coefficient $l_{ijkl} \in \mathbb{K}[X]$ of e_l in the Jacobi identity $J(e_i, e_j, e_k) = 0$, for all $i, j, k \leq n$.
- ii. The coefficient $m_{uijk} \in \mathbb{K}[X]$ of e_k in the Malcev identity $M(u, e_i, e_j) = 0$, for all $u \in \mathfrak{A}$ and $i, j \leq n$.
- iii. The coefficient $s_{ijklm} \in \mathbb{K}[X]$ of e_m in the Sagle identity $S(e_i, e_j, e_k, e_l) = 0$, for all $i, j, k, l \leq n$, whenever $\text{char}(\mathbb{K}) \neq 2$.

Finally, let us define the ideal

$$I = \langle c_{ij}^j \mid i, j \leq n \rangle + \langle c_{ij}^k - c_{ji}^k \mid i, j, k \leq n \rangle + \langle c_{ij}^k c_{ij}^{k'} \mid i, j, k, k' \leq n \text{ such that } k < k' \rangle \subset \mathbb{K}[X]. \tag{16}$$

Theorem 3.1 *The sets $\mathcal{L}_n(\mathbb{K})$ and $\mathcal{M}_n(\mathbb{K})$ are respectively identified with the algebraic sets defined by the zero-dimensional radical ideals in $\mathbb{K}[X]$*

$$I_L = \langle l_{ijkl} : i, j, k, l \leq n \rangle + I \quad \text{and} \quad I_M = \langle m_{uijk} : u \in \mathfrak{A}, i, j, k \leq n \rangle + I.$$

If $\text{char}(\mathbb{K}) \neq 2$, then the set $\mathcal{M}_n(\mathbb{K})$ is also identified with the algebraic set defined by the zero-dimensional radical ideal in $\mathbb{K}[X]$

$$I_S = \langle s_{ijklm} : i, j, k, l, m \leq n \rangle + I.$$

Besides, $|\mathcal{L}_n(\mathbb{K})| = \dim_{\mathbb{K}}(\mathbb{K}[X]/I_L)$ and $|\mathcal{M}_n(\mathbb{K})| = \dim_{\mathbb{K}}(\mathbb{K}[X]/I_M)$ ($= \dim_{\mathbb{K}}(\mathbb{K}[X]/I_S)$ if $\text{char}(\mathbb{K}) \neq 2$).

Proof. We prove the theorem for Malcev magma algebras and the ideal I_M . The other cases follow similarly. Observe that each zero $(c_{11}^1, \dots, c_{nn}^n) \in \mathcal{V}(I_M)$ constitutes the structure constants of an n -dimensional algebra A with basis $\{e_1, \dots, e_n\}$ such that $e_i e_j = \sum_{k=1}^n c_{ij}^k e_k$ and $M(u, e_i, e_j) = 0$, for all $i, j \leq n$ and $u \in A$. The generators of $I \subset I_M$ involve A to be a magma algebra such that $u^2 = 0$, for all $u \in A$. The linearity of $M(u, v, w)$ in the two last arguments involves A to be a Malcev algebra.

Since the base field \mathbb{K} is finite, the ideal I_M is zero-dimensional and the affine variety $\mathcal{V}(I_M)$ is a finite subset of \mathbb{K}^{n^3} . If \mathbb{K} is isomorphic to a Galois field $\text{GF}(q)$, then, from Proposition 2.7 of [39], the ideal I_M is also radical, because the unique monic generator of $I_M \cap \mathbb{K}[c_{ij}^k]$ is the polynomial $(c_{ij}^k)^q - c_{ij}^k$, which is intrinsically included in each one of the ideals and is square-free. As a consequence, the number of zeros in the algebraic set $\mathcal{V}(I_M)$ coincides with the Krull dimension of the quotient ring $\mathbb{K}[X]/I_M$ over \mathbb{K} . □

Corollary 3.2 Let \mathbb{K} be a finite field isomorphic to the Galois field $GF(q)$. The run time that is required by the Buchberger's algorithm in order to compute the reduced Gröbner bases of the ideals I_L , I_M and I_S in Theorem 3.1 are, respectively,

$$q^{O(n^3)} + O(n^8), \quad \max\{3, q\}^{O(n^3)} + O(q^{2n}) \quad \text{and} \quad q^{O(n^3)} + O(n^{10}).$$

Proof. The result follows straightforward from Proposition 4.1.1 exposed by Gao in [51], once we observe that all the generators of the three ideals in Theorem 3.1 are sparse in $\mathbb{K}[X]$. According to that result, the run time that is required by the Buchberger's algorithm in order to compute each reduced Gröbner basis is $q^{O(v)} + O(g^2l)$, where v , g and l are, respectively, the number of variables, the number of generators of the corresponding ideal that are not of the form $(c_{ij}^k)^q - c_{ij}^k$ and the longest length of these generators. \square

The previous result discards the use of the ideal I_M whenever $\text{char}(\mathbb{K}) \neq 2$. We have made use of the open computer algebra system for polynomial computations Singular [52] and the algorithm *slimgb* [48] to compute the reduced Gröbner bases of the ideals in Theorem 3.1 and hence, their respective algebraic sets and Krull dimensions. To this end, we have implemented four subprocedures called *Prod*, *Jacobild*, *Malcevd* and *Sagleld* and a main procedure called *MalcevAlg*. All of them have been included in the library *malcev.lib*, which is available online at <http://personales.us.es/raufalgan/LS/malcev.lib>. The subprocedure *Prod* outputs the list of polynomials corresponding to the coefficient of each basis vector in the product between two arbitrary vectors of the algebra A . Similar lists of polynomials are output by the rest of subprocedures. Their respective pseudocodes are exposed in Algorithms 1-4.

Algorithm 1 Polynomials related to the product of two vectors in A .

```

1: procedure Prod(u, v)
2:   for k ← 1, n do
3:     for i ← 1, n do
4:       for j ← i + 1, n do
5:         Lk ← Lk + (uivj - ujvi)cijk;
6:       end for
7:     end for
8:   end for
9:   return {L1, ..., Ln}
10: end procedure

```

Algorithm 2 Polynomials related to the Jacobi identity in A .

```

1: procedure Jacobild(u, v, w)
2:   L1 ← Prod(Prod(u, v), w);
3:   L2 ← Prod(Prod(v, w), u);
4:   L3 ← Prod(Prod(w, u), v);
5:   for i ← 1, n do
6:     L1i ← L1i + L2i + L3i;
7:   end for
8:   return L1
9: end procedure

```

Algorithm 3 Polynomials related to the Malcev identity in A .

```

1: procedure Malcevd(u, v, w)
2:   L1 ← Prod(Jacobild(u, v, w), u);
3:   L2 ← Jacobild(u, v, Prod(u, w));
4:   for i ← 1, n do
5:     L1i ← L1i - L2i;
6:   end for
7:   return L1
8: end procedure

```

Algorithm 4 Polynomials related to the Sagle identity in A .

```

1: procedure SagleId( $u, v, w, y$ )
2:    $L1 \leftarrow \text{Prod}(\text{Prod}(\text{Prod}(u, v), w), y)$ ;
3:    $L2 \leftarrow \text{Prod}(\text{Prod}(\text{Prod}(v, w), y), u)$ ;
4:    $L3 \leftarrow \text{Prod}(\text{Prod}(\text{Prod}(w, y), u), v)$ ;
5:    $L4 \leftarrow \text{Prod}(\text{Prod}(\text{Prod}(y, u), v), w)$ ;
6:    $L5 \leftarrow \text{Prod}(\text{Prod}(u, w), \text{Prod}(v, y))$ ;
7:   for  $i \leftarrow 1, n$  do
8:      $L1_i \leftarrow L1_i + L2_i + L3_i + L4_i - L5_i$ ;
9:   end for
10:  return  $L1$ 
11: end procedure

```

In case of being interested in computing an algebra containing a specific structure constant $c_{ij}^k \in \mathbb{K}$, we include the polynomial $c_{ij}^k - c_{ij}^k$ in the corresponding ideal of Theorem 3.1. This aspect has been incorporated in the main procedure *MalcevAlg*, whose pseudocode is exposed in Algorithm 5. This procedure receives as input

1. The dimension n of the algebra.
2. The order q of the base field \mathbb{K} .
3. A list C of tuples of the form (i, j, k, c_{ij}^k) , where $i, j, k \leq n$, $i < j$ and $c_{ij}^k \in \mathbb{K}$.
4. A positive integer $\text{opt1} \leq 2$ so that we use the ideal I_L if $\text{opt1} = 1$. Otherwise, we use the ideals I_M or I_S if $\text{char}(\mathbb{K}) = 2$ or $\text{char}(\mathbb{K}) \neq 2$, respectively.
5. A positive integer $\text{opt2} \leq 2$ that determines as output the Krull dimension of the ideal if $\text{opt2} = 1$. Otherwise, it determines the algebraic set defined by the ideal.

Algorithm 5 Computation of Lie and Malcev magma algebras.

```

1: procedure MalcevAlg( $n, q, C, \text{opt1}, \text{opt2}$ )
2:   Depending on the argument  $\text{opt1}$ , use the subprocedures JacobId, MalcevId or SagleId to initialize an ideal  $I$  as  $I_L$ ,  $I_M$  or  $I_S$ , respectively.
3:   for  $i \leftarrow 1, \text{size}(C)$  do
4:      $I = I + (c_{C_{i3}C_{i2}}^{C_{i3}} - C_{i4})$ ;
5:   end for
6:    $I = \text{slimgb}(I)$ ;
7:   if  $\text{opt1} = 1$  then
8:     return  $|\mathcal{V}(I)|$ 
9:   else
10:    return  $\mathcal{V}(I)$ 
11:  end if
12: end procedure

```

The correctness and termination of this main procedure is based on those of the algorithm *slimgb* [48] and hence, on those of Buchberger's algorithm [45]. This procedure has been implemented in a system with an *Intel Core i7-2600, with a 3.4 GHz processor and 16 GB of RAM*. Its effectiveness has been checked by computing the cardinalities that are exposed in Tables 1 and 2. The run time and memory usage are explicitly indicated in both tables. Both measures of computation efficiency fit positive exponential models even for dimension $n = 4$. Nevertheless, we are not interested in the computation of all Malcev magma algebras, but only in their distribution into isomorphism and isotopism classes. We indicate now how computational algebraic geometry can also be used to determine this distribution.

Let us consider the sets of variables

$$X = \{f_{11}, \dots, f_{nn}\} \quad \text{and} \quad Y = X \cup \{g_{11}, \dots, g_{nn}, h_{11}, \dots, h_{nn}\}.$$

q	$ \mathcal{M}_3(\text{GF}(q)) $	Run time		Used memory	
		I_L	$I_{M/S}$	I_L	$I_{M/S}$
2	32	0 s	0 s	0 MB	0 MB
3	123	0 s	0 s	0 MB	0 MB
5	581	0 s	0 s	0 MB	0 MB
7	1,567	0 s	0 s	0 MB	0 MB
11	5,891	0 s	0 s	0 MB	0 MB
13	9,613	0 s	0 s	0 MB	0 MB
17	21,137	0 s	0 s	0 MB	0 MB
⋮	⋮	⋮	⋮	⋮	⋮
499	498,492,523	0 s	0 s	0 MB	0 MB

Table 1. Computation of $|\mathcal{M}_3(\text{GF}(q))|$.

q	$ \mathcal{L}_4(\text{GF}(q)) $	Run time	Used memory	$ \mathcal{M}_4(\text{GF}(q)) $	Run time	Used memory
2	853	0 s	0 MB	897	6 s	0 MB
3	7,073	1 s	0 MB	7,073	8 s	0 MB
5	89,185	11 s	0 MB	89,377	41 s	0 MB
7	445,537	20 s	3 MB	445,969	55 s	2 MB
11	3,803,041	91 s	17 MB	3,804,241	154 s	427 MB
13	8,412,193	183 s	676 MB	8,413,921	258 s	859 MB
17	30,247,297	595 s	1,2 GB	30,250,369	752 s	1,5 GB

Table 2. Computation of $|\mathcal{L}_4(\text{GF}(q))|$ and $|\mathcal{M}_4(\text{GF}(q))|$.

These variables play the role of the entries in the nonsingular matrices F , G and H related to a possible isotopism (f, g, h) between two Malcev magma algebras of respective bases $\{e_1, \dots, e_n\}$ and $\{e'_1, \dots, e'_n\}$. Let p_{ijk} and q_{ijk} be the respective polynomials in $\mathbb{K}[X]$ and $\mathbb{K}[Y]$ that constitutes the coefficients of e_k in the expressions $f(e_i)f(e_j) - f(e_i e_j)$ and $f(e_i)g(e_j) - h(e_i e_j)$, for all $i, j, k \leq n$. The next result holds analogously to Theorem 3.1.

Theorem 3.3 *Let \mathbb{K} be a finite field isomorphic to the Galois field $\text{GF}(q)$ and let A and A' be two n -dimensional Malcev magma algebras over \mathbb{K} . The sets of isomorphisms and isotopisms between both algebras are respectively identified with the algebraic sets defined by the zero-dimensional radical ideals*

$$I_{A,A'} = \langle p_{ijk} : i, j, k \leq n \rangle + \langle \det(F)^{q-1} - 1 \rangle \subset \mathbb{K}[X].$$

$$J_{A,A'} = \langle q_{ijk} : i, j, k \leq n \rangle + \langle \det(F)^{q-1} - 1, \det(G)^{q-1} - 1, \det(H)^{q-1} - 1 \rangle \subset \mathbb{K}[Y].$$

Besides, $|\mathcal{V}(I_{A,A'})| = \dim_{\mathbb{K}}(\mathbb{K}[X]/I_{A,A'})$ $|\mathcal{V}(J_{A,A'})| = \dim_{\mathbb{K}}(\mathbb{K}[Y]/J_{A,A'})$. □

We have included the procedure *isoMalcev* in the previously mentioned library *malcev.lib* in order to compute the algebraic sets and the Krull dimensions in Theorem 3.3. Having as output the number of isomorphisms or that of isotopisms between two Malcev magma algebras A and A' , this procedure receives as input

1. The dimension n of both algebras.
2. The order q of the base field.
3. A list $L1$ formed by tuples (i, j, k, c_{ij}^k) that indicates the non-zero structure constants of the Malcev algebra A .
4. A list $L2$ formed by tuples (i, j, k, c'_{ij}^k) that indicates the non-zero structure constants of the Malcev algebra A' .
5. A positive integer $\text{opt} \leq 2$ that enable us to use the ideal $I_{A,A'}$ if $\text{opt} = 1$, or the ideal $J_{A,A'}$ if $\text{opt} = 2$.

Algorithm 6 shows how the procedure *isoMalcev* is implemented to determine the distribution of a set of Malcev magma algebras into isomorphism and isotopism classes.

Algorithm 6 Computation of the isomorphism (isotopism, resp.) classes of a set of Malcev algebras.

Require: A set M of Malcev magma algebras.

Ensure: S , the set of isomorphism (isotopism, resp.) classes of S

```

1:  $S = \emptyset$ .
2: while  $M \neq \emptyset$  do
3:   Take  $A \in M$ .
4:    $M := M \setminus \{A\}$ .
5:    $S := S \cup \{A\}$ .
6:   for  $A' \in M$  do
7:     if  $|\mathcal{V}(I_{A,A'})| > 0$  ( $|\mathcal{V}(J_{A,A'})| > 0$ , resp.) then
8:        $M := M \setminus \{A'\}$ .
9:     end if
10:  end for
11: end while
12: return  $S$ .
```

4. Classification of Malcev magma algebras of small dimensions

We have implemented the procedures and algorithms of the previous section to determine the distribution of 3- and 4-dimensional Malcev magma algebras into isomorphism and isotopism classes.

4.1. 3-dimensional Malcev magma algebras

The next result finishes the proof of Theorem 2.7 about the distribution of $\mathcal{M}_3(\mathbb{K})$ into isomorphism classes.

Proposition 4.1 *None of the algebras that are described in (9) is isomorphic to a Lie magma algebra.*

Proof. Let $a \in \mathbb{K} \setminus \{0\}$. Let f be an isomorphism between the algebra L_a^3 and a Malcev magma algebra A and let $F = (f_{ij})$ be the nonsingular matrix related to f . This algebra A should have a 2-dimensional derived algebra and its solvability index should be 2. The implementation of Algorithm 5 into a case study that takes into account Lemma 2.6 and the Jacobi identity involves A to be isomorphic to one of the next two magma algebras

a) $e_1 e_2 = e_3$ and $e_1 e_3 = \alpha e_2$.

b) $e_1 e_2 = e_2$ and $e_1 e_3 = \alpha e_3$.

In both cases, $\alpha \in \mathbb{K} \setminus \{0\}$. The implementation of the procedure *isoMalcev* enable us to compute for each case the corresponding reduced Gröbner basis in Theorem 3.3. In (a), the normal form of the polynomial that is related to the determinant of the corresponding matrix F modulo this reduced Gröbner basis is zero whatever the number α and the characteristic of the base field are. This means that F is a singular matrix, which is a contradiction with being f an isomorphism. Hence, L_a^3 is not isomorphic to a Malcev magma algebra of type (a). On the other hand, in (b), the previous normal form is the polynomial $(1 - \alpha)f_{11}f_{32}f_{33}$. The regularity of the matrix F involves that $\alpha \neq 1$. Besides, the reduced Gröbner basis involves the identity $\alpha a = a$. Since $\alpha \neq 1$, it must be $a = 0$, what is a contradiction with the hypothesis. Hence, L_a^3 is not isomorphic either to a Malcev magma algebra of type (b). \square

Even if not every 3-dimensional Malcev algebra is isomorphic to a Lie magma algebra, the next result shows that this statement is true in case of dealing with isotopisms instead of isomorphisms.

Proposition 4.2 *Every 3-dimensional Malcev algebra is isotopic to a Lie magma algebra.*

Proof. The result holds from the fact that, for all $a \in \mathbb{K} \setminus \{0\}$, the Lie algebras L_a^3 and L^2 are isotopic by means of the isotopism $(\text{Id}, \text{Id}, h)$, where $h(e_1) = e_1$, $h(e_2) = e_3$ and $h(e_3) = -\frac{1}{a}(e_3 - e_2)$. \square

The next result indicates the distribution of 3-dimensional Malcev magma algebras over a finite field into isotopism classes. In the proof and hereafter, any linear map between two n -dimensional algebras that is linearly defined from a permutation of the indices of their basis vectors is denoted by the corresponding permutation of the symmetric group S_n . Thus, for instance, given two 3-dimensional algebras of respective bases $\{e_1, e_2, e_3\}$ and $\{e'_1, e'_2, e'_3\}$, the permutation $(23) \in S_3$ maps e_1 to e'_1 , e_2 to e'_3 and e_3 to e'_2 . Besides, Id denotes the trivial linear map between algebras of the same dimension.

Theorem 4.3 *Let \mathbb{K} be a finite field. There exist four isotopism classes of $\mathcal{M}_3(\mathbb{K})$. They correspond to the abelian algebra and the algebras L^2 , L_0^3 and*

- $W(1; \underline{2})^{(1)}$ if $\text{char}(\mathbb{K}) = 2$.
- $\mathfrak{sl}(2, \mathbb{K})$, otherwise.

Proof. From Proposition 4.2, it is enough to study the distribution into isotopism classes of the Lie algebras in Theorem 2.7. Particularly, from Lemma 2.2, the algebras $W(1; \underline{2})^{(1)}$ and $\mathfrak{sl}(2, \mathbb{K})$, with a 3-dimensional derived algebra, are not isotopic to any other Malcev magma algebra of a distinct isomorphism class. Similarly, the algebra L^2 , with a 2-dimensional derived algebra, can only be isotopic to L_a^4 , with $a \in \mathbb{K} \setminus \{0\}$; whereas the algebra L_0^3 , with an 1-dimensional derived algebra, can only be isotopic to L_0^4 . Specifically, it is straightforward verified that the triple $(\text{Id}, \text{Id}, h)$ such that $h(e_2) = e_3$ and $h(e_3) = ae_2$ is an isotopism between L^2 and L_a^4 , whereas the triple $(\text{Id}, \text{Id}, (23))$ is an isotopism between L_0^3 and L_0^4 . \square

4.2. 4-dimensional Malcev magma algebras

The next result concludes the proof of Theorem 2.8 about the distribution of 4-dimensional Malcev magma algebras into isomorphism classes.

Proposition 4.4 *None of the 4-dimensional Lie algebras that are described in (10–14) is isomorphic to a Malcev magma algebra.*

Proof. The implementation of Algorithm 5 into a case study that takes into account the derived series and centers of a Malcev algebra, together with the Jacobi identity, enable us to assure that any possible 4-dimensional Malcev magma algebra with a 2-dimensional derived algebra and solvability index 2 is isomorphic to exactly one of the algebras M_0^3 , $M_{0,0}^6$, $M_{0,a}^7$ (with $a \in \mathbb{K}$) or M_0^{13} . Hence, the Lie algebras $M_{0,b}^6$ (with $b \in \mathbb{K} \setminus \{0\}$) and $M_{a_0}^9$ are not isomorphic to any Malcev magma algebra. Further, a similar case study enable us to assure that none of the algebras M_a^3 , $M_{a,b}^6$, $M_{a,a}^7$ and M_a^{13} , with $a, b \in \mathbb{K}$ such that $a \neq 0$, is isomorphic to a Malcev magma algebra. This is due to the fact that any possible 4-dimensional Malcev magma algebra with a 3-dimensional derived algebra is isomorphic to one of the algebras M^2 , M_{-1}^3 or $M_{a,0}^7$ if its solvability index is 2 or to $M_{1,0}^{11}$, M^{12} or M_a^{14} if its solvability index is 3. In both cases, $a \in \mathbb{K} \setminus \{0\}$. \square

We finish our study by focusing on the distribution of 4-dimensional Malcev magma algebras into isotopism classes.

Proposition 4.5 *Every 4-dimensional Malcev algebra over a finite field \mathbb{K} of characteristic distinct of two is isotopic to a Malcev magma algebra. If $\text{char}(\mathbb{K}) = 2$, then this assertion holds except for those Malcev algebras that are isomorphic to the Malcev algebra $M_{a_0}^9$.*

Proof. Let \mathbb{K} be a finite field. The implementation of the procedure *isoMalcev* in Algorithm 6 gives us the next isotopisms among the Malcev magma algebras described in Theorem 2.8 and those in (10–14). For each pair of isotopic algebras we show an isotopism (f, f, h) , which is described by means of those basis vectors that are not preserved by the transformations f and h .

a) $M^2 \simeq M_a^3 \simeq M_{b,c}^6$, for all $a, b, c \in \mathbb{K}$ such that $a \notin \{0, -1\}$ and $b \neq 0$. Here,

- The triple $(\text{Id}, \text{Id}, h)$ such that $h(e_3) = e_4$ and $h(e_4) = -ae_3 + (a+1)e_4$ is an isotopism between M^2 and M_a^3 , for all $a \in \mathbb{K} \setminus \{0, -1\}$.

- The triple $(\text{Id}, \text{Id}, h)$ such that $h(e_2) = e_2 + be_3 + ce_4$, $h(e_3) = e_4$ and $h(e_4) = e_2$ is an isotopism between M^2 and $M_{b,c}^6$, for all $b, c \in \mathbb{K}$ such that $b \neq 0$.

b) $M^0 \simeq M_a^{13}$, for all $a \in \mathbb{K} \setminus \{0\}$. Here, the triple (f, f, h) such that $f(e_2) = h(e_2) = e_3$, $f(e_3) = -e_4$, $f(e_4) = e_2 - e_4$, $h(e_3) = e_2$ and $h(e_4) = -ae_4$ is an isotopism between M^0 and M_a^{13} , for all $a \in \mathbb{K} \setminus \{0\}$.

A case study based on the same implementation of the procedure *isoMalcev* also enables us to assure that, if $\text{char}(\mathbb{K}) = 2$, then the Malcev algebra $M_{a_0}^9$ is not isotopic to any Malcev magma algebra with a 2-dimensional derived algebra and solvability index 3. \square

The proof of Proposition 4.5 enables us to assure that isotopisms do not preserve Jacobi anomalies. Specifically, we have proved that the non-Lie Malcev algebra M^0 is isotopic to any Lie algebra M_a^{13} , with $a \in \mathbb{K} \setminus \{0\}$. The next result holds straightforward.

Theorem 4.6 Any 4-dimensional Malcev algebra is isotopic to a Lie algebra. \square

The distribution of 4-dimensional Malcev algebras into isotopism classes is exposed in the next final result.

Theorem 4.7 Let \mathbb{K} be a finite field. There exist eight isotopism classes of $\mathcal{M}_3(\mathbb{K})$. They correspond to the abelian algebra and the algebras M^0 , M^2 , M_0^3 , M^4 , M^8 , M_0^{13} and M_1^{14} .

Proof. From Proposition 4.5, it is enough to study the distribution into isotopism classes of the Lie magma algebras in Theorem 2.8. This distribution has been obtained again from the implementation of the procedure *isoMalcev* in Algorithm 6 together with the results exposed in Subsection 2.1. For each pair of isotopic algebras we show an isotopism (f, f, h) , which is described by means of those basis vectors that are not preserved by the transformations f and h . Besides, each class is described according to the dimension of their derived algebras and centers and the isotopism invariants described in (6-7).

a) $\dim(\mathcal{C}_2(M)) = 1$: $M^4 \simeq M^5$. Here, the triple $(f, f, (23))$ such that $f(e_3) = e_2 + e_3$ is an isotopism between both algebras.

b) $\dim(\mathcal{C}_2(M)) = 2$:

- $\dim(Z(M)) = 0$:
 - $d_3(M) = 0$: M_0^{13} .
 - $d_3(M) = 1$: M^8 .
- $\dim(Z(M)) = 1$: $M_0^3 \simeq M_{0,a}^6 \simeq M_{0,b}^7$, for all $a, b \in \mathbb{K}$ such that $b \neq 0$. Here,
 - The triple (f, f, h) such that $f(e_4) = -e_2 + e_3 + e_4$ and $h(e_2) = e_2 - ae_4$ is an isotopism between M_0^3 and $M_{0,a}^6$, for all $a \in \mathbb{K}$.
 - The triple $(f, f, (23))$ such that $f(e_4) = e_3 + e_4$ is an isotopism between M_0^3 and $M_{0,b}^7$, for all $b \in \mathbb{K} \setminus \{0\}$.

c) $\dim(\mathcal{C}_2(M)) = 3$:

- $\dim(Z(M)) = 0$:
 - $D_3(M) = 1$: If $\text{char}(\mathbb{K}) = 2$, then $M^0 \simeq M_{1,0}^{11} \simeq W(1; \underline{2})$. Otherwise, $M^0 \simeq M^{12}$. Here,
 - * If $\text{char}(\mathbb{K}) = 2$, then the triple $((14)(23), (14)(23), \text{Id})$ is an isotopism between M^0 and $M_{1,0}^{11}$, whereas the triple $((1324), (1324), (143))$ is an isotopism between M^0 and $W(1; \underline{2})$.
 - * If $\text{char}(\mathbb{K}) \neq 2$, then the triple (f, f, h) such that $f(e_2) = e_3$, $f(e_3) = -2e_4$, $f(e_4) = e_2 + e_4$, $h(e_2) = 2e_3$, $h(e_3) = 2e_4$ and $h(e_4) = -e_2 - e_4$ is an isotopism between M^0 and M^{12} .
 - $D_3(M) = 3$: $M^2 \simeq M_{-1}^3 \simeq M_{a,a}^7 \simeq M_{c,0}^7$, for all $a, c \in \mathbb{K} \setminus \{0\}$. Here,
 - * The triple $(\text{Id}, \text{Id}, (34))$ is an isotopism between M^2 and M_{-1}^3 .
 - * The triple $(\text{Id}, \text{Id}, h)$ such that $h(e_2) = e_3$, $h(e_3) = e_4$ and $h(e_4) = ae_2 + ae_3$ is an isotopism between M^2 and $M_{a,a}^7$, for all $a \in \mathbb{K} \setminus \{0\}$.
 - * The triple $(\text{Id}, \text{Id}, h)$ such that $h(e_2) = e_3$, $h(e_3) = e_4$ and $h(e_4) = ce_2$ is an isotopism between M^2 and $M_{c,0}^7$, for all $c \in \mathbb{K} \setminus \{0\}$.

- $\dim(Z(M)) = 1$: $M_1^{14} \simeq M_a^{14}$, for all $a \in \mathbb{K} \setminus \{0\}$. If $\text{char}(\mathbb{K}) = 2$, then $M_1^{14} \simeq W(1; \underline{2})^{(1)} \oplus Z(L)$. Otherwise, $M_1^{14} \simeq \mathfrak{gl}(2, \mathbb{K})$. Here,
 - The triple $(\text{Id}, \text{Id}, h)$ such that $h(e_2) = ae_2$ is an isotopism between M_1^{14} and M_a^{14} , for all $a \in \mathbb{K} \setminus \{0\}$.
 - If $\text{char}(\mathbb{K}) = 2$, then the triple $(\text{Id}, \text{Id}, (1432))$ is an isotopism between M_1^{14} and $W(1; \underline{2})^{(1)} \oplus Z(L)$.
 - If $\text{char}(\mathbb{K}) \neq 2$, then the triple (f, f, h) such that $f(e_4) = e_1 + e_4$, $h(e_1) = e_4$, $h(e_2) = -e_3$, $h(e_3) = e_2$ and $h(e_4) = e_1$ is an isotopism between M_1^{14} and $\mathfrak{gl}(2, \mathbb{K})$. □

5. Conclusions and further studies

We have defined in this paper distinct zero-dimensional radical ideals whose related algebraic sets are uniquely identified with the set $\mathcal{M}_n(\mathbb{K})$ of n -dimensional Malcev magma algebras over a finite field \mathbb{K} . The computation of their reduced Gröbner bases together with the classification of Lie algebras over finite fields given by De Graaf and Strade have enabled us to determine the distribution of $\mathcal{M}_3(\mathbb{K})$ and $\mathcal{M}_4(\mathbb{K})$ not only into isomorphism classes, which is the usual criterion, but also into isotopism classes. Particularly, we have proved the existence of four isotopism classes in $\mathcal{M}_3(\mathbb{K})$ and eight isotopism classes in $\mathcal{M}_4(\mathbb{K})$. Besides, we have proved that every 3-dimensional Malcev algebra over any finite field and every 4-dimensional Malcev algebra over a finite field of characteristic distinct of two is isotopic to a Lie magma algebra. Keeping in mind the obtained results, the study of magma algebras by means of computational algebraic geometry constitutes a good first approach to the distribution of Malcev algebras over finite fields into isomorphism and isotopism classes. In this regard, the study of the sets $\mathcal{M}_5(\mathbb{K})$ and $\mathcal{M}_6(\mathbb{K})$ is established as a further work that complements the already known classification of 5- and 6-dimensional Malcev algebras in case of non-solvability [14] and nilpotency [15].

References

1. Löhmus J., Paal E, Sorgsepp L. About Nonassociativity in Mathematics and Physics. *Acta Applicandae Mathematicae* 1998; 50: 3–31.
2. Sabinin L., Sbitneva L., Shestakov I. *Non-Associative Algebra and Its Applications*. CRC Press: Boca Raton, FL; 2006.
3. Malcev AI. Analytic loops. *Mat. Sb. N.S.* 1955; , 36(78): 569–576.
4. Sagle AA. Malcev algebras. *Trans. Amer. Math. Soc.* 1961; 101:426–458.
5. Dirac PAM. *Lectures on Quantum Mechanics*. Yeshiva University: New York; 1964.
6. Lipkin HJ., Weisberger WI., Peshkin M., Magnetic Charge Quantization and Angular Momentum. *Ann. Phys.* 1969; 53: 203-214.
7. Günaydin M., Zumino B. Magnetic Charge And Nonassociative Algebras. *Proceedings of the Conference "Old and New Problems in Fundamental Physics: Symposium in honor of G. C. Wick"* 1986; 43–54.
8. Günaydin M. Exceptionality, supersymmetry and non-associativity in Physics. *Bruno Zumino Memorial Meeting*. CERN, Geneva; 2015.
9. Günaydin M., Minic D. Nonassociativity, Malcev algebras and string theory. *Fortschr. Phys.* 2013; 61: 873–892.
10. Kuzmin EN. Malcev algebras of dimension five over a field of zero characteristic. *Algebra Logika* 1971; 9: 691–700.
11. Kuzmin EN. Structure theory of Malcev algebras. *Radical theory* (Eger, 1982), North-Holland, Amsterdam, 1985; 38: 229–235.
12. Sagle AA. Simple Malcev algebras over fields of characteristic zero. *Pacific J. Math.* 1962; 12: 1057–1078.
13. De Graaf W. Classification of Solvable Lie Algebras *Exp. Math.* 2005; 14: 15–25.
14. Strade H. Lie algebras of small dimension. In: Lie algebras, vertex operator algebras and their applications, *Contemp. Math.* 2007; 442: 233–265.
15. Cicalo S., De Graaf W., Schneider C. Six-dimensional nilpotent Lie algebras. *Linear Algebra Appl.* 2012; 436: 163–189.
16. Albert AA. Non-Associative Algebras: I. Fundamental Concepts and Isotopy, *Ann. of Math., Second Series* 1942; 43
17. Bruck RH. Some results in the theory of quasigroups. *Trans. Amer. Math. Soc.* 1944; 55: 19–52.
18. Sandler R., Autotopism groups of some finite non-associative algebras, *Amer. J. Math.* 1962; 84: 239–264.
19. Petersson HP. Quasi composition algebras, *Abh. Math. Sem. Univ. Hamburg* 1971; 35: 215–222.
20. Benkart G. and Osborn JM. Real division algebras and other algebras motivated by physics, *Hadronic J.* 1980/81; 4: 392–443.
21. Dieterich E. Classification, automorphism groups and categorical structure of the two-dimensional real division algebras, *J. Algebra Appl.* 2005; 4: 517–538.
22. Babikov M. Isotopy and identities in alternative algebras, *Proc. Amer. Math. Soc.* 1997; 125: 1571–1575.
23. McCrimmon K. Homotopes of alternative algebras, *Math. Ann.* 1971; 191: 253–262.

24. Jiménez-Gestal C. and Pérez-Izquierdo JM. Ternary derivations of finite-dimensional real division algebras, *Linear Algebra Appl.* 2008; 428: 2192–2219.
25. Falcón OJ., Falcón RM., Núñez J. Isotopism and isomorphism classes of certain Lie algebras over finite fields *Results in Math.* 2015. In press. DOI 10.1007/s00025-015-0502-y.
26. Oehmke RH. and Sandler R. The collineation groups of division ring planes. I. Jordan division algebras, *J. Reine Angew. Math.* 1964; 216: 67–87.
27. Petersson HP. Isotopisms of Jordan algebras, *Proc. Amer. Math. Soc.* 1969; 20: 477–482.
28. McCrimmon K. Inner ideals in quadratic Jordan algebras. *Trans. Amer. Math. Soc.* 1971; 159: 445–468.
29. McCrimmon K. The generic norm of an isotope of a Jordan algebra, *Scripta Math.* 1973; 29: 229–236.
30. Petersson HP. The isotopy problem for Jordan matrix algebras, *Trans. Amer. Math. Soc.* 1978; 244: 185–197.
31. Thakur ML. Isotopy and invariants of Albert algebras, *Comment. Math. Helv.* 1999; 74: 297–305.
32. Pchelintsev SV. Isotopes of prime $(-1, 1)$ - and Jordan algebras, *Algebra Logika* 2010; 49: 388–423.
33. Holgate P. Genetic algebras associated with polyploidy, *Proc. Edinburgh Math. Soc.* 1966; 15: 1–9.
34. Campos TMM. and Holgate, P. Algebraic Isotopy in Genetics, *IMA J. Math. Appl. Med. Biol.* 1987; 4: 215–222.
35. Albert AA. Absolute valued real algebras, *Ann. of Math.* 1947; 48: 495–501.
36. Cuenca Mira JA., Darpö E. and Dieterich E. Classification of the finite dimensional absolute valued algebras having a non-zero central idempotent or a one-sided unity, *Bull. Sci. Math.* 2010; 134: 247–277.
37. Allison BN. and Hein W. Isotopes of some nonassociative algebras with involution, *J. Algebra* 1981; 69: 120–142.
38. Balanov Z. and Krasnov Y. Complex structures in real algebras. I. Two-dimensional commutative case, *Comm. Algebra* 2003; 31: 4571–4609.
39. Cox DA., Little JB., O’Shea D. *Using Algebraic Geometry*. Springer-Verlag: New York; 1998.
40. Cox DA., Little JB., O’Shea D. *Ideals, varieties, and algorithms. An introduction to computational algebraic geometry and commutative algebra*. Springer: New York; 2007.
41. Kuzmin EN. Binary Lie algebras of small dimensions, *Algebra and Logic* 1998; 37: 181–186.
42. Hillebrand D. *Triangulierung nulldimensionaler Ideale - Implementierung und Vergleich zweier Algorithmen*. Universitaet Dortmund: Fachbereich Mathematik; 1999.
43. Lazard D. Solving zero-dimensional algebraic systems. *J. Symbolic Comput.* 1992; 13: 117–131.
44. Möller HM. On decomposing systems of polynomial equations with finitely many solutions. *Appl. Algebra Engrg. Comm. Comput.* 1993; 4: 217–230.
45. Buchberger B. An algorithm for finding the basis elements of the residue class ring of a zero dimensional polynomial ideal. *J. Symbolic Comput.* 2006; 41: 475–511.
46. Faugère JC. A new efficient algorithm for computing Gröbner bases (F_4). *J. Pure Appl. Algebra* 1999; 139: 61–88.
47. Faugère JC. A new efficient algorithm for computing Gröbner bases without reduction to zero (F_5). *Proceedings of the 2002 International Symposium on Symbolic and Algebraic Computation*, ACM. New York; 2002: 75–83.
48. Brickenstein M. Slimgb: Gröbner bases with slim polynomials, *Rev. Mat. Complut.* 2010; 23: 453–466.
49. Hashemi A. Nullstellenstze for zero-dimensional Gröbner bases. *Comput. Complexity* 2009; 18: 155–168.
50. Hashemi A., Lazard D. Sharper complexity bounds for zero-dimensional Gröbner bases and polynomial system solving. *Internat. J. Algebra Comput.* 2011; 21: 703–713.
51. Gao S. *Counting Zeros over Finite Fields Using Gröbner Bases*. Carnegie Mellon University, 2009.
52. Decker W., Greuel GM., Pfister G., Schönemann H. Singular 4-0-2 — *A computer algebra system for polynomial computations* 2015.
53. Jacobson N. *Lie algebras*. Dover Publications: New York; 1979.