



# A hydrogen bond network in the active site of *Anabaena* ferredoxin-NADP<sup>+</sup> reductase modulates its catalytic efficiency

Q1 Ana Sánchez-Azqueta<sup>a,b</sup>, Beatriz Herguedas<sup>a,b</sup>, Ramón Hurtado-Guerrero<sup>a,b,c</sup>, Manuel Hervás<sup>d</sup>,  
 4 José A. Navarro<sup>d</sup>, Marta Martínez-Júlvez<sup>a,b</sup>, Milagros Medina<sup>a,b,\*</sup>

5 <sup>a</sup> Departamento de Bioquímica y Biología Molecular y Celular, Facultad de Ciencias, Universidad de Zaragoza, Zaragoza, Spain

6 <sup>b</sup> Institute of Biocomputation and Physics of Complex Systems (BIFI)-Joint Unit BIFI-IQFR (CSIC), Universidad de Zaragoza, Zaragoza, Spain

7 <sup>c</sup> Fundación ARAID, Gobierno de Aragón, Spain

8 <sup>d</sup> Instituto de Bioquímica Vegetal y Fotosíntesis, cicCartuja, Universidad de Sevilla & CSIC, Sevilla, Spain

## ARTICLE INFO

### Article history:

11 Received 19 July 2013

12 Received in revised form 13 October 2013

13 Accepted 23 October 2013

14 Available online xxxx

### Keywords:

20 Ferredoxin-NADP<sup>+</sup> reductase

21 Flavoenzyme

22 Isoalloxazine:nicotinamide interaction

23 Hydride transfer

24 Catalytically competent interaction

25 Charge-transfer complex

26 Site-directed mutagenesis

27 Kinetic isotope effect

## ABSTRACT

10 Ferredoxin-NADP<sup>+</sup> reductase (FNR) catalyses the production of NADPH in photosynthetic organisms, where its  
 11 FAD cofactor takes two electrons from two reduced ferredoxin (Fd) molecules in two sequential steps, and trans-  
 12 fers them to NADP<sup>+</sup> in a single hydride transfer (HT) step. Despite the good knowledge of this catalytic machin-  
 13 ery, additional roles can still be envisaged for already reported key residues, and new features are added to  
 14 residues not previously identified as having a particular role in the mechanism. Here, we analyse for the first  
 15 time the role of Ser59 in *Anabaena* FNR, a residue suggested by recent theoretical simulations as putatively in-  
 16 volved in competent binding of the coenzyme in the active site by cooperating with Ser80. We show that  
 17 Ser59 indirectly modulates the geometry of the active site, the interaction with substrates and the electronic  
 18 properties of the isoalloxazine ring, and in consequence the electron transfer (ET) and HT processes. Additionally,  
 19 we revise the role of Tyr79 and Ser80, previously investigated in homologous enzymes from plants. Our results  
 20 probe that the active site of FNR is tuned by a H-bond network that involves the side-chains of these residues  
 21 and that results to critical optimal substrate binding, exchange of electrons and, particularly, competent disposi-  
 22 tion of the C4n (hydride acceptor/donor) of the nicotinamide moiety of the coenzyme during the reversible HT  
 23 event.

© 2013 Elsevier B.V. All rights reserved.


## 1. Introduction

1 In the photosynthetic electron transfer (ET) chain of plants, algae  
 2 and cyanobacteria, the isoalloxazine ring of the FAD cofactor of  
 3 ferredoxin-NADP<sup>+</sup> reductase (FNR) gets reduced to its hydroquinone  
 4 state by sequentially accepting two electrons from two ferredoxin  
 5 (Fd) molecules. Subsequently, it transfers a hydride from the N5 atom  
 6 (N5Hi) of the isoalloxazine of its FAD cofactor to the nicotinamide C4  
 7 atom of NADP<sup>+</sup> (C4n) to provide the cell with reduction power in the  
 8 form of NADPH [1–3]. The overall ET process from Fd to NADP<sup>+</sup> is re-  
 9 versible, with transitory ternary complexes, Fd:FNR:NADP<sup>+</sup>, formed  
 10 during catalysis [4]. Structural, mutational and theoretical studies re-  
 11 vealed residues on the protein surface and in the isoalloxazine environ-  
 12 ment involved in the interaction and ET with the protein partner,  
 13 contributing to the optimal architecture of the active site for proton  
 14 and electron transfer, as well as playing key roles in the catalytic binding  
 15 of the nicotinamide moiety of the coenzyme (NMN) during the hydride  
 16 transfer (HT) event [5–13]. Among them, a particular role is proposed  
 17 for the C-terminal Tyr (Tyr303 in *Anabaena* FNR (AnFNR), numbering  
 18 used herein) (Fig. 1A). This residue stacks at the re-face of the isoallox-  
 19 azine ring of FAD, modulates its midpoint reduction potential and re-  
 20 duces the probability of a too strong stacking interaction between the

Q5 **Abbreviations:** FNR, FNR<sub>ox</sub>, FNR<sub>hq</sub>, FNR<sub>sq</sub>, ferredoxin-NADP<sup>+</sup> reductase and FNR in  
 the fully oxidised, anionic hydroquinone (fully reduced) and neutral semiquinone  
 (one-electron reduced) states, respectively; Fd, Fd<sub>rd</sub>, ferredoxin and in its the reduced  
 state; 2'-P, 2'-phosphate group of NADP<sup>+</sup>/H; dRf, 5-deazariboflavin; ET, electron  
 transfer; HT, hydride transfer; DT, deuteride transfer; WT, wild-type; CTC, charge-  
 transfer complex; CTC-1, FNR<sub>ox</sub>-NADPH CTC; CTC-2, FNR<sub>hq</sub>-NADP<sup>+</sup> CTC; NMN, nico-  
 tinamide nucleotide moiety of NADP<sup>+</sup>/H; 2'-P-AMP, 2'-P-AMP moiety of NADP<sup>+</sup>/H;  
 PP<sub>i</sub>, pyrophosphate; N5Hi, N5i, N5 hydride donor/acceptor of the FADH<sup>•</sup>/FAD iso-  
 lloxazine ring of FNR; C4n, C4Hn, C4 hydride acceptor/donor of the NADP<sup>+</sup>/H nico-  
 tinamide ring;  $k_A \rightarrow B$ ,  $k_B \rightarrow C$ , apparent/observed rate constants obtained by global  
 analysis of spectral kinetic data;  $k_{obsHT}$ ,  $k_{obsHT-1}$ ,  $k_{obsDT}$ ,  $k_{obsDT-1}$ , observed conversion  
 HT and DT rate constants for the forward and reverse reactions;  $k_{HT}$ ,  $k_{HT-1}$ , hydride  
 transfer first-order rate constants for the forward and reverse reactions, respectively;  
 $k_{DT}$ ,  $k_{DT-1}$ , deuteride transfer first-order rate constants for the forward and reverse re-  
 actions, respectively;  $K_d^{NADPH}$ ,  $K_d^{NADP^+}$ , dissociation constants for the intermediate  
 complexes in the reduction and reoxidation of FNR, respectively; KIE, kinetic isotopic  
 effect;  $A_H$ ,  $A_D$ , Arrhenius preexponential factors for hydrogen and deuteride, respec-  
 tively;  $E_{aH}$ ,  $E_{aD}$ , activation energies for hydride and deuteride transfer, respectively;  
 $k_{et}$ , first-order electron transfer rate;  $k_2$ , second-order rate constant for bimolecular  
 electron transfer;  $I$ , ionic strength

\* Corresponding author at: Departamento de Bioquímica y Biología Molecular y Celular,  
 Facultad de Ciencias, Universidad de Zaragoza, E-50009 Zaragoza, Spain. Tel.: +34 976  
 762476; fax: +34 976 762123.

E-mail address: mmedina@unizar.es (M. Medina).



**Fig. 1.** Key residues at the AnFNR active site. (A) Surface representation of the active site environment at the equilibrium of a MD simulation of a theoretical catalytically competent WT FNR<sub>100</sub>:NADP<sup>+</sup> complex [15]. NADP<sup>+</sup>, FAD, and selected key side-chains are shown in sticks with C in blue, orange and wheat, respectively. Selected water molecules at the active site are also shown as balls and sticks. (B) Sequence alignment of different members of the FNR superfamily (ClustalW2). Position of residues equivalent to those mutated in this work is shown in bold.

isoalloxazine and nicotinamide rings, thus contributing to the optimal geometry among the N5i, the C4n and the hydrogen that have to be transferred between them [5,12,14–18]. A second highly conserved aromatic side-chain, Tyr79, stacks at the isoalloxazine *si*-face with its hydroxyl H-bonding the 4'-ribityl hydroxyl of FAD, which is also connected through a complex H-bond network assisted by water molecules to the C2 of the isoalloxazine and to the side-chain Arg100 [8,19–22]. Other key highly conserved residues at the active site are the neighbours of Tyr303 at the *re*-face: Ser80, Cys261 and Glu301 [3,8,11,15,23–26]. They contribute to the

fine modulation of the FAD midpoint reduction potential, the affinity for Fd, the architecture of the catalytically competent complex, and/or the ET and HT rates [5,7,10,11,26–28]. Despite the fact that structural changes detected upon spinach FNR reduction are minor, they implicate a slight approach of the hydroxyl of Ser96 (Ser80 in AnFNR) to N5i that loses its H-bond with Glu312 (Glu301 in AnFNR) leading to the displacement of Tyr314 (Tyr303 in AnFNR) away from the flavin ring (decreasing the  $\pi$ - $\pi$  stacking with the reduced isoalloxazine), as well as the displacement of two highly conserved water molecules, W406 and W571, (W404 and W457 in AnFNR) located near the

ribityl of FAD [29]. Theoretical calculations for the HT process between the isoalloxazine and the nicotinamide of the coenzyme, also supported by experimental evidences, confirm that the sulphydryl group of Cys261 contributes to the approach of the N7n in the NMN amide to the isoalloxazine along the reaction path, while the O7n in the same amide H-bonds Ser80 (Fig. 1A) [18,28]. Ser80 is kept in the C4n and N5Hi atom environments along the reaction coordinate, contributing to a network of interactions involving the isoalloxazine ring, the nicotinamide ring, Cys261 and Ser80 itself (Fig. 1A). This network facilitates the approach of the reacting N5Hi and C4n atoms, and therefore, it is expected to contribute towards the adequate geometry for the chemical step of the reaction [6,10,11,15]. Thus, the architecture of the active site for the HT event must precisely contribute to the orientation of the N5Hi of the FAD<sub>hq</sub> isoalloxazine and the C4n of the coenzyme nicotinamide rings and, therefore, to the efficiency of the HT process [14–16]. Molecular dynamics (MD) simulations additionally indicated that H-bonds between the side-chains of Ser80 and Ser59 are highly populated (Fig. 1A) [15]. These side-chains also H-bond Glu301, proposed by theoretical and experimental evidences to switch positions in and out of the active site to provide a pathway for proton transfer between the external medium and N5i, via Ser80, during FNR reduction by Fd [14,27]. This route might also include Ser59, highly conserved in the plant type FNR family (Fig. 1B) [21,29].

In this study, we further analyse the roles of Ser59, Tyr79 and Ser80 in AnFNR during catalysis to better understand the function of the interacting network to which they contribute to within the active site. The presented results provide information about the role of this interacting network, indicating that it modulates the electronic environment of the isoalloxazine ring and influences the ET process from Fd, as well as the active site geometry during HT. Particularly, we show for the first time that Ser59 indirectly modulates the geometry of the active site, the interaction with substrates and the efficiency of the ET and HT processes. Additionally, the roles of these side-chains in the competent placement of the C4n (hydride acceptor/donor) atom and in the tunnelling contribution during the HT event have been analysed, being particularly relevant those of Ser80.

## 2. Materials and methods

### 2.1. Biological material

pET28a-AnFNR plasmids containing the S59A, S80A or Y79F mutations were obtained from the company Mutagenex® and used to produce and purify the corresponding proteins from *Escherichia coli* cultures as previously reported [30]. Samples were prepared in 50 mM Tris/HCl, pH 8.0. S80A FNR was further purified using a HiPrep TM 26/60 Sephacryl S-200 HR column (GE Healthcare). FNR<sub>hq</sub> variants were obtained by anaerobic photoreduction of the samples in the presence of 2 μM 5-deazariboflavin (dRf) and 3 mM EDTA in 50 mM Tris/HCl, pH 8.0, by irradiation from a 250 W light source [31]. Deuterated FNR<sub>hq</sub> (D-FNR<sub>hq</sub>) variants were produced by photoreduction with EDTA and dRf of the corresponding FNR<sub>ox</sub> previously dialysed in 50 mM Tris/DCl, pD ~ 8.0 in D<sub>2</sub>O. NADPD (4R-form, with the deuterium in the A face of the nicotinamide) was produced and purified as described [14,32]. *Anabaena* Fd (AnFd) was produced as previously described [12].

### 2.2. Spectroscopic assays

UV/vis spectra were recorded in a Cary-100 spectrophotometer. The molar absorption coefficient for each FNR variant was spectrophotometrically determined by thermal denaturation of the protein for 10 min at 90 °C, followed by centrifugation and separation of the precipitated apoprotein, and spectroscopic quantification of the FAD released to the supernatant [33]. Interaction parameters with NADP<sup>+</sup>, NAD<sup>+</sup> and Fd were determined by difference absorption spectroscopy

at 25 °C in 50 mM Tris/HCl, pH 8.0, as previously described [7,34]. Titrations were carried out by adding aliquots of 1 mM NADP<sup>+</sup> or Fd, and 50 mM NAD<sup>+</sup> to 20–80 μM FNR solutions. Errors in the determination of  $K_d$  and  $\Delta\epsilon$  were  $\pm 10\%$  and  $\pm 5\%$ , respectively.

### 2.3. Determination of midpoint reduction potentials of the FNR variants

Midpoint reduction potentials for the ox/hq couple ( $E_{ox/hq}$ , two-electron reduction process) of WT, S59A, Y79F, and S80A FNRs were determined at 25 °C by potentiometric titration under anaerobic conditions using a gold electrode and a calomel electrode as reference ( $E_m = +244.4$  mV at 25 °C). Due to the low degree of FNR semiquinone stabilisation it was not possible to measure the potential for the two one-electron steps. Typically, the solution contained ~20 μM FNR, 50 mM Tris/HCl buffer, pH 8.0, 3 mM EDTA and 2 μM dRf. 0.02% n-dodecyl-β-D-maltoside was also added to S80A FNR to increase its stability. Methylviologen ( $E_m = -446$  mV) and benzylviologen ( $E_m = -359$  mV) were additionally used as mediators. Solutions were made anaerobic over a 2–4 h period. Stepwise reduction of the protein was achieved by photoreduction using the equipment previously described [35]. The system was considered equilibrated when the potential ( $E$ ), measured with a Fluke 177 true-RMS multimeter, remained stable for at least 10 min. The UV/vis absorbance spectrum was then recorded and used to determine [FNR<sub>ox</sub>] and [FNR<sub>hq</sub>] at the equilibrium after each reduction step.  $E_{ox/hq}$  was calculated by linear regression analysis according to the Nernst equation. The values of each one-electron single step,  $E_{ox/sq}$  and  $E_{sq/hq}$ , were derived from Eqs. (1) and (2) using the experimentally determined  $E_{ox/hq}$  and the molar fraction of the maximum percentage of SQ stabilised.

$$E_{ox/sq} - E_{sq/hq} = 0.11 * \log \frac{2[SQ]}{1-[SQ]}; \quad (1)$$

$$\frac{E_{ox/sq} + E_{sq/hq}}{2} = E_{ox/hq}; \quad (2)$$

Error in the determined  $E_{ox/hq}$ ,  $E_{ox/sq}$  and  $E_{sq/hq}$  values was estimated to be  $\pm 5$  mV.

### 2.4. Steady-state kinetics measurements


The diaphorase activity of FNR was determined in a double beam Cary-100 spectrophotometer using either 2,6-dichlorophenolindophenol (DCPIP) ( $\Delta\epsilon_{620\text{ nm}} 21 \text{ mM}^{-1} \text{ cm}^{-1}$ ) or  $\text{K}_3\text{Fe}(\text{CN})_6$  ( $\Delta\epsilon_{420\text{ nm}} 1.05 \text{ mM}^{-1} \text{ cm}^{-1}$ ) as two- or one-electron acceptors, respectively. The final reaction mixture contained 4 nM FNR, 0.1 mM DCPIP or 1.5 mM  $\text{K}_3\text{Fe}(\text{CN})_6$ , and NADPH in the range 0–200 μM, while the reference cuvette contained 0.06 mM DCPIP when using this acceptor. Higher concentrations of FNR (1 μM) and/or nucleotide (0–5 mM) were required for the analysis of the reactions with NADH. The NADPH-dependent cytochrome c reductase activity was determined using AnFd, and horse heart cytochrome c (Cytc) as final electron acceptor. Reaction mixtures contained 4 nM FNR, 200 μM NADPH, 0.75 mg/ml Cytc and 0–15 μM AnFd. All measurements were carried out in 50 mM Tris/HCl, pH 8.0, at 25 °C.  $K_m$  and  $k_{cat}$  values were obtained by fitting the dependence of the observed initial rates on coenzyme concentration to the Michaelis–Menten equation. Estimated errors in  $K_m$  and  $k_{cat}$  were  $\pm 20\%$  and  $\pm 10\%$ , respectively.

### 2.5. Laser-flash induced kinetics

Laser-flash experiments were performed anaerobically at 25 °C in a 1 cm path-length cuvette using EDTA as electron donor and dRf as photosensitizer as previously described [7,36]. The standard reaction mixture contained, in a final volume of 1.5 mL, 4 mM sodium phosphate, 206



pH 7.0, 2 mM EDTA and 100  $\mu$ M dRf (low ionic strength (*I*) buffer). The laser-generated dRf triplet abstracts a hydrogen atom from EDTA which is present in large excess and produces the dRf semiquinone (dRfH•) which, in competition with its own disproportionation, reduces the oxidised protein. Direct reduction of AnFNR<sub>ox</sub> by the laser-flash photoreduced dRf (dRfH•) was followed by measuring the decrease of absorbance in the flavin band-I maxima at 458 nm. When both Fd and FNR are present simultaneously in the solution, the flash generated dRfH• reacts almost exclusively with free Fd<sub>ox</sub>, and thus the subsequent ET process from the generated Fd<sub>rd</sub> to FNR<sub>ox</sub> can be monitored [7,12,13]. FNR reduction by Fd<sub>rd</sub> was followed as the increase of absorbance at 600 nm, a wavelength at which the production of FNR<sub>sq</sub> can be monitored as FNR<sub>ox</sub> is reduced by Fd<sub>rd</sub>. Control experiments collected at 489–500 nm, an isosbestic point of the FNR<sub>ox</sub>/sq couple, allowed to monitor the oxidation of Fd<sub>rd</sub>, yielding rate constants that were the same, within experimental error, as those determined from the 600 nm data, as expected from the two step mechanism shown in Eq. (3). For these experiments, 40  $\mu$ M AnFd and AnFNR at varying concentrations were added to the standard reaction mixture, either in the absence or in the presence of 100 mM NaCl. For *I* dependence experiments, small amounts of a concentrated solution of 5 M NaCl were added to a reaction cuvette containing the low *I* buffer, 40  $\mu$ M AnFd and 30  $\mu$ M AnFNR. All experiments were performed under pseudo-first-order conditions, for which the amount of acceptor (FNR<sub>ox</sub>) was maintained well in excess over the amount of the generated Fd<sub>rd</sub> (<1  $\mu$ M). Each kinetic trace was the average of 8–15 measurements. All kinetic traces were fitted to monoexponential curves by using the Marquardt method to obtain the observed rate constants ( $k_{obs}$ ). Linear fittings of  $k_{obs}$  values on FNR concentration allowed obtaining second-order bimolecular rate constants ( $k_2$ ). Non-linear  $k_{obs}$  dependences on FNR concentration were adjusted to a two-step mechanism, given in Eq. (3) [37], to estimate minimal values of both the complex dissociation constant ( $K_d$ ) and the ET rate constant ( $k_{et}$ ).



Errors in the estimated values of  $K_d$  and  $k_{et}$  were  $\pm 20\%$  and  $\pm 10\%$ , respectively.

## 2.6. Stopped-flow pre-steady-state kinetic measurements

Transient charge transfer complex (CTC) formation and HT processes between the FNR<sub>hq/ox</sub> variants and NADP<sup>+</sup>/H were followed by stopped-flow in 50 mM Tris/HCl, pH 8.0, at 6 °C and under anaerobic conditions [14,38]. Final FNR concentrations were 25  $\mu$ M, while a 25–250  $\mu$ M range was used for the nucleotide. Reactions were followed by the evolution of the absorption spectra (400–1000 nm) using an Applied Photophysics SX17.MV stopped-flow equipment with a photodiode array detector (App. Photo. Ltd.). Typically, spectra were collected every 2.5 ms. Multiple wavelength absorption data were processed using the X-Scan software (App. Photo. Ltd.). Analysis of time dependent spectral changes was performed by global analysis and numerical integration methods using Pro-Kineticist (App. Photo. Ltd.). Data were fit to a single step model allowing estimation of the apparent conversion rate constants ( $k_{A \rightarrow B}$ ,  $k_{B \rightarrow C}$ ). In general, the first spectra after mixing show formation of some amount of CTC; this means a previous reaction,  $A \rightarrow B$ , has occurred in the instrumental dead time (2–3 ms in our conditions), then we correlate the observed reaction with a  $B \rightarrow C$  model. A, B and C are spectral species, reflecting a distribution of enzyme intermediates (reactants, CTCs, products, Michaelis-complexes) at a certain point along the reaction time course, and do not necessarily represent a single distinct enzyme intermediate. Moreover, none of them represents individual species, and their spectra cannot be included as fixed values in the global-fitting. Model validity was assessed by lack of systematic deviations from residual plots at different wavelengths,

inspection of calculated spectra and consistence among the number of significant singular values with the fit model. The apparent rate constants as a function of coenzyme concentration were globally fit to the reaction mechanisms including all the experimental data for processes in both directions (Figure S2C) [14]. In the simplest case the time-course of the reaction ( $k_{A \rightarrow B}$ ) will be equal to the sum of the rates for the forward ( $k_{HT}$ ) HT and reverse ( $k_{HT-1}$ ) HT at equilibrium [14,39]. In the presence of excess of coenzyme the dependence of  $k_{HT}$  and  $k_{HT-1}$  on substrate concentration is given by standard functions for substrate saturation and competitive inhibition (inhibition constant  $K_i$ ), respectively:

$$k_{A \rightarrow B} = \frac{[\text{NADPH}]k_{HT}}{[\text{NADPH}] + K_{\text{NADPH}}} + \frac{[\text{NADP}^+]k_{HT-1}}{[\text{NADP}^+] + K_{\text{NADP}^+}(1 + [\text{NADPH}]/K_i)} \quad (4)$$

The concentration of NADP<sup>+</sup>/H at equilibrium can also be estimated from the difference in the midpoint reduction potentials for NADP<sup>+</sup>/H and FNR<sub>ox/hq</sub> redox couples ( $\Delta E_m$ ) and the total concentration of enzyme ( $E_t$ ) (Eq. (5)):

$$[\text{NADP}^+] = [\text{NADPH}] \frac{(1 + 4\text{FNR}_t \cdot 10^{\Delta E_m/29.5} / [\text{NADPH}])^{1/2} - 1}{2 \cdot 10^{\Delta E_m/29.5}} \quad (5)$$

Errors in the determination of these kinetic constants were  $\pm 15\%$ –20%.

Single-wavelength kinetic traces were recorded for accurate estimation of the hydride or deuteride transfer (HT and DT, respectively) rate constants ( $k_{obsHT/HT-1}$  or  $k_{obsDT/DT-1}$ ) in the temperature dependence assays. In these cases traces at 458 nm were recorded with the single-wavelength monochromator using the SX18.MV software (App. Photo. Ltd.). A 1:1 enzyme:coenzyme concentration ratio was used for these experiments using temperatures between 5.3 and 17.3 °C, since, due to the reversibility of the process, this ratio relates with the maximal experimental values for  $k_{obsHT/HT-1}$  or  $k_{obsDT/DT-1}$ . Traces were fit to monoexponential decays to determine  $k_{obsHT}$ ,  $k_{obsHT-1}$ ,  $k_{obsDT}$  and  $k_{obsDT-1}$ . Errors in the determination of these kinetic constants were  $\pm 10\%$ . The kinetic isotope effects (KIEs) were calculated as:

$$\text{KIE} = \frac{k_{obsHT}}{k_{obsDT}} \quad \text{or} \quad \text{KIE} = \frac{k_{obsHT-1}}{k_{obsDT-1}} \quad (6)$$

Fitting the experimentally obtained rates to the Arrhenius equation allowed determining the Arrhenius pre-exponential factors ( $A_H$  and  $A_D$ ) and the Activation Energy values ( $E_{aH}$  and  $E_{aD}$ ). Combination of the Arrhenius equation with Eq. (6) leads to the graphical representation of the temperature dependence of the KIE.

## 2.7. Crystal growth, data collection and structure refinement

Crystals of S59A, Y79F and S80A AnFNR were produced under the same conditions previously reported for the WT [5,22], while those for the S80A FNR:NADP<sup>+</sup> complex were similarly obtained but without (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and adding 1  $\mu$ L of 10 mM NADP<sup>+</sup> to the drop. X-ray data sets for S59A and S80A AnFNR were collected on a Bruker-Incoatec 1  $\mu$ s microfocus generator with an Axiom detector. A microstar generator with an Image Plate detector was used to collect data for Y79F FNR, whereas data for the S80A FNR:NADP<sup>+</sup> complex were collected on the ID23-1 line at ESRF (Grenoble, France). Data were processed with Proteum Suite (Bruker) and XDS [40] and scaled and reduced with SCALA from CCP4 [41]. All structures were solved by MOLREP [42] from CCP4, using the structures of WT AnFNR (PDB ID: 1QUE) and the AnFNR:NADP<sup>+</sup> complex (PDB ID: 1GJR) as reference models. Refinement of all structures was performed with CCP4 and COOT [43] and SFCHECK [44]. PROCHECK [45] and MOLPROBITY [46] were used to

326 assess final structures. S59A, Y79F and S80A FNRs diffracted up to 327  
 328 1.92, 2.0 and 1.9 Å, respectively, and belonged to the P6<sub>5</sub> hexagonal 329  
 330 space group. Their  $V_m$  were 2.85, 2.86 and 2.68 Å<sup>3</sup>/Da with one FNR 331  
 332 molecule in their asymmetric units and 56.8, 57.0 and 53.7% solvent 333  
 334 contents, respectively. Each model comprised residues 9–303 (S59A 335  
 336 and Y79F FNRs) or 10–303 (S80A FNR), one FAD molecule, one 337  
 338 SO<sub>4</sub><sup>2-</sup> ion and water molecules. The S80A FNR:NADP<sup>+</sup> complex was 339  
 340 solved at 2.3 Å and crystals belonged to the I4 tetragonal space 341  
 342 group.  $V_m$  was 3.26 Å<sup>3</sup>/Da with two molecules in the asymmetric 343  
 344 unit and 61.93% solvent. The model included residues 9–303, one 345  
 346 FAD, one NADP<sup>+</sup> and waters. Data for collection and refinement pro- 347  
 348 cesses can be found in Table SP1. Coordinates and structure factors 349  
 350 were deposited in the Protein Data Bank with accession codes 3ZBT 351  
 352 for S59A FNR, 4BPR for Y79F FNR, 3ZBU for S80A FNR and 3ZC3 for 353  
 354 S80A FNR:NADP<sup>+</sup>. 355

### 341 3. Results


#### 342 3.1. Interaction with partners, and oxido-reduction properties of the FNR 343 344 variants

344 Purification of S59A, Y79F and S80A AnFNR variants produced 345  
 346 protein yields and spectral properties (including UV-vis spectral 347  
 348 shape, maxima position and  $A_{274\text{ nm}}/A_{458\text{ nm}}$  ratio) similar to the 349  
 350 WT, indicating that mutations prevented neither the assembly of 351  
 352 FAD nor the protein folding. Nevertheless, S80A UV-vis spectral max- 353  
 354 ima slightly shifted to longer wavelengths (to 276, 397 and 466 nm), 355  
 356 and its extinction coefficient in the flavin band-II was larger than that 357  
 of WT (Fig. 2A). This suggests that Ser80 directly influences the elec- 358  
 tronic environment of the FAD isoalloxazine. Titration of the FNR<sub>ox</sub> 359  
 variants with NADP<sup>+</sup> induced the typical difference spectra of 360  
 cyanobacterial FNRs (Fig. 2B) indicative of coenzyme binding, but 361  
 lacking the positive band at 509 nm detected in enzymes from plants 362  
 and related with direct stacking between the NMN and the isoalloxazine 363  
 [5,30,34,47].

357 When Ala substituted for Ser80 changes in the position 358  
 359 and intensities of spectral features were observed with respect to 360  
 WT FNR, this again suggests modification of the isoalloxazine envi- 361  
 ronment (Fig. 2B) [34,48]. Saturation of the difference spectra upon 362  
 increasing NADP<sup>+</sup> concentration allowed determination of  $K_d^{\text{NADP}^+}$  363  
 and  $\Delta\varepsilon$  (see Supplementary material, Figure S1). The affinity of S59A 364  
 and Y79F FNR<sub>ox</sub>s for NADP<sup>+</sup> was within a factor of two of that of WT, 365  
 but decreased up to 4-fold for the S80A variant (Table 1). Altogether 366  
 these observations suggest that removal of the Ser80 side-chain 367  
 modifies the nicotinamide disposition into the active site. Titration of 368  
 the different FNR variants with NAD<sup>+</sup> did not induce the appearance 369  
 of difference spectra. This indicated that the presence of NAD<sup>+</sup> does 370  
 not have any effect in the environment of the isoalloxazine ring and 371  
 suggested that binding of the coenzyme is not produced, similar to 372  
 that described for the WT [30,34]. 373

374 Difference spectra obtained upon titration of S59A and Y79F 375  
 376 FNR<sub>ox</sub> with Fd<sub>ox</sub> produced perturbations in the visible region very 377  
 378 similar to those reported for the WT FNR<sub>ox</sub> [7], while the S80A variant 379  
 380 showed a slight displacement of the maxima to shorter wavelengths 381  
 (not shown). All the variants showed interaction parameters in the 382  
 same range as for WT FNR<sub>ox</sub> with Fd<sub>ox</sub>, the only exception was S59A, 383  
 whose affinity for Fd<sub>ox</sub> increased by 3-fold (Table 1). 384

385 Photoreduction of S59A and Y79F FNRs took place following similar 386  
 387 spectral evolution and maximal percentage of semiquinone stabilisation 388  
 (ranging 16–19%) as for the WT (Fig. 2C), but reduction of the S80A 389  
 variant occurred with very little semiquinone stabilisation (<4%). Due 390  
 to the low degree of FNR<sub>sq</sub> stabilisation it was not possible to independ- 391  
 392 ently measure the potential for the two one-electron steps; therefore, mid- 393  
 394 point reduction potentials for the two electron processes ( $E_{\text{ox/hq}}$ ) were 395  
 396 determined for all of them.  $E_{\text{ox/hq}}$  for the mutants yielded values only 397  
 398 slightly less negative (8–12 mV) than for WT AnFNR (Fig. 2D, Table 2). 399  
 400 This was clearly the consequence of less negative  $E_{\text{sq/hq}}$  values for the 401  
 S59A and Y79F variants. The very little semiquinone stabilisation of 402  
 S80A FNR also prevented estimation of  $E_{\text{ox/sq}}$  and  $E_{\text{sq/hq}}$ . 403



**Fig. 2.** Spectroscopic properties of the AnFNR variants. (A) Absorbance spectra in the visible region of WT (—) and S80A (---) AnFNRs. (B) Difference absorbance spectra elicited upon addition of NADP<sup>+</sup> at saturating concentrations to WT (—), S59A (·····), Y79F (---), and S80A (— · —) AnFNR<sub>ox</sub> solutions (~20 μM). (C) Spectral evolution along step-wise photoreduction of Y79F AnFNR (20 μM). (D) Nernst plots for the reduction potential titrations of WT (●), Y79F (■), S59A (◆) and S80A (▲) AnFNRs.

**Table 1**  
Interaction parameters for complex formation of AnFNR<sub>ox</sub> variants with NADP<sup>+</sup> and AnFd<sub>ox</sub> as determined by difference spectroscopy in 50 mM Tris/HCl, pH 8.0, at 25 °C.

FNR variant	NADP <sup>+</sup>		AnFd <sub>ox</sub>	
	K <sub>d</sub> (μM)	Δε <sub>(482–390)</sub> (mM <sup>-1</sup> cm <sup>-1</sup> )	K <sub>d</sub> (μM)	Δε <sub>(462)</sub> (mM <sup>-1</sup> cm <sup>-1</sup> )
WT	4.0	1.15	6.7	1.95
S59A	8.5	1.28 <sup>a</sup>	2.2	1.60
Y79F	8.6	1.28	6.7	2.09
S80A	13.8	1.79 <sup>b</sup>	6.4	3.22 <sup>c</sup>

<sup>a</sup> Δε<sub>(458–396)</sub>.

<sup>b</sup> Δε<sub>(456–400)</sub>.

<sup>c</sup> Δε<sub>(449)</sub>.

### 3.2. Steady-state efficiency of the FNR variants

Kinetic parameters for the diaphorase activity of Y79F AnFNR reflected a very similar behaviour to the WT (Table 3). Replacement of Ser59 with Ala significantly increased the enzyme turnover ( $k_{cat}$ ) with both one- and two-electron acceptors, but the effect in the catalytic efficiency only resulted relevant when using the two-electron acceptor, due to the increase in  $K_m^{NADPH}$  when using the one-electron acceptor. On the contrary, when Ser80 was substituted by Ala the catalytic efficiency decreased by 2- to 4-fold, due to a decrease in the enzyme turnover. None of the mutations improved the AnFNR ability to catalyse the DCPIP diaphorase activity using NADH as electron donor. This activity was not detected at all in the S80A variant, making it even more specific towards the phosphorylated coenzyme.

Kinetic parameters for the FNR NADPH-dependent Cyt<sub>c</sub> reductase activity yielded lower  $k_{cat}$  values for the S59A and, particularly, Y79F variants with respect to WT FNR (37% and 5.5%, respectively), with unaltered  $K_m^{Fd}$  values (Table 3). No activity at all was detected for the S80A AnFNR, a fact probably related with its low ability to stabilise the semiquinone. These results indicate important deleterious effects in the ET from FNR<sub>hq</sub> to Fd<sub>ox</sub> by the introduced mutations.

### 3.3. Pre-steady-state kinetic analysis of the reduction of FNR<sub>ox</sub> by Fd<sub>rd</sub>

Reduction of the isoalloxazine of S59A, Y79F and S80A AnFNRS to the semiquinone state by the laser generated dRfH• followed a monoexponential absorbance decrease at 458 nm, as observed for the WT AnFNR.  $k_{obs}$  values for these reactions were linearly dependent on the FNR concentration with second-order rate constants indicating that all variants are as efficiently reduced by dRfH• as WT (Table 4) [7].

Fd/FNR ET reactions are optimised at relatively high  $I$ , as at very low salt concentrations the strong protein–protein charge interactions freeze the Fd:FNR complex in a non-optimal configuration [7]. FNR<sub>ox</sub> reduction by Fd<sub>rd</sub> has been here investigated in the presence of a moderately high salt concentration ( $I = 120$  mM). When an excess of Fd<sub>ox</sub> is additionally present in the cuvette, the laser-generated dRfH• causes its fast reduction, and a subsequent step of ET from Fd<sub>rd</sub> to WT FNR<sub>ox</sub> can be monitored by formation of neutral FNR<sub>sq</sub> [7,12]. The FNR<sub>sq</sub> formation observed at 600 nm is concomitant with Fd<sub>rd</sub> oxidation, as inferred from an absorbance increase at 498 nm (an

**Table 2**  
Midpoint reduction potentials of the AnFNR variants at 25 °C and pH 8.0<sup>a</sup>.

FNR variant	$E_m$ (mV)	% SQ	$E_{ox/sq}$ (mV)	$E_{sq/hq}$ (mV)
WT	-370	22	-384	-357
S59A	-364	19	-382	-346
Y79F	-363	16	-386	-340
S80A	-358	4	-	-

<sup>a</sup> Potentiometric titrations were carried out using 30 μM FNR, 1 μM methylviologen and benzylviologen, 2 μM dRf and 3 mM EDTA in 50 mM Tris/HCl, pH 8.0.

isobestic point for FNR<sub>ox/sq</sub>) in the same time frame. A similar kinetic behaviour was observed for S59A and Y79F AnFNRS. However, absorbance changes related with ET from Fd<sub>rd</sub> to S80A AnFNR<sub>ox</sub> were detected neither at 600 nm nor at the S80A isobestic point (500 nm), indicating that the S80A variant resulted in highly impaired accepting electrons from Fd<sub>rd</sub>. At  $I = 120$  mM the  $k_{obs}$  values for WT FNR<sub>sq</sub> formation present a hyperbolic dependence on FNR concentration, that can be related with the formation of a transient FNR<sub>ox</sub>:Fd<sub>rd</sub> complex prior to the ET step (Fig. 3A). Applying the formalism previously described [37], minimal values for  $k_{et}$ , and  $K_d$  can be estimated (Table 4), which are in agreement with previously reported data [7]. The two-step model here applied to estimate  $K_d$  and  $k_{et}$  values has largely demonstrated to be extremely useful in the characterization of ET in transient protein:protein reactions [13,35–37]. This model is based in the total amount of FNR<sub>ox</sub> in the sample, although under the experimental conditions used (an excess of Fd<sub>ox</sub>), part of the FNR molecules would be transiently complexed with Fd<sub>ox</sub>, and thus the free Fd<sub>rd</sub> generated by dRfH• has to replace FNR-bound Fd<sub>ox</sub>. However, it is widely accepted that, in transient protein complexes, protein association/dissociation is much faster than ET itself, and thus the ET process would be the rate limiting step rather than Fd exchange. It is worth to note that although the  $K_d$  values here described by difference spectroscopy upon titration refer to the Fd<sub>ox</sub>:FNR<sub>ox</sub> interaction, and the  $K_d$  values estimated by laser-flash experiments correspond to the interaction between Fd<sub>rd</sub> and FNR<sub>ox</sub> (which is only observable by kinetic methodology), both  $K_d$  values are of the same order of magnitude and comparable in both cases, thus supporting the validity of the kinetic models used. Regarding the kinetic properties of the FNR mutants, a similar behaviour was observed for Y79F AnFNR with  $k_{et}$  in the same range as for the WT, while the lower  $K_d$  suggested a stronger interaction between Fd<sub>rd</sub> and this FNR<sub>ox</sub> mutant. This later result is in agreement with the lower turnover for this variant in the Cyt<sub>c</sub> reductase activity. However, a linear dependence was observed on the enzyme concentration when analysing reduction of S59A AnFNR<sub>ox</sub> (Fig. 3A), allowing only estimation of a bimolecular second-order rate constant (Table 4). Decreasing the  $I$  of the medium, slowed  $k_{obs}$  and induced a hyperbolic dependence on FNR concentration (Fig. 3A, Table 4). This indicates the formation of a transient complex between S59A FNR<sub>ox</sub> and Fd<sub>rd</sub> at lower  $I$ , while more physiological salt concentrations make the system shift to a collisional ET mechanism. Changes in the concentration profile dependence have also been reported for the WT enzyme at different  $I$  (Fig. 3A, Table 4) [49].

The influence of  $I$  on the Fd/FNR interaction was further analysed to investigate the effects induced by the mutations on the electrostatics of the interaction. Biphasic dependences of  $k_{obs}$  with increasing NaCl concentration were observed with the WT, S59A and Y79F variants (Fig. 3B), despite the fact that for S59A FNR the  $k_{obs}$  maximum is shifted to higher  $I$ . The S80A variant did not show however significant reactivity at any salt concentration (not shown). The bell-shaped profile for the dependence of  $k_{obs}$  with  $I$  is related with the re-arrangement of the initial FNR<sub>ox</sub>:Fd<sub>rd</sub> interaction to achieve the optimal ET conformation, indicating the occurrence of protein–protein dynamic motions that are blocked by strong electrostatic interactions at very low  $I$  [7].

### 3.4. Transient kinetics of the hydride transfer reactions between FNR and the coenzyme. Kinetic isotopic effect (KIE) and dependence of the temperature

Stopped-flow analysis of the transient HT processes between S59A FNR<sub>hq/ox</sub> and NADP<sup>+</sup>/H followed very similar patterns to those reported for the WT enzyme: fast formation of two intermediate CTCs prior and after the HT event, whatever the direction of the reaction (CTC-1 and CTC-2, characterised by spectral bands centred at 600 nm and 800 nm, respectively (Figure S2)) [4,14,16,38]. Similar to the WT system, evolution of the initial CTC species for S59A also started in the instrumental dead time (Fig. 4A and D), but a slight increase in  $k_{HT}$  can be envisaged



t3.1 **Table 3**t3.2 Steady-state kinetic parameters of the different AnFNR variants for the diaphorase (with either DCPIP or ferricyanide as electron acceptors) and cytochrome c reductase activities in 50 mM  
t3.3 Tris/HCl, pH 8.0 at 25 °C.

FNR variant	DCPIP diaphorase						Fe(CN) <sub>6</sub> <sup>3-</sup> diaphorase			Cytochrome c reductase		
	NADPH			NADH			NADPH			$k_{\text{cat}}^{\text{Fd}}$ ( $\mu\text{M}^{-1}\text{s}^{-1}$ )	$k_{\text{cat}}$ ( $\text{s}^{-1}$ )	$k_{\text{cat}}/K_{\text{m}}$ ( $\mu\text{M}^{-1}\text{s}^{-1}$ )
	$K_{\text{m}}^{\text{NADPH}}$ ( $\mu\text{M}$ )	$k_{\text{cat}}$ ( $\text{s}^{-1}$ )	$k_{\text{cat}}/K_{\text{m}}$ ( $\mu\text{M}^{-1}\text{s}^{-1}$ )	$K_{\text{m}}^{\text{NADH}}$ ( $\mu\text{M}$ )	$k_{\text{cat}}$ ( $\text{s}^{-1}$ )	$k_{\text{cat}}/K_{\text{m}}$ ( $\mu\text{M}^{-1}\text{s}^{-1}$ )	$K_{\text{m}}^{\text{NADPH}}$ ( $\mu\text{M}$ )	$k_{\text{cat}}$ ( $\text{s}^{-1}$ )	$k_{\text{cat}}/K_{\text{m}}$ ( $\mu\text{M}^{-1}\text{s}^{-1}$ )			
WT	6.0	81.5	13.6	800	0.16	$2.0 \cdot 10^{-4}$	11	370	34	1	176	176
S59A	5.3	146.0	27.5	990	0.56	$5.6 \cdot 10^{-4}$	42	1000	24	0.99	65.7	66
Y79F	3.8	73.0	19.2	640	0.21	$3.3 \cdot 10^{-4}$	13.6	304	22	1.16	9.6	8.3
S80A	4.6	30.4	6.6	n.d. <sup>a</sup>	n.d. <sup>a</sup>	n.d. <sup>a</sup>	7.2	93.4	13	n.d. <sup>a</sup>	n.d. <sup>a</sup>	n.d. <sup>a</sup>

t3.11 <sup>a</sup> Activity was not detected.

for this variant with respect to WT (Table 5). Replacements at Tyr79 and Ser80 produced more evident effects (Fig. 4, Table 5). A decrease in the amplitude of the spectral band for CTC-2 was observed upon reduction of Y79F FNR<sub>ox</sub> by NADPH, while HT was slightly hampered in both directions (Fig. 4B and E, Table 5). More drastic effects were observed when Ala substituted for Ser80. Reduction of S80A FNR<sub>ox</sub> by NADPH occurred without spectral CTC-2 stabilisation (Fig. 4C), and no spectral features of CTC at all were detected for the reverse reaction (Fig. 4F). Additionally, both reactions showed less than 5% of the WT efficiency in HT (Table 5).

Due to the reversibility of the process producing the apparent decrease in the experimentally measured rate constants upon increasing coenzyme concentration and to the  $k_{\text{HT}}$  and  $k_{\text{HT-1}}$  rates being in close to the instrumental detection limit for some variants, equimolecular concentrations of enzyme and coenzyme, as well as the use of the single-wavelength detector, were selected to further investigate this mechanism by analysing KIEs on the HT processes. The main observable difference between HT and DT processes was the considerable decrease in  $k_{\text{obsDT}}$  and  $k_{\text{obsDT-1}}$  values with respect to the corresponding  $k_{\text{obsHT}}$  and  $k_{\text{obsHT-1}}$  ones for all the FNR variants (Fig. 5A and B, Table 6), therefore, inducing moderate to important KIEs. For each particular variant the KIE was slightly larger for the reduction of FNR<sub>ox</sub> by the coenzyme than for the reverse reaction. KIEs for the S59A and Y79F variants were in the same range as for WT, but processes with S80A FNR showed considerably larger KIEs (up to 6-fold for the FNR reductive process).  $k_{\text{obs}}$  values for the reduction of the different FNR<sub>ox</sub> forms by NADPH and NADPD, as well as for the reverse processes, resulted highly dependent on temperature (Fig. 5A and B), indicating high activation energies ( $E_{\text{a}}$ ) (Table 6), particularly for the reaction of S80A FNR<sub>ox</sub> with NADPD. The obtained parameters were evaluated within the environmentally coupled tunnelling model that distinguishes between two dynamics contributions to the protein motion: active (or gating) and passive (environmental reorganisation) [50–53]. Arrhenius plots for the HT and DT reactions of WT, S59A and Y79F FNRs corresponded to two almost parallel straight lines (Fig. 5A and B), which produced an almost temperature-independent KIE in the assayed range (Fig. 5C and D). In the reactions of WT FNR, the moderate KIEs and Arrhenius pre-exponential factors ratios ( $A_{\text{H}}/A_{\text{D}}$  and  $A_{\text{H-1}}/A_{\text{D-1}}$ ) just above

unity), together with the high  $E_{\text{a}}$  and small  $\Delta E_{\text{a}}$ , were interpreted as mainly tunnelling of the light isotope and contribution of both the environmental reorganisation (passive) and the vibrationally enhanced modulation (gating) to the tunnel reaction [14,54]. A similar behaviour might be predicted for S59A and Y79F FNRs in their non-photosynthetic reactions, with an apparent more predominant gating contribution, indicated by their slightly larger and lower, respectively,  $\Delta E_{\text{a}}$  and  $A_{\text{H}}/A_{\text{D}}$  values. Backward processes for these two variants showed slightly lower KIEs that were temperature-independent (Fig. 5C and D),  $\Delta E_{\text{a}}$  close to zero and larger  $A_{\text{H-1}}/A_{\text{D-1}}$  values with respect to the WT (Table 6). Altogether these parameters appear consistent with environmental heavy atom reorganisation contributing to the tunnel for the reaction of S59A and Y79F FNR<sub>ox</sub> with NADPH/D, and almost no gating contribution. Reduction of NADP<sup>+</sup> by S80A FNR<sub>hq</sub> behaved similarly to S59A and Y79F FNRs, but with larger KIE and  $A_{\text{H}}/A_{\text{D}}$  ratio, suggesting that a full tunnelling passive dynamics model with protein environmental reorganisation dominates the HT (without gating contribution). The most striking results were observed for reduction of S80A FNR<sub>ox</sub> by the coenzyme. This reaction occurred with a large KIE that considerably decreased with temperature (Fig. 5C), as consequence of a large  $E_{\text{aD}}$  that also produced a considerable increase in  $\Delta E_{\text{a}}$ . This, together with the much lower than unity  $A_{\text{H}}/A_{\text{D}}$ , suggests that vibrationally enhanced modulation of the tunnel probability (gating or active dynamics) is dominating this reaction [50–53].


### 3.5. The structural environment of the mutated positions

S59A, Y79F and S80A FNR<sub>ox</sub>s showed overall crystal structures similar to that of WT (r.m.s.d. ~0.26 on C $\alpha$  atoms aligned for all of them). Mutations did not lead to significant modification of the FAD isoalloxazine environment and were constricted to the interactions involving the modified side-chains (Fig. 6A). The larger differences resulted in the loop 105–111, flexible in all AnFNR structures so far described and suggested to accommodate the adenosine moiety of the FAD [21,22,55]. Y79F FNR showed the highest B factors for the residues contained in this loop; from 103 to 114 the B factor is higher than 40 Å<sup>2</sup> (being its overall B factor of 21.54 Å<sup>2</sup>). This might be due to the fact that the cofactor is not engaged through its ribityl motif with this residue and no water molecule is mimicking the removed hydroxyl group. Ala substitution for Ser59 provoked removal of two defined water molecules that interact with the OG atom of Ser59 and with the OE1 atom of Glu301, respectively, in WT FNR. Besides this, the FAD environment, especially the H-bond network involving Ser80 and Glu301, is not much affected by the mutation. Changes in the S80A FNR crystal structure are restricted to the H-bonds involving the removed hydroxyl group. Ala80 cannot H-bond the side-chain of Glu301, being only in contact with N5i through a H-bond with its main chain N. Again, no water molecule mimics the H-bond network established by the Ser80 side-chain. The highly conserved water molecule, proposed to act as a proton donor to the N5i of FAD, is found in the S80A FNR structure, as in that of WT, interacting with the hydroxyl of Tyr303, but it is at closer distance with the O4 atom of FAD (2.9 Å) than in the WT structure (3.17 Å)


t4.1 **Table 4**t4.2 Rate constants for the laser-flash induced reduction of the AnFNR<sub>ox</sub> variants by dRf and  
t4.3 AnFd.

FNR variant	Reduction by dRf <sup>a</sup>		Reduction by AnFd <sup>b</sup>	
	$k_2$ ( $\text{M}^{-1}\text{s}^{-1}$ )	$k_{\text{et}}$ ( $\text{s}^{-1}$ )	$k_2$ ( $\text{M}^{-1}\text{s}^{-1}$ )	$K_{\text{d}}$ ( $\mu\text{M}$ )
WT	$2.3 \times 10^8$	7780	$3.0 \times 10^7$ <sup>a</sup>	17.0
S59A	$3.5 \times 10^8$	7100 <sup>c</sup>	$2.5 \times 10^8$	26.0 <sup>c</sup>
Y79F	$3.5 \times 10^8$	10,870	–	4.0
S80A	$2.2 \times 10^8$	–	–	–

t4.10 <sup>a</sup> Reaction in 4 mM phosphate, pH 7.0 ( $I = 20$  mM).t4.11 <sup>b</sup> Reaction in 4 mM phosphate, pH 7.0 at 100 mM NaCl ( $I = 120$  mM), unless otherwise stated.t4.12 <sup>c</sup> Reaction in 4 mM phosphate, pH 7.0 at 20 mM NaCl ( $I = 40$  mM).



**Fig. 3.** Laser-flash induced transient kinetic analysis for the reduction of the AnFNR<sub>ox</sub> variants by AnFd<sub>red</sub>. (A) Dependence of  $k_{\text{obs}}$  on FNR concentration for the reduction by Fd<sub>red</sub> of WT (○) FNR<sub>ox</sub> at  $I = 20$  mM, and of WT (●), Y79F (■) and S59A (◆) FNR<sub>ox</sub> at  $I = 120$  mM, and of S59A (◇) FNR<sub>ox</sub> at  $I = 40$  mM. Reaction mixtures contained  $40 \mu\text{M}$  Fd<sub>red</sub>. (B) Dependence of  $k_{\text{obs}}$  on the square root of  $I$  for the reduction of WT (●), Y79F (■) and S59A (◆) FNR<sub>ox</sub> by Fd<sub>red</sub>. Reaction mixtures contained  $40 \mu\text{M}$  Fd<sub>red</sub> and  $30 \mu\text{M}$  FNR<sub>ox</sub>.



**Fig. 4.** Spectral evolution along the HT processes between AnFNR and the coenzyme followed by stopped-flow. Reduction by NADPH of (A) S59A FNR<sub>ox</sub> (spectra recorded at 1.28, 3.84, 6.4, 8.96 and 49.92 ms after mixing), (B) Y79F FNR<sub>ox</sub> (spectra recorded at 1.28, 3.84, 8.96, 24.32 and 49.92 ms), and (C) S80A FNR<sub>ox</sub> (spectra recorded at 3.84, 29.4, 80.6, 157.4 and 997.1 ms). Reoxidation by NADP<sup>+</sup> of photoreduced (D) S59A FNR<sub>hq</sub> (spectra recorded at 1.28, 3.84, 6.4, 8.96 and 49.92 ms), (E) Y79F FNR<sub>hq</sub> (spectra recorded at 1.28, 3.84, 8.96, 19.2 and 49.92 ms) and (F) S80A FNR<sub>hq</sub> (spectra recorded at 3.84, 29.4, 80.6, 157.4 and 997.1 ms). In all cases the thick line is the spectrum of the oxidised (for A, B and C) or reduced (D, E and F) protein before mixing. Insets show the time evolution of the absorption at 458 nm. Reactions were carried out with FNR at  $\sim 25 \mu\text{M}$  and coenzyme  $\sim 100 \mu\text{M}$  final concentrations in 50 mM Tris/HCl pH 8.0 at 6 °C.



t5.1 **Table 5**  
 t5.2 Transient-kinetics parameters for the HT processes between  $AnFNR_{\text{hq/ox}}$  and  $NADP^+/\text{H}$  at  
 t5.3  $6^\circ\text{C}$  in 50 mM Tris/HCl, pH 8.0.

t5.4	FNR variant	FNR <sub>ox</sub> and NADPH	FNR <sub>hq</sub> and NADP <sup>+</sup>
		$k_{\text{HT1}}$ (s <sup>-1</sup> )	$k_{\text{HT-1}}$ (s <sup>-1</sup> )
t5.6	WT <sup>a</sup>	300	285
t5.7	S59A	390	253
t5.8	Y79F	122.6	186
t5.9	S80A	13.2	10.1

t5.10 <sup>a</sup> Data from [14].

579 (Fig. 6A). Another common characteristic in these structures, and in  
 580 contrast with the WT, is the orientation of Arg264 towards the  
 581 C-terminal, its guanidinium H-bonding the C-terminal carboxylate  
 582 of Tyr303 (Fig. 6A). This interaction was already predicted by MD  
 583 simulations [15] and does not appear related with the introduced  
 584 mutations, further suggesting a contribution for Arg264 in the displac-  
 585 ment of Tyr303 to trigger the entrance of the nicotinamide into the  
 586 catalytic site.


587 Finally, in the S80A FNR:NADP<sup>+</sup> complex, the N atom of Ala80  
 588 H-bonds N5i and O4i. In this structure, NADP<sup>+</sup> binds in a similar  
 589 unproductive conformation to that previously reported for WT [22], al-  
 590 though the isoalloxazine-Tyr303 rings stacking distance gets slightly  
 591 larger (around 0.4 Å) and the nicotinamide lies slightly closer to  
 592 Tyr303, bridging its N7n atom and the OH of Tyr303.

#### 593 4. Discussion

594 Here we show that replacement of Ser59 with Ala improved the cat-  
 595 alytic efficiency of the diaphorase activity of *AnFNR* (particularly the  
 596  $k_{\text{cat}}$ ) with respect to the WT, as consequence of increasing the  $k_{\text{HT}}$

(Tables 3 and 5). This mutation also influences the affinity between 597  
 598 FNR and Fd as a function of their oxido-reduction states and of the  $I$  of  
 599 the media (Tables 1 and 4, Fig. 3). Thus, despite steady-state and pre-  
 600 steady-state kinetics indicate that S59A FNR<sub>ox</sub> is able to efficiently re-  
 601 ceive electrons from Fd<sub>rd</sub> (the photosynthetic process) once the interac-  
 602 tion is formed, its differential behaviour dependence on the ionic  
 603 strength with respect to the WT indicates that the mutation influences  
 604 the electrostatic and hydrophobic contributions to produce the most  
 605 productive complex for ET (Table 4, Fig. 3) [56]. Moreover, the intro-  
 606 duced mutation particularly favours the non-photosynthetic HT from  
 607 NADPH to FNR<sub>ox</sub>, as well as the ET from Fd<sub>rd</sub>, particularly at higher  
 608 ionic strengths than for the WT. This might correlate with the  $E_{\text{sq/hq}}$  of  
 609 the variant being slightly less negative. Additionally, KIE analyses for  
 610 the reactions of S59A FNR with the coenzyme (Fig. 5, Table 6) suggest  
 611 larger contribution of the vibrationally enhanced modulation to the tun-  
 612 nel reaction during HT for the reduction of the protein by the coenzyme.  
 613 These data indicate higher flexibility at the active site environment for  
 614 the non-photosynthetic process along the reaction coordinate for  
 615 S59A FNR, regarding both its photosynthetic reaction and the WT  
 616 behaviour. Therefore, Ser59 indirectly modulates the geometry of the  
 617 active site, the interaction with substrates and the electronic properties  
 618 of the isoalloxazine ring, and in consequence the ET and HT processes.  
 619 Altogether these data support the hypothesis derived from MD studies  
 620 on *AnFNR* [15], and confirm that the side-chain of Ser59 indirectly mod-  
 621 ulates the architecture of the reactive complexes during both ET with Fd  
 622 and HT with the coenzyme.

623 Previous substitutions at Tyr89 in *Pisum sativum* FNR (Tyr79 in  
 624 *AnFNR*) with Ser or Glu led to fatal consequences in terms of protein sta-  
 625 bility and FAD binding, while replacements by Phe and Trp considerably  
 626 reduced the catalytic efficiency and affinity for the coenzyme [8]. On the  
 627 contrary, replacement of Tyr79 by Phe in *AnFNR* only slightly modulated  
 628 the binding and diaphorase activity parameters with the coenzyme, as



**Fig. 5.** Temperature dependence of the HT rates and KIEs. Arrhenius plots ( $\ln k_{\text{obs}} = \ln A - E_a / RT$ ) of the kinetic data for the reactions of (A) FNR<sub>ox</sub> with NADPH (closed symbols) or NADPD (open symbols) and of (B) FNR<sub>hq</sub> or D-FNR<sub>hq</sub> (open symbols) with NADP<sup>+</sup> (closed symbols). Temperature dependence of the KIEs for (C) FNR<sub>ox</sub> reduction and (D) FNR<sub>hq</sub> reoxidation by the coenzyme. Data correspond to WT (●), S59A (◆), Y79F (■), and S80A (▲) *AnFNR*s.

**Table 6**  
KIEs for the HT processes catalysed by the different AnFNR variants.

FNR variant	HT (FNR <sub>ox</sub> + NADPH)			DT (FNR <sub>ox</sub> + NADPD)			KIE <sup>a</sup>	$\Delta E_a$ $E_{ad} - E_{ah}$ (kcal mol <sup>-1</sup> )	$A_H/A_D$
	$k_{obsHT}^a$ (s <sup>-1</sup> )	$E_{aH}$ (kcal mol <sup>-1</sup> )	$A_H$ (s <sup>-1</sup> )	$k_{obsDT}^a$ (s <sup>-1</sup> )	$E_{ad}$ (kcal mol <sup>-1</sup> )	$A_D$ (s <sup>-1</sup> )			
WT <sup>c</sup>	175	12.8	$1.8 \times 10^{12}$	27	13.5	$1.2 \times 10^{12}$	6.4	0.7	1.5
S59A	212	12.2	$8.2 \times 10^{11}$	29.2	14.2	$4.2 \times 10^{12}$	7.5	2.0	0.2
Y79F	90	10.2	$9.1 \times 10^9$	19	12.1	$6.4 \times 10^{10}$	4.8	1.9	0.14
S80A	6.8	14.4	$1.3 \times 10^{12}$	0.17	33.5	$4.2 \times 10^{25}$	39 <sup>b</sup>	19.2	$3 \times 10^{-14}$
FNR variant	HT-1 (FNR <sub>hq</sub> + NADP <sup>+</sup> )			DT-1 (D-FNR <sub>hq</sub> + NADP <sup>+</sup> )			KIE <sup>a</sup>	$\Delta E_a - 1$ $E_{ad-1} - E_{ah-1}$ (kcal mol <sup>-1</sup> )	$A_{H-1}/A_{D-1}$
	$k_{obsHT-1}^a$ (s <sup>-1</sup> )	$E_{aH-1}$ (kcal mol <sup>-1</sup> )	$A_{H-1}$ (s <sup>-1</sup> )	$k_{obsDT-1}^a$ (s <sup>-1</sup> )	$E_{ad-1}$ (kcal mol <sup>-1</sup> )	$A_{D-1}$ (s <sup>-1</sup> )			
WT <sup>c</sup>	229	11.6	$3.2 \times 10^{11}$	37	12.1	$1.1 \times 10^{11}$	5.7	0.5	2.8
S59A	196	11.5	$2.0 \times 10^{11}$	40.5	11.2	$2.6 \times 10^{10}$	4.8	-0.2	7.5
Y79F	73	15.1	$5.0 \times 10^{13}$	19.7	15.0	$1.3 \times 10^{13}$	3.7	-0.1	3.8
S80A	5.8	18.7	$2.8 \times 10^{15}$	0.36	18.4	$1.1 \times 10^{14}$	16.1	-0.3	15

<sup>a</sup> Values obtained in a stopped-flow equipment at 5.3 °C and with equimolar concentrations of protein and coenzyme. Evolution of the reaction was followed at the single wavelength of 458 nm.


<sup>b</sup> Temperature dependent KIEs.

<sup>c</sup> Data from [14].

well as the stabilisation of CTCs during HT, being major changes reduced to a decrease in  $k_{HT1}$  and  $k_{HT-1}$  within 2-fold regarding the WT (Tables 1, 3, 5 and 6). Regarding Fd, the mutation increases the affinity of the Fd<sub>rd</sub>:FNR<sub>ox</sub> complex and the ET rate between them (Table 4), while the reverse ET appears considerably hindered (Table 3). These effects with Fd might relate with the slightly less negative midpoint reduction potential of the variant, particularly  $E_{sq/hq}$ , that will favour its reduction regarding the WT (Table 2). Therefore, altogether the observed effects suggest that the H-bond between the hydroxyl group of Tyr79 and the ribityl portion of the cofactor, despite not being critical for ET and HT, indirectly contributes to the reactivity in productive complexes. These data agree with previous mutations in the Lys75–Leu78 peptide, where side-chains, despite not being in direct contact with the isoalloxazine ring, have been shown to slightly displace its midpoint reduction potentials to less negative values [57,58]. Thus, Tyr79 can be included among the side-chains tuning the flavin midpoint potential by creating a defined environment that modulates the FAD conformation.

The last residue here analysed, Ser80, is a key one in the active site of plastidic type FNRs where together with Cys261 and Glu301, it constitutes a highly conserved catalytic triad (Fig. 1A) [2,27,28]. Its main-chain directly H-bonds the isoalloxazine N5i atom as well as the O4i atom via a conserved water molecule, while its hydroxyl interacts with another conserved water molecule that H-bonds Glu301, Tyr303, and N5i (Fig. 6A) [21,59]. Mutations at the equivalent position in spinach leaves FNR, Ser96, [11] impaired catalysis for the S96V and S96G variants with respect to the WT, while it was not possible to produce the S96A variant. We have succeeded in producing the S80A mutant in AnFNR. This replacement slightly modifies the electronic environment of the FAD isoalloxazine ring regarding the WT, and has a deleterious effect in its semiquinone stabilisation (Fig. 2), suggesting that this later process is finely controlled by the H-bond network involving Ser80, Glu301, Tyr303 and N5i. This observation is also consistent with the mutant lacking the ability to accept a single electron from Fd<sub>rd</sub> in the Fd-mediated laser-flash induced ET experiments as well as to donate a single electron to Fd<sub>ox</sub> in the Cyt<sub>c</sub> reductase assay, given the formation of the FNR<sub>sq</sub> intermediate is necessary for both reactions. Noticeably, the S80A mutant shows a similar decrease in efficiency, regarding de WT, in the diaphorase assay when using either one-electron or two-electron acceptors. This suggests that the nature of the small, non-specific and non-physiological collisional K<sub>3</sub>Fe(CN)<sub>6</sub> one-electron acceptor allows it to extract a single electron from S80A FNR<sub>hq</sub> through the reduce amount of semiquinone that this mutant stabilizes (Table 2). On the contrary, processes with Fd include complex formation and dissociation rate limiting steps that modify FNR midpoint

reduction potentials [13]. The results here present suggest that in the case of S80A FNR interaction with Fd further decreases the low stability of its semiquinone. Regarding the coenzyme the mutation produced minor effects in its affinity. Despite the fact that its  $E_m$ , slightly less negative than in WT (Table 2), would apparently favour HT events from the coenzyme, we observed important deleterious effects in turnover, catalytic efficiency, HT rate constants, and the stabilisation of CTCs during the HT event for both the forward and reverse HT reactions (Fig. 4, Tables 3 and 5). All these data are in agreement with those reported for the Ser96 mutants in spinach FNR [11], further indicating that this Ser is critical to generate the architecture of the catalytically competent complex upon coenzyme binding in both the cyanobacterial and plant enzymes. Since the structures obtained for mutants at this Ser (both in AnFNR (Ser80) and the spinach enzyme (S96)) only show changes in the H-bond network involving this position (Fig. 6A), this network must be critical for the efficiency of the HT process. In agreement with this conclusion previous MD simulations suggested that the Ser80 side-chain might contribute to fix the position of the amide of NADP<sup>+</sup> and, as consequence, the position of the nicotinamide ring in the active site cavity [15]. The contribution of the Ser80 side-chain to the optimal architecture of the catalytically competent complex between FNR and the coenzyme is here further supported by the analysis of the active site dynamics during the HT event (Fig. 5 and Table 6). HT and DT reactions for WT FNR, in both the forward and reverse directions, have been explained applying a tunnelling model in which both environmental reorganisation (passive dynamics) and vibrational enhancement (active dynamics) contribute to the reaction [14,16]. However, KIE analyses for the processes of the S80A FNR mutant best fitted to the two extreme cases in which the tunnelling reaction is completely dominated by either environmental reorganisation or vibrational enhancement [60]. Thus, reduction of S80A FNR<sub>ox</sub> by NADPH/D was consistent with a full tunnelling model in which vibrational fluctuations of the active site (gating) are able to compress the hydrogen donor–acceptor distance during the HT event, making tunnelling more probable (specially for the light isotope) as long as the temperature increases. On the contrary, parameters measured for the oxidation of S80A FNR<sub>hq</sub> by NADP<sup>+</sup> suggest no gating contribution to the HT event, being the initial environmental thermal reorganisation of the whole active site the only dynamics contribution controlling the process. Therefore, modifications on the H-bond network caused by substitution of Ser80 with alanine entail important perturbations on the flexibility of the active site of FNR along the HT reaction coordinate. In the case of the reaction of S80A FNR<sub>ox</sub> with NADPH, the organisation of the close environment of the donor (C4n) and acceptor (N5i) atoms allows them to undergo



**Fig. 6.** Influence of the mutations in the active site geometry of AnFNR. (A) H-bond network at the active sites of the WT FNR (C in blue, PDB ID: 1QUE), S59A FNR (C in orange), Y79F FNR (C in yellow) and S80A FNR (C in green) crystal structures. Residues involved in the interactions are drawn in sticks. (B) Superposition of the active sites of WT FNR (C in white, PDB ID: 1QUE), the WT FNR:NADP<sup>+</sup> complex (C in blue, PDB ID: 1GJR) and the S80A FNR:NADP<sup>+</sup> complex (C in green).

717 important vibrational fluctuations to attain the optimal distance and  
 718 orientation for efficient HT. On the contrary, for the reverse reaction,  
 719 once the NADP<sup>+</sup> nicotinamide reaches the active site of S80A FNR<sub>hq</sub>  
 720 the reduced isoalloxazine and oxidised nicotinamide rings remain  
 721 “frozen”, without any further flexibility of the active site contrib-  
 722 uting to improve its relative distance and orientation to reach the  
 723 hydrogen tunnelling ready conformation. Therefore, the S80A muta-  
 724 tion highly compromises the active site environment fluctuations re-  
 725 quired in FNR to increase the HT probability, producing important

thermodynamic and kinetic consequences in the process that are 726  
 reflected in the enzyme efficiency, mechanism and reversibility of 727  
 the process, all of them particular facts of plant type FNRs. **Altogether** 728  
 these data confirm the importance of Ser80 in both the ET and HT 729  
 processes, adding information about its roles. Thus, this Ser side- 730  
 chain modulates the midpoint reduction potential of the flavin ring 731  
 and contributes to the stabilisation of its semiquinone state, a key 732  
 factor for efficient ET exchange between FNR and Fd. Additionally, 733  
 it also contributes to stabilise the nicotinamide ring in the optimal 734



geometric conformation regarding the isoalloxazine ring for an efficient HT event through a tunnelling process following similar dynamics for both the photosynthetic and non-photosynthetic reactions, thus ensuring the efficiency and reversibility of the process.

In conclusion, as new information becomes available additional roles might be envisaged for residues in the active site of FNR. Among these residues are Tyr79 and Ser80, previously analysed in FNRs from higher plants [8,11,26]. Additionally, other residues not considered previously, such as Ser59, might indirectly contribute to the efficiency of the HT with the coenzyme by modulating the architecture of the reactive complexes [15]. Here, we present proofs for the implication of H-bond connections between the flavin and the polypeptide chain, direct or via water molecules, in the FNR catalytic efficiency, showing that the isoalloxazine environment strongly influences FAD<sub>sq</sub> stabilisation and ET from Fd. Moreover, an additional key role is predicted for Ser80 during the HT step providing optimal active site geometry, including not only the final disposition between the reacting N5Hi and C4n atoms but also the active site motions required to achieve it.

## Acknowledgements

This work has been supported by the Ministerio de Ciencia e Innovación, Spain (BIO2010-14983 to M.M.), the Aragonian Government-FSE (B18) and the Andalusian Government-FEDER (PAIDI BIO-022 to J.A.N.A.). A.S.-A. holds a FPU fellowship from the Spanish Ministry of Education.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.bbabbio.2013.10.010>.

## References

- [1] M. Medina, Structural and mechanistic aspects of flavoproteins: photosynthetic electron transfer from photosystem I to NADP<sup>+</sup>, *FEBS J.* 276 (2009) 3942–3958.
- [2] A. Aliverti, V. Pandini, A. Pennati, M. de Rosa, G. Zanetti, Structural and functional diversity of ferredoxin-NADP<sup>+</sup> reductases, *Arch. Biochem. Biophys.* 474 (2008) 283–291.
- [3] E.A. Ceccarelli, A.K. Arakaki, N. Cortez, N. Carrillo, Functional plasticity and catalytic efficiency in plant and bacterial ferredoxin-NADP(H) reductases, *Biochim. Biophys. Acta* 1698 (2004) 155–165.
- [4] C.J. Batie, H. Kamin, Electron transfer by ferredoxin:NADP<sup>+</sup> reductase. Rapid-reaction evidence for participation of a ternary complex, *J. Biol. Chem.* 259 (1984) 11976–11985.
- [5] J. Tejero, I. Pérez-Dorado, C. Maya, M. Martínez-Júlvez, J. Sanz-Aparicio, C. Gómez-Moreno, J.A. Hermoso, M. Medina, C-terminal tyrosine of ferredoxin-NADP<sup>+</sup> reductase in hydride transfer processes with NAD(P)<sup>+</sup>/H, *Biochemistry* 44 (2005) 13477–13490.
- [6] M.A. Musumeci, A.K. Arakaki, D.V. Rial, D.L. Catalano-Dupuy, E.A. Ceccarelli, Modulation of the enzymatic efficiency of ferredoxin-NADP(H) reductase by the amino acid volume around the catalytic site, *FEBS J.* 275 (2008) 1350–1366.
- [7] M. Medina, M. Martínez-Júlvez, J.K. Hurlley, G. Tollin, C. Gómez-Moreno, Involvement of glutamic acid 301 in the catalytic mechanism of ferredoxin-NADP<sup>+</sup> reductase from *Anabaena* PCC 7119, *Biochemistry* 37 (1998) 2715–2728.
- [8] A.K. Arakaki, E.G. Orellano, N.B. Calcaterra, J. Ottado, E.A. Ceccarelli, Involvement of the flavin si-face tyrosine on the structure and function of ferredoxin-NADP<sup>+</sup> reductases, *J. Biol. Chem.* 276 (2001) 44419–44426.
- [9] A. Aliverti, T. Lubberstedt, G. Zanetti, R.G. Herrmann, B. Curti, Probing the role of lysine 116 and lysine 244 in the spinach ferredoxin-NADP<sup>+</sup> reductase by site-directed mutagenesis, *J. Biol. Chem.* 266 (1991) 17760–17763.
- [10] A. Aliverti, L. Piubelli, G. Zanetti, T. Lubberstedt, R.G. Herrmann, B. Curti, The role of cysteine residues of spinach ferredoxin-NADP<sup>+</sup> reductase as assessed by site-directed mutagenesis, *Biochemistry* 32 (1993) 6374–6380.
- [11] A. Aliverti, C.M. Bruns, V.E. Pandini, P.A. Karplus, M.A. Vanoni, B. Curti, G. Zanetti, Involvement of serine 96 in the catalytic mechanism of ferredoxin-NADP<sup>+</sup> reductase: structure–function relationship as studied by site-directed mutagenesis and X-ray crystallography, *Biochemistry* 34 (1995) 8371–8379.
- [12] I. Nogués, J. Tejero, J.K. Hurlley, D. Paladini, S. Frago, G. Tollin, S.G. Mayhew, C. Gómez-Moreno, E.A. Ceccarelli, N. Carrillo, M. Medina, Role of the C-terminal tyrosine of ferredoxin-nicotinamide adenine dinucleotide phosphate reductase in the electron transfer processes with its protein partners ferredoxin and flavodoxin, *Biochemistry* 43 (2004) 6127–6137.
- [13] J.K. Hurlley, R. Morales, M. Martínez-Júlvez, T.B. Brodie, M. Medina, C. Gómez-Moreno, G. Tollin, Structure–function relationships in *Anabaena* ferredoxin/ferredoxin-NADP<sup>+</sup>

- reductase electron transfer: insights from site-directed mutagenesis, transient absorption spectroscopy and X-ray crystallography, *Biochim. Biophys. Acta* 1554 (2002) 805–806.
- [14] J.R. Peregrina, A. Sánchez-Azqueta, B. Herguedas, M. Martínez-Júlvez, M. Medina, Role of specific residues in coenzyme binding, charge-transfer complex formation, and catalysis in *Anabaena* ferredoxin-NADP<sup>+</sup> reductase, *Biochim. Biophys. Acta* 1797 (2010) 1638–1646.
- [15] J.R. Peregrina, I. Lans, M. Medina, The transient catalytically competent coenzyme allocation into the active site of *Anabaena* ferredoxin-NADP<sup>+</sup> reductase, *Eur. Biophys. J.* (2012).
- [16] I. Lans, J.R. Peregrina, M. Medina, M. García-Viloca, A. González-Lafont, J.M. Lluch, Mechanism of the hydride transfer between *Anabaena* Tyr303Ser FNR<sub>rd</sub>/FNR<sub>ox</sub> and NADP<sup>+</sup>/H. A combined pre-steady-state kinetic/ensemble-averaged transition-state theory with multidimensional tunneling study, *J. Phys. Chem. B* 114 (2010) 817–818.
- [17] Z. Deng, A. Aliverti, G. Zanetti, A.K. Arakaki, J. Ottado, E.G. Orellano, N.B. Calcaterra, E.A. Ceccarelli, N. Carrillo, P.A. Karplus, A productive NADP<sup>+</sup> binding mode of ferredoxin-NADP<sup>+</sup> reductase revealed by protein engineering and crystallographic studies, *Nat. Struct. Biol.* 6 (1999) 847–853.
- [18] I. Lans, M. Medina, E. Rosta, G. Hummer, M. García-Viloca, J.M. Lluch, A. González-Lafont, Theoretical study of the mechanism of the hydride transfer between ferredoxin-NADP<sup>+</sup> reductase and NADP<sup>+</sup>: the role of Tyr303, *J. Am. Chem. Soc.* 134 (2012) 20544–20553.
- [19] P.A. Karplus, C.M. Bruns, Structure–function relations for ferredoxin reductase, *J. Bioenerg. Biomembr.* 26 (1994) 89–99.
- [20] M. Ingelman, S. Ramaswamy, V. Niviere, M. Fontecave, H. Eklund, Crystal structure of NAD(P)H:flavin oxidoreductase from *Escherichia coli*, *Biochemistry* 38 (1999) 7040–7049.
- [21] L. Serre, F.M. Vellieux, M. Medina, C. Gómez-Moreno, J.C. Fontecilla-Camps, M. Frey, X-ray structure of the ferredoxin:NADP<sup>+</sup> reductase from the cyanobacterium *Anabaena* PCC 7119 at 1.8 Å resolution, and crystallographic studies of NADP<sup>+</sup> binding at 2.25 Å resolution, *J. Mol. Biol.* 263 (1996) 20–39.
- [22] J.A. Hermoso, T. Mayoral, M. Faro, C. Gomez-Moreno, J. Sanz-Aparicio, M. Medina, Mechanism of coenzyme recognition and binding revealed by crystal structure analysis of ferredoxin-NADP<sup>+</sup> reductase complexed with NADP<sup>+</sup>, *J. Mol. Biol.* 319 (2002) 1133–1142.
- [23] M.B. Murataliev, R. Feyereisen, F.A. Walker, Electron transfer by diflavin reductases, *Biochim. Biophys. Acta* 1698 (2004) 1–26.
- [24] C.C. Correll, C.J. Batie, D.P. Ballou, M.L. Ludwig, Phthalate dioxygenase reductase: a modular structure for electron transfer from pyridine nucleotides to [2Fe–2S], *Science* 258 (1992) 1604–1610.
- [25] C.C. Correll, M.L. Ludwig, C.M. Bruns, P.A. Karplus, Structural prototypes for an extended family of flavoprotein reductases: comparison of phthalate dioxygenase reductase with ferredoxin reductase and ferredoxin, *Protein Sci.* 2 (1993) 2112–2133.
- [26] A. Aliverti, Z. Deng, D. Ravasi, L. Piubelli, P.A. Karplus, G. Zanetti, Probing the function of the invariant glutamyl residue 312 in spinach ferredoxin-NADP<sup>+</sup> reductase, *J. Biol. Chem.* 273 (1998) 34008–34015.
- [27] V.I. Dumit, T. Essigke, N. Cortez, G.M. Ullmann, Mechanistic insights into ferredoxin-NADP(H) reductase catalysis involving the conserved glutamate in the active site, *J. Mol. Biol.* 397 (2010) 814–825.
- [28] A. Sánchez-Azqueta, M.A. Musumeci, M. Martínez-Júlvez, E.A. Ceccarelli, M. Medina, Structural backgrounds for the formation of a catalytically competent complex with NADP(H) during hydride transfer in ferredoxin-NADP<sup>+</sup> reductases, *Biochim. Biophys. Acta* 1817 (2012) 1063–1071.
- [29] C.M. Bruns, P.A. Karplus, Refined crystal structure of spinach ferredoxin reductase at 1.7 Å resolution: oxidized, reduced and 2<sup>+</sup>-phospho-5<sup>+</sup>-AMP bound states, *J. Mol. Biol.* 247 (1995) 125–145.
- [30] J. Tejero, M. Martínez-Júlvez, T. Mayoral, A. Luquita, J. Sanz-Aparicio, J.A. Hermoso, J.K. Hurlley, G. Tollin, C. Gómez-Moreno, M. Medina, Involvement of the pyrophosphate and the 2<sup>+</sup>-phosphate binding regions of ferredoxin-NADP<sup>+</sup> reductase in coenzyme specificity, *J. Biol. Chem.* 278 (2003) 49203–49214.
- [31] M. Martínez-Júlvez, J. Hermoso, J.K. Hurlley, T. Mayoral, J. Sanz-Aparicio, G. Tollin, C. Gómez-Moreno, M. Medina, Role of Arg100 and Arg264 from *Anabaena* PCC 7119 ferredoxin-NADP<sup>+</sup> reductase for optimal NADP<sup>+</sup> binding and electron transfer, *Biochemistry* 37 (1998) 17680–17691.
- [32] V.V. Pollock, M.J. Barber, Kinetic and mechanistic properties of biotin sulfoxide reductase, *Biochemistry* 40 (2001) 1430–1440.
- [33] P. Macheroux, UV-visible spectroscopy as a tool to study flavoproteins, *Methods Mol. Biol.* 131 (1999) 1–7.
- [34] M. Medina, A. Luquita, J. Tejero, J. Hermoso, T. Mayoral, J. Sanz-Aparicio, K. Grever, C. Gómez-Moreno, Probing the determinants of coenzyme specificity in ferredoxin-NADP<sup>+</sup> reductase by site-directed mutagenesis, *J. Biol. Chem.* 276 (2001) 11902–11912.
- [35] S. Frago, I. Lans, J.A. Navarro, M. Hervás, D.E. Edmondson, M.A. De la Rosa, C. Gómez-Moreno, S.G. Mayhew, M. Medina, Dual role of FMN in flavodoxin function: electron transfer cofactor and modulation of the protein–protein interaction surface, *Biochim. Biophys. Acta* 1797 (2010) 262–271.
- [36] V. Rodríguez-Roldán, J.M. García-Heredia, J.A. Navarro, M. Hervás, B. De la Cerda, F.P. Molina-Heredia, M.A. De la Rosa, A comparative kinetic analysis of the reactivity of plant, horse, and human respiratory cytochrome c towards cytochrome c oxidase, *Biochem. Biophys. Res. Commun.* 346 (2006) 1108–1113.
- [37] T.E. Meyer, Z.G. Zhao, M.A. Cusanovich, G. Tollin, Transient kinetics of electron transfer from a variety of c-type cytochromes to plastocyanin, *Biochemistry* 32 (1993) 4552–4559.
- [38] J. Tejero, J.R. Peregrina, M. Martínez-Júlvez, A. Gutiérrez, C. Gómez-Moreno, N.S. Scrutton, M. Medina, Catalytic mechanism of hydride transfer between NADP<sup>+</sup>/H

- 890 and ferredoxin-NADP<sup>+</sup> reductase from *Anabaena* PCC 7119, Arch. Biochem. Biophys. 459 (2007) 79–90.
- 891
- 892 [39] S. Daff, An appraisal of multiple NADPH binding-site models proposed for cytochrome P450 reductase, NO synthase, and related diflavin reductase systems, Biochemistry 43 (2004) 3929–3932.
- 893
- 894
- 895 [40] W. Kabsch, XDS, Acta Crystallogr. D Biol. Crystallogr. 66 (2010) 125–132.
- 896 [41] M.D. Winn, C.C. Ballard, K.D. Cowtan, E.J. Dodson, P. Emsley, P.R. Evans, R.M. Keegan, E.B. Krissinel, A.G. Leslie, A. McCoy, S.J. McNicholas, G.N. Murshudov, N.S. Pannu, E.A. Potterton, H.R. Powell, R.J. Read, A. Vagin, K.S. Wilson, Overview of the CCP4 suite and current developments, Acta Crystallogr. D Biol. Crystallogr. 67 (2011) 235–242.
- 897
- 898 [42] A. Vagin, A. Teplyakov, MOLREP: an automated program for molecular replacement, J. Appl. Crystallogr. 30 (1997) 1022–1025.
- 899
- 900 [43] P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics, Acta Crystallogr. D Biol. Crystallogr. 60 (2004) 2126–2132.
- 901
- 902 [44] A.A. Vaguine, J. Richelle, S.J. Wodak, SFCHECK: a unified set of procedures for evaluating the quality of macromolecular structure-factor data and their agreement with the atomic model, Acta Crystallogr. D Biol. Crystallogr. 55 (1999) 191–205.
- 903
- 904 [45] R.A. Laskowski, M.W. MacArthur, D.S. Moss, J.M. Thornton, PROCHECK: a program to check the stereochemical quality of protein structures, J. Appl. Crystallogr. 26 (1993) 283–291.
- 905
- 906 [46] I.W. Davis, L.W. Murray, J.S. Richardson, D.C. Richardson, MOLPROBITY: structure validation and all-atom contact analysis for nucleic acids and their complexes, Nucleic Acids Res. 32 (2004) W615–W619.
- 907
- 908 [47] J. Sancho, C. Gómez-Moreno, Interaction of ferredoxin-NADP<sup>+</sup> reductase from *Anabaena* with its substrates, Arch. Biochem. Biophys. 288 (1991) 231–238.
- 909
- 910 [48] M. Martínez-Júlvez, J. Tejero, J.R. Peregrina, I. Nogués, S. Frago, C. Gómez-Moreno, M. Medina, Towards a new interaction enzyme:coenzyme, Biophys. Chem. 115 (2005) 219–224.
- 911
- 912 [49] J.K. Hurley, M. Fillat, C. Gómez-Moreno, G. Tollin, Structure–function relationships in the ferredoxin/ferredoxin:NADP<sup>+</sup> reductase system from *Anabaena*, Biochimie 77 (1995) 539–548.
- 913
- 914 [50] M.J. Knapp, J.P. Klinman, Environmentally coupled hydrogen tunneling. Linking catalysis to dynamics, Eur. J. Biochem. 269 (2002) 3113–3121.
- 915
- 916 [51] M.J. Knapp, K. Rickert, J.P. Klinman, Temperature-dependent isotope effects in soybean lipoxygenase-1: correlating hydrogen tunneling with protein dynamics, J. Am. Chem. Soc. 124 (2002) 3865–3874.
- 917
- 918 [52] Z.D. Nagel, J.P. Klinman, A 21st century revisionist's view at a turning point in enzymology, Nat. Chem. Biol. 5 (2009) 543–550.
- 919
- 920 [53] J. Basran, R.J. Harris, M.J. Sutcliffe, N.S. Scrutton, H-tunneling in the multiple H-transfers of the catalytic cycle of morphinone reductase and in the reductive half-reaction of the homologous pentaerythritol tetranitrate reductase, J. Biol. Chem. 278 (2003) 43973–43982.
- 921
- 922 [54] A. Kohen, Kinetic Isotope Effects as Probes for Hydrogen Tunneling in Enzyme Catalysis, Isotope Effects in Chemistry and Biology, CRC Press, 2005. 933
- 923
- 924 [55] J.R. Peregrina, B. Herguedas, J.A. Hermoso, M. Martínez-Júlvez, M. Medina, Protein motifs involved in coenzyme interaction and enzymatic efficiency in *Anabaena* ferredoxin-NADP<sup>+</sup> reductase, Biochemistry 48 (2009) 3109–3119. 936
- 925
- 926 [56] J.K. Hurley, J.L. Schmeits, C. Genzor, C. Gómez-Moreno, G. Tollin, Charge reversal mutations in a conserved acidic patch in *Anabaena* ferredoxin can attenuate or enhance electron transfer to ferredoxin:NADP<sup>+</sup> reductase by altering protein/protein orientation within the intermediate complex, Arch. Biochem. Biophys. 333 (1996) 243–250. 939
- 927
- 928 [57] M. Faro, C. Gomez-Moreno, M. Stankovich, M. Medina, Role of critical charged residues in reduction potential modulation of ferredoxin-NADP<sup>+</sup> reductase, Eur. J. Biochem. 269 (2002) 2656–2661. 940
- 929
- 930 [58] M. Martínez-Júlvez, I. Nogués, M. Faro, J.K. Hurley, T.B. Brodie, T. Mayoral, J. Sanz-Aparicio, J.A. Hermoso, M.T. Stankovich, M. Medina, G. Tollin, C. Gómez-Moreno, Role of a cluster of hydrophobic residues near the FAD cofactor in *Anabaena* PCC 7119 ferredoxin-NADP<sup>+</sup> reductase for optimal complex formation and electron transfer to ferredoxin, J. Biol. Chem. 276 (2001) 27498–27510. 941
- 931
- 932 [59] P.A. Karplus, M.J. Daniels, J.R. Herriott, Atomic structure of ferredoxin-NADP<sup>+</sup> reductase: prototype for a structurally novel flavoenzyme family, Science 251 (1991) 60–66. 942
- 933
- 934 [60] J.P. Klinman, Importance of protein dynamics during enzymatic C–H bond cleavage catalysis, Biochemistry (2013). 943
- 935
- 936
- 937
- 938
- 939
- 940
- 941
- 942
- 943
- 944
- 945
- 946
- 947
- 948
- 949
- 950
- 951
- 952
- 953
- 954
- 955
- 956