

SEMILINEAR FRACTIONAL DIFFERENTIAL EQUATIONS: GLOBAL SOLUTIONS, CRITICAL NONLINEARITIES AND COMPARISON RESULTS

BRUNO DE ANDRADE¹, ALEXANDRE N. CARVALHO², PAULO M. CARVALHO-NETO³,
AND PEDRO MARÍN-RUBIO⁴

ABSTRACT. In this work we study several questions concerning to abstract fractional Cauchy problems of order $\alpha \in (0, 1)$. Concretely, we analyze the existence of local mild solutions for the problem, and its possible continuation to a maximal interval of existence. The case of critical nonlinearities and corresponding regular mild solutions is also studied. Finally, by establishing some general comparison results, we apply them to conclude the global well-posedness of a fractional partial differential equation coming from heat conduction theory.

MSC 2010. Primary: 34A08, 34G20; Secondary: 35K05, 35B09.

Keywords: fractional differential equations; semilinear equations; comparison results; critical nonlinearities.

1. INTRODUCTION AND STATEMENT OF THE RESULTS

Origins of fractional calculus go back to a question posed by Leibniz to L'Hôpital about the meaning of the derivative of order $1/2$. Since then this theory was developed and nowadays it has generated a considerable amount of bibliography, see for example the monographs [11, 13, 14, 17] and the references therein.

Naturally the interest in fractional differential equations was considerably raised over the years; among engineers and scientists this is due to their vast potential of applications in several applied problems, such as fluid flow, rheology, diffusive transport akin to diffusion, electric networks, probability and statistical distribution theory, see [4, 18, 19, 20, 25] among others.

From the mathematical point of view, we observe that fractional differential equations is a rich and complex area and hence there exists a great interest in developing the theoretical analysis and numerical methods for these equations, see [7, 8, 10, 12,

¹Partially supported by CNPQ 100994/2011-3 and 478053/2013-4, and CAPES BEX 5549/11-6.

²Research partially supported by CNPq 305447/2005-0 and 451761/2008-1, CAPES/DGU 267/2008 and FAPESP 2008/53094-4, Brazil.

³Partially supported by FAPESP - Brazil, grant # 2008/58944-6, and CAPES - Brazil, grant # BEX 5537/11-8 .

⁴Partially supported by Ministerio de Educación–DGPU (Spain) project PHB2010-0002-PC, Ministerio de Ciencia e Innovación (Spain) project MTM2011-22411, and Junta de Andalucía grant P07-FQM-02468.

15, 16, 21, 22].

Motivated by this, we consider the abstract fractional Cauchy problem

$$\begin{cases} cD_t^\alpha u(t) = Au(t) + f(t, u(t)), & t > 0 \\ u(0) = u_0 \in X, \end{cases} \quad (1)$$

where X is a Banach space, $\alpha \in (0, 1)$, cD_t^α is Caputo's fractional derivative and $A : D(A) \subset X \rightarrow X$ is a sectorial operator, i.e., a closed operator with dense domain for which there are $\phi \in (\pi/2, \pi)$ and $N \geq 1$ such that

$$S_\phi := \{\lambda \in \mathbb{C} : |\arg(\lambda)| \leq \phi, \lambda \neq 0\} \subset \rho(A)$$

and

$$\|(\lambda - A)^{-1}\| \leq \frac{N}{|\lambda|} \quad \forall \lambda \in S_\phi.$$

In order to start our discussion, let us introduce some preliminaries. Consider $\{E_\alpha(t^\alpha A)\}_{t \geq 0}$ and $\{E_{\alpha, \alpha}(t^\alpha A)\}_{t \geq 0}$ the Mittag-Leffler families associated to A (see next section for details). By a global mild solution to (1) in $[0, \infty)$ we understand a continuous function $u : [0, \infty) \rightarrow X$ such that

$$u(t) = E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha, \alpha}((t-s)^\alpha A) f(s, u(s)) ds, \quad t \geq 0. \quad (2)$$

This concept is based upon the following (formal) construction. Suppose that $u : [0, \infty) \rightarrow X$ verifies (1). Then, applying the fractional integral operator J_t^α (see, e.g., [3, 12, 13, 14]) in both sides of the fractional differential equation (1), we obtain

$$\begin{aligned} u(t) &= u(0) + J_t^\alpha Au(t) + J_t^\alpha f(t, u(t)) \\ &= u_0 + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} Au(s) ds + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, u(s)) ds, \quad t \geq 0. \end{aligned}$$

Now, applying the Laplace transform \mathcal{L} in both sides of the above equality we have

$$\hat{u}(\lambda) = \frac{u_0}{\lambda} + \frac{1}{\lambda^\alpha} A \hat{u}(\lambda) + \frac{1}{\lambda^\alpha} (\widehat{f(u)})(\lambda) \Rightarrow \lambda^\alpha \hat{u}(\lambda) = u_0 \lambda^{\alpha-1} + A \hat{u}(\lambda) + (\widehat{f(u)})(\lambda),$$

where $\hat{u}(\lambda) = \mathcal{L}\{u(t)\}(\lambda)$ and $\widehat{f(u)}(\lambda) = \mathcal{L}\{f(t, u(t))\}(\lambda)$. Taking now $\lambda^\alpha \in \rho(A)$, it follows that

$$\hat{u}(\lambda) = \lambda^{\alpha-1} (\lambda^\alpha - A)^{-1} u_0 + (\lambda^\alpha - A)^{-1} (\widehat{f(u)})(\lambda),$$

and using the inverse Laplace transform, and Lemma 13 below, we deduce that

$$u(t) = E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha, \alpha}((t-s)^\alpha A) f(s, u(s)) ds, \quad t \geq 0.$$

Motivated by this discussion and by the previous related literature, we adopt the following concept for solution to the problem (1).

Definition 1. Let $\tau > 0$.

- i) A function $u : [0, \tau] \rightarrow X$ is said to be a local mild solution to (1) in $[0, \tau]$ if $u \in C([0, \tau]; X)$ and

$$u(t) = E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s)) ds, \quad t \geq 0.$$

- ii) A function $u : [0, \tau] \rightarrow X$ is said to be a local mild solution to (1) in $[0, \tau]$ if for any $\tau' \in [0, \tau]$, u is a local mild solution to (1) in $[0, \tau']$.

Our main purpose in this work is to ensure sufficient conditions for existence and uniqueness of mild solution to (1) and to establish some comparison results for this solution in ordered Banach spaces. As can be observed in (2), the notion of mild solution to (1) is very close to generalized Mittag-Leffler type functions. For this, in Section 2, among other things, we study some properties of these functions. In particular, we study the behavior of the families $\{E_\alpha(t^\alpha A)\}_{t \geq 0}$ and $\{E_{\alpha,\alpha}(t^\alpha A)\}_{t \geq 0}$ on the fractional power spaces associated to the linear operator A and we establish expressions for these families, similar to the second fundamental limit for semigroups (see Lemma 14 and Proposition 16 below).

Section 3 is devoted to study existence, uniqueness and continuation results to (1) when the nonlinear term is a locally Lipschitz continuous function $f : [0, \infty) \times X \rightarrow X$. In this direction we start with a proof of the following existence result.

Theorem 2. Let $f : [0, \infty) \times X \rightarrow X$ be a continuous function, and locally Lipschitz in the second variable, uniformly with respect to the first variable, that is, for each fixed $x \in X$, there exist an open ball B_x and a constant $L = L(B_x) \geq 0$ such that

$$\|f(t, z) - f(t, y)\| \leq L\|z - y\|$$

for all $z, y \in B_x$ and $t \in [0, \infty)$. Then, there exists $t_0 > 0$ such that (1) has a unique local mild solution in $[0, t_0]$.

In the remainder of Section 3 our goal is to ensure conditions for existence of global solutions of (1). For this purpose, we prove the following result on continuation of mild solutions.

Theorem 3. Let $f : [0, \infty) \times X \rightarrow X$ be as in Theorem 2. If $u : [0, t_0] \rightarrow X$ is a local mild solution to (1) in $[0, t_0]$, then there exists a unique continuation u^* of u in some interval $[0, t_0 + \tau]$ with $\tau > 0$.

We finish Section 3 with a result on global existence or non-continuation by blow-up. Namely, we have the following

Theorem 4. Let $f : [0, \infty) \times X \rightarrow X$ be a continuous function, locally Lipschitz in the second variable, uniformly with respect to the first variable, and bounded (i.e. it maps bounded sets onto bounded sets). Then the problem (1) has a global mild solution in $[0, \infty)$ or there exists $\omega \in (0, \infty)$ such that $u : [0, \omega) \rightarrow X$ is a local mild solution in $[0, \omega)$ and u does not have a continuation. Moreover, if u is the local mild solution to (1) in $[0, \omega)$, then $\limsup_{t \rightarrow \omega^-} \|u(t)\| = \infty$.

We observe that the assumption on the nonlinear term f is rather general but it does not allow to treat problems where the nonlinearity has critical growth. In Section 4 we treat the critical case, proving existence of ϵ -regular mild solution to (1).

In order to state this result, let us introduce some notation. For $\beta \geq 0$, let X^β be the fractional power spaces associated to the operator A . In this work we will use the following concepts.

Definition 5. A continuous function $u : [0, \tau] \rightarrow X^1$ is called an ϵ -regular mild solution to (1) if $u \in C((0, \tau]; X^{1+\epsilon})$ and verifies (2).

Definition 6. For $\epsilon > 0$ we say that a map g is an ϵ -regular map relative to the pair (X^1, X^0) if there exist $\rho > 1$, $\gamma(\epsilon)$ with $\rho\epsilon \leq \gamma(\epsilon) < 1$, and a positive constant c , such that $g : X^{1+\epsilon} \rightarrow X^{\gamma(\epsilon)}$ and

$$\|g(x) - g(y)\|_{X^{\gamma(\epsilon)}} \leq c(1 + \|x\|_{X^{1+\epsilon}}^{\rho-1} + \|y\|_{X^{1+\epsilon}}^{\rho-1})\|x - y\|_{X^{1+\epsilon}},$$

for all $x, y \in X^{1+\epsilon}$.

We recall from [2] the following classification for a map f which is ϵ -regular, for $\epsilon \in I$, relative to the pair (X^1, X^0) .

- If $I = [0, \epsilon_1]$ for some $\epsilon_1 > 0$ and $\gamma(0) > 0$, we say that f is a *subcritical* map relative to (X^1, X^0) .
- If $I = [0, \epsilon_1]$ for some $\epsilon_1 > 0$ with $\gamma(\epsilon) = \rho\epsilon$, $\epsilon \in I$, and if f is not subcritical, then we say that f is a *critical* map relative to (X^1, X^0) .
- If $I = (0, \epsilon_1]$ for some $\epsilon_1 > 0$ with $\gamma(\epsilon) = \rho\epsilon$, $\epsilon \in I$, and f is not subcritical or critical, then we say that f is a *double-critical* map relative to (X^1, X^0) .
- If $I = [\epsilon_0, \epsilon_1]$ for some $\epsilon_1 > \epsilon_0 > 0$ with $\gamma(\epsilon_0) > \rho\epsilon_0$ and f is not subcritical, critical or double critical, then we say that f is an *ultra-subcritical* map relative to (X^1, X^0) .
- If $I = [\epsilon_0, \epsilon_1]$ for some $\epsilon_1 > \epsilon_0 > 0$ with $\gamma(\epsilon) = \rho\epsilon$, $\epsilon \in I$, and if f is not subcritical, critical, double critical or ultra-subcritical, then we say that f is an *ultra-critical* map relative to (X^1, X^0) .

Next theorem is the main result of Section 4 (where the class $\mathcal{F}(\epsilon, \rho, \gamma(\epsilon), c, \nu(\cdot), \xi)$ is specified).

Theorem 7. Let $f \in \mathcal{F}(\epsilon, \rho, \gamma(\epsilon), c, \nu(\cdot), \xi)$. If $v_0 \in X^1$, there exist positive values r and τ_0 such that for any $u_0 \in B_{X^1}(v_0, r)$ there exists a continuous function $u(\cdot, u_0) : [0, \tau_0] \rightarrow X^1$ with $u(0) = u_0$, which is an ϵ -regular mild solution to the problem

$$\begin{cases} cD_t^\alpha u = Au + f(t, u(t)), & t > 0, \quad \alpha \in (0, 1), \\ u(0) = u_0. \end{cases} \quad (3)$$

This solution satisfies

$$\begin{aligned} u &\in C((0, \tau_0]; X^{1+\theta}), \quad 0 \leq \theta < \gamma(\epsilon), \\ \lim_{t \rightarrow 0^+} t^{\alpha\theta} \|u(t, u_0)\|_{X^{1+\theta}} &= 0, \quad 0 < \theta < \gamma(\epsilon). \end{aligned}$$

Moreover, for each $\theta_0 < \gamma(\epsilon) + \epsilon - \rho\epsilon$ there exists a constant $C > 0$ such that if $u_0, w_0 \in B_{X^1}(v_0, r)$, then

$$t^{\alpha\theta} \|u(t, u_0) - u(t, w_0)\|_{X^{1+\theta}} \leq C \|u_0 - w_0\|_{X^1} \quad \forall t \in [0, \tau_0], \quad 0 \leq \theta \leq \theta_0.$$

In particular we obtain an existence theorem in X^1 without the nonlinearity being defined on X^1 . The main motivation to consider situations as in Theorem 7 is the fact that if the only requirement on the nonlinear term is that $f : X^1 \rightarrow X^0$ be locally Lipschitz, we cannot ensure that problem (1) is well-posed in an ϵ -regular sense. For example, taking $f(u) = -2Au$, which satisfies $f : X^1 \rightarrow X^0$ and is globally Lipschitz, we will have $cD_t^\alpha u = -Au$, which is not locally well-posed, in general. Hence, some extra conditions should be imposed on f to guarantee the existence of solutions of the above problem.

As application we consider our abstract result in the framework of fractional heat equations.

Example 8. Let $\Omega \subset \mathbb{R}^N$ be a bounded domain with smooth boundary. We will treat the fractional equation (with $\alpha \in (0, 1)$)

$$\begin{cases} \partial_t^\alpha u = \Delta u + u|u|^{\rho-1} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (4)$$

in the L^q theory, $1 < q < \infty$ and $q = \frac{N(\rho-1)}{2}$. Observe that this equation for the case $\alpha = 1$ is well-known (e.g. cf. [2, 5, 23, 24]).

Let $L = \Delta$ with Dirichlet boundary conditions in Ω . Then, L can be seen as an unbounded operator in $E_q^0 = L^q(\Omega)$ with domain

$$D(A) := W^{2,q}(\Omega) \cap W_0^{1,q}(\Omega).$$

It is well-known that the scale of fractional power spaces $\{E_q^\alpha\}_{\alpha \in \mathbb{R}}$ associated to L verifies (see [1, 2])

$$\begin{aligned} E_q^\alpha &\hookrightarrow H_q^{2\alpha}(\Omega), \quad \alpha \geq 0, \quad 1 < q < \infty, \\ E_q^{-\alpha} &\hookrightarrow (E_{q'}^\alpha)', \quad \alpha \geq 0, \quad 1 < q < \infty, \quad q' = \frac{q}{q-1}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} E_q^\alpha &\hookrightarrow L^r(\Omega) \text{ for } r \leq \frac{Nq}{N-2q\alpha}, \quad 0 \leq \alpha \leq \frac{N}{2q} \\ E_q^0 &= L^q(\Omega), \\ E_q^\alpha &\hookrightarrow L^r(\Omega) \text{ for } r \geq \frac{N}{N-2q\alpha}, \quad -\frac{N}{2q'} \leq \alpha \leq 0. \end{aligned}$$

Let L_α be the realization of L in E_q^α . Then we have that L_α is an isometry from $E_q^{\alpha+1}$ into E_q^α and

$$L_\alpha : D(L_\alpha) = E_q^{\alpha+1} \subset E_q^\alpha \rightarrow E_q^\alpha$$

is a sectorial operator. Denote $X_q^\alpha := E_q^{\alpha-1}$, $\alpha \in \mathbb{R}$. The fractional power spaces associated to $L_{-1} : X_q^1 \subset X_q^0 \rightarrow X_q^0$ satisfy

$$\begin{aligned} X_q^\alpha &\hookrightarrow L^r(\Omega) \text{ for } r \leq \frac{Nq}{N+2q-2q\alpha}, \quad 1 \leq \alpha < \frac{2q+N}{2q}, \\ X_q^1 &= L^q(\Omega), \\ X_q^\alpha &\hookrightarrow L^r(\Omega) \text{ for } r \geq \frac{Nq}{N+2q-2q\alpha}, \quad \frac{2q'-N}{2q'} < \alpha \leq 1. \end{aligned}$$

Following [2, Lemma 8] and with the above constructions, we may classify the non-linear term of (4) in the following way.

- If $\rho > \frac{N}{N-2}$, then f is an ϵ -regular map relative to (X_q^1, X_q^0) for $0 \leq \epsilon < \frac{N}{N+2q}$ and $\gamma(\epsilon) = \rho\epsilon$. Therefore f is a critical map.
- If $\rho = \frac{N}{N-2}$, then f is an ϵ -regular map relative to (X_q^1, X_q^0) for $0 < \epsilon < \frac{N}{N+2q}$ and $\gamma(\epsilon) = \rho\epsilon$. Therefore f is a double-critical map.
- If $\frac{N+2}{N} < \rho < \frac{N}{N-2}$, then f is an ϵ -regular map relative to (X_q^1, X_q^0) for $0 < \epsilon_0(\rho) < \epsilon < \frac{1}{\rho}$, with $\epsilon_0(\rho) = \frac{1}{\rho} \left(1 - \frac{N}{2} \left(1 - \frac{2}{N(\rho-1)} \right) \right) > 0$, and $\gamma(\epsilon) = \rho\epsilon$. Therefore f is an ultra-critical map.

Applying Theorem 7, for each $u_0 \in L^q(\Omega)$ we ensure the existence of an ϵ -regular solution to the above problem, starting at u_0 , for any $\epsilon \in \left(\epsilon_0(\rho), \frac{N}{N+2q} \right)$. Furthermore, for any $0 < \theta < \gamma\left(\frac{N}{N+2}\right) = 1$ this solution verifies

$$t^{\alpha\theta} \|u(t)\|_{X^{1+\theta}} \rightarrow 0 \text{ as } t \rightarrow 0^+,$$

$$t^{\alpha\theta} \|u(t, u_0) - u(t, v_0)\|_{X^{1+\theta}} \leq C \|u_0 - v_0\|_{L^q}, \quad 0 < t < \tau(u_0, v_0).$$

In Section 5 we consider the problem of comparison and positive solutions. We start this with the linear version of (1); particularly, we establish the equivalence between the positivity of the families $\{E_\alpha(t^\alpha A)\}_{t \geq 0}$ and $\{E_{\alpha, \alpha}(t^\alpha A)\}_{t \geq 0}$ with the positivity of the function $\lambda \mapsto (\lambda^\alpha - A)^{-1}$ (see Proposition 30 and Proposition 32 below). With respect to the semilinear problem (1), we establish the following result on positive solutions.

Theorem 9. *Let (X, \leq_X) be an ordered Banach space and suppose that the families $\{E_\alpha(t^\alpha A)\}_{t \geq 0}$ and $\{E_{\alpha, \alpha}(t^\alpha A)\}_{t \geq 0}$ are increasing. If $0 \leq_X f(t, x)$ a.e. $t \in [0, t_0]$ and for all $x \in X$ with $0 \leq_X x$, then $0 \leq_X u_0$ implies that the local mild solution $u_f(t, u_0)$ is positive, i.e., $0 \leq_X u_f(t, u_0)$ for all $t \in [0, t_0]$.*

This allows us to conclude the following comparison result.

Theorem 10. *Let (X, \leq_X) be an ordered Banach space and suppose that the families $\{E_\alpha(t^\alpha A)\}_{t \geq 0}$ and $\{E_{\alpha, \alpha}(t^\alpha A)\}_{t \geq 0}$ are increasing.*

- (i) *Let $u_0, u_1 \in X$ be given, and assume that there exist $t_0, t_1 \in [0, \infty)$ such that $\{u_f(t, u_i)\}_{i=0,1}$ are local mild solutions in $[0, t_i]$, $i \in \{0, 1\}$, to*

$$\begin{cases} cD_t^\alpha u(t) = Au(t) + f(t, u(t)), & t > 0, \\ u(0) = u_i \in X. \end{cases}$$

Then, if $t^ = \min\{t_0, t_1\}$, $f(t, \cdot)$ is increasing a.e. $t \in [0, t^*)$, and $u_1 \leq_X u_0$, it holds that $u_f(t, u_1) \leq_X u_f(t, u_0)$ for all $t \in [0, t^*]$.*

- (ii) *Consider functions f_0 and f_1 , and $u_0 \in X$, and assume that there exist $t_0, t_1 \in [0, \infty)$ such that $\{u_{f_i}(t, u_0)\}_{i=0,1}$ are local mild solutions in $[0, t_i]$, $i \in \{0, 1\}$, to*

$$\begin{cases} cD_t^\alpha u(t) = Au(t) + f_i(t, u(t)), & t > 0, \\ u(0) = u_0 \in X. \end{cases}$$

Then, if $t^ = \min\{t_0, t_1\}$ and $f_0(t, x) \leq_X f_1(t, x)$ a.e. $t \in [0, t^*]$ and for all $x \in X$, it holds that $u_{f_0}(t, u_0) \leq_X u_{f_1}(t, u_0)$ for all $t \in [0, t^*]$.*

- (iii) Consider functions f_0 and f_1 , and $u_0, u_1 \in X$, and assume that there exist $t_0, t_1 \in [0, \infty)$ such that $\{u_{f_i}(t, u_i)\}_{i=0,1}$ are local mild solutions in $[0, t_i]$, $i \in \{0, 1\}$, to

$$\begin{cases} cD_t^\alpha u(t) = Au(t) + f_i(t, u(t)), & t > 0, \\ u(0) = u_i \in X. \end{cases}$$

Then, if $t^* = \min\{t_0, t_1\}$, and $x \leq_X y$ imply $f_0(t, x) \leq_X f_1(t, y)$ a.e. $t \in [0, t^*]$, and $u_0 \leq_X u_1$, it holds that $u_{f_0}(t, u_0) \leq_X u_{f_1}(t, u_1)$ for all $t \in [0, t^*]$.

Next result uses the comparison of solutions to establish the existence of a unique mild solution in a certain interval. As a consequence, when the solutions of the auxiliary problems are globally defined in time, from the continuation results we obtained previously, we will conclude the existence of a unique global mild solution (see Example 12 below).

Corollary 11. Let (X, \leq_X) be an ordered Banach space and $C > 0$ such that

$$x \leq_X y \leq_X z \Rightarrow \|y\|_X \leq C(\|x\|_X + \|z\|_X) \quad \forall x, y, z \in X.$$

Suppose that the families $\{E_\alpha(t^\alpha A)\}_{t \geq 0}$ and $\{E_{\alpha, \alpha}(t^\alpha A)\}_{t \geq 0}$ are increasing. Let $u_0 \in X$ and consider f_1, f_2 , and $g : [0, \infty) \times X \rightarrow X$ as in Theorem 2, and that g is bounded. Moreover, assume that $f_1(t, x) \leq_X g(t, x) \leq_X f_2(t, x)$ a.e. $t \geq 0$ and for all $x \in X$. Suppose that the problems

$$\begin{cases} cD_t^\alpha u(t) = Au(t) + f_i(t, u(t)), & t > 0, \\ u(0) = u_0 \in X. \end{cases}$$

for $i = \{1, 2\}$ have unique local mild solutions in $[0, t^*]$ for some $t^* > 0$. Then, the problem

$$\begin{cases} cD_t^\alpha u(t) = Au(t) + g(t, u(t)), & t > 0, \\ u(0) = u_0 \in X. \end{cases}$$

has a unique local mild solution in $[0, t^*]$.

Finally, as an application of this last result (and some of Section 3) we ensure existence of global mild solution for the following fractional partial differential equation coming from heat conduction theory:

Example 12. Let $\Omega \subset \mathbb{R}^N$, $N \in \mathbb{N}$, be a bounded open subset with smooth boundary $\partial\Omega$. Consider the problem

$$\begin{cases} c\partial_t^\alpha u(t, x) = D\Delta u(t, x) + F(u(t, x)), & t > 0, & x \in \Omega, \\ \partial u / \partial \vec{v}(t, x) = 0, & t > 0, & x \in \partial\Omega, \\ u(0, x) = u_0(x), & & x \in \Omega. \end{cases} \quad (5)$$

where $c\partial_t^\alpha$ is Caputo's fractional derivative, $u = (u_1, \dots, u_n)^\top$, $n \geq 1$, $\frac{\partial u}{\partial \vec{v}} = (\langle \nabla u_1, \vec{v} \rangle, \dots, \langle \nabla u_n, \vec{v} \rangle)^\top$, where \vec{v} is the outward normal vector, and D is the diagonal matrix

$$D = \begin{bmatrix} d_1 & 0 & 0 & \dots & 0 \\ 0 & d_2 & 0 & \dots & 0 \\ 0 & 0 & d_3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & d_n \end{bmatrix}_{n \times n}$$

with $d_i > 0$, for all $i \in \{0, 1, \dots, n\}$. The nonlinearity

$$F = (F_1, \dots, F_n) : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

is assumed to be locally Lipschitz.

To treat this problem we set $X = (L^q(\Omega))^n$, for $1 < q < \infty$, with the usual order \leq_X and we consider $A_D = \text{diag}(A_1, \dots, A_n)$, where for each $i \in \{0, \dots, n\}$ we define $A_i : D(A_i) \subset L^q(\Omega) \rightarrow L^q(\Omega)$ by

$$\begin{aligned} D(A_i) &= \{\phi \in W^{2,q}(\Omega); \partial\phi/\partial\vec{\nu} = 0 \text{ on } \partial\Omega\} =: W_{\mathcal{N}}^{2,q}(\Omega); \\ A_i\phi &= -d_i\Delta\phi \end{aligned}$$

and suppose that $\inf\{\text{Re}(\sigma(A_D))\} = \tilde{d}_1 < 0$. In [6, Proposition 2.8] it was shown that this operator is sectorial and has positive resolvent. Then, rewriting (5) in the abstract form

$$\begin{cases} cD_t^\alpha u(t) = -A_D u(t) + F(u(t)), & t > 0, \\ u(0) = u_0 \in (L^q(\Omega))^n, \end{cases}$$

we may ensure, by Theorem 2, existence and uniqueness of a local mild solution u . If there exist constants $c_0 \in [0, -\tilde{d}_1)$ and $c_1 \geq 0$ such that

$$-c_0|u_i| - c_1 \leq F_i(u) \leq c_0|u_i| + c_1 \quad \forall i \in \{1, \dots, n\},$$

then problem (5) has a global mild solution.

Indeed, we conclude that

$$-c_0\|u\| - C_1 \leq_X F(u) \leq_X c_0\|u\| + C_1 \quad \forall u \in (L^q(\Omega))^n,$$

where $\|u\| := (|u_1|, \dots, |u_n|)$ and $C_1 = (c_1, \dots, c_1)$.

Since the problems

$$\begin{cases} cD_t^\alpha u(t) = -A_D u(t) - c_0\|u(t)\| - C_1, & t > 0, \\ u(0) = u_0 \in (L^q(\Omega))^n, \end{cases}$$

and

$$\begin{cases} cD_t^\alpha u(t) = -A_D u(t) + c_0\|u(t)\| + C_1, & t > 0, \\ u(0) = u_0 \in (L^q(\Omega))^n, \end{cases}$$

have global mild solutions (by using Theorem 4), we conclude by Corollary 11 and Theorems 3 and 4 that (5) possesses a unique global mild solution.

2. PRELIMINARY RESULTS

Let $(X, \|\cdot\|)$ be a complex Banach space. As usual, for a linear operator A , we denote by $D(A)$ the domain of A , by $\sigma(A)$ its spectrum, while $\rho(A) := \mathbb{C} - \sigma(A)$ is the resolvent set of A . Moreover, we denote by $\mathcal{L}(Y, Z)$ the space of all bounded linear operators between two normed spaces Y and Z with the operator norm $\|\cdot\|_{\mathcal{L}(Y, Z)}$. We abbreviate this notation to $\mathcal{L}(Y)$ when $Y = Z$. When no confusion arise, we write $\|T\|_{\mathcal{L}(Y)}$ as $\|T\|$ for every $T \in \mathcal{L}(Y)$. Finally, for $\beta \geq 0$ we denote by X^β the fractional power spaces associated to the operator A , being $X^0 = X$ and $X^1 = D(A)$.

The aim of this section is to present the basic tools we will use in this manuscript. Our results include the classical theory of the Mittag-Leffler function that is closely related to the solutions of the fractional differential equations.

2.1. Mittag-Leffler functions and sectorial operators. We start with a generalization of the Cauchy representation for semigroups associated to sectorial operators. To this end we recall that given $\epsilon > 0$ and $\theta \in (\pi/2, \pi)$, the Hankel path $Ha = Ha(\epsilon, \theta)$ is the path given by $Ha = Ha_1 + Ha_2 - Ha_3$, where Ha_i are such that

$$Ha_1 := \{te^{i\theta} : t \in [\epsilon, \infty)\}, \quad Ha_2 := \{\epsilon e^{it} : t \in [-\theta, \theta]\}, \quad Ha_3 := \{te^{-i\theta} : t \in [\epsilon, \infty)\}. \quad (6)$$

Lemma 13. *Consider $\alpha \in (0, 1)$ and suppose that $A : D(A) \subset X \rightarrow X$ is a sectorial operator. Then, the functions*

$$E_\alpha(t^\alpha A) := \frac{1}{2\pi i} \int_{Ha} e^{\lambda t} \lambda^{\alpha-1} (\lambda^\alpha - A)^{-1} d\lambda, \quad t \geq 0,$$

and

$$E_{\alpha,\alpha}(t^\alpha A) := \frac{t^{1-\alpha}}{2\pi i} \int_{Ha} e^{\lambda t} (\lambda^\alpha - A)^{-1} d\lambda, \quad t \geq 0$$

(where Ha is given by (6)) are well defined. Furthermore, there exists a constant $M > 0$ such that

$$\|E_\alpha(t^\alpha A)\| \leq M \quad \text{and} \quad \|E_{\alpha,\alpha}(t^\alpha A)\| \leq M \quad \text{for all } t \geq 0.$$

Proof. Let $\phi \in (\pi/2, \pi)$, S_ϕ be the sector associated with the sectorial operator A and choose arbitrary values $\epsilon > 0$ and $\theta \in (\pi/2, \phi]$. We will estimate the function $\|E_\alpha(t^\alpha A)\|$ on each Ha_i , according to definition (6), for any $t > 0$. Just observe that for each fixed $t \neq 0$, if we assume that $\epsilon = 1/t$, then

- On Ha_1 , it holds that

$$\left\| \frac{1}{2\pi i} \int_{Ha_1} e^{\lambda t} \lambda^{\alpha-1} (\lambda^\alpha - A)^{-1} d\lambda \right\| \leq \frac{1}{2\pi} \left\| \int_\epsilon^\infty e^{tse^{i\theta}} (se^{i\theta})^{\alpha-1} ((se^{i\theta})^\alpha - A)^{-1} e^{i\theta} ds \right\|$$

and using that if $\lambda = se^{i\theta} \in Ha(\epsilon, \theta) \subset S_\phi$, then $\lambda^\alpha \in S_\phi$, we obtain by the sectorial property that

$$\begin{aligned} \left\| \frac{1}{2\pi i} \int_{Ha_1} e^{\lambda t} \lambda^{\alpha-1} (\lambda^\alpha - A)^{-1} d\lambda \right\| &\leq \frac{N}{2\pi} \int_\epsilon^\infty e^{ts \cos(\theta)} |(se^{i\theta})|^{-1} ds \\ &\leq \frac{N}{2\pi \epsilon} \int_\epsilon^\infty e^{ts \cos(\theta)} ds = \frac{N e^{\cos(\theta)}}{-2\pi \cos(\theta)}. \end{aligned}$$

- On Ha_2 , we have that

$$\begin{aligned} \left\| \frac{1}{2\pi i} \int_{Ha_2} e^{\lambda t} \lambda^{\alpha-1} (\lambda^\alpha - A)^{-1} d\lambda \right\| &= \frac{1}{2\pi} \left\| \int_{-\theta}^\theta e^{t\epsilon e^{is}} (\epsilon e^{is})^{\alpha-1} ((\epsilon e^{is})^\alpha - A)^{-1} i\epsilon e^{is} ds \right\| \\ &\leq \frac{N}{2\pi} \int_{-\theta}^\theta e^{t\epsilon \cos(s)} ds \leq \frac{\theta N e}{\pi}. \end{aligned}$$

- On Ha_3 we proceed in the same way as in Ha_1 .

Taking M as the maximum over all the bounds obtained above, we deduce that $\|E_\alpha(At^\alpha)\|$ is well defined for each $t \geq 0$.

Now we shall seek a uniform bound for the function $\|E_\alpha(At^\alpha)\|$ for all $t \geq 0$. For this, just observe that if $\epsilon < \epsilon'$ and $\pi/2 < \theta' < \theta < \phi$ and we take $Ha = Ha(\epsilon, \theta)$ and

$Ha' = Ha(\epsilon', \theta')$, we obtain the equality

$$\frac{1}{2\pi i} \int_{Ha} e^{\lambda t} \lambda^{\alpha-1} (\lambda^\alpha - A)^{-1} d\lambda = \frac{1}{2\pi i} \int_{Ha'} e^{\lambda t} \lambda^{\alpha-1} (\lambda^\alpha - A)^{-1} d\lambda \quad \forall t \geq 0.$$

This is enough to justify that the estimate obtained before is uniform on $t > 0$. To verify the boundedness for $t = 0$, we observe that making a change of variables

$$E_\alpha(t^\alpha A) = \frac{1}{2\pi i} \int_{Ha} e^\lambda \lambda^{\alpha-1} (\lambda^\alpha - At^\alpha)^{-1} d\lambda, \quad t \geq 0,$$

and therefore, using the dominated convergence theorem, we conclude that $E_\alpha(0^\alpha A)$ is the identity operator.

A similar procedure proves that $\|E_{\alpha,\alpha}(At^\alpha)\| \leq M$ for all $t \in [0, \infty)$. \square

Next result give us information on the Mittag-Leffler families in the fractional power spaces X^β , $\beta \geq 0$, associated to the sectorial operator A .

Lemma 14. *Consider $\alpha \in (0, 1)$, $0 \leq \beta \leq 1$, and suppose that $A : D(A) \subset X^0 \rightarrow X^0$ is a sectorial operator. Then, there exists a constant $M > 0$ such that*

$$\|E_\alpha(t^\alpha A)x\|_{X^\beta} \leq Mt^{-\alpha\beta} \|x\|_{X^0} \quad \text{and} \quad \|t^{\alpha-1} E_{\alpha,\alpha}(t^\alpha A)x\|_{X^\beta} \leq Mt^{\alpha(1-\beta)-1} \|x\|_{X^0}$$

for all $t > 0$.

Proof. We have that

$$\begin{aligned} \|E_\alpha(t^\alpha A)x\|_{X^\beta} &= \left\| \frac{1}{2\pi i} \int_{Ha} e^{\lambda t} \lambda^{\alpha-1} A^\beta (\lambda^\alpha - A)^{-1} x d\lambda \right\|_{X^0} \\ &\leq \frac{t^{-\alpha}}{2\pi} \left\| \int_{Ha} e^\lambda \lambda^{\alpha-1} A^\beta ((\lambda/t)^\alpha - A)^{-1} x d\lambda \right\|_{X^0} \\ &\leq t^{-\alpha\beta} \left(\frac{C}{2\pi} \int_{Ha} |e^\lambda \lambda^{\alpha\beta-1}| |d\lambda| \right) \|x\|_{X^0}. \end{aligned}$$

Hence, it is sufficient to choose $M > 0$ such that $\frac{C}{2\pi} \int_{Ha} |e^\lambda \lambda^{\alpha\beta-1}| |d\lambda| \leq M$. By a similar procedure one may prove the second estimate. \square

Our goal now is to establish an expression for Mittag-Leffler functions associated to sectorial operators similar to the second fundamental limit for semigroups. For this, we recall the Post-Widder inversion formula, which can be found in [26]. This expression will be very useful to our following results.

Lemma 15 (Post-Widder). *Let $u : [0, \infty) \rightarrow X$ be a continuous function such that $u(t) = O(\exp(\omega t))$ as $t \rightarrow \infty$ for some $\omega \in \mathbb{R}$, and let \hat{u} be the Laplace transform of u . Then*

$$u(t) = \lim_{n \rightarrow \infty} \frac{(-1)^n}{n!} \left(\frac{n}{t}\right)^{n+1} \left(\frac{d^n}{d\lambda^n} \hat{u}\right)(n/t),$$

uniformly on compact sets of $(0, \infty)$.

Proposition 16. Consider $A : D(A) \subset X \rightarrow X$ a sectorial operator and $\alpha \in (0, 1)$. Let $\{E_\alpha(t^\alpha A)\}_{t \geq 0}$ and $\{E_{\alpha, \alpha}(t^\alpha A)\}_{t \geq 0}$ be the Mittag-Leffler families associated to A . Then, for each $x \in X$ we have

$$E_\alpha(t^\alpha A)x = \lim_{n \rightarrow \infty} \frac{1}{n!} \sum_{k=1}^{n+1} b_{k, n+1}^\alpha \left(\left(\frac{n}{t} \right)^\alpha \left(\left(\frac{n}{t} \right)^\alpha - A \right)^{-1} \right)^k x, \quad (7)$$

and

$$t^{\alpha-1} E_{\alpha, \alpha}(t^\alpha A)u_0 = \lim_{n \rightarrow \infty} \frac{1}{n!} \sum_{k=1}^n \alpha k b_{k, n}^\alpha \left(\frac{n}{t} \right)^{\alpha k+1} \left(\left(\frac{n}{t} \right)^\alpha - A \right)^{-(k+1)} x, \quad (8)$$

uniformly on compact sets of $(0, \infty)$, where the positive real numbers $b_{k, n}^\alpha$ are defined by

$$\begin{cases} b_{1, 1}^\alpha = 1, \\ b_{k, n}^\alpha = (n-1-\alpha k)b_{k, n-1}^\alpha + \alpha(k-1)b_{k-1, n-1}^\alpha, & \text{if } 1 \leq k \leq n \text{ and } n = 2, 3, \dots \\ b_{k, n}^\alpha = 0, & \text{if } k > n \text{ and } n = 1, 2, \dots \end{cases} \quad (9)$$

Proof. By Lemma 13 there exists $M \geq 1$ such that $\|E_\alpha(t^\alpha A)\| \leq M$ and $\|E_{\alpha, \alpha}(t^\alpha A)\| \leq M$ for all $t \geq 0$. By induction in n , we obtain for $Re \lambda > 0$ that

$$\frac{d^n}{d\lambda^n} (\lambda^{\alpha-1} (\lambda^\alpha - A)^{-1}) = (-1)^n \lambda^{-(n+1)} \sum_{k=1}^{n+1} b_{k, n+1}^\alpha [\lambda^\alpha (\lambda^\alpha - A)^{-1}]^k, \quad n = 1, 2, \dots$$

Furthermore, for each $x \in X$ the function $u(t) = E_\alpha(t^\alpha A)x$ is such that

$$\hat{u}(\lambda) = \lambda^{\alpha-1} (\lambda^\alpha - A)^{-1} x.$$

Hence, (7) follows by Lemma 15.

On the other hand, to prove (8), observe that

$$\frac{d^n}{d\lambda^n} (\lambda^\alpha - A)^{-1} = (-1)^n \lambda^{-n} \sum_{k=1}^n \alpha k b_{k, n}^\alpha \lambda^{\alpha k} (\lambda^\alpha - A)^{-(k+1)}, \quad n = 1, 2, \dots$$

and for each $x \in X$, the function $v(t) = t^{\alpha-1} E_{\alpha, \alpha}(t^\alpha A)x$ verifies $\hat{v}(\lambda) = (\lambda^\alpha - A)^{-1} x$. Using again Lemma 15, the proof is finished. \square

2.2. Ordered Banach spaces. We recall briefly the basic properties of an order on a Banach space X , in order to develop our comparison results.

Definition 17. An ordered Banach space is a couple (X, \leq_X) where X is a Banach space and \leq_X is an order relation in X such that for every $x, y, z \in X$ and for any scalar $\lambda \geq 0$, it holds that

- (i) $x \leq_X y$ implies $x + z \leq_X y + z$;
- (ii) $x \leq_X y$ implies $\lambda x \leq_X \lambda y$;
- (iii) the positive cone $C = \{x \in X : 0 \leq_X x\}$ is closed in X .

Remark 18. Observe that $x \leq_X y$ is equivalent to $0 \leq_X y - x$. Furthermore, $x \leq_X 0$ if and only if $0 \leq_X -x$.

Example 19. Consider $1 \leq p \leq \infty$. The Banach spaces $L^p(\Omega)$ and $C(\bar{\Omega})$ with the order " $f \leq g$ if and only if $f(x) \leq g(x)$ a.e." are ordered Banach spaces.

Definition 20. Suppose that (X, \leq_X) and (Y, \leq_Y) are ordered Banach spaces. A function $T : D(T) \subset X \rightarrow Y$ is called increasing if $x \leq_X y$ implies $Tx \leq_Y Ty$.

T is called positive if $0 \leq_X x$ implies $0 \leq_Y Tx$, for all $x, y \in D(T)$.

Lemma 21. Let (X, \leq_X) be an ordered Banach space and consider $f \in L^1(t_0, t_1; X)$ such that $0 \leq_X f(t)$ a.e. $t \in (t_0, t_1)$. Then

$$0 \leq_X \int_{t_0}^{t_1} f(s) ds.$$

3. LOCAL WELL POSEDNESS

We start this section with a proof of our result on local existence and uniqueness of mild solutions to (1).

Proof of Theorem 2. Given $u_0 \in X$, let $B_{u_0}(r)$ be the open ball with center u_0 and radius $r > 0$ and $L = L(B_{u_0}(r))$ be the Lipschitz constant of f associated to $B_{u_0}(r)$. Fix $\beta \in (0, r)$ and choose $t_0 > 0$ such that

$$(M/\alpha)(L\beta + C)t_0^\alpha \leq \beta/2 \quad \text{and} \quad \|E_\alpha(At^\alpha)u_0 - u_0\| \leq \beta/2 \quad \forall t \in [0, t_0].$$

where $M = \sup_{t \in [0, \infty)} \|E_{\alpha, \alpha}(At^\alpha)\|$ and $C = \sup_{t \in [0, t_0]} \|f(s, u_0)\|$. Consider

$$K := \{u \in C([0, t_0]; X) : u(0) = u_0 \quad \text{and} \quad \|u(t) - u_0\| \leq \beta \quad \forall t \in [0, t_0]\}$$

and define the operator T on K by

$$T(u(t)) = E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha, \alpha}((t-s)^\alpha A) f(s, u(s)) ds.$$

If $u \in K$, then $T(u(0)) = u_0$ and $T(u(t)) \in C([0, t_0]; X)$. Furthermore, we have that

$$\begin{aligned} \|T(u(t)) - u_0\| &\leq \|E_\alpha(At^\alpha)u_0 - u_0\| \\ &\quad + \int_0^t (t-s)^{\alpha-1} M (\|f(s, u(s)) - f(s, u_0)\| + \|f(s, u_0)\|) ds \\ &\leq \|E_\alpha(At^\alpha)u_0 - u_0\| + (M/\alpha)(L\beta + C)t_0^\alpha \\ &\leq \beta/2 + \beta/2 = \beta, \end{aligned}$$

that is, $T(K) \subset K$. Now, if $u, v \in K$, we see that

$$\begin{aligned} \|T(u(t)) - T(v(t))\| &\leq \int_0^t (t-s)^{\alpha-1} M \|f(s, u(s)) - f(s, v(s))\| ds \\ &\leq \frac{LMt_0^\alpha}{\alpha} \sup_{s \in [0, t_0]} \|u(s) - v(s)\| \\ &\leq \frac{1}{2} \sup_{s \in [0, t_0]} \|u(s) - v(s)\|. \end{aligned}$$

So, by the Banach contraction principle we have that T has a unique fixed point in K . We will denote this fixed point by w .

Uniqueness of local mild solution to (1) follows from the Singular Gronwall Inequality (see [9, Lemma 7.1.1, page 188]). Therefore, we obtain that the solution ω is the unique continuous function that satisfies the integral equation and this finishes the proof. \square

Definition 22. Let $u : [0, t_0] \rightarrow X$ be a local mild solution in $[0, t_0]$ to (1). If $t_1 > t_0$ and $v : [0, t_1] \rightarrow X$ is a local mild solution to (1) in $[0, t_1]$, then we say that v is a continuation of u in $[0, t_1]$.

The remainder of this paragraph will be devoted to the problem of continuation of local mild solutions and existence of global mild solutions of (1).

Proof of Theorem 3. Let $u : [0, t_0] \rightarrow X$ be the local mild solution to (1) in $[0, t_0]$. Since f is locally Lipschitz, there exist $r > 0$, an open ball $B = B(u(t_0))$ with center in $u(t_0)$ and radius r , and $L = L_{u(t_0)}$ the Lipschitz constant of f associated to B . Fix $\beta \in (0, r)$ and choose $\tau > 0$ such that the following conditions are satisfied

- $\|E_\alpha(At^\alpha)u_0 - E_\alpha(At_0^\alpha)u_0\| \leq \beta/4$,
- $(M/\alpha)(L\beta + C)\tau^\alpha \leq \beta/4$, $(MD/\alpha)[t^\alpha - (t - t_0)^\alpha - t_0^\alpha] \leq \beta/4$,
- $\int_0^{t_0} (t_0 - s)^{\alpha-1} \|[E_{\alpha,\alpha}(A(t-s)^\alpha) - E_{\alpha,\alpha}(A(t_0 - s)^\alpha)]f(s, u(s))\| ds \leq \beta/4$

for all $t \in [t_0, t_0 + \tau]$, where

$$\begin{aligned} C &= \sup_{s \in [0, t_0]} \|f(s, u(t_0))\|, \\ D &= \sup_{s \in [0, t_0]} \|f(s, u(s))\|, \\ M &= \max\left\{ \sup_{t \in [0, \infty)} \|E_{\alpha,\alpha}(At^\alpha)\|, \sup_{t \in [0, \infty)} \|E_\alpha(At^\alpha)\| \right\}. \end{aligned}$$

Consider

$$K := \left\{ w \in C([0, t_0 + \tau]; X) : \begin{array}{l} w(t) = u(t) \text{ for all } t \in [0, t_0] \text{ and} \\ \|w(t) - u(t_0)\| \leq \beta \text{ for all } t \in [t_0, t_0 + \tau] \end{array} \right\}$$

and $T : K \rightarrow C([0, t_0 + \tau]; X)$ given by

$$T(w(t)) = E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, w(s)) ds.$$

We check that $T(K) \subset K$.

- i) If $w \in K$, then $w(t) = u(t)$ in $[0, t_0]$ with u the local mild solution to (1) in $[0, t_0]$. So, if $t \in [0, t_0]$,

$$\begin{aligned} T(w(t)) &= E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, w(s)) ds \\ &= E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s)) ds \\ &= u(t) \end{aligned}$$

ii) If $t \in [t_0, t_0 + \tau]$, then

$$\begin{aligned} & \|T(w(t)) - u(t_0)\| \\ & \leq \|E_\alpha(t^\alpha A)u_0 - E_\alpha(t_0^\alpha A)u_0\| + \left\| \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, w(s)) ds \right. \\ & \quad \left. - \int_0^{t_0} (t_0-s)^{\alpha-1} E_{\alpha,\alpha}((t_0-s)^\alpha A) f(s, w(s)) ds \right\| \\ & \leq \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3 + \mathcal{I}_4 \\ & \leq \beta, \end{aligned}$$

where

$$\begin{aligned} \mathcal{I}_1 &= \|E_\alpha(t^\alpha A)u_0 - E_\alpha(t_0^\alpha A)u_0\| \leq \beta/4, \\ \mathcal{I}_2 &= \left\| \int_{t_0}^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, w(s)) ds \right\| \leq \beta/4, \\ \mathcal{I}_3 &= \left\| \int_0^{t_0} [(t-s)^{\alpha-1} - (t_0-s)^{\alpha-1}] E_{\alpha,\alpha}((t-s)^\alpha A) f(s, w(s)) ds \right\| \leq \beta/4, \\ \mathcal{I}_4 &= \left\| \int_0^{t_0} (t_0-s)^{\alpha-1} [E_{\alpha,\alpha}((t-s)^\alpha A) - E_{\alpha,\alpha}((t_0-s)^\alpha A)] f(s, w(s)) ds \right\| \leq \beta/4. \end{aligned}$$

By similar computations, we conclude that for every $t \in [0, t_0 + \tau]$

$$\begin{aligned} \|T(\omega(t)) - T(v(t))\| &\leq \frac{LM\tau^\alpha}{\alpha} \sup_{s \in [0, t_0 + \tau]} \|\omega(s) - v(s)\| \\ &\leq \frac{1}{2} \sup_{s \in [0, t_0 + \tau]} \|\omega(s) - v(s)\| \quad \forall u, v \in K. \end{aligned}$$

Therefore, by the Banach contraction principle, we conclude that there exists a unique fixed point $u^* \in K$ of the integral equation. As in the proof of Theorem 2, it is not difficult to see that u^* is the unique continuous continuation of u in $[0, t_0 + \tau]$. \square

The following result will be useful in the proof of Theorem 4.

Lemma 23. *Consider $\omega \in (0, \infty)$, $u : [0, \omega) \rightarrow X$ a bounded continuous function, and $f : [0, \infty) \times X \rightarrow X$ continuous and bounded. If $\{t_n\} \subset [0, \omega)$ satisfies $\lim_n t_n = \omega$, then*

$$\lim_n \int_0^{t_n} (t_n - r)^{\alpha-1} \| [E_{\alpha,\alpha}(A(t_n - r^\alpha)) - E_{\alpha,\alpha}(A(\omega - r^\alpha))] f(r, u(r)) \| dr = 0.$$

Proof. Let $M = \sup_{t \in [0, \infty)} \|E_{\alpha,\alpha}(At^\alpha)\|$ and $K = \sup_{s \in [0, \omega)} \|f(s, u(s))\|$. Given $\epsilon > 0$, fix $\gamma \in (0, \omega)$ such that

$$\frac{(\omega - \gamma)^\alpha}{\alpha} MK \leq \epsilon/2.$$

Now, choose $N \in \mathbb{N}$ such that

- $t_n > \gamma$ for all $n \geq N$;
- $\int_0^\gamma (t_n - r)^{\alpha-1} \| [E_{\alpha,\alpha}(A(t_n - r^\alpha)) - E_{\alpha,\alpha}(A(\omega - r^\alpha))] f(r, u(r)) \| dr < \epsilon/2$ for all $n \geq N$.

Hence, we conclude that for $n \geq N$,

$$\begin{aligned} & \int_0^{t_n} (t_n - r)^{\alpha-1} \| [E_{\alpha,\alpha}(A(t_n - r^\alpha)) - E_{\alpha,\alpha}(A(w - r^\alpha))] f(r, u(r)) \| dr \\ & \leq \epsilon/2 + \int_\gamma^{t_n} (t_n - r)^{\alpha-1} MK dr \\ & \leq \epsilon/2 + \frac{(\omega - \gamma)^\alpha}{\alpha} MK \leq \epsilon. \end{aligned}$$

□

Proof of Theorem 4. Consider

$$H := \{ \tau \in [0, \infty) : \exists u_\tau : [0, \tau] \rightarrow X \text{ unique local mild solution to (1) in } [0, \tau] \}.$$

If $\sup H = w$, we can consider a continuous function $u : [0, w) \rightarrow X$ that is a local mild solution to (1) in $[0, w)$. If $w = \infty$, then u is a global mild solution in $[0, \infty)$. Otherwise, if $w < \infty$ we will prove that $\limsup_{t \rightarrow w} |u(t)| = \infty$. By contradiction, suppose that there exists $K < \infty$ such that $\|u(t)\| \leq K$ for all $t \in [0, w)$. Therefore, it follows from Lemma 23 that if $\{t_n\} \subset [0, w)$ is a sequence that converges to w , given $\epsilon > 0$, there exists $N \in \mathbb{N}$, such that, if $m, n \geq N$, we have

$$\|E_\alpha(t_n^\alpha A)u_0 - E_\alpha(t_m^\alpha A)u_0\| \leq \epsilon/5, \quad |t_n - t_m|^\alpha \frac{MK}{\alpha} \leq \epsilon/5, \quad |t_n^\alpha - (t_n - t_m)^\alpha - t_m^\alpha| \frac{MK}{\alpha} \leq \epsilon/5,$$

$$\int_0^{t_n} (t_n - r)^{\alpha-1} \| [E_{\alpha,\alpha}((t_n - r)^\alpha A) - E_{\alpha,\alpha}((w - r)^\alpha A)] f(r, u(r)) \| dr \leq \epsilon/5,$$

$$\int_0^{t_m} (t_m - r)^{\alpha-1} \| [E_{\alpha,\alpha}((t_m - r)^\alpha A) - E_{\alpha,\alpha}((w - r)^\alpha A)] f(r, u(r)) \| dr \leq \epsilon/5,$$

where

$$K = \sup_{t \in [0, w)} \|f(t, u(t))\|, \quad M = \max\left\{ \sup_{t \in [0, \infty)} \|E_{\alpha,\alpha}(At^\alpha)\|, \sup_{t \in [0, \infty)} \|E_\alpha(At^\alpha)\| \right\}.$$

Hence, for $n, m \geq N$ and assuming, without loss of generality, that $t_n > t_m$, it follows from the estimate

$$\|u(t_n) - u(t_m)\| \leq \|E_\alpha(t_n^\alpha A)u_0 - E_\alpha(t_m^\alpha A)u_0\| + \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3,$$

where

$$\begin{aligned} \mathcal{I}_1 &= \int_{t_m}^{t_n} (t_n - r)^{\alpha-1} \|E_{\alpha,\alpha}((t_n - r)^\alpha A) f(r, u(r))\| dr \\ &\leq |t_n - t_m|^\alpha \frac{MK}{\alpha}, \end{aligned}$$

$$\begin{aligned}
\mathcal{I}_2 &= \int_0^{t_m} (t_n - r)^{\alpha-1} \| [E_{\alpha,\alpha}((t_n - r)^\alpha A) - E_{\alpha,\alpha}((t_m - r)^\alpha A)] f(r, u(r)) \| dr \\
&\leq \int_0^{t_m} (t_n - r)^{\alpha-1} \| [E_{\alpha,\alpha}((t_n - r)^\alpha A) - E_{\alpha,\alpha}((\omega - r)^\alpha A)] f(r, u(r)) \| dr \\
&\quad + \int_0^{t_m} (t_n - r)^{\alpha-1} \| [E_{\alpha,\alpha}((t_m - r)^\alpha A) - E_{\alpha,\alpha}((\omega - r)^\alpha A)] f(r, u(r)) \| dr \\
&\leq \int_0^{t_n} (t_n - r)^{\alpha-1} \| [E_{\alpha,\alpha}((t_n - r)^\alpha A) - E_{\alpha,\alpha}((\omega - r)^\alpha A)] f(r, u(r)) \| dr \\
&\quad + \int_0^{t_m} (t_m - r)^{\alpha-1} \| [E_{\alpha,\alpha}((t_m - r)^\alpha A) - E_{\alpha,\alpha}((\omega - r)^\alpha A)] f(r, u(r)) \| dr
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{I}_3 &= \int_0^{t_m} |[(t_n - s)^{\alpha-1} - (t_m - s)^{\alpha-1}]| \| E_{\alpha,\alpha}((t_m - s)^\alpha A) f(s, u(s)) \| ds \\
&\leq |t_n^\alpha - (t_n - t_m)^\alpha - t_m^\alpha| \frac{MK}{\alpha},
\end{aligned}$$

that

$$\|u(t_n) - u(t_m)\| \leq \epsilon.$$

This computation shows that $\{u(t_n)\}$ is a Cauchy sequence and therefore it has a limit, $u_t \in X$. Then, we may extend u over $[0, \omega]$ obtaining the equality

$$u(t) = E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s)) ds$$

for all $t \in [0, \omega]$. With this, by Theorem 3, we can extend the solution to some bigger interval, which is a contradiction with the definition of ω . \square

Corollary 24. *Let $f : [0, \infty) \times X \rightarrow X$ be as in Theorem 4 and suppose that there exists a constant $K > 0$ such that if $\|u_0\| \leq K$, then the solutions to (1), while exists, are bounded by K . Then (1) possesses a unique global mild solution.*

4. CRITICAL NONLINEARITIES

Following the notation of [2], we consider the following class of nonlinearities: let ϵ , $\gamma(\epsilon)$, ξ , ζ , c , and δ be positive constants, and a function ν with values in $[0, \delta)$ and $\lim_{t \rightarrow 0^+} \nu(t) = 0$. Define $\mathcal{F}(\epsilon, \rho, \gamma(\epsilon), c, \nu(\cdot), \xi, \zeta)$ as the family of functions f such that, for $t \geq 0$, $f(t, \cdot)$ is an ϵ -regular map relative to the pair (X^1, X^0) , satisfying

$$\|f(t, x) - f(t, y)\|_{X^{\gamma(\epsilon)}} \leq c(\|x\|_{X^{1+\epsilon}}^{\rho-1} + \|y\|_{X^{1+\epsilon}}^{\rho-1} + \nu(t)t^{-\zeta})\|x - y\|_{X^{1+\epsilon}},$$

$$\|f(t, x)\|_{X^{\gamma(\epsilon)}} \leq c(\|x\|_{X^{1+\epsilon}}^\rho + \nu(t)t^{-\xi}),$$

for all $x, y \in X^{1+\epsilon}$.

Without loss of generality we may assume that the function ν is non-decreasing. We will suppose that $0 \leq \zeta \leq \alpha(\gamma(\epsilon) - \epsilon)$ and $0 \leq \xi \leq \alpha\gamma(\epsilon)$, $\alpha \in (0, 1)$. In most cases

in the arguments below we will fix the parameters ϵ , $\gamma(\epsilon)$, ρ , ξ , ζ and c , and we will denote the class \mathcal{F} defined above by $\mathcal{F}(\nu(\cdot))$.

As immediate consequence of Lemma 14 we have that if $0 \leq \theta, \beta \leq 1$, then

$$t^{\alpha(1+\theta-\beta)} \|E_\alpha(t^\alpha A)x\|_{X^{1+\theta}} \leq M \|x\|_{X^\beta} \quad (10)$$

and

$$t^{\alpha(\theta-\beta)+1} \|t^{\alpha-1} E_{\alpha,\alpha}(t^\alpha A)x\|_{X^{1+\theta}} \leq M \|x\|_{X^\beta}.$$

To prove Theorem 7 we need some previous results.

Lemma 25. *Consider $\alpha \in (0, 1)$, $\theta \in [0, 1]$, and a sectorial operator A . The operators $\{t^{\alpha\theta} E_\alpha(t^\alpha A) : X^1 \rightarrow X^{1+\theta}\}_{t>0}$ are bounded linear operators satisfying*

$$\|t^{\alpha\theta} E_\alpha(t^\alpha A)\|_{\mathcal{L}(X^1, X^{1+\theta})} \leq M,$$

with $M > 0$ independent of t . Moreover, given a compact subset J of X^1 , we have

$$\limsup_{t \rightarrow 0^+} \sup_{x \in J} \|t^{\alpha\theta} E_\alpha(t^\alpha A)x\|_{X^{1+\theta}} = 0.$$

Proof. The fact that $\|t^{\alpha\theta} E_\alpha(t^\alpha A)\|_{\mathcal{L}(X^1, X^{1+\theta})} \leq M$ follows from (10). For the remaining part it suffices to observe that the operators $t^\alpha E_\alpha(t^\alpha A) : X^1 \rightarrow X^{1+\alpha}$ are uniformly bounded in t , that

$$\lim_{t \rightarrow 0^+} \|t^\alpha E_\alpha(t^\alpha A)x\|_{X^{1+\alpha}} = 0,$$

for $x \in X^{1+\alpha}$, and that X^α is a dense subset of X^1 . \square

Let $\mathbf{B} : (0, \infty) \times (0, \infty) \rightarrow (0, \infty)$ be the beta function given by

$$\mathbf{B}(a, b) = \int_0^1 (1-x)^{a-1} x^{b-1} dx.$$

For $\zeta \geq 0$, define

$$\mathbf{B}_\epsilon^\theta(\zeta) = \sup_{0 \leq \eta \leq \theta} \{\mathbf{B}(\alpha(\gamma(\epsilon) - \eta), 1 - \zeta), \mathbf{B}(\alpha(\gamma(\epsilon) - \eta), 1 - \alpha\rho\epsilon)\}.$$

Lemma 26. *With the above notation, let $f \in \mathcal{F}(\nu(\cdot))$. If $u \in C((0, \tau]; X^{1+\epsilon})$, then, for all $0 \leq \theta < \gamma(\epsilon)$, we have that*

$$\begin{aligned} & t^{\alpha\theta} \left\| \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s)) ds \right\|_{X^{1+\theta}} \\ & \leq M c \mathbf{B}_\epsilon^\theta(\xi) (\nu(t) t^{\alpha\gamma(\epsilon)-\xi} + \lambda(t)^\rho t^{\alpha(\gamma(\epsilon)-\rho\epsilon)}) \end{aligned}$$

for $0 < t \leq \tau$, where $\lambda(t) = \sup_{s \in (0, t]} \{s^{\alpha\epsilon} \|u(s)\|_{X^{1+\epsilon}}\}$.

Proof. Indeed, we have that

$$\begin{aligned}
& t^{\alpha\theta} \int_0^t \|(t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s))\|_{X^{1+\theta}} ds \\
& \leq Mt^{\alpha\theta} \int_0^t (t-s)^{\alpha(\gamma(\epsilon)-\theta)-1} c(\nu(s) s^{-\xi} + \|u(s)\|_{X^{1+\epsilon}}^\rho) ds \\
& \leq Mc\nu(t) t^{\alpha\theta} \int_0^t (t-s)^{\alpha(\gamma(\epsilon)-\theta)-1} s^{-\xi} ds \\
& \quad + Mct^{\alpha\theta} \int_0^t (t-s)^{\alpha(\gamma(\epsilon)-\theta)-1} s^{-\alpha\rho\epsilon} (s^{\alpha\epsilon} \|u(s)\|_{X^{1+\epsilon}})^\rho ds \\
& \leq Mc\nu(t) t^{\alpha\theta-\xi+\alpha(\gamma(\epsilon)-\theta)} \int_0^1 (1-s)^{\alpha(\gamma(\epsilon)-\theta)-1} s^{-\xi} ds \\
& \quad + Mc\lambda(t)^\rho t^{\alpha\theta-\alpha\rho\epsilon+\alpha(\gamma(\epsilon)-\theta)} \int_0^1 (t-s)^{\alpha(\gamma(\epsilon)-\theta)-1} s^{-\alpha\rho\epsilon} ds \\
& \leq Mc\mathbf{B}_\epsilon^\theta(\xi) (\nu(t) t^{\alpha\gamma(\epsilon)-\xi} + \lambda(t)^\rho t^{\alpha(\gamma(\epsilon)-\rho\epsilon)}),
\end{aligned}$$

which concludes the proof. \square

Lemma 27. *With the above notation, let $f \in \mathcal{F}(\nu(\cdot))$ and consider $u, v \in C((0, \tau]; X^{1+\epsilon})$ such that $t^{\alpha\epsilon} \|u(t)\|_{X^{1+\epsilon}} \leq \mu$ and $t^{\alpha\epsilon} \|v(t)\|_{X^{1+\epsilon}} \leq \mu$ for some $\mu > 0$. Then, for all $0 \leq \theta < \gamma(\epsilon) < 1$, we have*

$$\begin{aligned}
& t^{\alpha\theta} \left\| \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) (f(s, u(s)) - f(s, v(s))) ds \right\|_{X^{1+\theta}} \\
& \leq Mc\mathbf{B}_\epsilon^\theta(\zeta + \alpha\epsilon) [\nu(t) t^{\alpha(\gamma(\epsilon)-\theta)-\zeta} + 2\mu^{\rho-1} t^{\alpha(\gamma(\epsilon)+\epsilon-\theta-\rho\epsilon)}] \sup_{0 < s \leq \tau} s^{\alpha\epsilon} \|u(s) - v(s)\|_{X^{1+\epsilon}}.
\end{aligned}$$

Proof. It follows from the ϵ -regularity property of f that

$$\begin{aligned}
& t^{\alpha\theta} \int_0^t \|(t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) (f(s, u(s)) - f(s, v(s)))\|_{X^{1+\theta}} ds \\
& \leq Mt^{\alpha\theta} \int_0^t (t-s)^{\alpha(\gamma(\epsilon)-\theta)-1} \|f(s, u(s)) - f(s, v(s))\|_{X^{\gamma(\epsilon)}} ds \\
& \leq Mct^{\alpha\theta} \int_0^t (t-s)^{\alpha(\gamma(\epsilon)-\theta)-1} \|u(s) - v(s)\|_{X^{1+\epsilon}} (\|u(s)\|_{X^{1+\epsilon}}^{\rho-1} + \|v(s)\|_{X^{1+\epsilon}}^{\rho-1} + \nu(s) s^{-\zeta}) ds \\
& \leq Mc\nu(t) t^{\alpha\theta} \int_0^t (t-s)^{\alpha(\gamma(\epsilon)-\theta)-1} s^{-\zeta-\alpha\epsilon} s^{\alpha\epsilon} \|u(s) - v(s)\|_{X^{1+\epsilon}} ds \\
& \quad + Mct^{\alpha\theta} \int_0^t (t-s)^{\alpha(\gamma(\epsilon)-\theta)-1} s^{-\alpha\rho\epsilon} [(s^{\alpha\epsilon} \|u(s)\|_{X^{1+\epsilon}})^{\rho-1} + (s^{\alpha\epsilon} \|v(s)\|_{X^{1+\epsilon}})^{\rho-1}] \\
& \quad \quad \times s^{\alpha\epsilon} \|u(s) - v(s)\|_{X^{1+\epsilon}} ds \\
& \leq [\nu(t) t^{\alpha(\gamma(\epsilon)-\theta)-\zeta} + 2\mu^{\rho-1} t^{\alpha(\gamma(\epsilon)+\epsilon-\theta-\rho\epsilon)}] Mc\mathbf{B}_\epsilon^\theta(\zeta + \alpha\epsilon) \sup_{0 \leq s \leq \tau} s^{\alpha\epsilon} \|u(s) - v(s)\|_{X^{1+\epsilon}}.
\end{aligned}$$

\square

Now we may prove the main result of this section.

Proof of Theorem 7. Define μ by

$$Mc\mathbf{B}_\epsilon^\epsilon \mu^{\rho-1} = \frac{1}{8},$$

where $\mathbf{B}_\epsilon^\theta := \max\{\mathbf{B}_\epsilon^\theta(\xi), \mathbf{B}_\epsilon^\theta(\zeta + \alpha\epsilon)\}$ and choose $r = r(\mu, M) > 0$ such that

$$r = \frac{\mu}{4M} = \frac{1}{4M(8cM\mathbf{B}_\epsilon^\epsilon)^{\frac{1}{\rho-1}}}.$$

For v_0 fixed, choose $\tau_0 \in (0, 1]$ such that $\nu(t) < \delta$ for all $t \in (0, \tau_0]$,

$$\|t^{\alpha\epsilon} E_\alpha(t^\alpha A)v_0\|_{X^{1+\epsilon}} \leq \frac{\mu}{2} \quad \text{if } 0 \leq t \leq \tau_0,$$

and

$$Mc\delta\mathbf{B}_\epsilon^\epsilon = \min\left\{\frac{\mu}{8}, \frac{1}{4}\right\}.$$

Consider

$$K(\tau_0) = \left\{ u \in C((0, \tau_0]; X^{1+\epsilon}) : \sup_{t \in (0, \tau_0]} t^{\alpha\epsilon} \|u(t)\|_{X^{1+\epsilon}} \leq \mu \right\}$$

with norm

$$\|u\|_{K(\tau_0)} = \sup_{t \in (0, \tau_0]} t^{\alpha\epsilon} \|u(t)\|_{X^{1+\epsilon}}.$$

Suppose that $u_0 \in X^1$ with $\|u_0 - v_0\|_{X^1} < r$ and define on $K(\tau_0)$ the map

$$Tu(t) = E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s)) ds.$$

Our purpose is to show that for any $u_0 \in B_{X^1}(v_0, r)$, $K(\tau_0)$ is T -invariant and T is a contraction.

Initially, let us prove that $T : K(\tau_0) \rightarrow K(\tau_0)$ is well defined.

Claim 1: if $u \in K(\tau_0)$, then $Tu \in C((0, \tau_0]; X^{1+\theta})$ for all $\theta \in [0, \gamma(\epsilon))$.

For this, let $t_1, t_2 \in (0, \tau_0]$, $t_1 > t_2$; then, for every $0 \leq \theta < \gamma(\epsilon)$, we have that

$$\begin{aligned} & \|Tu(t_1) - Tu(t_2)\|_{X^{1+\theta}} \\ & \leq \|(E_\alpha(t_1^\alpha A) - E_\alpha(t_2^\alpha A))u_0\|_{X^{1+\theta}} \\ & \quad + \left\| \int_0^{t_2} ((t_1-s)^{\alpha-1} E_{\alpha,\alpha}((t_2-s)^\alpha A) - (t_1-s)^{\alpha-2} E_{\alpha,\alpha}((t_2-s)^\alpha A)) f(s, u(s)) ds \right\|_{X^{1+\theta}} \\ & \quad + \left\| \int_{t_2}^{t_1} (t_1-s)^{\alpha-1} E_{\alpha,\alpha}((t_1-s)^\alpha A) f(s, u(s)) ds \right\|_{X^{1+\theta}}. \end{aligned}$$

The first term trivially goes to 0 as $t_1 \rightarrow t_2^+$. A similar procedure to Theorem 4 shows that the second term also goes to 0 as $t_1 \rightarrow t_2^+$. Let us consider the third term. We

have

$$\begin{aligned}
& \left\| \int_{t_2}^{t_1} (t_1 - s)^{\alpha-1} E_{\alpha,\alpha}((t_1 - s)^\alpha A) f(s, u(s)) ds \right\|_{X^{1+\theta}} \\
& \leq M \int_{t_2}^{t_1} (t_1 - s)^{\alpha(\gamma(\epsilon)-\theta)-1} \|f(s, u(s))\|_{X^{\gamma(\epsilon)}} ds \\
& \leq Mc \int_{t_2}^{t_1} (t_1 - s)^{\alpha(\gamma(\epsilon)-\theta)-1} (\nu(s) s^{-\xi} + \|u(s)\|_{X^{1+\epsilon}}^{\rho-1}) ds \\
& \leq Mc\delta \int_{t_2}^{t_1} (t_1 - s)^{\alpha(\gamma(\epsilon)-\theta)-1} s^{-\xi} ds + Mc \int_{t_2}^{t_1} (t_1 - s)^{\alpha(\gamma(\epsilon)-\theta)-1} \|u(s)\|_{X^{1+\epsilon}} ds \\
& \leq Mc\delta t_1^{\alpha(\gamma(\epsilon)-\theta)-\xi} \int_{t_2/t_1}^1 (1-s)^{\alpha(\gamma(\epsilon)-\theta)-1} s^{-\xi} ds \\
& \quad + Mc\mu^\rho t^{\alpha(\gamma(\epsilon)-\theta)-\rho\epsilon} \int_{t_2/t_1}^1 (1-s)^{\alpha(\gamma(\epsilon)-\theta)-1} s^{-\alpha\rho\epsilon} ds,
\end{aligned}$$

which goes to 0 as $t_1 \rightarrow t_2^+$.

The case $t_1 < t_2$ is analogous.

Claim 2: $t^{\alpha\epsilon} \|Tu(t)\|_{X^{1+\epsilon}} \leq \mu$ for all $t \in (0, \tau_0]$.

Indeed, we may estimate as follows.

$$\begin{aligned}
& t^{\alpha\epsilon} \|Tu(t)\|_{X^{1+\epsilon}} \\
& \leq \|t^{\alpha\epsilon} E_\alpha(At^\alpha)u_0\|_{X^{1+\epsilon}} + Mt^{\alpha\epsilon} \int_0^t (t-s)^{\alpha(\gamma(\epsilon)-\epsilon)-1} \|f(s, u(s))\|_{X^{\gamma(\epsilon)}} ds \\
& \leq \|t^{\alpha\epsilon} E_\alpha(At^\alpha)(u_0 \mp v_0)\|_{X^{1+\epsilon}} + Mct^{\alpha\epsilon} \int_0^t (t-s)^{\alpha(\gamma(\epsilon)-\epsilon)-1} (\nu(s) s^{-\xi} + \|u(s)\|_{X^{1+\epsilon}}^\rho) ds \\
& \leq Mr + \|t^{\alpha\epsilon} E_\alpha(At^\alpha)v_0\|_{X^{1+\epsilon}} + Mc\delta t^{\alpha\gamma(\epsilon)-\xi} \int_0^1 (1-s)^{\alpha(\gamma(\epsilon)-\epsilon)-1} s^{-\xi} ds \\
& \quad + Mc\mu^\rho t^{\alpha(\gamma(\epsilon)-\rho\epsilon)} \int_0^1 (1-s)^{\alpha(\gamma(\epsilon)-\epsilon)-1} s^{-\alpha\rho\epsilon} ds \\
& \leq Mr + \|t^{\alpha\epsilon} E_\alpha(At^\alpha)v_0\|_{X^{1+\epsilon}} + Mc\delta \mathbf{B}_\epsilon^\epsilon + Mc\mu^\rho \mathbf{B}_\epsilon^\epsilon \leq \mu.
\end{aligned}$$

This shows that $T(K(\tau_0)) \subset K(\tau_0)$.

Finally, by taking $\theta = \epsilon$, it follows from Lemma 27 that T is a strict contraction in $K(\tau_0)$ and that

$$\|Tu(t) - Tv(t)\|_{K(\tau_0)} \leq \frac{1}{2} \|u - v\|_{K(\tau_0)}.$$

By the Banach contraction principle we have that T has a unique fixed point in $K(\tau_0)$, which will be denoted by $u(\cdot, u_0)$. It is defined for $\|u_0 - v_0\|_{X^1} < r$, $0 \leq t \leq \tau_0$.

Observe that $u(\cdot, u_0) \in C((0, \tau_0]; X^{1+\theta})$ for all $0 \leq \theta < \gamma(\epsilon)$. Furthermore, we will prove that

Claim 3: $t^{\alpha\theta} \|u(t, u_0)\|_{X^{1+\theta}} \rightarrow 0$ as $t \rightarrow 0$, for all $0 < \theta < \gamma(\epsilon)$.

Indeed, from Lemma 26 we have that

$$\begin{aligned} & t^{\alpha\theta} \|u(t, u_0)\|_{X^{1+\theta}} \\ & \leq t^{\alpha\theta} \|E_\alpha(t^\alpha A)u_0\|_{X^{1+\theta}} + t^{\alpha\theta} \left\| \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s, u_0)) ds \right\|_{X^{1+\theta}} \\ & \leq t^{\alpha\theta} \|E_\alpha(t^\alpha A)u_0\|_{X^{1+\theta}} + Mc\mathbf{B}_\epsilon^\theta(\xi)\nu(t) + Mc\mathbf{B}_\epsilon^\theta(\xi)\mu^{\rho-1} \sup_{0 < s \leq t} \{s^{\alpha\epsilon} \|u(s, u_0)\|_{X^{1+\epsilon}}\}. \end{aligned}$$

Therefore, if $\theta = \epsilon$ we deduce

$$t^{\alpha\epsilon} \|u(t, u_0)\|_{X^{1+\epsilon}} \leq t^{\alpha\epsilon} \|E_\alpha(t^\alpha A)u_0\|_{X^{1+\epsilon}} + Mc\mathbf{B}_\epsilon^\epsilon \nu(t) + \frac{1}{8} \sup_{0 < s \leq t} \{s^{\alpha\epsilon} \|u(s, u_0)\|_{X^{1+\epsilon}}\},$$

from which we obtain

$$\sup_{0 < s \leq t} \{s^{\alpha\epsilon} \|u(s, u_0)\|_{X^{1+\epsilon}}\} \leq \frac{8}{7} \left(\sup_{0 < s \leq t} \{s^{\alpha\epsilon} \|E_\alpha(s^\alpha A)u_0\|_{X^{1+\epsilon}}\} + Mc\mathbf{B}_\epsilon^\epsilon \nu(t) \right) \rightarrow 0 \text{ as } t \rightarrow 0.$$

From above we also conclude that Claim 3 holds.

Claim 4: $\lim_{t \rightarrow 0^+} \|u(t, u_0) - u_0\|_{X^1} = 0$.

For this observe that from Lemma 26,

$$\begin{aligned} & \|u(t, u_0) - u_0\|_{X^1} \\ & \leq \|E_\alpha(t^\alpha A)u_0 - u_0\|_{X^1} + \int_0^t \|(t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s, u_0))\|_{X^1} ds \\ & \leq \|E_\alpha(t^\alpha A)u_0 - u_0\|_{X^1} + Mc\mathbf{B}_\epsilon^0(\xi)(\nu(t) + [\sup_{0 < s \leq t} \{s^{\alpha\epsilon} \|u(s, u_0)\|_{X^{1+\epsilon}}\}]^\rho). \end{aligned}$$

Therefore, the claim follows and we have that $u(t, u_0)$ is an ϵ -regular mild solution starting at u_0 and it is the unique ϵ -regular mild solution, starting at u_0 , in the set $K(\tau_0)$. We will call it the K -solution starting at u_0 .

Moreover, if $u_0, w_0 \in B_{X^1}(v_0, r)$, it follows from Lemma 27 and the choice of τ_0 that

$$\begin{aligned} & t^{\alpha\theta} \|u(t, u_0) - u(t, w_0)\|_{X^{1+\theta}} \\ & \leq t^{\alpha\theta} \|E_\alpha(t^\alpha A)(u_0 - w_0)\|_{X^{1+\theta}} \\ & \quad + t^{\alpha\theta} \left\| \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) (f(s, u(s, u_0)) - f(s, u(s, w_0))) ds \right\|_{X^{1+\theta}} \\ & \leq M \|u_0 - w_0\|_{X^1} + Mc\mathbf{B}_\epsilon^\theta(\zeta + \alpha\epsilon)\nu(t) t^{\alpha(\gamma(\epsilon)-\theta)-\zeta} \sup_{0 < s \leq \tau_0} \{s^{\alpha\theta} \|u(s, u_0) - u(s, w_0)\|_{X^{1+\epsilon}}\} \\ & \quad + Mc\mathbf{B}_\epsilon^\theta(\zeta + \alpha\epsilon) 2\mu^{\rho-1} t^{\alpha(\gamma(\epsilon)+\epsilon-\theta-\rho\epsilon)} \sup_{0 < s \leq \tau_0} \{s^{\alpha\theta} \|u(s, u_0) - u(s, w_0)\|_{X^{1+\epsilon}}\}. \quad (11) \end{aligned}$$

For $\theta = \epsilon$ we obtain

$$t^{\alpha\epsilon} \|u(t, u_0) - u(t, w_0)\|_{X^{1+\epsilon}} \leq M \|u_0 - w_0\|_{X^1} + \frac{1}{2} \sup_{0 < s \leq \tau_0} \{s^{\alpha\epsilon} \|u(t, u_0) - u(t, w_0)\|_{X^{1+\epsilon}}\},$$

which implies

$$t^{\alpha\epsilon} \|u(t, u_0) - u(t, w_0)\|_{X^{1+\epsilon}} \leq 2M \|u_0 - w_0\|_{X^1}.$$

For $0 \leq \theta \leq \theta_0 < \gamma(\epsilon) + \epsilon - \rho\epsilon$ we have from (11) that

$$\begin{aligned} & t^{\alpha\theta} \|u(t, u_0) - u(t, w_0)\|_{X^{1+\theta}} \\ & \leq M \|u_0 - w_0\|_{X^1} + 2M^2 c\mathbf{B}_\epsilon^\theta (\zeta + \alpha\epsilon) [\nu(t) t^{\alpha(\gamma(\epsilon)-\theta)-\zeta} + 2\mu^{\rho-1} t^{\alpha(\gamma(\epsilon)+\epsilon-\theta-\rho\epsilon)}] \|u_0 - w_0\|_{X^1} \\ & \leq C(\theta_0) \|u_0 - w_0\|_{X^1}, \end{aligned}$$

where

$$C(\theta_0) = M \left(1 + 2M \sup_{t \in [0, \tau_0], 0 \leq \theta \leq \theta_0} \{c\mathbf{B}_\epsilon^\theta (\zeta + \alpha\epsilon) [\nu(t) t^{\alpha(\gamma(\epsilon)-\epsilon)-\zeta} + 2\mu^{\rho-1} t^{\alpha(\gamma(\epsilon)+\epsilon-\theta-\rho\epsilon)}] \} \right).$$

This concludes the proof of the theorem. \square

We have the following two consequences of Theorem 7. The first one is that if f is independent of time, we obtain the same result. Observe that this is not just by an application of the theorem, since now we do not have a time-dependent class \mathcal{F} any more. However, it is not difficult to readapt its proof.

Corollary 28. *With the above notation, assume that f is independent of time and it is an ϵ -regular map, for some $\epsilon > 0$, relative to the pair (X^1, X^0) . Then if $v_0 \in X^1$, there exist $r = r(v_0) > 0$ and $\tau_0 = \tau_0(v_0) > 0$ such that for every $u_0 \in B_{X^1}(v_0, r)$ there is a continuous function $u(\cdot, u_0) : [0, \tau_0] \rightarrow X^1$ with $u(0) = u_0$, which is an ϵ -regular mild solution to the problem (3) starting at u_0 . Furthermore, this solution satisfies*

$$u \in C((0, \tau_0]; X^{1+\theta}), \quad 0 \leq \theta < \gamma(\epsilon),$$

$$\lim_{t \rightarrow 0^+} t^{\alpha\theta} \|u(t, u_0)\|_{X^{1+\theta}} = 0, \quad 0 < \theta < \gamma(\epsilon).$$

Moreover, for each $\theta_0 < \gamma(\epsilon) + \epsilon - \rho\epsilon$ there exists a constant $C > 0$ such that if $u_0, w_0 \in B_{X^1}(v_0, r)$, then

$$t^{\alpha\theta} \|u(t, u_0) - u(t, w_0)\|_{X^{1+\theta}} \leq C \|u_0 - w_0\|_{X^1} \quad \forall t \in [0, \tau_0], 0 \leq \theta \leq \theta_0 < \gamma(\epsilon) + \epsilon - \rho\epsilon.$$

The second consequence of Theorem 7 is the following

Corollary 29. *Let f be as in Theorem 7 and K a relatively compact set in X^1 , then there exists $\tau_0 = \tau_0(K)$ such that the ϵ -regular solution starting at u_0 exists in the time interval $[0, \tau_0]$ for any $u_0 \in K$.*

5. COMPARISON AND POSITIVITY RESULTS

5.1. Linear equations. The following result establishes the equivalence between the positivity of the resolvent operator of A and the positivity of the Mittag-Leffler families.

Proposition 30. *Let (X, \leq_X) be an ordered Banach space and consider $\alpha \in (0, 1)$ and $A : D(A) \subset X \rightarrow X$ a sectorial operator. Suppose that there exists $\lambda_0 > 0$ such that $(\lambda^\alpha - A)^{-1}$ is increasing for any $\lambda > \lambda_0$. Then, $E_\alpha(t^\alpha A)$ is increasing for all $t \geq 0$. Conversely, if $X \ni u_0 \mapsto E_\alpha(t^\alpha A)u_0 \in C([0, \infty); X)$ is increasing for all $t \geq 0$, then $(\lambda^\alpha - A)^{-1}$ is increasing for any $\lambda > 0$.*

Proof. The case $t = 0$ is easily verified.

To prove the case $t > 0$, consider $0 \leq_X u_0 \in X$. From Proposition 16, write

$$E_\alpha(t^\alpha A)u_0 = \lim_{n \rightarrow \infty} \frac{1}{n!} \sum_{k=1}^{n+1} b_{k,n+1}^\alpha \left(\left(\frac{n}{t} \right)^\alpha \left(\left(\frac{n}{t} \right)^\alpha - A \right)^{-1} \right)^k u_0, \quad t > 0. \quad (12)$$

If $n \in \mathbb{N}$ is such that $n/t > \lambda_0$, then

$$0 \leq \left(\left(\frac{n}{t} \right)^\alpha - A \right)^{-1} u_0.$$

Since the values $b_{k,n+1}^\alpha$, given by (9), are positive real numbers and the positive cone is closed, it follows from (12) that $0 \leq_X E_\alpha(t^\alpha A)u_0$.

Conversely, since the integral is a positive operator and

$$\lambda^{\alpha-1}(\lambda^\alpha - A)^{-1} = \int_0^\infty e^{-\lambda t} E_\alpha(t^\alpha A) dt \quad \forall \lambda > 0,$$

we conclude that $(\lambda^\alpha - A)^{-1}$ is a positive operator whenever $\lambda > 0$. \square

Remark 31. (i) *Proposition 30 is equivalent to the following result: if $0 \leq_X u_0$, then the solution of the linear homogeneous problem*

$$\begin{cases} cD_t^\alpha u(t) = Au(t), \\ u(0) = u_0 \in X, \end{cases}$$

verifies $0 \leq_X u(t)$ for all $t \geq 0$. Equivalently, if $u_1 \leq_X u_0$, then $v(t) \leq_X u(t)$ for all $t \geq 0$, where u and v are the corresponding solutions to the problem with initial conditions $u(0) = u_0$ and $v(0) = u_1$, respectively.

(ii) *The assumption on the operator A is equivalent to the following positivity result for elliptic problems: for any $\lambda > \lambda_0$ and for any f with $0 \leq_X f$, the solution to the problem $\lambda^\alpha u - Au = f$ satisfies $0 \leq_X u$.*

A similar result may be stated in terms of the family $\{E_{\alpha,\alpha}(t^\alpha A)\}_{t \geq 0}$.

Proposition 32. *Let (X, \leq_X) be an ordered Banach space and consider $\alpha \in (0, 1)$ and $A : D(A) \subset X \rightarrow X$ a sectorial operator. Suppose that there exists $\lambda_0 > 0$ such that $(\lambda^\alpha - A)^{-1}$ is increasing for any $\lambda > \lambda_0$. Then, $E_{\alpha,\alpha}(t^\alpha A)$ is increasing for all $t \geq 0$. Conversely, if $X \ni u_0 \mapsto E_{\alpha,\alpha}(t^\alpha A)u_0 \in C([0, \infty); X)$ is increasing for all $t \geq 0$, then $(\lambda^\alpha - A)^{-1}$ is increasing for any $\lambda > 0$.*

Proof. It suffices to observe that if $u_0 \in X$, then

$$t^{\alpha-1} E_{\alpha,\alpha}(t^\alpha A)u_0 = \lim_{n \rightarrow \infty} \frac{1}{n!} \sum_{k=1}^n \alpha k b_{k,n}^\alpha \left(\frac{n}{t} \right)^{\alpha k+1} \left(\left(\frac{n}{t} \right)^\alpha - A \right)^{-(k+1)} u_0, \quad t \geq 0.$$

Now, we may proceed as in the proof of Proposition 30. Conversely, we have that

$$(\lambda^\alpha - A)^{-1} = \int_0^\infty e^{-\lambda t} t^{\alpha-1} E_{\alpha,\alpha}(t^\alpha A) dt,$$

for any $\lambda > 0$, which concludes the proof. \square

Corollary 33. *Let (X, \leq_X) be an ordered Banach space and consider $\alpha \in (0, 1)$ and $A : D(A) \subset X \rightarrow X$ a sectorial operator. Suppose that there exists $\lambda_0 > 0$ such that $(\lambda^\alpha - A)^{-1}$ is increasing for any $\lambda > \lambda_0$. Let $u_f(t, u_0)$ be the local mild solution in $[0, \tau]$, for some $\tau > 0$, to the problem*

$$\begin{cases} cD_t^\alpha u(t) = Au(t) + f(t), & t > 0, \\ u(0) = u_0 \in X, \end{cases}$$

and suppose that $f \in L^1(0, \tau; X)$. Assume that $u_1 \leq_X u_0$ and $f_1(t) \leq_X f_0(t)$ a.e. $t \in (0, \tau)$. Then $u_{f_1}(t, u_0) \leq_X u_{f_0}(t, u_0)$ for all $t \in [0, \tau]$. In particular, if $0 \leq_X u_0$ and $0 \leq_X f(t)$ a.e. $t \in (0, \tau)$, then $0 \leq_X u_f(t, u_0)$ for all $t \in [0, \tau]$.

Proof. Observe that the corresponding solutions ($i = 0, 1$) are given by

$$u_i(t) = E_\alpha(t^\alpha A)u_i + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f_i(s) ds.$$

Since $E_\alpha(t^\alpha A)u_1 \leq_X E_\alpha(t^\alpha A)u_0$ and, for any $0 < s < t < \tau$,

$$E_{\alpha,\alpha}((t-s)^{\alpha-1} A) f_1(s) \leq_X E_{\alpha,\alpha}((t-s)^{\alpha-1} A) f_0(s),$$

the result follows from the fact that the integral is a positive operator. \square

5.2. Semilinear equations. In this paragraph we deal with comparison results and positivity for the semilinear problem (1) where $f : [0, \tau] \times X \rightarrow X$ a continuous function, locally lipschitz in the second variable (uniformly with respect to the first variable), and, eventually, bounded.

Proof of Theorem 9. Let T be the operator defined by

$$T(u)(t) = E_\alpha(t^\alpha A)u_0 + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s)) ds.$$

We proved in Theorem 2 that if $\tau \in [0, t_0]$ and $\beta > 0$ are small enough, then T is a contraction in $K = \{u \in C([0, \tau]; X) : \|u(t) - u_0\| \leq \beta\}$ and it has a unique fixed point. Consider $y_0(s) = u_0$, $s \in [0, \tau]$. Consequently, $0 \leq_X y_0(s)$ and $0 \leq_X f(s, y_0(s))$ a.e. $(0, \tau)$. Hence, $y_1(t) = T(y_0(t))$, $t \in [0, \tau]$ satisfies $0 \leq_X y_1(t)$. Iterating, we obtain

$$0 \leq_X y_n(t) = T(y_{n-1}(t)), \quad t \in [0, \tau].$$

Since $\{y_n\}_{n \in \mathbb{N}}$ converges to u in $C([0, \tau]; X)$ we have that $0 \leq_X u(t)$ for all $t \in [0, \tau]$. Now, combining this with some continuation arguments from Section 3, we may conclude that $0 \leq_X u(t)$ for all $t \in [0, t_0]$. \square

Proof of Theorem 10. (i) For $i = 0, 1$, we know that $u_i(t) = u_f(t, u_i)$, is a fixed point of the operator

$$T(u)(t) = E_\alpha(t^\alpha A)u_i + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}((t-s)^\alpha A) f(s, u(s)) ds$$

in $[0, t_i]$.

Consider initially the function $y_i^0(t) = u_i$, $t \in [0, t_i]$, $i = 0, 1$. Iterating, we have $y_i^n(t) = T(y_i^{n-1})(t)$ and this sequence converges to $u_i(t)$ in $C([0, t_i]; X)$. Furthermore, we have $y_1^0(t) \leq_X y_0^0(t)$ a.e. $t \in (0, t^*)$ and hence

$$f(t, y_1^0(t)) \leq_X f(t, y_0^0(t)) \quad \text{a.e. } t \in (0, t^*).$$

Then, $y_1^1(t) \leq_X y_0^1(t)$ in $[0, t^*]$. Iterating, we obtain $y_1^n(t) \leq_X y_0^n(t)$ and consequently $u_1(t) \leq_X u_0(t)$ in $[0, t^*]$.

Statements (ii) and (iii) can be proved analogously. \square

Acknowledgments. This paper was started during a P.M.-R.'s visit to the Instituto de Ciências Matemáticas e de Computação (ICMC) at the Universidade de São Paulo, in São Carlos (Brazil), and finished while B.A. and P.M.C.-N. were visiting the Departamento de Ecuaciones Diferenciales y Análisis Numérico (EDAN) at the Universidad de Sevilla in 2011-12. The authors would like to thanks to the people of these two institutions for their hospitality during their respective stays, and in particular P.M.-R. would also like to thank to Ministerio de Educación–DGPU (Spain) for its support, under project PHB2010-0002-PC.

REFERENCES

- [1] H. Amann, Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems, 9–126, in: Schmeisser/Triebel: *Function Spaces, Differential Operators and Nonlinear Analysis*, **133**, Teubner, Stuttgart, 1993.
- [2] J. M. Arrieta and A. N. Carvalho, Abstract parabolic problems with critical nonlinearities and applications to Navier-Stokes and Heat equations, *Trans. Amer. Math. Soc.* **352** (2000), 285–310.
- [3] E. Bazhikova, *Fractional Evolution Equations in Banach Spaces*, Ph.D. Thesis, Eindhoven University of Technology, 2001.
- [4] M. N. Berberan-Santos, Properties of the Mittag-Leffler relaxation function, *J. Math. Chem.* **38** (2005), 629–635.
- [5] H. Brezis and T. Cazenave, A nonlinear heat equation with singular initial data, *J. Anal. Math.* **68** (1996), 277–304.
- [6] A. N. Carvalho and M. R. T. Primo, Spatial homogeneity in parabolic problems with nonlinear boundary conditions, *Commun. Pure Appl. Anal.* **3** (2004), 637–651.
- [7] S.D. Eidelman and A. N. Kochubei, Cauchy problem for fractional diffusion equations, *J. Differential Equations* **199** (2004), 211–255.
- [8] L. Galeone and R. Garrappa, Explicit methods for fractional differential equations and their stability properties, *J. Comput. Appl. Math.* **228** (2009), 548–560.
- [9] D. Henry, *Geometric Theory of Semilinear Parabolic Equations*, Lecture Notes in Mathematics **840**, Springer-Verlag, Berlin, 1981.
- [10] H. J. Haubold, A. M. Mathai, and R. K. Saxena, Mittag-Leffler Functions and Their Applications, *J. Appl. Math.* **2011** (2011), 51 pages.
- [11] A. A. Kilbas; H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, 2006.
- [12] A. A. Kilbas, T. Pierantozzi, J. J. Trujillo, and L. Vazquez, On the solution of fractional evolution equations, *J. Phys. A: Math. Gen.* **37** (2004), 3271–3283.
- [13] F. Mainardi, *Fractional calculus and waves in linear viscoelasticity*, Imperial College Press, London, 2010.
- [14] K. B. Oldham and J. Spanier, *The Fractional Calculus*, Academic Press, London, 1974.
- [15] I. V. Ostrovskii and I. N. Peresolkova, Nonasymptotic results on distribution of zeros of the function $E_\rho(z, \mu)$, *Anal. Math.* **23** (1997), 283–296.

- [16] J. Peng and K. Li A note on property of the Mittag-Leffler function, *J. Math. Anal. Appl.* **370** (2010), 635–638.
- [17] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [18] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives. Theory and Applications*, Gordon & Breach Sci. Publishers, Yverdon, 1993.
- [19] W. R. Schneider, Fractional diffusion, *Lecture Notes in Phys.* **355** (1990), 276–286.
- [20] W. R. Schneider and W. Wyss, Fractional Diffusion and wave equations, *J. Math. Phys.* **30** (1989), 134–144.
- [21] R. N. Wang, D. H. Chen, and T. J. Xiao, Abstract fractional Cauchy problems with almost sectorial operators *J. Differential Equations*, **252** (2012), 202–235.
- [22] R. N. Wang, T. J. Xiao and J. Liang, A note on the fractional Cauchy problems with nonlocal initial conditions, *Appl. Math. Lett.* **24** (2011), 1435–1442.
- [23] F. B. Weissler, Semilinear evolution equations in Banach spaces, *J. Funct. Anal.* **32** (1979), 277–296.
- [24] F. B. Weissler, Local existence and nonexistence for semilinear parabolic equations in L^p , *Indiana Univ. Math. J.* **29** (1980), 79–102.
- [25] W. Wyss, The fractional diffusion equation, *J. Math. Phys.* **27** (1986), 2782–2785.
- [26] K. Yosida, *Functional Analysis*, Springer-Verlag, New York, 1978.

(B. de Andrade, A. N. Carvalho, and P. M. Carvalho-Neto) INSTITUTO DE CIÊNCIAS MATEMÁTICAS E DE COMPUTAÇÃO, UNIVERSIDADE DE SÃO PAULO-CAMPUS DE SÃO CARLOS, CAIXA POSTAL 668, 13560-970 SÃO CARLOS SP, BRAZIL

E-mail address: bruno00luis@gmail.com, andcarva@icmc.usp.br, pmat@icmc.usp.br

(P. Marín Rubio) DEPARTAMENTO DE ECUACIONES DIFERENCIALES Y ANÁLISIS NUMÉRICO, UNIVERSIDAD DE SEVILLA, APDO. DE CORREOS 1160, 41080-SEVILLA, SPAIN

E-mail address: pmr@us.es