

UNIFORMLY SUMMING SETS OF OPERATORS ON SPACES OF CONTINUOUS FUNCTIONS

J. M. DELGADO and CÁNDIDO PIÑEIRO

Received 30 March 2004

Let X and Y be Banach spaces. A set \mathcal{M} of 1-summing operators from X into Y is said to be *uniformly summing* if the following holds: given a weakly 1-summing sequence (x_n) in X , the series $\sum_n \|Tx_n\|$ is uniformly convergent in $T \in \mathcal{M}$. We study some general properties and obtain a characterization of these sets when \mathcal{M} is a set of operators defined on spaces of continuous functions.

2000 Mathematics Subject Classification: 47B38, 47B10.

1. Introduction. Throughout this paper, X and Y will be Banach spaces. If X is a Banach space, $B_X = \{x \in X : \|x\| \leq 1\}$ will denote its closed unit ball and X^* will be the topological dual of X . Given a real number $p \in [1, \infty)$, a (linear) operator $T : X \rightarrow Y$ is said to be *p-summing* if there exists a constant $C > 0$ such that

$$\left(\sum_{i=1}^n \|Tx_i\|^p \right)^{1/p} \leq C \cdot \sup \left\{ \left(\sum_{i=1}^n |\langle x^*, x_i \rangle|^p \right)^{1/p} : x^* \in B_{X^*} \right\} \quad (1.1)$$

for every finite set $\{x_1, \dots, x_n\} \subset X$. The least C for which the above inequality always holds is denoted by $\pi_p(T)$ (the *p-summing norm* of T). The linear space of all *p*-summing operators from X into Y is denoted by $\Pi_p(X, Y)$ which is a Banach space endowed with the *p*-summing norm.

As usual, $\ell_w^p(X)$ will be the Banach space of weakly *p*-summable sequences in X , that is, the sequences $(x_n) \subset X$ satisfying $\sum_n |\langle x^*, x_n \rangle|^p < \infty$ for all $x^* \in X^*$; the norm in $\ell_w^p(X)$ is $\epsilon_p(x_n) = \sup \{ (\sum_n |\langle x^*, x_n \rangle|^p)^{1/p} : x^* \in B_{X^*} \}$. The set of all strongly *p*-summable sequences in X is denoted by $\ell_a^p(X)$; the norm in this space is $\pi_p(x_n) = (\sum_n \|x_n\|^p)^{1/p}$. If $T \in \Pi_p(X, Y)$, the correspondence $\hat{T} : (x_n) \mapsto (Tx_n)$ always induces a bounded operator from $\ell_w^p(X)$ into $\ell_a^p(Y)$ with $\|\hat{T}\| = \pi_p(T)$ [5, Proposition 2.1].

Families of operators arise in different applications: equations containing a parameter, homotopies of operators, and so forth. In these applications, it may be very interesting to know that, given a set $\mathcal{M} \subset \Pi_p(X, Y)$ and $(x_n) \in \ell_w^p(X)$, the series $\sum_n \|Tx_n\|^p$ is uniformly convergent in $T \in \mathcal{M}$. The main purpose of this paper is to study *uniformly p-summing* sets, that is, those sets $\mathcal{M} \subset \Pi_p(X, Y)$ for which, given $(x_n) \in \ell_w^p(X)$, the series $\sum_n \|Tx_n\|^p$ is uniformly convergent in $T \in \mathcal{M}$. These sets also enjoy some properties that justify their study; the next proposition lists some of them.

PROPOSITION 1.1. (a) *Let (T_k) be a sequence in $\Pi_p(X, Y)$. Then, $\widehat{T}_k \xrightarrow{k} 0$ pointwise if and only if $T_k \xrightarrow{k} 0$ pointwise and (T_k) is uniformly p -summing.*

(b) *Let $\mathcal{M} \subset \Pi_p(X, Y)$ be a uniformly p -summing set. If \mathcal{M} is endowed with the strong operator topology, then the map $T \in \mathcal{M} \mapsto \sum_n \|Tx_n\|^p \in \mathbb{R}$ is continuous for every $(x_n) \in \ell_w^p(X)$.*

A basic argument shows that uniformly p -summing sets are bounded for the p -summing norm. In fact, if X does not contain any copy of c_0 , bounded sets and uniformly 1-summing sets are the same. That is the reason for which we only consider operators defined on a $\mathcal{C}(\Omega)$ -space, Ω being a compact Hausdorff space. We recall that every weakly compact operator $T : \mathcal{C}(\Omega) \rightarrow Y$ has a representing measure $m_T : \Sigma \rightarrow Y$ defined by $m_T(B) = T^{**}(\chi_B)$ for all $B \in \Sigma$, where Σ denotes the Borel σ -field of subsets of Ω and χ_B denotes the characteristic function of B . The vector measure m_T is regular and countably additive [6, Theorem VI.2.5 and Corollary VI.2.14]. If we denote by \tilde{T} the operator T^{**} restricted to $B(\Sigma)$ (the space of all bounded Borel-measurable scalar-valued functions defined on Ω), then

$$\tilde{T}\varphi = \int_{\Omega} \varphi dm_T, \tag{1.2}$$

for all $\varphi \in B(\Sigma)$ (the integral is the elementary Bartle integral [6, Definition I.1.12]).

It is well known that every p -summing operator defined on a Banach space X is weakly compact. In Section 2, we consider 1-summing operators T defined on $\mathcal{C}(\Omega)$; these operators are characterized as those with representing measure m_T having finite variation and $\pi_1(T) = |m_T|(\Omega)$ [6, Theorem VI.3.3]. We show that a set $\mathcal{M} \subset \Pi_1(\mathcal{C}(\Omega), Y)$ is uniformly 1-summing if and only if the family of all variation measures $\{|m_T| : T \in \mathcal{M}\}$ is uniformly bounded and there is a countably additive measure $\mu : \Sigma \rightarrow [0, \infty)$ such that $\{|m_T| : T \in \mathcal{M}\}$ is uniformly μ -continuous.

In Section 3, we mention a special class of uniformly p -summing operators: *uniformly dominated sets*. The relationship between uniformly summing sets and relatively weak compactness is also studied. Finally, we give some examples and open problems.

2. Uniformly 1-summing sets in $\Pi_1(\mathcal{C}(\Omega), Y)$. Before facing our main theorem, we include three results which correspond to the vector measure theory. These results will be usually invoked along the following lines.

PROPOSITION 2.1 [6, Proposition I.1.17]. *The following statements about a collection $\{m_i : i \in I\}$ of Y -valued measures defined on a σ -field Σ are equivalent:*

- (a) *the set $\{m_i : i \in I\}$ is uniformly countably additive, that is, if (E_n) is a sequence of pairwise disjoint members of Σ , then $\lim_n \|\sum_{k \geq n} m_i(E_k)\| = 0$ uniformly in $i \in I$,*
- (b) *the set $\{\gamma^* \circ m_i : i \in I, \gamma^* \in B_{Y^*}\}$ is uniformly countably additive,*
- (c) *if (E_n) is a sequence of pairwise disjoint members of Σ , then $\lim_n \|m_i(E_n)\| = 0$ uniformly in $i \in I$,*
- (d) *if (E_n) is a sequence of pairwise disjoint members of Σ , then $\lim_n \|m_i\|(E_n) = 0$ uniformly in $i \in I$, where $\|m_i\|$ denotes the semivariation of m_i ,*
- (e) *the set $\{\gamma^* \circ m_i : i \in I, \gamma^* \in B_{Y^*}\}$ is uniformly countably additive.*

THEOREM 2.2 [6, Theorem I.2.4]. *Let $\{m_i : \Sigma \rightarrow Y : i \in I\}$ be a uniformly bounded (with respect to the semivariation) family of countably additive vector measures on a σ -field Σ . The family $\{m_i : i \in I\}$ is uniformly countably additive if and only if there exists a positive real-valued countably additive measure μ on Σ such that $\{m_i : i \in I\}$ is uniformly μ -continuous, that is,*

$$\lim_{\mu(E) \rightarrow 0} \|m_i(E)\| = 0 \tag{2.1}$$

uniformly in $i \in I$.

If Ω is a compact Hausdorff space and Σ denotes the σ -field of the Borel subsets of Ω , a vector measure m on Σ is regular if for each Borel set E and $\varepsilon > 0$ there exists a compact set K and an open set O such that $K \subset E \subset O$ and $\|m\|(O \setminus K) < \varepsilon$.

PROPOSITION 2.3 [6, Lemma VI.2.13]. *Let \mathcal{K} be a family of regular (countably additive) scalar measures defined on Σ . Each of the following statements implies all the others:*

- (a) *for each pairwise disjoint sequence (O_n) of open subsets of Ω , $\lim_n \mu(O_n) = 0$ uniformly in $\mu \in \mathcal{K}$,*
- (b) *for each pairwise disjoint sequence (O_n) of open subsets of Ω , $\lim_n |\mu|(O_n) = 0$ uniformly in $\mu \in \mathcal{K}$,*
- (c) *\mathcal{K} is uniformly countably additive,*
- (d) *\mathcal{K} is uniformly regular, that is, if $E \in \Sigma$ and $\varepsilon > 0$, then there exists a compact set K and an open set O such that $K \subset E \subset O$ and $\sup_{\mu \in \mathcal{K}} |\mu|(O \setminus K) < \varepsilon$.*

Now, we are able to show our main result. In the proof, we will use the fact that $|m_T|$ is regular when $T : \mathcal{C}(\Omega) \rightarrow Y$ is 1-summing [7, Proposition 15.21].

THEOREM 2.4. *Let $\mathcal{M} \subset \Pi_1(\mathcal{C}(\Omega), Y)$ be a bounded set. The following statements are equivalent:*

- (a) *\mathcal{M} is uniformly 1-summing,*
- (b) *the family of nonnegative measures $\{|m_T| : T \in \mathcal{M}\}$ is uniformly countably additive,*
- (c) *given $\varepsilon > 0$ and a disjoint sequence (E_n) of Borel subsets of Ω , there exists $n_0 \in \mathbb{N}$ such that*

$$\sum_{n \geq n_0} \|m_T(E_n)\| < \varepsilon, \tag{2.2}$$

for all $T \in \mathcal{M}$.

PROOF. (a) \Rightarrow (b). According to [6, Lemma VI.2.13], it suffices to show that $\lim_{n \rightarrow \infty} |m_T|(O_n) = 0$ uniformly in $T \in \mathcal{M}$, for all disjoint sequences (O_n) of open subsets of Ω . By contradiction, suppose that there exists $\varepsilon > 0$, a sequence (T_n) in \mathcal{M} , and a strictly increasing sequence (k_n) of natural numbers such that

$$|m_{T_n}|(O_{k_n}) > 2\varepsilon, \quad \forall n \in \mathbb{N}. \tag{2.3}$$

Now we consider the operators $S_n : \mathcal{C}(\Omega, O_{k_n}) \rightarrow Y$ defined by

$$S_n \varphi = \int_{O_{k_n}} \varphi \, d\mathbf{m}_{T_n}, \tag{2.4}$$

for all $\varphi \in \mathcal{C}(\Omega, O_{k_n})$, where $\mathcal{C}(\Omega, O_{k_n})$ is the closed subspace of $\mathcal{C}(\Omega)$ formed by all continuous functions φ on Ω such that φ vanishes in $\Omega \setminus O_{k_n}$. It is known that $\pi_1(S_n) = |\mathbf{m}_{T_n}|(O_{k_n})$, for all $n \in \mathbb{N}$ [7, Theorem 19.3]. For each $n \in \mathbb{N}$, we can choose a finite set $\{\varphi_1^n, \dots, \varphi_{p_n}^n\} \subset \mathcal{C}(\Omega, O_{k_n})$ satisfying $\epsilon_1(\varphi_i^n)_{i=1}^{p_n} \leq 1$ and

$$\sum_{i=1}^{p_n} \|S_n \varphi_i^n\| > \pi_1(S_n) - \epsilon. \tag{2.5}$$

Since the open sets O_{k_n} are disjoint, it follows that the sequence $(\varphi_1^1, \dots, \varphi_{p_1}^1, \varphi_1^2, \dots, \varphi_{p_2}^2, \dots)$ belongs to $\ell_w^1(\mathcal{C}(\Omega))$. Nevertheless, for all $n \in \mathbb{N}$, we have

$$\sum_{m \geq n} \sum_{i=1}^{p_m} \|T_n \varphi_i^m\| \geq \sum_{i=1}^{p_n} \|T_n \varphi_i^n\| = \sum_{i=1}^{p_n} \|S_n \varphi_i^n\| > \pi_1(S_n) - \epsilon = |\mathbf{m}_{T_n}|(O_{k_n}) - \epsilon > \epsilon. \tag{2.6}$$

This denies (a) and proves that (a) implies (b).

(b) \Rightarrow (c). Again we proceed by contradiction. Suppose (E_n) is a disjoint sequence of Borel subsets of Ω for which there exists $\epsilon > 0$, a sequence (T_n) in \mathcal{M} , and a strictly increasing sequence (k_n) of natural numbers so that

$$\sum_{i=k_n+1}^{k_{n+1}} \|\mathbf{m}_{T_n}(E_i)\| > \epsilon, \quad \forall n \in \mathbb{N}. \tag{2.7}$$

If we put $B_n = \bigsqcup_{i=k_n+1}^{k_{n+1}} E_i$, the above inequality yields $|\mathbf{m}_{T_n}|(B_n) > \epsilon$. So, in view of [6, Proposition I.1.17], the family $\{|\mathbf{m}_T| : T \in \mathcal{M}\}$ is not uniformly countably additive.

(c) \Rightarrow (b). We need to prove

$$\lim_{n \rightarrow \infty} |\mathbf{m}_T|(E_n) = 0 \quad \text{uniformly in } T \in \mathcal{M}, \tag{2.8}$$

for all disjoint sequences (E_n) of Borel subsets of Ω . Suppose (b) fails. Then, there exists $\epsilon > 0$, a sequence (T_n) in \mathcal{M} , and a strictly increasing sequence (k_n) of natural numbers satisfying

$$|\mathbf{m}_{T_n}|(E_{k_n}) > \epsilon, \quad \forall n \in \mathbb{N}. \tag{2.9}$$

For each $n \in \mathbb{N}$, we choose a finite partition $\{E_1^n, \dots, E_{p_n}^n\}$ of E_{k_n} for which

$$\sum_{i=1}^{p_n} \|\mathbf{m}_{T_n}(E_i^n)\| > \epsilon. \tag{2.10}$$

Then, the disjoint sequence $(E_1^1, \dots, E_{p_1}^1, E_1^2, \dots, E_{p_2}^2, \dots)$ does not satisfy (c).

(b)⇒(a). According to [6, Theorem I.2.4] there exists a countably additive measure $\mu : \Sigma \rightarrow [0, \infty)$ so that

$$\lim_{\mu(E) \rightarrow 0} |m_T|(E) = 0 \quad \text{uniformly in } T \in \mathcal{M}. \tag{2.11}$$

Hence, given $\varepsilon > 0$, there exists $\delta > 0$ such that, if $E \in \Sigma$ verifies $\mu(E) < \delta$, then $|m_T|(E) < \varepsilon/2$, for all $T \in \mathcal{M}$.

Next, given $(\varphi_n) \in \ell_w^1(\mathcal{C}(\Omega))$ with $\epsilon_1(\varphi_n) \leq 1$, notice that the series $\sum_{n=1}^\infty |\varphi_n(t)|$ is convergent for all $t \in \Omega$. Put $f_n(t) = \sum_{k=1}^n |\varphi_k(t)|$ and $f(t) = \lim_{n \rightarrow \infty} f_n(t)$. By Egorov’s theorem, the sequence (f_n) is quasi-uniformly convergent to f . Then, there exists $E \in \Sigma$ such that $\mu(E) < \delta$ and

$$f_n|_{\Omega \setminus E} \rightarrow f|_{\Omega \setminus E} \tag{2.12}$$

uniformly. If $C = \sup\{|m_T|(\Omega) : T \in \mathcal{M}\}$, there exists $n_0 \in \mathbb{N}$ so that

$$\sum_{n \geq n_0} |\varphi_n(t)| < \frac{\varepsilon}{2C}, \quad \forall t \in \Omega \setminus E. \tag{2.13}$$

Now,

$$\begin{aligned} \sum_{n \geq n_0} \|T\varphi_n\| &= \sum_{n \geq n_0} \left\| \int_{\Omega} \varphi_n(t) dm_T \right\| \\ &\leq \sum_{n \geq n_0} \left\| \int_E \varphi_n(t) dm_T \right\| + \sum_{n \geq n_0} \left\| \int_{\Omega \setminus E} \varphi_n(t) dm_T \right\| \\ &\leq \sum_{n \geq n_0} \int_E |\varphi_n(t)| d|m_T| + \sum_{n \geq n_0} \int_{\Omega \setminus E} |\varphi_n(t)| d|m_T| \\ &= \int_E \left(\sum_{n \geq n_0} |\varphi_n(t)| \right) d|m_T| + \int_{\Omega \setminus E} \left(\sum_{n \geq n_0} |\varphi_n(t)| \right) d|m_T| \\ &\leq |m_T|(E) + \frac{\varepsilon}{2C} |m_T|(\Omega \setminus E) \\ &< \varepsilon. \end{aligned} \tag{2.14} \quad \square$$

We denote by $\mathcal{V}(X, Y)$ the class of completely continuous operators from X into Y , that is, the class of operators which map weakly convergent sequences in X into norm-convergent sequences in Y . A set $\mathcal{M} \subset \mathcal{V}(X, Y)$ is said to be *uniformly completely continuous* if, given a weakly convergent sequence (x_n) in X , (Tx_n) is norm convergent uniformly in $T \in \mathcal{M}$. The following result gives some characterizations of uniformly completely continuous sets in $\mathcal{V}(\mathcal{C}(\Omega), Y)$. Recall that an operator T defined on $\mathcal{C}(\Omega)$ is completely continuous if and only if T is weakly compact [6, Corollary VI.2.17], so m_T is countably additive and regular, too.

THEOREM 2.5. *Let $\mathcal{M} \subset \mathcal{V}(\mathcal{C}(\Omega), Y)$ be a bounded set for the operator norm. The following statements are equivalent:*

- (a) \mathcal{M} is uniformly completely continuous,
- (b) the family $\{m_T : T \in \mathcal{M}\}$ is uniformly countably additive,

(c) $\mathcal{M}^* = \{T^* : T \in \mathcal{M}\}$ is collectively weakly compact, that is, the set $\bigcup_{T \in \mathcal{M}} T^*(B_{Y^*})$ is relatively weakly compact in $\mathcal{C}(\Omega)^*$.

PROOF. (a) \Rightarrow (b). By [6, Proposition I.1.17], the family $\{m_T : T \in \mathcal{M}\}$ is uniformly countably additive if and only if $\mathcal{N} = \{\gamma^* \circ m_T : T \in \mathcal{M}, \gamma^* \in B_{Y^*}\}$ is. According to [6, Lemma VI.1.13], we have to prove that

$$\lim_{n \rightarrow \infty} \gamma^* \circ m_T(O_n) = 0 \quad \text{uniformly in } \mathcal{N}, \tag{2.15}$$

for all disjoint sequences (O_n) of open subsets of Ω . By contradiction, suppose there exists such a sequence (O_n) for which $\lim_{n \rightarrow \infty} \gamma^* \circ m_T(O_n) = 0$ but not uniformly in \mathcal{N} . Then, there exists $\varepsilon > 0$ and sequences $(\gamma_n^*) \subset B_{Y^*}$, $(T_n) \in \mathcal{M}$, and $(O_{k_n}) \subset (O_n)$ such that

$$|\gamma_n^* \circ m_{T_n}(O_{k_n})| > \varepsilon, \quad \forall n \in \mathbb{N}. \tag{2.16}$$

Now, using the regularity of each m_{T_n} , we can find a sequence of compact sets (H_n) with $H_n \subset O_{k_n}$ and

$$\|m_{T_n}\|(O_{k_n} \setminus H_n) < \frac{\varepsilon}{2}, \quad \forall n \in \mathbb{N}, \tag{2.17}$$

($\|m\|$ denotes the semivariation of m , that is, $\|m\|(E) = \sup\{|\gamma^* \circ m|(E) : \gamma^* \in B_{Y^*}\}$). By Urysohn’s lemma, for every $n \in \mathbb{N}$ there exists a continuous function $\varphi_n : \Omega \rightarrow [0, 1]$ such that $\varphi_n(H_n) = 1$ and $\varphi_n(\Omega \setminus O_{k_n}) = 0$. Obviously, the series $\sum_{n=1}^\infty \varphi_n$ is unconditionally convergent in $\mathcal{C}(\Omega)$. Since \mathcal{M} is uniformly completely continuous, there exists $n_0 \in \mathbb{N}$ such that

$$\|T\varphi_n\| < \frac{\varepsilon}{2}, \quad \forall n \geq n_0, \forall T \in \mathcal{M}. \tag{2.18}$$

Then, we have

$$\begin{aligned} \|m_{T_n}(O_{k_n})\| &\leq \|m_{T_n}(O_{k_n}) - T_n\varphi_n\| + \|T_n\varphi_n\| \\ &= \left\| \int_{\Omega} \chi_{O_{k_n}} dm_{T_n} - \int_{\Omega} \varphi_n dm_{T_n} \right\| + \|T_n\varphi_n\| \\ &= \left\| \int_{O_{k_n}} (1 - \varphi_n) dm_{T_n} \right\| + \|T_n\varphi_n\| \\ &= \left\| \int_{O_{k_n} \setminus H_n} (1 - \varphi_n) dm_{T_n} \right\| + \|T_n\varphi_n\| \\ &\leq \|m_{T_n}\|(O_{k_n} \setminus H_n) + \|T_n\varphi_n\| \\ &< \varepsilon, \end{aligned} \tag{2.19}$$

for all $n \geq n_0$. This is in contradiction with (2.16).

(b) \Rightarrow (a). By [6, Theorem I.2.4], there exists a scalar countably additive measure $\mu : \Sigma \rightarrow [0, \infty)$ such that $\{m_T : T \in \mathcal{M}\}$ is uniformly μ -continuous. Then, if (φ_n) is a sequence

that tends to zero weakly in $\mathcal{C}(\Omega)$, it is obvious that zero is the pointwise limit of the sequence $(\varphi_n(t))$. Now, using Egorov's theorem and proceeding along similar lines as the proof of (b) \Rightarrow (a) in [Theorem 2.4](#), the proof concludes.

(b) \Leftrightarrow (c). The set $\bigcup_{T \in \mathcal{M}} T^*(B_{Y^*}) = \{\gamma^* \circ m_T : T \in \mathcal{M}, \gamma^* \in B_{Y^*}\} \subset \mathcal{C}(\Omega)^*$ is relatively weakly compact if and only if it is bounded and uniformly countably additive [[4](#), Theorem VII.13]. A call to [[6](#), Proposition I.1.17] makes clear that $\bigcup_{T \in \mathcal{M}} T^*(B_{Y^*})$ is uniformly countably additive if and only if condition (b) is satisfied. \square

COROLLARY 2.6. *If $\mathcal{M} \subset \Pi_1(\mathcal{C}(\Omega), Y)$ is uniformly 1-summing, then \mathcal{M} is uniformly completely continuous.*

The converse of the last result is not true in general.

PROPOSITION 2.7. *Suppose that the cardinal of Ω is infinite. The following statements are equivalent:*

- (a) *each subset of $\Pi_1(\mathcal{C}(\Omega), Y)$ uniformly completely continuous is uniformly 1-summing,*
- (b) *Y is finite-dimensional.*

PROOF. (a) \Rightarrow (b). By contradiction, suppose there is an unconditionally summable serie $\sum_k \gamma_k$ in Y such that $\sum_k \|\gamma_k\| = \infty$. Let (ω_k) be a sequence in Ω with $\omega_k \neq \omega_l$ when $k \neq l$. For each $m \in \mathbb{N}$ consider the operator $T_m : \mathcal{C}(\Omega) \rightarrow Y$ defined by

$$T_m \varphi = \sum_{k=1}^m \varphi(\omega_k) \gamma_k. \tag{2.20}$$

It is not difficult to show that $\mathcal{M} = (T_m)$ is uniformly completely continuous. Nevertheless,

$$\pi_1(T_m) = \sum_{k=1}^m \|\gamma_k\| \xrightarrow{m} \infty, \tag{2.21}$$

so \mathcal{M} cannot be uniformly 1-summing because it is not π_1 -bounded.

(b) \Rightarrow (a). This follows easily in view of conditions (b) in [Theorems 2.4](#) and [2.5](#). \square

We have showed that the converse of [Corollary 2.6](#) is not true in general. However, a direct argument using [Theorems 2.4](#) and [2.5](#) leads up to conclude that every uniformly completely continuous set $\mathcal{M} \subset \Pi_1(\mathcal{C}(\Omega), Y)$ verifying the following condition is uniformly 1-summing:

- (i) given $T \in \mathcal{M}$ and a finite subset $\{(\varphi_1, \gamma_1^*), \dots, (\varphi_m, \gamma_m^*)\}$ of $\mathcal{C}(\Omega) \times B_{Y^*}$, there exist $S \in \mathcal{M}$ and $z^* \in B_{Y^*}$ such that $|\langle \gamma_n^*, T \varphi_n \rangle| \leq |\langle z^*, S \varphi_n \rangle|$, $n = 1, \dots, m$.

3. Final notes and examples. The Grothendieck-Pietsch domination theorem states that an operator $T: X \rightarrow Y$ is p -summing if and only if there exists a positive Radon measure μ defined on the (weak*) compact space B_{X^*} such that

$$\|Tx\|^p \leq \int_{B_{X^*}} |\langle x^*, x \rangle|^p d\mu(x^*), \tag{3.1}$$

for all $x \in X$ [5, Theorem 2.12]. Since the appearance of this theorem, there is a great interest in finding out the structure of uniformly p -dominated sets. A subset \mathcal{M} of $\Pi_p(X, Y)$ is *uniformly p -dominated* if there exists a positive Radon measure μ such that the inequality (3.1) holds for all $x \in X$ and all $T \in \mathcal{M}$. In [3, 8, 9], the reader can find some of the most recent steps given on this subject. Now we are going to show that these sets are uniformly p -summing.

PROPOSITION 3.1. *If $\mathcal{M} \subset \Pi_p(X, Y)$ is a uniformly p -dominated set, then $\mathcal{M}^{**} = \{T^{**} : T \in \mathcal{M}\}$ is uniformly p -summing.*

PROOF. Let μ be a measure for which \mathcal{M} is uniformly p -dominated. In a similar way as in the Pietsch factorization theorem [5, Theorem 2.13], we can obtain, for all $T \in \mathcal{M}$, operators $U_T: L_p(\mu) \rightarrow \ell_\infty(B_{Y^*})$, $\|U_T\| \leq \mu(B_{X^*})^{1/p}$, and an operator $V: X \rightarrow L_\infty(\mu)$ such that the following diagram is commutative:

$$\begin{array}{ccc}
 X & \xrightarrow{T} & Y \\
 \downarrow V & & \searrow i_Y \\
 & & \ell_\infty(B_{Y^*}) \\
 & \nearrow U_T & \\
 L^\infty(\mu) & \xrightarrow{i_p} & L^p(\mu)
 \end{array} \tag{3.2}$$

Here, i_p is the canonical injection from $L_\infty(\mu)$ into $L_p(\mu)$ and i_Y is the isometry from Y into $\ell_\infty(B_{Y^*})$ defined by $i_Y(y) = (\langle y^*, y \rangle)_{y^* \in B_{Y^*}}$. Notice that i_p^{**} can be viewed as i_p composed with the canonical projection $P: L_\infty(\mu)^{**} \rightarrow L_\infty(\mu)$ which is simply the adjoint of the usual embedding $L_1(\mu) \rightarrow L_1(\mu)^{**}$. By weak compactness, we may and do consider T^{**} as a map from X^{**} into Y for which

$$i_Y \circ T^{**} = U_T \circ i_p \circ P \circ V^{**}. \tag{3.3}$$

Given $\varepsilon > 0$ and $(x_n^{**}) \in \ell_w^p(X^{**})$, we can choose $n_0 \in \mathbb{N}$ so that

$$\sum_{n \geq n_0} \|i_p \circ P \circ V^{**}(x_n^{**})\|^p < \frac{\varepsilon}{\mu(B_{X^*})}, \tag{3.4}$$

because $i_p \circ P \circ V^{**}$ is p -summing. Then, we have

$$\begin{aligned}
 \sum_{n \geq n_0} \|T^{**}x_n^{**}\|^p &= \sum_{n \geq n_0} \|i_Y \circ T^{**}(x_n^{**})\|^p = \sum_{n \geq n_0} \|U_T \circ i_p \circ P \circ V^{**}(x_n^{**})\|^p \\
 &\leq \mu(B_{X^*}) \sum_{n \geq n_0} \|i_p \circ P \circ V^{**}(x_n^{**})\|^p < \varepsilon,
 \end{aligned} \tag{3.5}$$

for all $T \in \mathcal{M}$. So, \mathcal{M}^{**} is uniformly p -summing. □

It is easy to show that the study of uniformly p -summing sets can be reduced to the behavior of its sequences. Indeed, a bounded set \mathcal{M} in $\Pi_p(X, Y)$ is uniformly p -summing if and only if every sequence (T_n) in \mathcal{M} admits a uniformly p -summing subsequence. Thus, it seems to be interesting to make clear the relationship between uniformly p -summing sets and relatively weakly compact sets. For $p = 1$, we have the following result.

PROPOSITION 3.2. *Every relatively weakly compact set in $\Pi_1(X, Y)$ is uniformly 1-summing.*

PROOF. Let \mathcal{M} be a relatively weakly compact set in $\Pi_1(X, Y)$. Given $\hat{x} = (x_n) \in \ell_w^1(X)$, consider the (weak-weak) continuous operator $U_{\hat{x}} : \Pi_1(X, Y) \rightarrow \ell_a^1(Y)$ defined by $U_{\hat{x}}(T) = (Tx_n)$. Then, $U_{\hat{x}}(\mathcal{M})$ is relatively weakly compact in $\ell_a^1(Y)$; according to [2, Theorem 2], we can conclude that \mathcal{M} is uniformly 1-summing. \square

Proposition 3.2 does not remain true if $p = 2$. For example, for each $\beta = (\beta_n) \in \ell_2$ consider the operator $T_\beta : c_0 \rightarrow \ell_2$ defined by $T(\alpha_n) = (\alpha_n \cdot \beta_n)$ and put $\mathcal{M} = \{T_\beta : \beta \in B_{\ell_2}\} \subset \Pi_2(c_0, \ell_2)$ [5, Theorem 3.5]. If we consider ℓ_2 as a subspace of $\Pi_2(c_0, \ell_2)$, the set $\mathcal{M} = B_{\ell_2}$ is relatively weakly compact. Nevertheless, no matter how we choose $k \in \mathbb{N}$,

$$\sum_{n \geq k} \|T_{e_k} e_n\|^2 = 1, \tag{3.6}$$

so \mathcal{M} cannot be uniformly 2-summing.

Now we show that there are uniformly p -summing sets failing to be relatively weakly compact.

PROPOSITION 3.3. *If every uniformly p -summing set is relatively weakly compact in $\Pi_p(X, Y)$, then Y is reflexive.*

PROOF. Fixing $x_0^* \in X^*$ with $\|x_0^*\| = 1$, the isometry $y \in Y \mapsto x_0^* \otimes y \in x_0^* \otimes Y$ allows us to see Y as a subspace of $\Pi_p(X, Y)$. If $\varepsilon > 0$ and $(x_n) \in \ell_w^p(X)$, there exists $n_0 \in \mathbb{N}$ so that

$$\sum_{n \geq n_0} |\langle x_0^*, x_n \rangle|^p < \varepsilon; \tag{3.7}$$

hence, for every $y \in B_Y$,

$$\sum_{n \geq n_0} \|(x_0^* \otimes y)(x_n)\|^p = \sum_{n \geq n_0} |\langle x_0^*, x_n \rangle|^p \|y\|^p < \varepsilon. \tag{3.8}$$

This yields that B_Y is uniformly p -summing and, by hypothesis, weakly compact. \square

The converse of Proposition 3.3 is not always true. By contradiction, suppose every uniformly 1-summing set in $\Pi_1(\ell_1, \ell_2)$ is relatively weakly compact. Because ℓ_1 does not contain any copy of c_0 , every bounded set in $\Pi_1(\ell_1, \ell_2)$ is relatively weakly compact. Then, we conclude that $\Pi_1(\ell_1, \ell_2)$ is reflexive, which is not possible since ℓ_1^* , viewed as a subspace of $\Pi_1(\ell_1, \ell_2)$, is not.

However, if $p = 1$ and $X = \mathcal{C}(\Omega)$, the reflexivity of Y is a sufficient condition for a uniformly 1-summing set to be relatively weakly compact. Indeed, if $rbvca(\Sigma, Y)$ denotes

the set of all regular, countably additive, Y -valued measures m on Σ with bounded variation, recall that relatively weakly compact sets \mathcal{M} in $rbvca(\Sigma, Y)$ are those verifying the following conditions: (i) \mathcal{M} is bounded; (ii) the family of nonnegative measures $\{|m| : m \in \mathcal{M}\}$ is uniformly countably additive; and (iii) for each $E \in \Sigma$, the set $\{m(E) : m \in \mathcal{M}\}$ is relatively weakly compact in Y [6, Theorem IV.2.5]. Having in mind the identification between $\Pi_1(\mathcal{C}(\Omega), Y)$ and $rbvca(\Sigma, Y)$, and making use of the characterization of uniformly 1-summing sets obtained in Theorem 2.4, we conclude the next characterization.

COROLLARY 3.4. *The following statements are equivalent:*

- (a) Y is reflexive,
- (b) every set \mathcal{M} in $\Pi_1(\mathcal{C}(\Omega), Y)$ is uniformly 1-summing if and only if \mathcal{M} is relatively weakly compact.

It is well known that a linear operator T is 1-summing if and only if T^{**} is. So, it is natural to ask if a set \mathcal{M} is uniformly 1-summing whenever $\mathcal{M}^{**} = \{T^{**} : T \in \mathcal{M}\}$ is. Unfortunately, we are going to show that this is not true in general. It suffices to take X as the separable \mathcal{L}_∞ -space of Bourgain and Delbaen [1]. This space has the Radon-Nikodym property, so it does not contain any copy of c_0 . Nevertheless, X^* is isomorphic to ℓ_1 and, therefore, X^{**} contains a copy of c_0 . Let (e_n) be the canonical basis of ℓ_1 and $J : \ell_1 \rightarrow X^*$ an isomorphism. Put $T_n = J e_n \in \Pi_1(X, \mathbb{R})$; the set $\mathcal{M} = \{T_n : n \in \mathbb{N}\}$ is uniformly 1-summing since it is bounded and X does not contain any copy of c_0 . Notice that the elements of \mathcal{M}^{**} are the linear forms $x^{**} \in X^{**} \mapsto \langle x^{**}, J e_n \rangle \in \mathbb{R}$, for all $n \in \mathbb{N}$. If (e_n^*) is the canonical basis of c_0 , then $((J^*)^{-1}(e_n^*)) \in \ell_w^1(X^{**})$; hence, no matter how we choose $k \in \mathbb{N}$, it turns out that

$$\sum_{n \geq k} |T_k^{**}((J^*)^{-1}(e_n^*))| = \sum_{n \geq k} |\langle (J^*)^{-1}(e_n^*), J e_k \rangle| = \sum_{n \geq k} |\langle e_n^*, e_k \rangle| = 1, \tag{3.9}$$

and \mathcal{M}^{**} cannot be uniformly 1-summing.

Nevertheless, if \mathcal{M} is a set of operators defined on c_0 , then it is true that \mathcal{M} is uniformly 1-summing if and only if \mathcal{M}^{**} is too. To see this, notice that for a 1-summing operator $T : (\alpha_n) \in c_0 \mapsto \sum_{n=1}^\infty \alpha_n x_n \in X$, the second adjoint $T^{**} : \ell_\infty \rightarrow X$ is defined by $T^{**}(\beta_n) = \sum_{n=1}^\infty \beta_n x_n$, for all $(\beta_n) \in \ell_\infty$.

When \mathcal{M} is a set of operators defined on a $\mathcal{C}(\Omega)$ -space, we do not know whether \mathcal{M}^{**} inherits the property or not. Anyway, we are going to prove the following weaker result. We inject isometrically $B(\Sigma)$ into $\mathcal{C}(\Omega)^{**}$ in the natural way.

PROPOSITION 3.5. *If $\mathcal{M} \subset \Pi_1(\mathcal{C}(\Omega), X)$ is uniformly 1-summing, then $\tilde{\mathcal{M}} = \{\tilde{T} : B(\Sigma) \rightarrow X : T \in \mathcal{M}\}$ is uniformly 1-summing too.*

PROOF. The argument is similar to the one used in the proof of (b) \Rightarrow (a) in Theorem 2.4. □

Finally, we give an example to show that Corollary 2.6 is not true if $\mathcal{C}(\Omega)$ is replaced by a general Banach space X . It suffices to take $X = \ell_2$ and $\mathcal{M} = \{e_n^* : n \in \mathbb{N}\}$, where (e_n^*) is the unit basis of $\ell_2^* \simeq \ell_2$. The set \mathcal{M} is bounded in $\Pi_1(\ell_2, \mathbb{R})$ and, therefore, uniformly 1-summing but it is not uniformly completely continuous.

REFERENCES

- [1] J. Bourgain and F. Delbaen, *A class of special \mathcal{L}_∞ -spaces*, Acta Math. **145** (1980), no. 3-4, 155-176.
- [2] J. K. Brooks and N. Dinculeanu, *Weak compactness in spaces of Bochner integrable functions and applications*, Adv. Math. **24** (1977), no. 2, 172-188.
- [3] J. M. Delgado and C. Piñeiro, *A note on uniformly dominated sets of summing operators*, Int. J. Math. Math. Sci. **29** (2002), no. 5, 307-312.
- [4] J. Diestel, *Sequences and Series in Banach Spaces*, Graduate Texts in Mathematics, vol. 92, Springer-Verlag, New York, 1984.
- [5] J. Diestel, H. Jarchow, and A. Tonge, *Absolutely Summing Operators*, Cambridge Studies in Advanced Mathematics, vol. 43, Cambridge University Press, Cambridge, 1995.
- [6] J. Diestel and J. J. Uhl, Jr., *Vector Measures*, Mathematical Surveys, vol. 15, American Mathematical Society, Rhode Island, 1977.
- [7] N. Dinculeanu, *Vector Measures*, International Series of Monographs in Pure and Applied Mathematics, vol. 95, Pergamon Press, Oxford; VEB Deutscher Verlag der Wissenschaften, Berlin, 1967.
- [8] R. Khalil and M. Hussain, *Uniformly dominated sets of p -summing operators*, Far East J. Math. Sci., Special Volume (1998), no. Part I, 59-68.
- [9] B. Marchena and C. Piñeiro, *Bounded sets in the range of an X^{**} -valued measure with bounded variation*, Int. J. Math. Math. Sci. **23** (2000), no. 1, 21-30.

J. M. Delgado: Departamento de Matemáticas, Facultad de Ciencias Experimentales, Campus Universitario del Carmen, Avda. de las Fuerzas Armadas, 21071 Huelva, Spain

E-mail address: jmde1ga@uhu.es

Cándido Piñeiro: Departamento de Matemáticas, Facultad de Ciencias Experimentales, Campus Universitario del Carmen, Avda. de las Fuerzas Armadas, 21071 Huelva, Spain

E-mail address: candido@uhu.es



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

