

SUBDIVISIONS OF ROTATIONALLY SYMMETRIC PLANAR CONVEX BODIES MINIMIZING THE MAXIMUM RELATIVE DIAMETER

ANTONIO CAÑETE, UWE SCHNELL, AND SALVADOR SEGURA GOMIS

ABSTRACT. In this work we study subdivisions of k -rotationally symmetric planar convex bodies that minimize the maximum relative diameter functional. For some particular subdivisions called k -partitions, consisting of k curves meeting in an interior vertex, we prove that the so-called *standard k -partition* (given by k equiangular inradius segments) is minimizing for any $k \in \mathbb{N}$, $k \geq 3$. For general subdivisions, we show that the previous result only holds for $k \leq 6$. We also study the optimal set for this problem, obtaining that for each $k \in \mathbb{N}$, $k \geq 3$, it consists of the intersection of the unit circle with the corresponding regular k -gon of certain area. Finally, we also discuss the problem for planar convex sets and large values of k , and conjecture the optimal k -subdivision in this case.

1. INTRODUCTION

The study of the classical geometric magnitudes associated to a planar compact convex set (perimeter, area, inradius, circumradius, diameter and minimal width) is one of the main points of interest for Convex Geometry, and these magnitudes have been considered since ancient times. They appear in the origin of this mathematical field, and in fact, there is a large variety of related problems, providing a more complete understanding of these geometric measures.

One of the most common problems involving these magnitudes consists of finding precise relations between some of them (and not only in the convex framework). For instance, the well-known isoperimetric inequality relates the perimeter and the area functionals, stating that, among all planar sets with fixed enclosed area, the circle is the one with the least possible perimeter [15, 14]. In the same direction, the isodiametric inequality asserts that the circle is also the planar compact convex set of prescribed area with the minimum possible diameter [2]. Another example is given by the equilateral triangle, which is the planar compact convex set with minimum enclosed area for fixed minimal width [16]. We point out that these relations are usually established by means of appropriate general inequalities, with the characterization of the equality case providing then the optimal set for the corresponding functionals

Date: September 25, 2018.

2010 Mathematics Subject Classification. 52A10, 52A40.

Key words and phrases. partitioning problems, k -rotationally symmetric planar convex body, maximum relative diameter.

(that is, the set attaining the minimum possible value for the functional). A nice detailed description of the relations between two (and three) of these classical magnitudes can be found in [17].

Apart from the above approach, these geometric functionals can be studied in a large range of different related problems. One of these problems, which has been considered in several works during the last years, is the so-called *fencing problem*, or more generally, the *partitioning problem*. For these problems, the main interest is determining the division of the given set into k connected subsets (of equal or unequal areas) which minimizes (or maximizes) a particular geometric magnitude [7]. The relative isoperimetric problem is one of these questions, where the goal is minimizing the perimeter of the division curves for fixed enclosed areas. For this problem, the general geometric properties satisfied by the minimizing divisions can be deduced by using a variational approach, and particular results characterizing the solutions have been obtained for some regular polygons and for the circle, see [18, 1, 4, 5].

Regarding the diameter functional, the partitioning problem can be studied by means of the *maximum relative diameter* functional. We recall that for a planar compact convex set C and a decomposition of C into k connected subsets C_1, \dots, C_k , the maximum relative diameter associated to the decomposition is defined as the maximum of the diameters of the subsets C_i , for $i = 1, \dots, k$ (this means that it measures the largest distance in the subsets provided by the decomposition). In this setting, the partitioning problem seeks the decomposition of C into k subsets with the least possible value for the maximum relative diameter. This problem has been recently studied for $k = 2$ and $k = 3$ under some additional symmetry hypotheses [11, 3], and the results therein constitute the main motivation for our work.

In [11], the previous problem is treated for *centrally symmetric* planar compact convex sets, and for decompositions into two subsets of equal areas, proving that the minimizing division for the maximum relative diameter is always given by a line segment passing through the center of symmetry of the set [11, Prop. 4]. We point out that a more precise characterization of such a minimizing segment is not provided. Furthermore, the optimal set for this problem (that is, the centrally symmetric planar compact convex set attaining the minimum possible value for the maximum relative diameter) is also determined up to dilations, consisting of a certain intersection of two unit circles and a strip bounded by two parallel lines, see [11, Th. 5].

In [3], the problem is considered for *3-rotationally symmetric* planar compact convex sets, and for divisions into three subsets of equal areas. The main result establishes that the so-called *standard trisection*, determined by three equiangular inradius segments, minimizes the maximum relative diameter [3, Th. 3.5 and Prop. 5.1] for any set of that class. In addition, the optimal set for this problem is also characterized, up to dilations, as the intersection of the unit circle with a certain equilateral triangle [3, Th. 4.7]. We remark that, although the results in [3] are stated for divisions into three subsets of *equal areas*, all of them also hold in the case of unequal areas.

This paper is inspired precisely in these last two references involving the maximum relative diameter [11, 3]. As a natural continuation, we shall focus on the class \mathcal{C}_k of k -rotationally symmetric planar convex bodies (that is, compact convex sets) for each $k \in \mathbb{N}$, $k \geq 3$. And for each set in \mathcal{C}_k , we shall investigate the divisions into k connected subsets minimizing the maximum relative diameter functional. In this setting we shall consider two different types of divisions, namely k -subdivisions and k -partitions. For a given set $C \in \mathcal{C}_k$, a k -partition of C will be a decomposition of C into k connected subsets by k curves starting at different points of ∂C , and with all of them meeting at an interior point in C . On the other hand, a k -subdivision of C will be a general decomposition of C into k connected subsets, with no additional restrictions.

The goal of this paper is studying the minimizing k -partitions and minimizing k -subdivisions for the maximum relative diameter, for any k -rotationally symmetric planar convex body. In fact, we shall see that the so-called *standard k -partition* (consisting of k inradius segments symmetrically placed) is a minimizing k -partition for the maximum relative diameter, for any $k \in \mathbb{N}$, $k \geq 3$ (Theorem 4.5), being also minimizing among k -subdivisions only when $k \leq 6$ (Theorem 4.6). For $k > 7$ we shall provide some examples showing that the standard k -partition is not a minimizing k -subdivision, which suggests that, in those cases, a complete characterization is a difficult question. In addition, we point out that the uniqueness of the minimizing k -partition does not hold for this problem, as explained in Subsection 4.1, since proper slight deformations of a minimizing k -partition are also minimizing.

Another interesting result in this paper is the characterization of the *optimal bodies* for this problem. For each $k \in \mathbb{N}$, $k \geq 3$, we are able to determine the k -rotationally symmetric planar convex body attaining the minimum possible value for the maximum relative diameter, when considering k -partitions. Taking into account the previous Theorem 4.5, for each set of this class, that minimum value will be provided by the corresponding standard k -partition. In Section 5 we give the description of these optimal bodies for each $k \geq 3$, which, in general, consist of the intersections of the unit circle with a certain regular k -gon, up to dilations.

We remark that the case $k = 2$ (treating the decompositions into two subsets for centrally symmetric planar convex bodies and previously studied in [11]) presents some differences with respect to the other situations, as explained in Remark 3.6, and cannot be treated as for $k \geq 3$. The main reason for this particular behaviour lies in the following technical fact: the associated standard 2-partition of a set of this class, which is provided by two inradius segments placed symmetrically, is just a line segment passing through the center of symmetry of the set, and so the computation of the maximum relative diameter associated to this standard 2-partition cannot be done by using Lemma 3.2, which is fundamental when $k \geq 3$.

We have organized this paper as follows. Section 2 contains the precise definitions and the setting of the problem, and in Section 3 we describe the standard k -partition for any k -rotationally symmetric planar convex body.

This k -partition is constructed by applying successively the existing rotational symmetry to an inradius segment of the set. We remark that Lemma 3.2 establishes how to compute the maximum relative diameter for the standard k -partition.

In Section 4 we obtain the main results of this paper. Due to Lemmata 3.2, 4.1 and 4.4, we prove our Theorem 4.5:

For any k -rotationally symmetric planar convex body, the associated standard k -partition is a minimizing k -partition for the maximum relative diameter.

Furthermore, regarding general k -subdivisions, we obtain Theorem 4.6:

For any k -rotationally symmetric planar convex body, the associated standard k -partition is a minimizing k -subdivision for the maximum relative diameter, when $k \leq 6$.




FIGURE 1. The standard 3-partition for the equilateral triangle, the standard 4-partition for the square, and the standard 8-partition for the regular octagon

When $k \geq 7$, Examples 4.7 and 4.8 show that the standard k -partition is not a minimizing k -subdivision in general, suggesting that the characterization of such minimizing k -subdivisions is a difficult task which will depend on the particular considered set. We also point out that, for $k = 3$, the minimizing 3-partition and 3-subdivision in this setting have been previously obtained, see [3, Th. 3.5 and Prop. 5.1].

Moreover, in Subsection 4.1 we discuss the uniqueness of the solutions for this problem. We will show with some examples that slight modifications of a minimizing k -partition in a proper way (preserving the value of the maximum relative diameter) will produce minimizing k -partitions as well, and so we do not have uniqueness in this setting (as in most of the problems involving the diameter functional). This subsection concludes by posing some necessary conditions for a given k -partition to be minimizing, derived from Lemma 3.2 and Theorem 4.5.

In Section 5 we study the *optimal bodies* for this problem. More precisely, for each $k \in \mathbb{N}$, $k \geq 3$, we characterize the k -rotationally symmetric planar convex body for fixed area attaining the lowest possible value for the maximum relative diameter when considering k -partitions. Notice that, in view of Theorem 4.5, we have to focus on the standard k -partitions, and as a consequence of Lemma 3.2 we are able to prove our Theorem 5.1:

For any $k \in \mathbb{N}$, $k \geq 3$, the minimum value for the maximum relative diameter for k -partitions in the class of k -rotationally symmetric planar convex bodies is uniquely attained by the standard k -partition associated to the intersection of the unit circle and the regular k -gon with inradius equal to $1/(2 \sin(\pi/k))$, up to dilations.

The previous Theorem 5.1 characterizes the optimal body, for each $k \in \mathbb{N}$, $k \geq 3$, as the intersection of the unit circle with a particular regular k -gon. In Subsection 5.1 we analyze these intersections in each case, obtaining the precise optimal sets: for $k = 3$ and $k = 5$, the optimal bodies are respectively an equilateral triangle and a regular pentagon *with rounded vertices*, for $k = 4$ we have a square, and for $k \geq 6$ the best set in this setting is the unit circle. We remark that the optimal body for $k = 3$ was already characterized in [3, Th. 4.7] by means of a constructive procedure.




FIGURE 2. Optimal sets for $k = 3$, $k = 4$, $k = 5$, and $k \geq 6$

Finally, in Section 6 we complete our study by considering large values of $k \in \mathbb{N}$, discussing our problem for k -subdivisions (focusing on planar convex bodies). Recall that Theorem 4.5 establishes that the standard k -partitions are minimizing k -partitions for any $k \in \mathbb{N}$, $k \geq 3$, and minimizing k -subdivisions only when $k \leq 6$, for any k -rotationally symmetric planar convex body. In Corollary 6.4 we obtain a lower bound for the maximum relative diameter functional when considering k -subdivisions of a planar convex body, involving a negligible addend when k is large enough. However, we think that such lower bound can be improved, and some reasonings in this section lead us to Conjecture 6.8: for large values of $k \in \mathbb{N}$, a minimizing k -subdivision will be given by a decomposition induced by a planar tiling of regular hexagons.

2. PRELIMINARIES

In this paper we will consider, for any given $k \in \mathbb{N}$, $k \geq 3$, the class \mathcal{C}_k of k -rotationally symmetric planar convex bodies (assuming therefore the compactness of the sets). We recall that, in this setting, each set C in the class \mathcal{C}_k has a center of symmetry p , and then, the k -rotationally symmetry means that C is invariant under the rotation centered at p with angle $\varphi_k := 2\pi/k$. Some examples of sets belonging to \mathcal{C}_k are the regular k -gon and the Reuleaux k -gon.

Given a planar compact set C , we shall denote by $r(C)$ the *inradius* of C (that is, the radius of the largest ball contained in C), and by $R(C)$ the




FIGURE 3. Some examples of k -rotationally symmetric planar convex bodies: the equilateral and Reuleaux triangles are 3-rotationally symmetric, the regular octagon is 8-rotationally symmetric (and also 4-rotationally symmetric), and the circle is k -rotationally symmetric for any $k \in \mathbb{N}$.

circumradius of C (radius of the smallest ball containing C). If no confusion may arise, we shall simply denote them by r and R , respectively. For k -rotationally symmetric planar convex bodies, we have the following Lemma, which provides a metric characterization for the inradius and the circumradius.

Lemma 2.1. *Let $C \in \mathcal{C}_k$, and let p be its center of symmetry. Then, the inradius r coincides with the minimal Euclidean distance between p and a point in ∂C , and the circumradius R coincides with the maximal Euclidean distance between p and a point in ∂C .*

Proof. It is known that the set I of centers of all largest balls contained in C is a convex set with empty interior [13]. As C is k -rotationally symmetric, this implies that $I = \{p\}$ (note that for any $c \in I$, $c \neq p$, the images of c by the existing rotations would belong to I , and so p would be an interior point of I). Then there is a unique largest ball contained in C , which is centered at p . This immediately yields the statement for the inradius.

On the other hand, as the smallest ball containing C is unique, its center has to coincide with p (otherwise, an analogous rotation argument will contradict that uniqueness). This implies trivially the statement for the circumradius. \square




FIGURE 4. The inradius r and the circumradius R of an equilateral triangle.

We now define the notions of k -subdivision and k -partition of a planar compact set, which will be considered along this work. Notice that these definitions can be generally done without assuming additional hypotheses regarding convexity or symmetry.

Definition 2.2. Let C be a planar compact set. A k -subdivision S of C is a decomposition of C into k connected subsets $\{C_1, \dots, C_k\}$, not necessarily of equal areas, satisfying:

- i) $C = \bigcup_{i=1}^k C_i$,
- ii) $\text{int}(C_i) \cap \text{int}(C_j) = \emptyset$, for $i, j \in \{1, \dots, k\}$, $i \neq j$.

In addition, if a k -subdivision S is given by k curves meeting at a point $c \in \text{int}(C)$, and the other endpoints of the curves are different points in ∂C , then S will be called a k -partition of C . In this case, c is called the common point of the k -partition and the endpoints of the curves meeting ∂C will be called the endpoints of the k -partition.

We stress that k -subdivisions are general decompositions of the original set C into k connected subsets, and that k -partitions are k -subdivisions with a particular topological structure: they consist of k curves converging to an interior point, each of them meeting the boundary of C at a different point.




FIGURE 5. Three different 5-subdivisions for the regular pentagon.




FIGURE 6. Three different 5-partitions for the regular pentagon.

Remark 2.3. In this work, we are focusing on the class \mathcal{C}_k of k -rotationally symmetric planar convex bodies since the lack of the rotational symmetry prevents a general study of this question. Moreover, for any set $C \in \mathcal{C}_k$, its center of symmetry is a remarkable point in the interior of C , and so it seems natural to consider k -partitions, which are precisely the k -subdivisions determined by interior points of C .

We now recall the definition of the maximum relative diameter for a k -subdivision of a planar compact set.

Definition 2.4. Let C be a planar compact set, and let S be a k -subdivision of C into subsets $\{C_1, \dots, C_k\}$. We define the maximum relative diameter associated to S as

$$d_M(S, C) = \max\{D(C_i) : i = 1, \dots, k\},$$

where $D(C_i) = \max\{d(x, y) : x, y \in C_i\}$ denotes the Euclidean diameter of C_i .

Remark 2.5. Along this work, we shall usually write $d_M(S)$ instead of $d_M(S, C)$, if no confusion may arise.

Remark 2.6. For a given planar compact set C and a k -subdivision S of C , the existence of $d_M(S, C)$ is assured due to the continuity of the Euclidean distance $d : \mathbb{R}^2 \rightarrow \mathbb{R}$, and the compactness of C .

Remark 2.7. By a standard argument it is clear that $d_M(S, C)$ is attained by points lying in the boundary of one of the subsets C_i determined by the k -subdivision S . Moreover, it is well-known that the diameter of a convex polygon is given by the distance between two of its vertices.

In the forthcoming sections, we shall investigate the k -partitions and the k -subdivisions of a given k -rotationally symmetric planar convex body which minimize the maximum relative diameter functional. In other words, for a given set $C \in \mathcal{C}$, we shall look for

$$\min\{d_M(P, C) : P \text{ is a } k\text{-partition of } C\},$$

and

$$\min\{d_M(S, C) : S \text{ is a } k\text{-subdivision of } C\}.$$

The k -partitions and k -subdivisions attaining those minimum values will be called *minimizing*. As usual in this kind of optimization problems involving the diameter functional, we point out that the uniqueness of the minimizing k -subdivision is not expected, as we shall see in Subsection 4.1.

3. STANDARD k -PARTITION

In this section we shall describe a particular k -partition for the sets of our class \mathcal{C}_k , called the *standard k -partition*, by means of a simple geometrical intrinsic construction. This k -partitions will play an important role along this work.

For $C \in \mathcal{C}_k$, let $x_1 \in \partial C$ be a point in ∂C with minimal distance to the center of symmetry p . In view of Lemma 2.1, this implies that $d(p, x_1)$ is the inradius of C , and so the segment $\overline{px_1}$ will be referred to as *an inradius segment* of C (notice that for regular polygons, such a segment is called apothem). By applying $k - 1$ times the rotation centered at p with angle $\varphi_k = 2\pi/k$ to the point x_1 , we shall finally obtain a set of points $\{x_1, \dots, x_k\} \subset \partial C$, all of them minimizing the distance to p . If we join these points to p , we shall get k inradius segments dividing C into k connected congruent subsets. This k -partition will be called *standard k -partition* of C , and will be denoted by $P_k(C)$, or simply P_k .

Remark 3.1. In general, the previous construction of the standard k -partition associated to a k -rotationally symmetric planar convex body C is not uniquely determined. In fact, if there are more than k points in ∂C attaining the minimal distance to the center of symmetry of C , then the associated standard k -partition is not unique (for instance, this happens for the standard 3-partition




FIGURE 7. Standard k -partitions for different k -rotationally symmetric planar bodies

of the regular hexagon, or for the standard k -partition of the circle for any $k \geq 2$). In any case, all the standard k -partitions associated to C will be k -rotationally congruent by construction.

For the standard k -partition of a given set C in \mathcal{C}_k , the associated maximum relative diameter can be easily computed in terms of the inradius and the circumradius of C , as proved in the following Lemma 3.2.

Lemma 3.2. *Let C be a k -rotationally symmetric planar convex body, with inradius r and circumradius R . Let P_k be the standard k -partition associated to C . Then,*

$$d_M(P_k, C) = \max\{R, 2r \sin(\varphi_k/2)\}.$$

Remark 3.3. Recall that $\varphi_k/2 = \pi/k$.

Proof. We can assume that $d_M(P_k, C) = D(C_1)$, where C_1 is one of the congruent subsets determined by P_k . Call $x_1, x_2 \in \partial C_1$ the endpoints of the two inradius segments bounding C_1 . Any pair of points realizing $D(C_1)$ cannot be contained in the relative interior of an inradius segment (in that case, the diameter could be clearly increased by considering some proper point along the segment). As the maximal distance between two points on $\partial C \cap \partial C_1$ is $d(x_1, x_2) = 2r \sin(\pi/k)$, and the maximal distance between p and ∂C equals R , the statement follows. \square

The following examples show that both possibilities from Lemma 3.2 may occur.

Example 3.4. (Circle) Any circle is a k -rotationally symmetric planar convex body for any $k \in \mathbb{N}$, $k \geq 2$, whose inradius r coincides with its circumradius R . Consequently, in view of Lemma 3.2, it is immediate checking that $d_M(P_k) = 2r \sin(\varphi_k/2)$ for $k \in \{3, 4, 5, 6\}$, and $d_M(P_k) = R$ for $k \geq 6$.

Example 3.5. (Regular polygons) For a regular k -gon E_k , Lemma 3.2 yields that $d_M(P_k, E_k) = R(E_k)$, for $k \in \mathbb{N}$, $k \geq 3$, due to the well-known relation $r(E_k) = R(E_k) \cos(\varphi_k/2)$.

Remark 3.6. Lemma 3.2 does not hold for $k = 2$ (it can be easily checked by considering the square or the circle). The reason of this fact is that the standard 2-partition P_2 will divide any centrally symmetric planar convex body C into two subsets whose boundaries do not contain the center of symmetry p as a *vertex*. Notice that p will belong to the line segment $\overline{x_1 x_2}$ (being x_1, x_2

the endpoints of P_2), and so p will not be relevant for computing $d_M(P_2, C)$. In fact, it can be checked that

$$d_M(P_2, C) = \max\{d(x_1, x) : x \in \partial C\}.$$

The main difficulty in this case is that the quantity $d(x_1, x)$, when $x \in \partial C$, cannot be estimated in general, which prevents using the arguments from this paper for $k = 2$. We recall that this case was studied in [11], where it is proved that the minimizing 2-partition for d_M is given by a line segment passing through the center of symmetry of the set, without a more precise description [11, Prop. 4].

4. MAIN RESULTS

In this section, we aim to compare the maximum relative diameters associated to the standard k -partition P_k and an arbitrary k -partition P of a given k -rotationally symmetric planar convex body C , when $k \geq 3$. By means of Lemmata 4.1 and 4.4, we will conclude that $d_M(P_k, C) \leq d_M(P, C)$, and so the standard k -partition always provides a minimizing k -partition for the maximum relative diameter functional. Furthermore, if we consider general k -subdivisions, we will see that the same previous result holds only when $k \leq 6$ (Theorem 4.6), showing also some counterexamples for $k \geq 7$. Finally, in Subsection 4.1 we will see that the uniqueness of the minimizing k -partition cannot be expected for this problem.

Lemma 4.1. *Let C be a k -rotationally symmetric planar convex body, with circumradius R , for $k \geq 3$. Let P be a k -partition of C . Then $d_M(P) \geq R$.*

Proof. Let c be the common point of P , and C_1, \dots, C_k the corresponding subsets of C given by P . Note that $c \in \bigcap_{i=1}^k C_i$. Call p the center of symmetry of C , and let y_1, \dots, y_k be points in ∂C such that $d(p, y_i) = R$ (we can assume they are placed k -symmetrically along ∂C). Then, it is not difficult to check that $p = \sum_{i=1}^k y_i/k$ is the unique minimum for the functional $S : C \rightarrow \mathbb{R}$ given by

$$S(x) = \sum_{i=1}^k d(x, y_i)^2,$$

and so

$$S(c) = \sum_{i=1}^k d(c, y_i)^2 \geq S(p) = kR^2.$$

This necessarily implies that some y_j satisfies $d(c, y_j) \geq R$, yielding $d_M(P) \geq d(c, y_j) \geq R$, as desired. \square

The above Lemma 4.1 also holds for k -subdivisions, when $k \leq 6$. Notice that the proof is different from the one of Lemma 4.1, since k -subdivisions do not have a common interior point as k -partitions.

Lemma 4.2. *Let C be a k -rotationally symmetric planar convex body, with circumradius R , for $3 \leq k \leq 6$. Let S be a k -subdivision of C . Then $d_M(S) \geq R$.*

Proof. Let y_1, \dots, y_k be points in ∂C such that $d(p, y_i) = R$. We can assume they are placed k -symmetrically along ∂C , and so they will determine a regular k -gon E_k with edge length equal to $2R \sin(\pi/k) \geq 2R \sin(\pi/6) = R$, since $k \leq 6$. Then we have that p, y_1, \dots, y_k are $k+1$ points with mutual distances greater than or equal to R . Since at least two of these points will be contained in a same subset given by the subdivision S , it follows that $d_M(S) \geq R$. \square

Remark 4.3. The previous Lemma 4.2 does not hold when $k > 6$. For instance, for $k = 7$, consider a regular heptagon H with circumradius $R = 1$ and center of symmetry p , and the 7-subdivision S shown in Figure 8 (which is a modification of the 7-partition of H whose endpoints are the vertices of H with p as common point). This 7-subdivision is given by eleven line segments meeting in threes at five inner points (placed at the same distance from p), with congruent subsets H_2, H_7 , congruent subsets H_3, H_4, H_5, H_6 , and $p \in H_1$. In this case, we have checked that $d_M(S) = D(H_2) = d(a, b) = 0.9892 < R = 1$, and so Lemma 4.2 is not satisfied.




FIGURE 8. A 7-subdivision of the regular heptagon with maximum relative diameter strictly less than the circumradius

The following Lemma 4.4 is true not only for k -partitions, but also for general k -subdivisions.

Lemma 4.4. *Let C be a k -rotationally symmetric planar convex body, with in-radius r , for $k \geq 3$. Let S be a k -subdivision of C . Then $d_M(S) \geq 2r \sin(\pi/k)$.*

Proof. Let x_1, \dots, x_k be points in ∂C with $d(p, x_i) = r$. We can assume that they are placed k -symmetrically, and so they determine a regular k -gon with edge length $2r \sin(\pi/k)$. Call C_1, \dots, C_k the subsets of C given by the k -subdivision S .

If two (consecutive) points x_α, x_β belong to a same subset C_i , then $d_M(S) \geq D(C_i) \geq d(x_\alpha, x_\beta) = 2r \sin(\pi/k)$, and the result follows. So we will assume that each point x_i only belongs to C_i , for $i = 1, \dots, k$.

Let p be the center of symmetry of C , and let K be the circle with center p and radius r . Clearly $x_1, \dots, x_k \in \partial K$. Since any pair of consecutive points x_{i-1}, x_i are contained in different subsets C_{i-1}, C_i , respectively, there must be a point $q_i \in \partial K$ between them, with $q_i \in C_{i-1} \cap C_i$. Let α_i denote the angle

at p determined by the segments $\overline{pq_i}$ and $\overline{pq_{i+1}}$. We have that $\alpha_i \in (0, 2\varphi_k) = (0, 4\pi/k)$, and $\alpha_1 + \dots + \alpha_k = 2\pi$. Then it follows that at least one angle α_j satisfies $2\pi/k \leq \alpha_j < 4\pi/k$. This implies that $d(q_j, q_{j+1}) = 2r \sin(\alpha_j/2) \geq 2r \sin(\pi/k)$, and so $d_M(S) \geq D(C_j) \geq d(q_j, q_{j+1}) \geq 2r \sin(\pi/k)$, which proves the statement. \square

We can now state the main results of this section, asserting that for any k -rotationally symmetric planar convex body, the associated standard k -partition is a minimizing k -partition for the maximum relative diameter functional, and also a minimizing k -subdivision when $k \leq 6$. Both results are immediate consequences of the previous Lemmata.

Theorem 4.5. *Let C be a k -rotationally symmetric planar convex body, and P_k the associated standard k -partition of C . Let P be any k -partition of C . Then, $d_M(P) \geq d_M(P_k)$.*

Proof. From Lemma 3.2, we have that $d_M(P_k) = \max\{R, 2r \sin(\pi/k)\}$, where R and r are the circumradius and the inradius of C . By using Lemmata 4.1 and 4.4, we conclude that $d_M(P) \geq d_M(P_k)$. \square

Theorem 4.6. *Let C be a k -rotationally symmetric planar convex body, and P_k the associated standard k -partition of C , with $k \leq 6$. Let S be any k -subdivision of C . Then $d_M(S) \geq d_M(P_k)$.*

Proof. Taking into account Lemmata 3.2, 4.2 and 4.4, the result follows. \square

We point out that the previous Theorem 4.6 is the best result we can obtain for minimizing k -subdivisions, since it does not hold for $k \geq 7$, as shown in the following examples.

Example 4.7. In Remark 4.3 we considered a regular heptagon H with circumradius $R = 1$, and described a 7-subdivision S with $d_M(S) = 0.9892$. As $d_M(P_7) = R = 1$ for the standard 7-partition P_7 associated to H (see Example 3.5), it follows that P_7 is not a minimizing 7-subdivision for the maximum relative diameter.

Example 4.8. Consider the unit circle C and the 8-subdivision S shown in Figure 9, given by an inner subset C_1 bounded by a circle of radius 0.43, and seven congruent subsets C_2, \dots, C_8 determined by 7-symmetric segments joining ∂C and ∂C_1 . Straightforward computations yield that $D(C_1) = 0.86$ and $D(C_2) = 0.86$, and so $d_M(S) = 0.86$. Thus the corresponding standard 8-partition P_8 is not a minimizing 8-subdivision for the circle, since $d_M(P_8) = R(C) = 1$, see Example 3.4.

Example 4.9. The following idea suggests that when $k \in \mathbb{N}$ is large enough, the corresponding standard k -partition will not be a minimizing k -subdivision for the maximum relative diameter. Consider the unit circle, which satisfies $d_M(P_k) \geq 1$ for any $k \in \mathbb{N}$, $k \geq 3$, where P_k is the standard k -partition, see Example 3.4. It is clear that, when $k \geq k_0$ for $k_0 \in \mathbb{N}$ large enough, we can divide the unit circle into k *small* subsets, each one with diameter strictly




FIGURE 9. A 8-subdivision of the circle

less than 1, thus obtaining a k -subdivision with maximum relative diameter strictly less than 1. This implies that P_k is not minimizing for $k \geq k_0$. This reasoning can be probably applied for regular k -gons when k is large enough (and then close to be a circle). In Section 6 we shall study the minimizing k -subdivision for a planar convex body, for large values of $k \in \mathbb{N}$.

Remark 4.10. We remark that Theorems 4.5 and 4.6 were already obtained in [3] for the case $k = 3$.

4.1. Uniqueness of the minimizing k -partition. The uniqueness of the solution is also an interesting question for these kinds of optimization problems. In our setting, where the considered functional is the maximum relative diameter, the uniqueness of the minimizing k -partition does not hold, as we can see in the example from Figure 10. For the regular hexagon H , we exhibit three different minimizing 6-partitions: the associated standard 6-partition P_6 (which is minimizing by Theorem 4.5), and two others 6-partitions obtained by slight modifications of P_6 , keeping invariant the value of the maximum relative diameter $d_M(P_6, H)$. Other similar examples can be constructed for the regular hexagon and, in general, for any k -rotationally symmetric planar convex body (see Figure 11). This shows that the minimizing k -partition for any set of our class is not unique, being not possible to obtain a complete classification of all of them.




FIGURE 10. Three different minimizing 6-partitions for the regular hexagon. The first one is the corresponding standard 6-partition P_6 , and the others are slight modifications.

However, some necessary conditions for being a minimizing k -partition can be derived from a deeper analysis of Lemmata 4.1 and 4.4. More precisely, let C be a k -rotationally symmetric planar convex body, with center of symmetry p , and let P be a minimizing k -partition. From Theorem 4.5 and Lemma 3.2 we have that $d_M(P) = d_M(P_k) = \max\{R, 2r \sin(\pi/k)\}$, where P_k is the standard




FIGURE 11. Three different minimizing 4-partitions for the circle. The first one is the corresponding standard 4-partition P_4 , and the others are slight modifications.

k -partition associated to C , and R and r are the circumradius and the inradius of C , respectively. By discussing the two different possibilities, we shall obtain conditions that the minimizing k -partition P must satisfy. If $d_M(P_k) = R$, from the proof of Lemma 4.1 we can deduce that the common point of P has to coincide necessarily with p (otherwise, $d_M(P)$ will be *strictly greater* than R). On the other hand, if $d_M(P_k) = 2r \sin(\pi/k)$, an analysis of the equality case in the proof of Lemma 4.4 will yield that the endpoints of P have to coincide with the endpoints of P_k .

5. OPTIMAL SETS

The search for the optimal sets in geometric optimization problems is, in general, a hard question. For some problems, it is only possible to obtain some properties of the solutions, but not a complete description of them. However, for the problem considered in this paper, we will see that we can give a precise characterization of the optimal set, which in this setting corresponds with the k -rotationally symmetric planar convex body attaining the minimum value for the maximum relative diameter.

Our initial motivation resides in the following question: given a k -rotationally symmetric planar convex body C , we know from Theorem 4.5 that the lowest value for the maximum relative diameter among all k -partitions of C is achieved by the standard k -partition $P_k(C)$ associated to C , and it is equal to $d_M(P_k(C), C)$. Then, a natural further step is investigating for the minimum of such lowest values, among all sets in the class \mathcal{C}_k . Equivalently, we are interested in finding

$$\min\{d_M(P_k(C), C) : C \in \mathcal{C}_k\},$$

which will consequently determine the optimal body in \mathcal{C}_k .

However, in this setting it is more convenient to find

$$\min\left\{\frac{d_M(P_k(C), C)^2}{A(C)} : C \in \mathcal{C}_k\right\},$$

where $A(C)$ denotes the area of C , since the quotient

$$(5.1) \quad \frac{d_M(P_k(C), C)^2}{A(C)}$$

is invariant under dilations about the center of symmetry of C , and therefore the optimal body will be independent from the enclosed area.

The following Theorem 5.1 provides the characterization of the optimal body in \mathcal{C}_k , for each $k \in \mathbb{N}$, $k \geq 3$.

Theorem 5.1. *Among all k -rotationally symmetric planar convex bodies for fixed $k \in \mathbb{N}$, $k \geq 3$, the minimum value in (5.1) is uniquely attained by the intersection of the unit circle with the regular k -gon with inradius $1/(2 \sin(\pi/k))$, up to dilations.*

Proof. In order to obtain the minimum value in (5.1), we can fix either the value of the maximum relative diameter or the area of the set. We shall proceed by fixing the maximum relative diameter, and so we shall *maximize* the area of the set.

Consider an arbitrary set C in \mathcal{C}_k with $d_M(P_k(C), C) = 1$, where $P_k(C)$ is the standard k -partition associated to C . From Lemma 3.2, it follows that $R(C) \leq 1$ and $2r(C) \sin(\pi/k) \leq 1$, where $R(C)$ and $r(C)$ are the circumradius and the inradius of C , respectively. The first inequality implies that C is contained in the unit circle, and the second one implies that $r(C) \leq 1/(2 \sin(\pi/k))$, which gives that C is contained in the regular k -gon with inradius $1/(2 \sin(\pi/k))$, due to the convexity of C . Therefore, C will be contained in the corresponding intersection, and the set with maximal area under these conditions will be necessarily that intersection. \square

Remark 5.2. The previous Theorem 5.1 characterizes the optimal body in \mathcal{C}_k for our problem, when considering k -partitions, for any $k \in \mathbb{N}$, $k \geq 3$. We can also study the same question for k -subdivisions. In that case, taking into account Theorem 4.6, analogous reasonings will yield to the same characterization of the optimal bodies only when $k \leq 6$.

5.1. Description of the optimal bodies. We shall now determine the optimal bodies for each $k \in \mathbb{N}$, $k \geq 3$, by analyzing the corresponding intersection described in Theorem 5.1 in each particular case. The precise optimal bodies are depicted in Figure 12.

For $k \in \{3, 5\}$, we have that the value of the inradius $1/(2 \sin(\pi/k))$ of the corresponding regular k -gon is strictly less than 1, and its circumradius is strictly greater than 1 (see Example 3.5). This implies that the regular k -gon actually intersects the unit circle, obtaining in these cases a sort of regular k -gons with *rounded corners* as optimal bodies.

For $k = 4$, the square with inradius $1/(2 \sin(\pi/4))$ has circumradius equal to 1. Hence, that square is completely contained in the unit circle and provides the optimal body in this case.

Finally, for $k \geq 6$, the inradius $1/(2 \sin(\pi/k))$ of the regular k -gon E_k is greater than or equal to 1, which implies that the unit circle is entirely contained in E_k . Therefore, the optimal bodies coincide with the unit circle for $k \geq 6$.




FIGURE 12. Optimal bodies for $k = 3$, $k = 4$, $k = 5$, and $k \geq 6$.

Remark 5.3. The optimal body for $k = 3$ was already obtained in [3, Th. 4.7] by means of a constructive procedure, and is a solution for some others geometric optimization problems, see [9, §. 4].

6. k -SUBDIVISIONS FOR LARGE VALUES OF k

In Section 4 we have proved that, for each k -rotationally symmetric planar convex body, the associated standard k -partition is a minimizing k -partition for the maximum relative diameter, for any $k \in \mathbb{N}$, $k \geq 3$ (Theorem 4.5), and also a minimizing k -subdivision for $k \in \{3, 4, 5, 6\}$ (Theorem 4.6). In this section, we will go further with the study of the k -subdivisions problem, by considering the asymptotic situation for large values of $k \in \mathbb{N}$, and searching for a minimizing k -subdivision for the maximum relative diameter.

We have already seen that, in general, the standard k -partition is not minimizing when $k \geq 7$ (for instance, see Examples 4.7 and 4.8). The reasonings in this section will lead us to conjecture that an optimal k -subdivision, when the number of regions k is large enough, is provided by a *hexagonal tiling* (up to some subsets touching the boundary of the original set). This kind of tilings have already appeared in the literature during the last years, specially from the isoperimetric point of view (see [10] for the least-perimeter subdivision of the plane, or [6] for the structure of the solutions of the planar soap bubble problem for a large number of regions). We point out that the assumption of rotational symmetry can be removed, and so we shall focus on general planar convex bodies.

The following lemma, whose proof relies on the classical isodiametric inequality [2], provides a first lower bound for the maximum relative diameter, when considering k -subdivisions of an arbitrary planar convex body.

Lemma 6.1. *Let C be a planar convex body, and let S be a k -subdivision of C , for $k \in \mathbb{N}$. Then,*

$$(6.1) \quad d_M(S, C) \geq \sqrt{\frac{A(C)}{k}} \sqrt{\frac{4}{\pi}}.$$

Proof. Let C_1, \dots, C_k be the subsets of C given by S , and call $d = d_M(S, C)$. By using the isodiametric inequality [2], since $D(C_i) \leq d$, we have that

$A(S_i) \leq \frac{\pi}{4} d^2$, for $i = 1, \dots, k$. Then,

$$A(C) = \sum_{i=1}^k A(C_i) \leq k \frac{\pi}{4} d^2,$$

which yields the statement. \square

Remark 6.2. We recall that the equality in the classical isodiametric inequality holds for circles, and so, the equality in (6.1) cannot be attained, since it is not possible to consider a subdivision given by circles due to the overlappings of the corresponding subsets. Therefore, (6.1) actually provides a *strict* lower bound for the maximum relative diameter functional.

The following Lemma 6.3 will allow to improve the lower bound obtained in the previous Lemma 6.1, for large values of k .

Lemma 6.3. *Let C be a planar convex body, and let S be a k -subdivision of C with $d_M(S, C) = d$. Then,*

$$(6.2) \quad A(C) \leq 0.688452 k d^2 + d P(C),$$

where $P(C)$ denotes the perimeter of C .

Proof. Call C_1, \dots, C_k the subsets of C given by S . Let \widetilde{C}_i be the convex hull of C_i . Then $D(\widetilde{C}_i) = D(C_i)$ for $i = 1, \dots, k$, and C is covered by the union of these convex hulls. Since the possible overlappings can be eliminated by dividing the intersection regions by proper line segments, we can then assume that C_1, \dots, C_k are convex polygons up to some arcs contained in ∂C . By replacing each of these arcs by the line segment determined by its endpoints, we shall obtain convex polygons $C'_i \subset C_i$, $i = 1, \dots, k$, whose union is a convex polygon C' contained in C . It is clear that $C \setminus C'$ is contained in the union of stripes parallel to $\partial C'$ and breadth d . This implies

$$(6.3) \quad A(C) \leq A(C') + d P(C') \leq A(C') + d P(C) = \sum_{i=1}^k A(C'_i) + d P(C).$$

The convex polygons C'_1, \dots, C'_k determine a planar graph with $k + 1$ faces (together with the unbounded component C'_0). If m and n denote the number of edges and vertices, by Euler's theorem we have $(k + 1) - m + n = 2$.

Since each edge is incident to two vertices and the degree of each vertex is at least 3 we obtain $2m \geq 3n$. It follows $1 = k - m + n \leq k - m + 2/3m = k - m/3$ and so

$$m \leq 3k - 3.$$

Let f_j be the number of polygons with j edges in $\{C'_1, \dots, C'_k\}$, and s the number of edges of C_0 . Then

$$2m = s + \sum_j f_j j$$

and so

$$\sum_j f_j j < s + \sum_j f_j j = 2m \leq 6k - 6 \leq 6k.$$

On the other hand, let m_j be the maximal area of a polygon of j edges with diameter one. It is known (see [12]) that m_j is attained by regular polygons for odd j , with

$$m_j = j/2 \cos(\pi/j) \tan(\pi/(2j)).$$

Furthermore, the values for m_4 , m_6 and m_8 are also numerically known (see [12]), so that $m_3 = 0.433012$, $m_4 = 0.5$, $m_5 = 0.657163$, $m_6 = 0.6749814429$, $m_7 = 0.719740$, $m_8 = 0.7268684828$, $m_9 = 0.745619$. Moreover, notice that $m_j \leq \pi/4$, for any $j \geq 3$, as the corresponding polygon will be contained in the circle of diameter one. Then, if C'_i is a polygon with j edges, $A(C'_i) \leq d^2 m_j$, and so

$$(6.4) \quad \sum_{i=1}^k A(C'_i) \leq \sum_j f_j m_j d^2 = d^2 \sum_j f_j m_j \leq d^2 \left(\sum_{j=3}^9 f_j m_j + \pi/4 \sum_{j \geq 10} f_j \right).$$

We are interested in finding an upper bound for (6.4). By replacing $\sum_{j \geq 10} f_j$ by \tilde{f} and calling $\tilde{m} = \pi/4$, and using linear programming, it is straightforward checking that the maximal value for $f_3 m_3 + \dots + f_9 m_9 + \tilde{f} \tilde{m}$ under the restrictions $f_3 + \dots + f_9 + \tilde{f} = k$, $3f_3 + \dots + 9f_9 + 10\tilde{f} \leq 6k$, and $f_3, \dots, f_9, \tilde{f} \geq 0$ is given by $k(m_5 + m_7)/2 = 0.688452k$, being attained by $f_5 = f_7 = k/2$ and $f_3 = \dots = f_9 = \tilde{f} = 0$.

Using that bound in (6.4), we conclude that $\sum_{i=1}^k A(C'_i) \leq 0.6884k d^2$, which yields the statement taking into account (6.3). \square

Corollary 6.4. *Let C be a planar convex body, and let S be a k -subdivision of C . Then*

$$(6.5) \quad d_M(S, C) \geq \sqrt{\frac{A(C)}{k}} \sqrt{\frac{1}{0.688452}} - O(1/k).$$

Proof. The statement is a direct consequence of (6.2). \square

Remark 6.5. For large values of $k \in \mathbb{N}$, (6.5) gives an improvement of the lower bound in (6.1). However, such a bound is *strict*, since a planar tiling by regular pentagons and regular heptagons cannot be realized.

In the following Lemma 6.6 we shall consider a particular k -subdivision for any planar convex body and any $k \in \mathbb{N}$, induced by a proper tiling of the plane by regular hexagons, see Figure 13. We shall estimate the maximum relative diameter d_M for these k -subdivisions, obtaining an upper bound for the minimal value of d_M , which we conjecture it is the optimal one for large values of $k \in \mathbb{N}$.

Lemma 6.6. *Let C be a planar convex body C . For each $k \in \mathbb{N}$, there exists a k -subdivision S_k such that*

$$(6.6) \quad d_M(S_k, C) \leq \sqrt{\frac{A(C)}{k}} \sqrt{\frac{8}{3\sqrt{3}}} + O\left(\frac{1}{k}\right).$$

Proof. Call H the regular hexagon with unit diameter, and so $A(H) = 3\sqrt{3}/8$. For any $d > 0$, let L_d be the planar lattice with fundamental cell dH . For any $k \in \mathbb{N}$, let d_k be the unique positive solution of

$$(6.7) \quad k A(dH) = A(C) + dP(C) + \pi d^2,$$

where $P(C)$ denotes the perimeter of C . It is clear that C is covered by the union of the fundamental cells of L_{d_k} with nonempty intersection with C . Call n the number of such fundamental cells, which will be contained in the outer parallel body $C + d_k B$, where B is the planar unit ball. Then, Steiner's formula and (6.7) give

$$n A(d_k H) \leq A(C + d_k B) = A(C) + d_k P(C) + \pi d_k^2 = k A(dH),$$

and so $n \leq k$. This means that the lattice L_{d_k} induces a k -subdivision S_k of C (if $n < k$, we can divide a fundamental cell until having a decomposition of C into k connected subsets) with $d_M(S_k, C) \leq d_k$. This fact, together with a straightforward estimate of d_k by using (6.7), proves the statement. \square

Remark 6.7. We stress that the k -subdivision S_k from Lemma 6.6 consists of regular hexagons of diameter d_k in the interior of the set, and portions of such hexagons when meeting the boundary of the set.




FIGURE 13. A subdivision of the circle given by regular hexagons

The previous Lemma 6.6 gives an upper bound for the minimum possible value of the maximum relative diameter for large enough values of $k \in \mathbb{N}$. However, we think that such a bound is the optimal one due to the following fact: taking into account Lemma 6.1 and using the covering density [8] (recall that the subdivision by circles would produce overlappings of the subsets), it is easy to check that the bound from (6.6) can be obtained from the bound given in (6.1), when k is large enough. This means, in some sense, that the hexagonal tiling corrects the defects appearing in the decomposition by circles. In view of this, we finish this section with the following Conjecture 6.8, regarding the minimizing k -subdivisions for a planar convex set and large values of $k \in \mathbb{N}$.

Conjecture 6.8. Let C be a planar convex set, and let S be any k -subdivision of C . For large enough values of $k \in \mathbb{N}$, the minimal value for $d_M(S, C)$ is provided by the k -subdivision described in Lemma 6.6, given by regular hexagons of fixed diameter in the interior of C and portions of such hexagons when meeting the boundary of C .

Acknowledgements. The authors would like to thank the Instituto de Matemáticas de la Universidad de Sevilla (IMUS), where part of this work was done. The first author is partially supported by the research projects MTM2010-21206-C02-01 and MTM2013-48371-C2-1-P, and by Junta de Andalucía grants FQM-325 and P09-FQM-5088. The third author is partially supported by MINECO (Ministerio de Economía y Competitividad) and FEDER (Fondo Europeo de Desarrollo Regional) project MTM2012-34037.

REFERENCES

- [1] M. Bleicher, *Isoperimetric divisions into several cells with natural boundary*, In Intuitive geometry (Siofok, 1985), volume 48 of Colloq. Math. Soc. Janos Bolyai (1987), 63-84.
- [2] L. Bieberbach, *Über eine Extremaleigenschaft des Kreises*, Jber. Deutsch. Math.-Vereinig **24** (1915), 247–250.
- [3] A. Cañete, C. Miori, S. Segura Gomis, *Trisections of a 3-rotationally symmetric planar convex body minimizing the maximum relative diameter*, J. Math. Anal. Appl. **418** (2014), no. 2, 1030–1046.
- [4] A. Cañete, M. Ritoré, *Least-perimeter partitions of the disk into three regions of given areas*, Indiana Univ. Math. J. **53** (2004), no. 3, 883904.
- [5] S. Cox, *Calculations of the minimal perimeter for n deformable cells of equal area confined in a circle*, Phil. Mag. Letts. **86** (2006), 569–578.
- [6] S. J. Cox, F. Morgan, F. Graner, *Are large perimeter-minimizing two-dimensional clusters of equal-area bubbles hexagonal or circular?*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **469** (2013), no. 2149, 20120392, 10 pp.
- [7] H. T. Croft, K. J. Falconer, R. K. Guy, *Unsolved problems in Geometry*, Springer-Verlag, Berlin, 1991.
- [8] G. Fejes Tóth, W. Kuperberg, *Packing and covering with convex sets, Chapter 3.3 in: Handbook of Convex Geometry*, Elsevier 1993, 799–860.
- [9] M. A. Hernández Cifre, G. Salinas, S. Segura Gomis, *Complete systems of inequalities*, J. Inequal. Pure Appl. Math. **2** (2001), no. 1, article 10.
- [10] T. C. Hales, *The honeycomb conjecture*, Discrete Comput. Geom. **25** (2001), no. 1, 1–22.
- [11] C. Miori, C. Peri, S. Segura Gomis, *On fencing problems*, J. Math. Anal. Appl. **300** (2004), no. 2, 464–476.
- [12] M. J. Mossinghoff, *Isodiametric problems for polygons*, Discrete Comput. Geom. **36** (2006), no. 2, 363–379.
- [13] M. Moszyńska, E. Saorín Gómez, *On the kernel centre of a convex body*, Beitr. Algebra Geom. **55** (2014), 595–602.
- [14] R. Osserman, *The isoperimetric inequality*, Bull. Amer. Math. Soc. **84** (1978), no. 6, 1182-1238.
- [15] E. Schmidt, *Über eine neue Methode zur Behandlung einer Klasse isoperimetrischer Aufgaben im Grossen*. Math. Z., 47 (1942), 489?-642.
- [16] J. Pál, *Ein Minimumproblem für Ovale*, Math. Ann. **83** (1921), no. 3–4, 311–319.
- [17] P. R. Scott, P. W. Awyong, *Inequalities for convex sets*, J. Inequal. Pure Appl. Math. **1** (2000), no. 1, article 6 (electronic).
- [18] Y. Tomonaga, *Geometry of length and area. 1*. Dept. of Mathematics, Utsunomiya University, Utsunomiya, 1974.

DEPARTAMENTO DE MATEMÁTICA APLICADA I, UNIVERSIDAD DE SEVILLA

E-mail address: `antonioc@us.es`

FACULTY OF MATHEMICS AND SCIENCES, UNIVERSITY OF APPLIED SCIENCES ZITTAU/GÖRLITZ

E-mail address: `uschnell@hszg.de`

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE ALICANTE

E-mail address: `salvador.segura@ua.es`