



## Use of polar and nonpolar fractions as additional information sources for studying thermoxidized virgin olive oils by FTIR

N. Tena, R. Aparicio-Ruiz and D.L. García-González<sup>✉</sup>

Instituto de la Grasa (CSIC), Padre García Tejero 4, E-41012 Sevilla, Spain  
<sup>✉</sup>Corresponding author: [dluisg@cica.es](mailto:dluisg@cica.es)

Submitted: 2 December 2013; Accepted: 7 February 2014

**SUMMARY:** Fourier transform infrared (FTIR) spectroscopy has been proposed to study the degradation of virgin olive oils (VOO) in samples undergoing thermoxidation. The polar and nonpolar fractions of oxidized oils have been analyzed by FTIR to provide further information on the minor spectral changes taking place during thermoxidation. This information assists in the interpretation of the spectra of the samples. For this purpose polar and nonpolar fractions of 47 VOO samples thermoxidized (190 °C) in a fryer were analyzed by FTIR. The time-course change of the band area assigned to single *cis* double bonds was explained by their correlation with the decrease in oleic acid (adjusted- $R^2=0.93$ ). The bands assigned to the hydroxyl groups and the first overtone of ester groups was better studied in the spectra collected for the polar and nonpolar fractions, respectively. The bands assigned to peroxide, epoxy, tertiary alcohols and fatty acids were clearly observed in the spectra of the polar fraction while they are not noticeable in the spectra of the oils.

**KEYWORDS:** FTIR spectroscopy; Polar and nonpolar fractions; Thermoxidation; Virgin olive oil

**RESUMEN:** *Uso de las fracciones polares y no polares como fuente de información adicional al estudio de aceites de oliva vírgenes por FTIR.* La espectroscopía de infrarrojos por transformada de Fourier (FTIR) se ha propuesto para estudiar la degradación de los aceites de oliva vírgenes (AOV) sujetas a termoxidación. Las fracciones polares y no polares de aceites oxidados se analizaron mediante FTIR para obtener más información sobre los cambios espectrales menores que tienen lugar durante la termoxidación. Esa información ayuda en la interpretación de los espectros de las muestras puras. Con este objetivo, fracciones polares y no polares de 47 AOV termoxidados (190 °C) en una freidora se analizaron mediante FTIR. La banda asignada a dobles enlaces *cis* se explica por su correlación con la disminución de ácido oleico ( $R^2$ -ajustado=0,93). Las bandas asignadas a los grupos hidroxilos y del primer sobretono de los grupos éster se estudió mejor en los espectros recogidos para la fracción polar y no polar, respectivamente. Grupos asignados a peróxidos, epoxi, alcoholes terciarios y los ácidos grasos se observan claramente en los espectros de la fracción polar.

**PALABRAS CLAVE:** Aceite de oliva virgen; Espectroscopia FTIR; Fracciones polares y no polares; Termoxidación

**Citation/Cómo citar este artículo:** Tena N, Aparicio-Ruiz R, García-González DL. 2014. Use of polar and nonpolar fractions as additional information sources for studying thermoxidized virgin olive oils by FTIR. *Grasas Aceites* 65 (3): e030. doi: <http://dx.doi.org/10.3989/gya.121913>.

**Copyright:** © 2014 CSIC. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial (by-nc) Spain 3.0 Licence.

## 1. INTRODUCTION

The consumption of frying products and fast food has dramatically increased over the last decades. The high concern for the toxic hazard of fried products (Friedman and Levin, 2008; Billek, 2000) and the economical implications of the oil stability during frying (Piper, 2001) have led to a major focus on virgin olive oil (VOO) as a frying oil. In fact, deep- and pan-frying is the second most important application of olive oil, which is a millenarian culinary practice of the Mediterranean countries (García-González *et al.*, 2009). The high stability of olive oil is mainly due to its low content in polyunsaturated fatty acids and its high content in antioxidants (e.g. phenols).

Although the official method of percentage of total polar compounds (TPC) is well correlated with the thermal degradation of the oil and oxidized fatty acids formed at frying temperatures (Sanibal and Mancini-Filho, 2004), this methodology is lengthy, expensive and laborious. Some alternative methods (determination of free fatty acids, percentage of polymers, peroxide value, Lovibond color, etc) have the general drawback of providing only partial information that is not necessarily correlated to the quality and safety decay of the frying oil (Stier, 2001). For that reason, several rapid spectroscopic methods are being proposed because they provide information on the representative markers of the chemical processes that occur during frying (hydrolysis and oxidation reactions). Fourier transform infrared spectroscopy (FTIR) has already been proven to be an appropriate technique in the quantitative determination of some parameters of edible oils during frying, such as free fatty acids (Li *et al.*, 2008), quantification of *trans* fatty acids (van de Voort *et al.*, 2008), peroxide values (Yu *et al.*, 2007) and iodine value (Sedman *et al.*, 1998; van de Voort *et al.*, 2008a). In one of our publications (Tena *et al.*, 2009) we described a method based on attenuated total reflectance (ATR)-FTIR to predict the deterioration degree of virgin olive oil during frying. The optimum infrared region to predict the TPC of thermoxidized oil was assigned to *trans* isomers ( $978\text{--}960\text{ cm}^{-1}$ ) and it is highly correlated with the percentage of elaidic acid (*trans* isomer of oleic acid). This compound rises as the frying process progresses at the same pace as TPC and the FTIR band assigned to single *trans* double bonds. However, during the thermoxidation process other kinds of degradation reactions occur, most of them being related to hydrolysis, oxidation and other reactions favored by high temperatures. The compounds formed or degraded in these reactions may not follow the same trend as TPC and *trans* double bonds, and consequently their control needs a further study beyond a simple correlation with single spectral regions. The ability of FTIR to study chemical groups associated with these reactions also

makes this technique optimal for studying these chemical processes.

Although ATR-FTIR provides information about the entire medium IR spectral region with an easy and rapid output, this technique has less sensitivity for small spectral changes. On the basis of Beer's law, this limitation means that the use of ATR for sample handling may cause difficulties in tracking low concentrations of chemical species. Alternatively, the use of transmission cells permits studying bands of low intensity that require higher path lengths ( $\sim 100\text{ }\mu\text{m}$  vs. few  $\mu\text{m}$  for ATR). A higher path length allows for maximizing the signal intensity of bands, hence they are recorded at a higher signal-to-noise ratio compared to the spectra collected with ATR. Thus, using transmission cells for sample handling in FTIR analysis provides better results (higher sensitivity and precision) for low intensity bands, although it has the problem of saturation of the detector signal for the most intense bands. Thus, transmission cells of several path lengths (100–350  $\mu\text{m}$ ) and window materials (KBr, CaF<sub>2</sub>, etc.) have been used to study hydroperoxides, the unsaturation degree and fatty acids, among other chemical species, in oxidized edible oils (van de Voort *et al.*, 2008; Moya Moreno *et al.*, 1999).

An approach to increasing the intensity of certain spectral bands is to extract the polar and nonpolar fractions of the oil. Separating the oil into two fractions additionally entails simplifying the spectra, avoiding, to some extent, the overlapping bands. Thus, the individual study of fractions obtained from complex matrices with solvents or solid-extraction cartridges has been addressed for a further interpretation of spectroscopic properties (Christophoridou and Dais, 2009; Scano *et al.*, 2008). Most of the fractionation methods are based on polarity or solubility, and extraction of the unsaponifiable matter. Thus, the examination of unsaponifiable matter of olive oil with FTIR and Raman spectroscopies allowed for the development of authentication methods with a better performance compared to the same methods applied to the oils (Baeten *et al.*, 2005).

To our knowledge fractionation into polar and nonpolar fractions has not yet been employed in the FTIR studies of edible oils, although it has been successfully applied to other kinds of spectroscopies. Thus, Valavanidis *et al.*, (2004) studied the antioxidant activity of methanol soluble and non soluble phases by ultraviolet-visible (UV-Vis) and by electron paramagnetic resonance (EPR) spectroscopies. Zamora *et al.*, (2002a, 2002b) also studied the polar fractions of several edible oils with <sup>13</sup>C-NMR spectroscopy to characterize the oxidative and hydrolytic states of the samples, as well as to assure their genuineness with an improved degree of certainty.

The selective removal of the triacylglycerol base of thermoxidized oils by chromatography leads to the separation of polar and nonpolar fractions. The individual study of these fractions permits a better characterization of the chemical changes catalyzed by high temperatures. The nonpolar fraction contains the unaltered triacylglycerols and consequently it has a composition similar to the original unaltered oils. Most of the newly formed compounds during thermoxidation are expected to be included in the polar fraction so the study of this fraction reveals the most relevant chemical changes taking place. Further chromatographic analysis of the polar fraction has classified the polar compounds into 4 major groups: polymerized triacylglycerols, oxidized triacylglycerols, diacylglycerols and free fatty acids (Dobarganes *et al.*, 2000). The quantitative determination of these polar compounds by high-performance high-exclusion chromatography (HPSEC) (Marmesat *et al.*, 2007) allows for studying in detail the degradation of thermoxidized oils and the progress of oxidative, hydrolytic and thermal reactions. This information can be a useful tool for a proper interpretation of FTIR spectra.

The aim of the present study is to examine the major and minor spectral changes occurring in virgin olive oil during thermoxidation by means of FTIR spectroscopic analysis of the samples, and their polar and nonpolar fractions of the samples. Special attention is given to the FTIR spectra of polar fractions and the spectral bands that are not observed in the spectra of the oils, or they are barely noticed. The polar fractions were analyzed by HPSEC and the results served to explain the time-course change in the different bands and the chemical reactions associated with them. Special attention is given to the FTIR spectra of polar fractions and the spectral bands that are not observed in the spectra of the oils, or they are barely noticed.

## 2. MATERIALS AND METHODS

### 2.1. Sample Preparation

Forty-seven samples of thermoxidized virgin olive oil were obtained from a discontinuous domestic thermoxidation process. Commercial Spanish virgin olive oil, purchased at a local supermarket for frying purposes, was used as a model system. Although the oxidative protection of virgin olive oils greatly depends on their cultivar due to differences in composition (e.g. fatty acids, phenols, etc.), this oil was considered to be representative enough for tracking and assessing oxidation through a qualitative observation of the spectral changes. The thermoxidation process was carried out in a 4 L domestic fryer with an automatic

temperature controller (Heidolph EKT 3001, Schwabach, Germany). Three liters of virgin olive oil were heated at 190 °C for 94 hours in cycles of 8 hours per day. A sample of 40 mL was collected every 2 hours until the end of the heating process. Samples were kept in brown glass vials at 4 °C until further chemical and spectroscopic analyses. This experiment was repeated with another commercial virgin olive oil to validate the results obtained with the first sequence of samples. In this second experiment a sample was collected every 10 hours until obtaining 10 samples of 40 mL. These samples were used to validate the regression analysis in those bands that allowed for a linear mathematical modeling and to check the consistency of the trends observed for each spectral change.

### 2.2. Analysis of Total Polar Compounds

The percentage of total polar compounds (TPC) was determined gravimetrically according to the IUPAC Standard Method N° 2.507 (IUPAC, 1992). Nonpolar and polar fractions were separated from 1 g of oil by silica gel column chromatography (20 g silica adjusted to a water content of 5%, w/w) hexane/diethyl ether (90:10, v/v) and 150 mL diethyl ether as elution systems, respectively. The nonpolar fraction was eluted with 150 mL of n-hexane/diethyl ether (90:10, v/v), while the polar fraction was eluted with 150 mL of diethyl ether. Efficiency of the separation was confirmed by TLC using n-hexane/diethyl ether/acetic acid (80:20:1, v/v/v) and visualized with iodine vapor. The percentage of polar fraction was calculated by weighing both fractions after the evaporation of the solvents.

### 2.3. Distribution of Polar Compounds by High-Performance Size-Exclusion Chromatography (HPSEC)

This method determines the contents of polymerized triacylglycerols, oxidized triacylglycerols, and diacylglycerols in oils. It was applied to the polar fraction of heated virgin olive oil.

A solution of sample polar fraction in hexane (15 mg/mL) was analyzed by high performance liquid chromatography (Agilent Technologies 1200) on two columns Hewlett Packard PL gel, 30 cm×0.75 cm i.d., packed with high-performance spherical gel made of styrene-divinylbenzene co-polymer, particle size 5 µm, pore size 100 Å and 500 Å, respectively (Agilent Technol., Madrid, España), connected in a series in the order 500 Å and 100 Å, eluting with tetrahydrofuran at a flow rate of 1 mL/min. The injection volume was 10 µL. A refractive index detector (Agilent Technologies 1200) was used. The quantification was done according to Dobarganes *et al.*, 2000. The calculation is achieved assuming that all compounds of the sample are eluted. The contents

of the different groups of compounds were expressed as percentages in the oil.

#### 2.4. Fatty Acid Analysis

Total fatty acids were transmethylated according to ISO 5509. About 0.2 g of commercial virgin olive oil were dissolved in 4 mL of hexane and 0.4 mL of a solution of 2 N de KOH in methanol. The mixture was vigorously shaken for 10–15 seconds. The sample (0.2  $\mu$ L) was injected into a gas chromatograph (Varian 3900) equipped with a split-splitless injector and a flame ionization detector. An SP-2380, 60 $\times$ 0.25 mm ID and 0.25  $\mu$ m column (Supelco) was used. The injector and detector temperatures were set at 225 °C and 250 °C respectively. The oven temperature was held at 170 °C for 10 min, then programmed from 170 °C to 210 °C at 1.5 °C $\cdot$ min<sup>-1</sup> and finally held at the last temperature for 8 min. Hydrogen was used as carrier gas at a flow rate of 1.0 mL $\cdot$ min<sup>-1</sup>. Each sample was analyzed in duplicate.

The results were expressed as g fatty acid $\cdot$ 100 g<sup>-1</sup> total fatty acids (%). The results were normalized according to Dobarganes and Pérez-Camino (1988).

#### 2.5. Analysis of Phenols

A standard solution (0.5 mL) made of *p*-hydroxyphenylacetic (0.12 mg $\cdot$ mL<sup>-1</sup>) and *o*-coumaric acids (0.01 mg $\cdot$ mL<sup>-1</sup>) in methanol was added to a sample of filtered virgin olive oil (2.5 g). A rotary evaporator at 40 °C under vacuum was used to evaporate the solvent and the oily residue was dissolved in 6 mL of hexane.

The diol-bonded phase cartridge was conditioned according to Mateos *et al.*, (2001). After the sample loading, the final residue was extracted with 500  $\mu$ L of methanol-water (1:1 v/v) at 40 °C and a filtrated aliquot (20  $\mu$ L) of the final colorless solution was injected into the HPLC system (LaChrom Elite Tokio, Japan), equipped with a diode array detector. The column was a Lichrospher 100RP-18 column (4.0 mm i.d. $\times$ 250 mm; 5  $\mu$ m, particle size) (Darmstadt, Germany) maintained at 30 °C. The gradient elution, at a flow rate of 1.0 mL $\cdot$ min<sup>-1</sup>, was created using the following mobile phases: A mixture of water/phosphoric acid (95.5:0.5 v/v) (solvent A) and methanol-acetonitrile (50:50 v/v) (solvent B). The change in solvent gradient was programmed as follows: from 95% (A)–5% (B) to 70% (A)–30% (B) in 25 min; 62% (A)–38% (B) in 10 min; 62% (A)–38% (B) in 5 min; 55% (A)–45% (B) in 5 min; 47.5% (A)–52.5% (B) in 5 min and 100% (B) in 5 min, followed by 5 min of maintenance. The chromatographic signals were obtained at 235, 280, and 335 nm. The response factors and recoveries were based on the procedure developed by Mateos *et al.*, (2001).

#### 2.6. FTIR Spectroscopy

Spectral data were collected using a Bruker 55 Equinox S FTIR spectrometer equipped with a DGTS detector (Bruker Optics, Ettlingen, Germany). The sampling station was equipped with a support of transmission cell. The cell used to collect the spectra was a removable cell model 20510 (Specac, Orpington, UK), with windows of KBr and an optical path length of 150  $\mu$ m. Each spectrum was recorded at room temperature in the region of 4000–400 cm<sup>-1</sup> from an average of 22 scans at a resolution of 4 cm<sup>-1</sup>. Each sample (20  $\mu$ L) was injected into the interior of the cell using a micropipette and analyzed in duplicate. Between samples, the cell was thoroughly cleaned by using a membrane vacuum pump (Vacuubrand MD 4C NT, Wertheim, Germany). The sample was removed from the cell by aspiration and by passing hexane through the cell. The cell was later dried with nitrogen gas and a spectrum was scanned to test the cleanliness of the windows and the total absence of sample.

The spectra were manipulated with OPUS version 4.0 (Bruker Optics, Ettlingen, Germany). Peak areas were computed on the raw spectra and the results were exported as ASCII data for further statistical analysis.

The spectra of oleic, linoleic and linolenic acids and trioleine were acquired from pure standards (Sigma-Aldrich, St. Louis, MO).

#### 2.7. Statistical Analysis

Univariate and multivariate algorithms have been applied by means of Statistica version 6.0 (Statsoft, Tulsa OK). Stepwise multiple linear regression analysis (SMLRA) was the statistical procedure for the selection of IR bands that better correlate with the polar compound percentage and the percentage of the different groups of compounds identified in the polar fraction. The objective of this statistical procedure was to select the wavenumbers that could predict which kind of degradation reaction is happening in the frying oil during the process, using the TPC and the percentage of the different groups of compounds in the polar fraction as reference for the alteration degrees of the samples. The F-to-enter and F-to-remove values specified a priori determined the wavenumbers included or excluded from the model (Tabachnick and Fidell, 1983). In order to assure the significance of the results and avoid a good result by chance the F-to-Enter/Remove values were selected under the strictest conditions (Fisher-distribution table at p=0.05). The concentration of some compounds did not show a linear trend during the thermoxidation process, and for this reason no linear regression model was used in those cases.

### 3. RESULTS AND DISCUSSION

#### 3.1. FTIR spectra of neat thermoxidized virgin olive oils

The FTIR spectra of the virgin olive oils (VOOs) thermoxidized from 0 to 94 hours show remarkable changes in the bands assigned to (i) hydroxyl groups (-OH) indicative of the oxidation reactions ( $3584\text{--}3508\text{ cm}^{-1}$ ), (ii) first overtone of the stretching vibration of the ester group of triacylglycerols ( $\text{C}=\text{O}$ ) ( $3472\text{ cm}^{-1}$ ), (iii) single *cis* double bonds ( $3005\text{ cm}^{-1}$ ), (iv) changes in the band assigned to single *trans* double bonds, the band at  $967\text{ cm}^{-1}$  ( $976\text{--}963\text{ cm}^{-1}$ ). All these spectral regions provide information about the chemical structures that are involved in the complex degradation reactions occurring in the oil during the frying process. The band that better correlated with the percentage of total polar compounds (TPC) was the band assigned to single *trans* double bonds, the

band centered at  $967\text{ cm}^{-1}$  ( $976\text{--}963\text{ cm}^{-1}$ ), which was already observed in a previous work carried out with ATR (Tena *et al.*, 2009). The equation resulting from regressing TPC against the area of the single *trans* double bonds has a regression coefficient adjusted- $R^2$  of 0.99 and a standard deviation in the prediction of  $\text{RSD}=1.44\%$ :

$$\text{TPC} (\%) = -3.78 + 124.06 \times [\text{FTIR } \textit{trans} \text{ band area}]$$

The regression line points out a linear relationship when TPC is lower than 30% (Figure 1A), and the goodness of the regression model allowed for a successful validation with the other 10 thermoxidized olive oil samples collected every 10 hours from a domestic fryer for 90 hours. However, the prediction value is overestimated from 30% onwards (Figure 1B). This overestimation is due to the heterocedastic behavior of the regression, where the lower values of band areas have more error than the higher values, probably due to a problem of lack of sensitivity.

As a consequence of the formation of *trans* double bonds during thermoxidation (Tena *et al.*, 2009), the band assigned to *cis* double bonds ( $3013\text{--}2991\text{ cm}^{-1}$ ) decreases as the thermal degradation progresses. The area of this band in the initial non thermoxidized sample is higher than any value of band area of the single *trans* double bonds quantified in the rest of the samples, and it decreases up to 50% of the initial value at the end of the experiment. This band area is correlated with TPC by linear regression (adjusted- $R^2=0.98$ ;  $\text{RSD}=2.48\%$ ). However, the behavior of the band area assigned to the single *cis* double bonds is not completely linear, unlike the band area assigned to *trans* double bonds. The lack of linearity explains that the former band was not selected as a control parameter of the thermoxidation process in a previous study carried out with ATR-FTIR (Tena *et al.*, 2009). Thus, the band area of *cis* double bonds ( $3013\text{--}2991\text{ cm}^{-1}$ ) explains TPC with an adjusted- $R^2$  of 0.99 ( $\text{RSD}=0.96\%$ ) by means of the following non linear regression model:

$$\text{TPC} (\%) = 56.22 + 42.78 \times [\text{FTIR } \textit{cis} \text{ band area}] - 25.45 \times [\text{FTIR } \textit{cis} \text{ band area}]^2$$

The content of *cis* double bonds follows two approximately linear behaviors with intersection at around 45% of the total polar compounds (60 h of thermoxidation). The two lines intersect forming a five degree angle between them.

The composition of fatty acids in the samples was determined by gas chromatography to provide a chemical explanation for the observed changes in the band assigned to single *cis* double bonds. Oleic acid accounts for 75.93% of the total fatty acids present in the unaltered oil (non thermoxidized sample), while linoleic and linolenic acids account for 4.55%




FIGURE 1. Linear regression between the area of the single *trans* double bonds band ( $976\text{--}963\text{ cm}^{-1}$ ) and the percentage of total polar compounds for percentages lower than 30% (A), and for the entire experiment (B). Validation samples are labelled with the letter "v" and a number corresponding to the hours of thermoxidation.




FIGURE 2. Time-course change in the bands assigned to hydroxyl groups ( $3544\text{ cm}^{-1}$ ), hydroperoxides ( $3425\text{ cm}^{-1}$ ) and the first overtone of esters groups ( $3472\text{ cm}^{-1}$ ) during the thermoxidation process in virgin olive oil.

and 0.67%, respectively (Table 1). The relatively high percentage of oleic acid suggests that a significant amount of single *cis* double bonds, quantified by the FTIR band area ( $3013\text{--}2991\text{ cm}^{-1}$ ), is due to oleic acid. This hypothesis was verified by regressing the percentage of oleic acid using the band area assigned to single *cis* double bonds ( $3013\text{--}3991\text{ cm}^{-1}$ ). The linear model that explains the percentage of oleic acid has an adjusted- $R^2$  of 0.93 (RSD=1.84%):

$$\text{Oleic acid (\%)} = 19.95 + 22.32 \times [\text{FTIR } cis \text{ band area}]$$

In addition to *cis/trans* transformation during thermoxidation, the use of FTIR spectroscopy is also particularly useful for monitoring hydroperoxides and hydroxyl derivatives that are produced during oxidation (van de Voort *et al.*, 2001). Thus, the spectral region of  $3600\text{--}3410\text{ cm}^{-1}$  was examined to extract information about the oxidation process involved in a thermal degradation (Figure 2). This region includes the band assigned to hydroperoxide moieties due to the  $\text{--OO--H}$  stretching vibration ( $\sim 3425\text{ cm}^{-1}$ ) and the band associated to alcohol groups ( $\sim 3544\text{ cm}^{-1}$ ) (van de Voort *et al.*, 2001; Russin *et al.*, 2003; van de Voort *et al.*, 1994). The locations of these bands strongly depend on sample polarity and they gradually shift as the oxidation progresses (Russin *et al.*, 2003). The first overtone of stretching vibration of ester group  $\text{C=O}$  of triacylglycerols, whose fundamental band absorbs at  $\sim 1746\text{ cm}^{-1}$ , is also located in this region ( $\sim 3472\text{ cm}^{-1}$ ) (van de Voort *et al.*, 2001; Russin *et al.*, 2003; van de Voort *et al.*, 1994).

Although the concentration of hydroperoxides in oils undergoing thermoxidation is expected to be low due to the breakdown of these compounds at high temperatures (Velasco and Dobarganes, 2002), it is important to note that hydroperoxides may be formed shortly after taking each sample and they

are cooled at room temperature before being frozen. Furthermore, the thermoxidation process was carried out emulating a common discontinued frying process, where the oil is kept at room temperature between frying cycles and hydroperoxides are accumulated before the next frying session. Although the FTIR absorption of hydroperoxides is observable in this region, the band assigned to the first overtone of the ester groups  $\text{C=O}$  ( $\sim 3472\text{ cm}^{-1}$ ) clearly interferes in the surrounding regions and makes any study of this area difficult to carry out (Figure 2). Thus, this overlapping negatively affects the quantification of the hydroperoxide band ( $\sim 3425\text{ cm}^{-1}$ ) and the overtone of the ester group, which provides information on the hydrolysis of triacylglycerols during thermoxidation. Furthermore, the small concentrations of hydroperoxydes  $\sim 12\text{ meq of O}_2\cdot\text{kg}^{-1}$  of oil (Moya Moreno *et al.*, 1999) – prevent them from being quantified accurately (van de Voort *et al.*, 1994). The extraction of the polar and nonpolar fractions of the samples and their subsequent study by FTIR can resolve the overlapping problem of the hydroxyl region by simplifying the whole spectrum and would provide the added advantage of supplying more details about the chemical changes that take place.

### 3.2. FTIR spectra of polar and non polar fractions of thermo-oxidized virgin olive oils

Figure 3 shows the variance spectra computed from the whole set of spectra collected for the neat oil, polar and nonpolar fractions. These spectra highlight the most significant changes during the whole thermoxidation process.

Comparing the variance spectra of polar and nonpolar fractions, it is noticeable that most of the spectral changes are observed in the former (Figure 3). The near absence of relevant spectral changes in the nonpolar fraction spectrum is explained by the fact




FIGURE 3. Variance spectra computed from the whole set of spectra of 47 thermoxidation samples, polar and nonpolar fractions.




FIGURE 4. Polar and nonpolar fraction spectra in the region (3600–3200  $\text{cm}^{-1}$ ) of a thermoxidized virgin olive oil (VOO) samples.

that most of the new products generated in thermal processes are polar (Marmesat *et al.*, 2007). Thus, the compounds containing hydroxyl groups are exclusively present in the polar fraction, and the high intensity of the band assigned to hydroperoxides allows for their quantification. Furthermore, this

band is free of the interference of the overtone band associated to the ester linkage (Figure 2). Figure 4 shows the spectral region between 3600–3200  $\text{cm}^{-1}$  collected from the polar fraction and nonpolar fraction of a thermoxidized virgin olive oil sample. The spectrum of the nonpolar fraction only shows a

single band centered within  $3510\text{--}3430\text{ cm}^{-1}$ , which is due to the first overtone of the stretching vibration of the ester group in triacylglycerols (van de Voort *et al.*, 2001). By contrast, in the polar fraction, the spectrum has a higher absorption, which can be divided into three bands: (i) a weak band located within  $3510\text{--}3430\text{ cm}^{-1}$ , assigned to the first overtone of the ester group of triacylglycerols; (ii) a shoulder at higher wavenumbers assigned to hydroxyl groups of alcohols ( $\sim 3544\text{ cm}^{-1}$ ) (van de Voort *et al.*, 2001; Russin *et al.*, 2003; van de Voort *et al.*, 1994); and (iii) a shoulder at lower wavenumbers assigned to the hydroxyl groups of hydroperoxides ( $3425\text{ cm}^{-1}$ ) (van de Voort *et al.*, 2001). The hydroperoxides, already present in the non thermoxidized sample, are decomposed into secondary oxidation products during the whole thermoxidation process, reducing their concentration until 20 h of thermoxidation (14% of TPC). At this moment, the intensity of this band varies slightly with time and tends to reach a plateau.

One of the most relevant bands in the variance spectrum of the polar fraction is located around  $\sim 1167\text{ cm}^{-1}$  (Figure 3). This band is assigned to tertiary alcohols (C-O stretching) (Coates, 2000), which they can be formed as secondary oxidation products. This band rises during thermoxidation following a trend opposite to that of the hydroperoxide band (Figure 5). Thus, this band reaches a maximum intensity at 20 hours of thermoxidation (14% of TPC), probably due to the end of the hydroperoxide breakdown. This band may also be assigned to phenols ( $\sim 1200\text{ cm}^{-1}$ , C-O stretching) (Coates, 2000), although their contribution to this

band may be irrelevant since its intensity rises during the process while the concentration of phenols decreases (Table 1).

The moment in which the hydroperoxide and the tertiary alcohols reach a plateau matches with a trend change observed for the different polar chemical groups analyzed by HPSEC. Table 1 shows the percentages of TPC, polymerized triacylglycerols, oxidized triacylglycerols, and diacylglycerols throughout the experiment. Polar polymers and oxidized triacylglycerols are the main compounds produced by oxidation at high temperatures (Dobarganes and Marquez-Ruiz, 1998; Caldwell *et al.*, 2011), and they accounted for 45.61% and 21.91% in most of the thermoxidized oil. The concentration of these compounds rises more rapidly after 20 h of the process. This inflection at 20 hours is not observed for other polar compounds resulting from hydrolysis (diacylglycerols and free fatty acids), whose formation is not directly related to hydroperoxide breakdown. The observed inflection at 20 h also coincides with the moment in which the concentration of *o*-diphenols abruptly decreases to  $8.71\text{ mg}\cdot\text{kg}^{-1}$ , which means 6.5% of the initial concentration (Table 1). After that moment, the oil is hardly protected against oxidation by means of antioxidants, and the oil undergoes oxidation at a higher rate, which depends on its composition of fatty acids (Barrera-Arellano *et al.*, 2002). Similar inflections in the increasing trend of polar polymers and oxidized triacylglycerols are observed after 20 h (Table 1).

In addition to the O-H vibration observed for hydroperoxides at  $\sim 3425\text{ cm}^{-1}$ , the C-O-O stretching of peroxides and epoxy produce a weak absorption




FIGURE 5. Time-course trend in the the band assigned to hydroperoxides ( $\sim 3400\text{ cm}^{-1}$ ) and tertiary alcohols ( $\sim 1167\text{ cm}^{-1}$ ) in the spectra of polar fractions during the thermoxidation process.



TABLE 1. Percentages of total polar compounds, polar polymers, oxidized triacylglycerols, diacylglycerols, oleic, linoleic and linolenic acids, and the content of *o*-diphenols (mg/kg) in the virgin olive oil samples during the thermoxidation experiment.  
Note: TAG, Triacylglycerols; DAG, Diacylglycerols; tr, trace levels.

Thermoxidation time (hours)	Total polar compounds	Polar polymers	Oxidized TAG	DAG	C18:1	C18:2	C18:3	<i>o</i> -diphenols
0	3.5	0.2	1.4	1.3	75.9	4.6	0.7	134.1
6	5.1	1.5	1.7	1.4	72.9	4.0	0.6	70.0
10	7.1	2.7	2.5	1.5	72.4	3.9	0.5	29.8
16	9.4	3.9	3.5	1.6	72.2	3.4	0.5	11.6
20	13.0	5.9	4.9	1.8	71.4	3.4	0.5	8.7
26	18.6	8.9	7.2	2.0	70.8	3.1	0.5	7.4
30	20.8	10.9	7.9	1.6	69.5	3.0	0.5	7.2
36	31.0	16.3	12.7	2.2	69.3	2.7	0.4	tr
40	36.1	18.9	15.9	2.0	67.3	2.3	0.2	tr
46	37.9	21.6	14.5	2.6	65.2	2.0	0.2	tr
50	45.7	25.9	18.3	2.6	64.0	1.8	0.2	–
56	50.6	30.5	18.8	3.0	63.0	1.6	0.2	–
60	53.9	32.8	18.8	4.0	62.2	1.6	0.2	–
66	60.5	37.2	21.7	4.0	60.8	1.4	0.2	–
70	61.9	39.7	21.6	3.8	58.4	1.2	0.2	–
76	63.8	40.3	21.9	4.9	58.0	1.2	0.1	tr
80	64.8	42.1	22.1	4.6	55.8	1.0	0.1	–
86	66.0	42.8	22.3	5.1	53.8	0.8	0.1	–
90	67.7	44.7	22.2	6.0	51.3	0.7	tr	–
94	68.1	45.6	21.9	5.5	50.5	0.7	tr	–




FIGURE 6. Time-course change in the peak area and the absorption in the region ( $890\text{--}800\text{ cm}^{-1}$ ) assigned to peroxide and epoxy groups for the polar fraction during the thermoxidation process.

in the range of 890–800. Although this absorption is not as evident in the variance spectra as the hydroperoxide band (Figure 3), it is present in the spectra of the polar fractions and absent in the spectra for oils and their nonpolar fraction (Figure 6). Since it is a complex region characterized by weak bands along the wavenumbers of the region, the area under the spectra was considered the best parameter for studying the trend, instead of the peak height. Figure 6 shows the time-course change of the peak area and the spectra in the region under study (890–800  $\text{cm}^{-1}$ ). The area rapidly increases during the first hours of the experiment. However, unlike the trends observed for hydroperoxide and tertiary alcohols, the epoxy and peroxide band rises at a slower pace after 6 hours of thermoxidation (9% of TPC). The epoxy groups responsible for this absorption are formed during thermoxidation and they attach to the triacylglycerol molecules, resulting in oxidized monomeric triacylglycerols (Velasco *et al.*, 2004). Thus, an oxidation study carried out with methyl oleate and triolein prove the occurrence of *trans*-9,10- and *cis*-9,10-epoxystearate, while the same experiment carried out with methyl linoleate and trilinolein produced a series of monounsaturated epoxides (epoxyoleates) (Bordeaux *et al.*, 1999). In this study the epoxides coming from monounsaturated fatty acids showed a higher stability, and thereby a higher concentration.

In the variance spectrum for the polar fraction the band located at 1450  $\text{cm}^{-1}$  is also noteworthy. This band is assigned to  $\text{CH}_2$  scissoring deformation (Ismail *et al.*, 1999). However, the intensity of

this band is close to the cut-off limit (Figure 7) and consequently the signal is out of the working range for quantitative studies. On the contrary, the nearby band at 1417  $\text{cm}^{-1}$  (Figure 7) is on-scale and can be quantified. The observable magnitude of this band in the variance spectra is likely due to the hydrolysis of triacylglycerols to produce free acids, although in this case the hydrolysis process is much less significant compared to oxidation since no food was fried (Zhang *et al.*, 2012; Romero *et al.*, 1998). The absorption at 1417  $\text{cm}^{-1}$  is assigned to the out-of-plane hydroxyl group bending mode  $\delta$  (O-H---O) (Smith and Victoria, 1968). Thus, the spectrum of pure oleic acid shows this band, unlike that of their corresponding triacylglycerols (triolein) (Figure 7). This difference between the free fatty acids and their corresponding triacylglycerols has also been found for other fatty acids (linoleic and linolenic acids). Figure 6 shows the trend of this band (1417  $\text{cm}^{-1}$ ) over time. It rises for 24 hours, and then suddenly reaches stabilization.

The rise in free fatty acids is also noticeable in the variance spectrum of the polar fraction in the range of 1780–1670  $\text{cm}^{-1}$  (Figure 3). Thus, the ester linkage band is doubled into two bands. One of them is centered at 1743  $\text{cm}^{-1}$ , which is assigned to the decrease in ester linkages because of the hydrolysis of triacylglycerols. This band is off-scale and cannot be quantified. The other band is located at 1713  $\text{cm}^{-1}$  and is assigned to the carbonyl group of the carboxylic acid (Coates, 2000). Although this band is more noticeable in the spectra of the polar fraction, it is not clear that this band is only




FIGURE 7. FTIR Spectra (1600–1300  $\text{cm}^{-1}$ ) of virgin olive oil thermoxidized for 94 hours, pure triolein and oleic acid. The time-course change in the band assigned to hydroxyl groups (1417  $\text{cm}^{-1}$ ) in the polar fraction spectra during the thermoxidation process is shown as an insert.




FIGURE 8. The time-course change in the peak height at  $1713\text{ cm}^{-1}$  computed in the spectra of virgin olive oils and polar fractions during the thermoxidation process. The spectra of the oils ( $1850\text{--}1650\text{ cm}^{-1}$ ) are shown as an insert.

due to free fatty acids, since it decreases at the beginning of the process, until it starts rising from 30 hours onwards (22% of TPC) (Figure 8). The dramatic shrink of this band at the beginning of the process is probably due to the carbonyl group of ketones, which also absorb in this region ( $1725\text{--}1705\text{ cm}^{-1}$ ) (Coates, 2000), together with the thermal deterioration of the existing free fatty acids

of the unaltered oil. The increasing trend shown in the peak height computed in the spectra of the oil corresponds to the results obtained by other authors analyzing free fatty acids in frying oils (Innawong *et al.*, 2004; Casal *et al.*, 2010) as well as to the determined concentrations of diacylglycerols resulting from hydrolysis (Table 1), which rise from 1.32 % to 5.51 %.




FIGURE 9. (A) Time-course change in the band area of the first overtone of ester linkage absorption ( $3493\text{--}3452\text{ cm}^{-1}$ ) in the nonpolar fraction of the samples during the thermoxidation process. (B) Relationship between diacylglycerol content and the percentage of total polar compounds.

A reduction in the band at  $\sim 1746\text{ cm}^{-1}$  assigned to the stretching vibration of ester linkage (C=O) was also observed, which means a decline in the concentration of triacylglycerols. This band is off-scale in all the spectra because of its strong intensity. Thus, the time-course change of the ester groups of triacylglycerols has been studied through the first overtone of ester groups ( $\sim 3472\text{ cm}^{-1}$ ). Although this band interferes with the rest of absorptions of the hydroxyl region ( $3600\text{--}3410\text{ cm}^{-1}$ ) (Figure 2), in the spectra of the nonpolar fractions this band is isolated and consequently can be easily measured (Figure 4).

Figure 9A shows the trend of the band area of the first overtone of ester linkage absorption ( $3493\text{--}3452\text{ cm}^{-1}$ ) (Figure 4) in the nonpolar fraction of the samples *adversus* the percentage of total polar compounds. The band area is almost constant with a slight tendency to decrease until the moment that the oil reaches  $\sim 26\%$  of total polar compounds ( $\sim 36$  hours of thermoxidation). At this moment, the concentration of esters starts decreasing linearly. This trend matches that of the band assigned to free fatty acids in the oils ( $1713\text{ cm}^{-1}$ ) (Figure 8), whose intensity increases at a faster rate from 30 hours (TPC=22%) onwards. The decline in the ester linkage absorption associated to triacylglycerols is explained by the hydrolysis reactions that take place during thermoxidation, the principal by-products being free fatty acids and diacylglycerols. Figure 9B shows the relationship between the diacylglycerol content and the total polar compounds. The time-course change in the concentration of diacylglycerols is consistent with the decrease in intensity of the overtone band of the ester linkage band (Figure 9A), which corroborates the relationship of the latter with hydrolysis.

#### 4. CONCLUSION

The analysis of thermoxidized oils with transmission cell allows for the tracking of weak bands that evolve during thermoxidation with a sufficient signal to noise ratio. Unlike the single *trans* double bonds, the band assigned to *cis* double bonds ( $3010\text{ cm}^{-1}$ ) has a non linear trend and it can only be modeled by the quadratic regression model to predict TPC (%). Regarding the rest of the chemical species evolving during thermoxidation, their time-course change does not match the total polar compounds contents, and they show different trends depending on the chemical reaction responsible for their formation (either oxidation or hydrolysis). Since most of the chemical changes take place in the polar fraction, the FTIR analysis of polar and nonpolar fractions leads to a better interpretation of the oil spectra. Thus, the analysis of the hydroxyl region is simplified by isolating the hydroperoxide and alcohol bands in the polar fraction, and taking apart the first overtone of the ester linkage band in the nonpolar fraction. The spectra of polar fractions also provides valuable

information on the bands associated with free fatty acids, epoxy and peroxide compounds since their absorption are more intense compared to the spectra of the oil. This study improves the interpretation of the spectral changes occurring during thermoxidation of virgin olive oil, and establishes an advanced knowledge in support of the use of FTIR spectroscopy for monitoring thermoxidized oils.

#### 5. ACKNOWLEDGEMENT

This research has been partially funded by the Spanish State Secretary for Research (Ramón y Cajal Program, AGL2008-01411 and AGL2011-30371-C02-02).

#### REFERENCES

- Baeten V, Fernandez Pierna JA, Dardenne P, Meurens M, García-González DL, Aparicio-Ruiz R. 2005. Detection of the presence of hazelnut oil in olive oil by FT-Raman and FT-MIR spectroscopy. *J. Agric. Food Chem.* **53**, 6201–6206. <http://dx.doi.org/10.1021/jf050595n>
- Barrera-Arellano D, Ruiz-Méndez V, Velasco J, Márquez-Ruiz G, Dobarganes C. 2002. Loss of tocopherols and formation of degradation compounds at frying temperatures in oils differing in degree of unsaturation and natural antioxidant content. *J. Sci. Food Agric.* **82**, 1696–1702. <http://dx.doi.org/10.1002/jsfa.1245>
- Billek G. 2000. Health aspects of thermoxidized oils and fats. *Eur. J. Lipid Sci. Technol.* **102**, 587–593. [http://dx.doi.org/10.1002/1438-9312\(200009\)102:8/9<587::AID-EJLT587>3.3.CO;2-R](http://dx.doi.org/10.1002/1438-9312(200009)102:8/9<587::AID-EJLT587>3.3.CO;2-R)
- Bordeaux O, Márquez-Ruiz G, Dobarganes MC. 1999. Characterization, quantitation and evolution of mono-epoxy compounds formed in model systems of fatty acid methyl esters and monoacid triglycerides heated at high temperature. *Grasas Aceites* **50**, 53–59. <http://dx.doi.org/10.3989/gya.1999.v50.i1.636>
- Caldwell JD, Cooke BS, Greer MK. 2011. High performance liquid chromatography-size exclusion chromatography for rapid analysis of total polar compounds in used frying oils. *J. Am. Oil Chem. Soc.* **88**, 1669–1674. <http://dx.doi.org/10.1007/s11746-011-1845-5>
- Casal S, Malheiro R, Sendas A, Oliveira BPP, Pereira JA. 2010. Olive oil stability under deep-frying conditions. *Food Chem. Toxicol.* **48**, 2972–2979. <http://dx.doi.org/10.1016/j.fct.2010.07.036>
- Christophoridou S, Dais P. 2009. Detection and quantification of phenolic compounds in olive oil by high resolution  $^1\text{H}$  nuclear magnetic resonance spectroscopy. *Anal. Chim. Acta* **633**, 283–292. <http://dx.doi.org/10.1016/j.aca.2008.11.048>
- Coates J. 2000. Interpretation of infrared spectra, a practical approach. In: R.A. Meyers (Ed.) *Encyclopedia of Analytical Chemistry*. John Wiley & Sons Ltd.: Chichester, U.K. pp. 1–23.
- Dobarganes MC, Márquez-Ruiz G. 1998. Analytical evaluation of fats and oils by size-exclusion chromatography. *Analysis* **26**, 61–65. <http://dx.doi.org/10.1051/analysis:199826030061>
- Dobarganes MC, Pérez-Camino MC. 1988. Fatty acid composition: a useful tool for the determination of alteration level in heated fats. *Rev. Fr. Corps Gras* **35**, 67–70.
- Dobarganes MC, Velasco J, Dieffenbacher A. 2000. Determination of polar compounds, polymerized and oxidized triacylglycerols, and diacylglycerols in oils and fats – Results of collaborative studies and the standardized method (Technical Report). *Pure Appl. Chem.* **72**, 1563–1575. <http://dx.doi.org/10.1351/pac200072081563>

- Friedman M, Levin CE. 2008. Review of methods for the reduction of dietary content and toxicity of acrylamide. *J. Agric. Food Chem.* **56**, 6113–6140. <http://dx.doi.org/10.1021/jf0730486>
- García-González DL, Aparicio-Ruiz R, Aparicio R. 2009. Olive oil. In *Gourmet and Health-Promoting Oils*; Kamal-Eldin A, Moreaou R (Ed.) AOCS press: Champaign, IL, pp. 33–72.
- Innawong B, Mallikarjunan P, Marcy JE. 2004. The determination of frying oil quality using a chemosensory system. *Food Sci. Technol.-Leb.* **37**, 35–41.
- Ismail AA, Nicodemo A, Sedman J, van de Voort F, Holzbaure IE. 1999. Infrared spectroscopy of lipids: Principles and applications. In *Spectral Properties of Lipids*; Hamilton, RJ, Cast, J (Ed.) CRC Press: Boca Raton, FL. pp. 235–269.
- IUPAC 2.507. 1992. International Union of Pure and Applied Chemistry (IUPAC). Determination of polar compounds in frying fats. In *Standard Methods for the Analysis of Oils, Fats and Derivatives*, (7th ed.). Blackwell: Oxford, U.K.
- Li Y, García-González DL, Yu X, van de Voort FR. 2008. Determination of free fatty acids in edible oils with the use of a variable filter array IR spectrometer. *J. Am. Oil Chem. Soc.* **85**, 599–604. <http://dx.doi.org/10.1007/s11746-008-1232-z>
- Marmesat S, Velasco J, Marquez-Ruiz G, Dobarganes MC. 2007. A rapid method for determination of polar compounds in used frying fats and oils. *Grasas Aceites* **58**, 179–184.
- Mateos R, Espartero JL, Trujillo M, Ríos JJ, León-Camacho M, Alcudia F, Cert A. 2001. Determination of phenols, flavones, and lignans in virgin olive oils by solid-phase extraction and high-performance liquid chromatography with diode array ultraviolet detection. *J. Agric. Food Chem.* **49**, 2185–2192. <http://dx.doi.org/10.1021/jf0013205>
- Moya Moreno MCM, Mendoza Olivares D, Amézquita López FJ, Gimeno Adelantado JV, Bosch Reig F. 1999. Analytical evaluation of polyunsaturated fatty acids degradation during thermal oxidation of edible oils by Fourier transform infrared spectroscopy. *Talanta* **50**, 269–275. [http://dx.doi.org/10.1016/S0039-9140\(99\)00034-X](http://dx.doi.org/10.1016/S0039-9140(99)00034-X)
- Piper R. 2001. The market for fried food. In *Frying: Improving Quality*; Rossell JB. (Ed.) CRC Press: Boca Raton, FL, pp. 7–16. <http://dx.doi.org/10.1533/9781855736429.1.7>
- Romero A, Cuesta C, Sánchez-Muniz FJ. 1998. Effect of oil replenishment during deep-fat frying of frozen foods in sunflower oil and high-oleic acid sunflower oil. *J. Am. Oil Chem. Soc.* **75**, 161–167. <http://dx.doi.org/10.1007/s11746-998-0028-5>
- Russin TA, van de Voort FR, Sedman J. 2003. Novel method for rapid monitoring of lipid oxidation by FTIR spectroscopy using disposable IR cards. *J. Am. Oil Chem. Soc.* **80**, 635–641. <http://dx.doi.org/10.1007/s11746-003-0751-8>
- Sanibal EAA, Mancini-Filho J. 2004. Frying oil and fat quality measured by chemical, physical, and test kit analyses. *J. Am. Oil Chem. Soc.* **81**, 847–852. <http://dx.doi.org/10.1007/s11746-004-0990-8>
- Scano P, Rosa A, Cesare Marincola F, Locci E, Melis MP, Dessi MA, Lai A. 2008. <sup>13</sup>C NMR, GC and HPLC characterization of lipid components of the salted and dried mullet (*Mugil cephalus*) roe “bottarga”. *Chem. Phys. Lipids* **151**, 69–76. <http://dx.doi.org/10.1016/j.chemphyslip.2007.10.001>
- Sedman J, van de Voort FR, Ismail AA, Maes P. 1998. Industrial validation of FTIR trans and iodine value analyses of fats and oils. *J. Am. Oil Chem. Soc.* **75**, 33–39. <http://dx.doi.org/10.1007/s11746-998-0006-y>
- Smith JW, Victoria MC. 1968. Infrared spectroscopic investigations of acid-base interactions in aprotic solvents. Part I. The interaction of tri-n-propylamine and some carboxylic acids. *J. Chem. Soc. A*, 2468–2474. <http://dx.doi.org/10.1039/j19680002468>
- Stier RF. 2001. The measurement of frying oil quality and authenticity. In *Frying: Improving Quality*; Rossell JB (Ed.) CRC Press: Boca Raton, FL, pp. 7–16. <http://dx.doi.org/10.1533/9781855736429.2.165>
- Tabachnick BG, Fidell LS. 1983. Using multivariate Statistics. New York: Harper & Row.
- Tena N, Aparicio R, García-González DL. 2009. Thermal deterioration of virgin olive oil monitored by ATR-FTIR analysis of trans content. *J. Agric. Food Chem.* **57**, 9997–10003. <http://dx.doi.org/10.1021/jf9012828>
- Valavanidis A, Nisiotou C, Papageorgiou Y, Kremli I, Satravelas N, Zinieris N, Zygali H. 2004. Comparison of the radical scavenging potential of polar and lipidic fractions of olive oil and other vegetable oils under normal conditions and after thermal treatment. *J. Agric. Food Chem.* **52**, 2358–2365. <http://dx.doi.org/10.1021/jf030491h>
- van de Voort FR, Sedman J, Russin T. 2001. Lipid analysis by vibrational spectroscopy. *Eur. J. Lipid Sci. Technol.* **103**, 815–840. [http://dx.doi.org/10.1002/1438-9312\(200112\)103:12<815::AID-EJLT1111815>3.3.CO;2-G](http://dx.doi.org/10.1002/1438-9312(200112)103:12<815::AID-EJLT1111815>3.3.CO;2-G)
- van de Voort FR, Ghetler A, García-González DL, Li Y. 2008a. Perspectives on Quantitative Mid-FTIR spectroscopy in relation to edible oil and lubricant analysis: evolution and integration of analytical methodologies. *Food Anal. Meth.* **1**, 153–163. <http://dx.doi.org/10.1007/s12161-008-9031-6>
- van de Voort FR, Ismail AA, Sedman J, Dubois J, Nicodemo T. 1994. The determination of peroxide value by Fourier transform infrared spectroscopy. *J. Am. Oil Chem. Soc.* **71**, 921–926. <http://dx.doi.org/10.1007/BF02542254>
- van de Voort FR, Sedman J, Sherazi STH. 2008. Correcting for underlying absorption interferences in Fourier transform infrared trans analysis of edible oils using two-dimensional correlation techniques. *J. Agric. Food Chem.* **56**, 1532–1537. <http://dx.doi.org/10.1021/jf0725068>
- Velasco, J, Dobarganes, MC. 2002. Oxidative stability of virgin olive oil. *Eur. J. Lipid Sci. Technol.* **104**, 661–676. [http://dx.doi.org/10.1002/1438-9312\(200210\)104:9/10<661::AID-EJLT661>3.0.CO;2-D](http://dx.doi.org/10.1002/1438-9312(200210)104:9/10<661::AID-EJLT661>3.0.CO;2-D)
- Velasco J, Marmesat S, Márquez-Ruiz G, Dobarganes MC. 2004. Formation of short-chain glycerol-bound oxidation products and oxidised monomeric triacylglycerols during deep-frying and occurrence in used frying fats. *Eur. J. Lipid Sci. Technol.* **106**, 728–735. <http://dx.doi.org/10.1002/ejlt.200401032>
- Yu X, van de Voort FR, Sedman J. 2007. Determination of peroxide value of edible oils by FTIR spectroscopy with the use of the spectral reconstitution technique. *Talanta* **74**, 241–246. <http://dx.doi.org/10.1016/j.talanta.2007.06.004>
- Zhang Q, Saleh ASM, Chen J, Shen Q. 2012. Chemical alterations taken place during deep-fat frying based on certain reaction products: A review. *Chem. Phys. Lipids* **165**, 662–681. <http://dx.doi.org/10.1016/j.chemphyslip.2012.07.002>
- Zamora R, Gómez G, Dobarganes MC, Hidalgo FJ. 2002a. Oil fractionation as a preliminary step in the characterization of vegetable oils by high-resolution <sup>13</sup>C NMR spectroscopy. *J. Am. Oil Chem. Soc.* **79**, 261–266. <http://dx.doi.org/10.1007/s11746-002-0472-z>
- Zamora R, Gómez G, Hidalgo FJ. 2002b. Classification of vegetable oils by high-resolution <sup>13</sup>C NMR spectroscopy using chromatographically obtained oil fractions. *J. Am. Oil Chem. Soc.* **79**, 267–272. <http://dx.doi.org/10.1007/s11746-002-0472-z>