

# Polychaete-parasitizing copepods from the deep-sea Kuril-Kamchatka Trench (Pacific Ocean), with the description of a new *Ophelicola* species and comments on the currently known annelidicolous copepods

Mercedes Conradi\*, M<sup>a</sup> Eugenia Bandera\*, Ivan Marin\*\* & Daniel Martin\*\*\*

(\*) Laboratorio de Biología Marina, Departamento de Zoología, Facultad de Biología. Universidad de Sevilla. Reina Mercedes 6, 41012 - Sevilla, Spain. Fax: 954233480,

(\*\*) A.N. Severtzov Institute of Ecology and Evolution of the Russian Academy of Science, Leninsky prosp., 33, Moscow, 117071, Russia.

(\*\*\*) Centre d'Estudis Avancats de Blanes (CEAB - CSIC). Carrer d'accés a la Cala Sant Francesc 14, 17300 Blanes (Girona), Catalunya (Spain). E-mail: [dani@ceab.csic.es](mailto:dani@ceab.csic.es), Corresponding author.

## ABSTRACT

The annelid associated copepods, collectively called annelidicolous, were placed in 21 families. Some genera, such as *Ophelicola*, are considered phylogenetically isolated and are placed into the order Cyclopoida as *incertae sedis*. In this paper, we describe *Ophelicola kurambia*, the second species recorded for the genus and the first for the Pacific Ocean. The single known specimen, a female, was found during the German-Russian deep-sea expedition KuramBio at the deep-sea Kuril-Kamchatka Trench. The new species differs from *Ophelicola drachi* (known from the Gulf of Biscay, Atlantic Ocean) in being attached to the host through the mandibles instead of maxillae and, specially, in the formula of the antennular armature. The study of the new species contributes to clarify the diagnosis of the genus, which clearly differs from *Notomasticola* (another *incertae sedis* genus), and resembles both the most modified clausiids (in the mandibular shape and antennular segmentation) and the clausidiids (in the shape of maxilla). However, it does not contribute to clarify the position of *Ophelicola* within the order Cyclopoida. The paper includes a list of the known annelidicolous copepods (excluding Monstrilloidae) and summarizes the main trends shown in terms of diversity, distribution and relationships. Currently, 168 species of copepods from 74 genera and 22 families and 7 *incertae sedis* (excluding Monstrilloida) are known to be involved in 235 parasitic relationships (mostly ectoparasitic) with polychaetes. Host polychaetes include 156 species belonging to 104 genera from 22 families (plus 14 unknown). About 50% of these relationships are known from European waters, mainly from shallow depths.

**Key words:** KuramBio, Pacific Ocean, deep-sea, Copepoda, *Ophelicola*, parasitic, Polychaeta, Opheliidae.

## 1. INTRODUCTION

More than 120 species of Copepoda belonging to at least 21 families, the so called annelidicolous copepods, were reported to be associated with annelids, most of them being external or internal parasites of polychaetes. Some families include annelid symbionts together with free-living members and/or associates of other invertebrates. However, some others are known as exclusive parasites of polychaete hosts (Boxshall and Halsey, 2004; Humes, 1994).

The symbiotic relationships with polychaetes might have evolved independently from various copepod ancestors (Björnberg and Radashevsky, 2011). A comprehensive hypothesis about the relationships involving parasitic copepods has not yet been developed and therefore, placing annelidicolous species into genera and even families is often problematic (Kim et al., 2013). In fact, the definition of some families is rather nebulous and the boundaries among families are sometimes poorly defined, such as those among Clausidiidae Embleton, 1901, Clausiidae Giesbrecht, 1895 and Anomoclausidiidae Gotto, 1964 (Boxshall and Halsey, 2004; Humes and Ho, 1967; Kim et al., 2013). This also caused some genera to be phylogenetically isolated due to their unusual features

In 1978, Laubier described a new genus of annelidicolous copepod collected from an unidentified ophelid polychaete found between 4,706 and 4,475 m depth in the Atlantic coast of France. The genus *Ophelicola* Laubier, 1978 was considered as phylogenetically isolated due to its unusual features. Thus, it was placed into the order Cyclopoida as *incertae sedis* (Boxshall and Halsey, 2004).

During the German-Russian deep-sea expedition KuramBio (Kuril-Kamchatka Biodiversity Study) to the Kuril-Kamchatka trench and abyssal plain, two specimens of moderately transformed copepods associated with polychaete worms were collected. Parasitic copepods from polychaete hosts are seldom reported, likely because their very low prevalence. Usually, these parasites are found only after observing large numbers of potential hosts (Kim et al., 2013), which is a particularly difficult task in the deep-sea. Accordingly, in this paper we describe one of the specimens as a new species of *Ophelicola*, despite having found a single female, and discuss whether this new discovery provides new insights in the relationship of *Ophelicola* within the cyclopoid families. Unfortunately, the second annelidicolous copepod, belonging to the genus *Anomopsyllus* G.O. Sars, 1921, was in very poor conditions and, thus, it could not be formally described. This paper also includes a list of the known annelidicolous copepods (excluding Monstrilloidae) and summarizes all known characteristics in terms of diversity of both the symbionts and the hosts, type of relationship and bathymetrical and geographical distribution.

## 2. MATERIAL AND METHODS

The polychaete hosts were collected during the KuramBio Expedition 2012 to the Kuril-Kamchatka Trench and abyssal plain, with the help of the supranet of the epibenthic sledge EBS-S or the box corer GKG, both operated from the R/V SONNE-223. Infested host were extracted from sediments collected in stations 223-3-9 (4987 - 4991 m depth) and 2-5A (4869 m depth), carefully washed on board, photographed alive, and then fixed in 70% ethanol.

In the laboratory, the copepods were extracted from the hosts, dissected in lactic acid prior to staining with Chlorazol black E (Sigma® C-1144), examined as temporary mounts in lactophenol, and finally sealed with Entellan as permanent mounts. Drawings were made with the help of a *camera lucida* attached to a Leica DMLB differential interference microscope. Body length was measured from the anterior margin of the rostrum to the posterior margin of the caudal rami. All appendage segments and setation elements are named and numbered according to Huys and Boxshall (Huys and Boxshall, 1991).

The dissected holotype is deposited in the Museo Nacional de Ciencias Naturales of Madrid (MNCN), Spain.

## 3. RESULTS AND DISCUSSION

### 3.1. Taxonomic account

Subclass Copepoda  
Order Cyclopoida  
*Incertae sedis*

Genus *Ophelicola* Laubier, 1978

Diagnosis (redefined): Body of adult female transformed by swelling and fusion of free pedigerous somites. Prosome comprising cephalothorax incorporating 1st pedigerous somite and swollen 2nd to 4th pedigerous somites. Urosome distinct, comprising partly swollen 5th pedigerous somite fused to genital somite, and 4 free abdominal somites. Genital apertures paired, located posterolaterally on genital somite. Caudal rami with 6 setae.

Rostrum weakly developed. Antennule 5-segmented, distal 3 segments homologous, with XXI-XXIV, XXV and XXVI-XXVIII; armature 4(5), 16(14), 4+aesthetasc, 2+aesthetasc, 7+aesthetasc. Antenna uniramous, 4-segmented with coxa and basis fused to form coxobasis bearing single seta; 1st endopodal segment with 1 mid-margin seta, 2nd with 4 elements sometimes including 1 claw, 3rd with 7 elements; exopod lacking. Entognathous, with mouthparts arranged in perioral depression. Mandible small, consisting of a strongly sclerotized gnathobase with articulated distal portion denticulate or plumose. Maxillule lobate, with 5-9 setae. Maxilla 2-segmented, comprising large unarmed syncoxa and basis; basis with basal naked seta, produced into trifid claw-like process and articulated bifid claw. Maxilliped reduced, sometimes located in transverse groove on surface of cephalothorax, indistinctly 3-segmented, 1st segment unarmed, 2nd with setulose seta, and 3rd smallest, bearing short naked seta and small spine.

Swimming legs 1 to 4 ventrally on somites. Intercoxal sclerite in leg 1 only. Legs 1 to 4 biramous, with 3-segmented rami. Spine and seta formula as in Table 1.

Inner basal seta absent on leg 1. Inner coxal setae absent in all legs. 5th leg small, located laterally on somite; 2-segmented with protopodal segment more or less separate from somite and bearing outer seta: exopodal segment with 3 setae. 6th legs represented by paired opercula in female, sometimes with 1 seta. Egg sacs unknown.

Type species: *Ophelicola drachi* Laubier, 1978.

Remarks: The original description of *Ophelicola* pointed out the similarities with the family Clausidiidae in the general structure of the maxillae and swimming legs. However, *Ophelicola* lacks armature in the maxillar syncoxa, which is armed in clausidiid genera (except for *Conchylurus* Bocquet & Stock, 1957 and *Hippomolgus* G.O. Sars, 1917). Except for *Hyphalion* Humes, 1987, *Conchylurus*, and *Hermadona* Ho & Kim, 2004, most female clausidiids have well-developed 4-segmented maxillipeds (Ho and Kim, 2003), contrarily to the rudimentary limbs of *Ophelicola* females. Furthermore, the antennules of the Clausidiidae, with the exception of *Hermadona*, *Conchylirus*, and *Hersilioides* Canu 1888, are 7-segmented (Boxshall and Humes, 1987; Ho and Kim, 1990, 2003, 2004). Moreover, the derived structure of the mandible excludes *Ophelicola* from the Clausidiidae and indicates a possible relationship with the Clausiidae (Boxshall and Halsey, 2004). In fact, the mandible and the antennule segmentation of *Ophelicola* resemble that of the most modified genera of this family (such as *Boreoclausia* Kim et al. 2013, *Vivgottoia*, Kim et al. 2013, and *Sheaderia* Kim et al. 2013). However, these genera have also very reduced and modified legs 1-4, present the typical clausiid maxillule and maxilla, and have a single free abdominal segment. The clausiid genus *Spionicola* Björnberg & Radashevsky, 2009 shares the 5-segmented antennules with *Ophelicola* (Björnberg and Radashevsky, 2009), but clearly differs in all remaining characteristics. *Ophelicola* resembles the clausiid genus *Rhodinicola* Levinsen, 1878 in having 3-segmented rami of legs 1-4 and in lacking posterior median element at the basis of leg 1 (Björnberg and Radashevsky, 2011). However, most oral appendages of *Ophelicola* (i.e. mandible and maxilla) differ from the typical clausiid form (Boxshall and Halsey, 2004).

Finally, the *incertae sedis* genus *Notomasticola* Kim et al. 2013 clearly differs from *Ophelicola* in having 1-segmented abdomen, 4-segmented antennule, antennal armature,

reduced oral appendages lacking maxilliped, 2-segmented rami of legs 1-3, reduced leg 4, and 2-segmented leg 5 (Kim et al., 2013).

Therefore, nearly forty years later from its description, the genus *Ophelicola*, cannot be placed with confidence in any existing family, and should still be considered as *incertae sedis* within Cyclopoida until a full cladistic analysis of the annelidicolous families, ideally incorporating molecular data as they become available, could be carried out.

*Ophelicola kurambia* sp. nov.  
(Figures 1-4)

Material examined: MNCN 20.04/10007: 1 female holotype from KuramBio, r/v SONNE-223, station 223-3-9, gear EBS-S, date: 05-08-2012, depth 4987-4991 m, 47°14.6'N 154°42.88'E, 47°14.86'N 154°43.18'E.

Female: Body slightly transformed cyclopiform, by swelling and partial fusion of prosomal somites (Figures 1A, 1B). Total body length of female 1557  $\mu\text{m}$  (measured from anterior margin of cephalic somite to posterior margin of caudal rami on holotype in lactic acid), maximum width 586  $\mu\text{m}$ . Prosome typically comprising cephalothorax incorporating first pedigerous somite and free 2nd to 4th pedigerous somites. Prosome length/width ratio = 1.78:1 Prosome/urosome length ratio = 2.25:1. Urosome 5-segmented (Figure 1C) comprising 5th pedigerous somite, genital double-somite, and 3 free abdominal somites. 5th pedigerous somite much smaller than preceding pedigers, wider than long 92x304  $\mu\text{m}$ . Genital double somite (Figures 1A, 1C) nearly rounded and somewhat inflated, 453  $\mu\text{m}$  x 489.6  $\mu\text{m}$ , wider in middle. Genital areas located ventrolaterally at end of genital double somite. Each genital area (Figure 1F) with 1 short plumose seta. Egg sacs not seen. 3 free abdominal somites, each wider than long, 79x263, 45x226.4, 124,4x187  $\mu\text{m}$  (Figure 1C). Caudal ramus (Figures 1A, 1C), 85  $\mu\text{m}$  long, twice longer than wide, with 6 terminal setae. Outer lateral and dorsal setae naked, similar in length. Outermost and two median terminal setae broken; innermost terminal seta very small, naked. Urosome with minute setules (Figure 1C).

Rostrum (Figure 2B) broad, with truncate anterior margin. Antennule (Figure 1E) about 220  $\mu\text{m}$  long, with 5 segments measuring (along posterior, non-setiferous margin): 32 (68  $\mu\text{m}$  along anterior margin), 99, 29, 22, and 34.6  $\mu\text{m}$ , respectively. Formula for armature: 5, 14, 4 + aesthetasc, 2 + aesthetasc and 7 + aesthetasc. Except for 1 barbed seta on third segment, all setae naked.

Antenna (Figure 2A) uniramous and 4-segmented, 184  $\mu\text{m}$  long (terminal setae excluded), with 1st segment longest. 1st segment with 1 long seta, spinulose on inner margin and setulose on outer. 2nd segment with 1 weakly setulose seta, 3 setules and tiny setules close to insertion of 3rd segment. Outer corner of 3rd segment with patch of setules, inner corner with 5 elements: 1 setule, 1 barbed spine, 1 naked seta, 1 strong claw with strong curved spines near tip, and 1 long naked seta. Segment 3 smaller, articulating with segment 2 proximally, somewhat displaced on lateral side, with 7 setae ornamented as figured, 1 of them longer than total length of antenna. Labrum (Figures 2E, 2F) with patch of surface setules, with 1 pair of curved digitiform processes in middle of posterior margin, and membranous areas on each side of process, each membranous area with rounded process entirely covered with setules.

Mandible (Figure 2G) reduced and small, consisting of a strongly sclerotized gnathobase on which inserts a distal portion; articulation clearly visible. Basal part conical-shaped, with membranous flange along medial (posterior) margin. Distal portion short dagger-shaped, no sclerotized but densely plumose.

Maxillule (Figures 2C, 2D) complex, lobate but without clear distinction between lobes (1 setulose), tapering towards apex. Armature of maxillule with 10 elements, 9 setae with length and ornamentation as figured and 1 rounded element.

Maxilla (Figure 2H) 2-segmented, comprising large unarmed syncoxa. Segment 2 sclerotized with basal naked seta, produced into trifid claw-like process, articulated bifid claw-like process (widest claw with long setules on surface), and 1 naked seta.

Maxilliped (Figure 2I) small, 54  $\mu\text{m}$  long; 3-segmented with segment 1 unarmed, segment 2 with 1 spinulose seta, and segment 3 smallest, bearing 1 short naked seta and 1 small spine.

Swimming legs 1-4 (Figures 3A-C), located ventrally on somites, biramous, with 3-segmented rami. Spine and seta formula listed in Table 2.

Inner basal seta absent on leg 1. Basis with outer naked seta in legs 1-4. Both, endopodal and exopodal segments with spinules at outer corner. Setae and spines very long.

Leg 5 (Figure 1D) 2-segmented but proximal protopodal segment incorporated into somite, with 1 posterolateral seta. Free distal segment (exopod) small, nearly as long as wide, 20.3 x 27  $\mu\text{m}$ ; armed with 1 subterminal seta and 2 terminal setae, the outer, the largest.

Male: Unknown

Etymology: The specific name derives from “KuramBio”, the acronym of the expedition during which the copepod was collected. Gender feminine.

Distribution: Known only from the type locality at the Northwest Pacific, abyssal Kuril-Kamchatka Trench area.

Ecology: The observed specimen was attached to the skin of a non identified opheliid polychaete, using its left and right maxillae together as pincers.

Remarks: The genus *Ophelicola* was erected by Laubier (1978) to include a species parasitizing an opheliid polychaete, *O. drachi*, found in the abyssal plain of the Gulf of Biscay (Atlantic coast of France) at about 4.500 m depth. To date, no other species of this genus has been discovered. *Ophelicola kurambia* sp. nov. is, thus, the second known species and was also found deeper than 4,000 m but in the abyssal plain of the Kuril-Kamchatka Trench (Northwest Pacific Ocean).

Both species are ectoparasites of an unidentified opheliid, however, *O. kurambia* sp. nov. was attached to the host through its maxillae, while *O. drachi* was attached through the mandibles (Laubier, 1978).

Both species also differ in the formula for the antennular armature (5, 14, 4 + aesthetasc, 2 + aesthetasc and 7 setae + aesthetasc in *O. kurambia* sp. nov.; 4, 16, 4 + aesthetasc, 2 + aesthetasc, 7 + aesthetasc in *O. drachi*). As for the antenna, both species present the same number of elements per segment (1, 1, 4, 7), but the 3rd segment has 3 setae plus 1 claw in *O. kurambia* sp. nov. and 4 setae in *O. drachi*. The formula proposed by Boxshall and Halsey (2004) for the genus (1, 1, 3 + 1 claw, 4 + 3 claws) do not match with the two species studied. Both species show a mandible strongly sclerotized, but in the new species the gnathobase has a short dagger-shaped densely plumose, instead of the large blade distally denticulated of *O. drachi*.

The maxillule of *O. kurambia* sp. nov. has 9 setae plus 1 rounded distal element, and 1 setulose lobe tapering towards the apex, while *O. drachi* possesses 5 setae and a rounded distal lobe.

The maxilla of *O. kurambia* sp. nov. possesses 1 bifid, pincer-like element articulated at base, 1 trifold claw-like process and 2 naked setae. In turn, the maxillar distal segment of *O. drachi* presents a basal seta and a complex system of claws, one of them bifid and the other one represented by a truncated stump with a pointed process distally according to Laubier (1978), while Boxshall and Halsey (2004) described the maxilla of *Ophelicola* with a basis produced into a trifold claw-like process bearing 1 seta.

The original description of *O. drachi* points out that maxillipeds are located in a transverse groove on the surface of the cephalothorax, as confirmed by Boxshall and Halsey (2004). However, during the dissection of *O. kurambia* sp. nov., this circumstance has not been observed.

The armature formula for legs 1-4 is similar in both species. Since the table showing the legs' ornamentation (Table 1 in Laubier, 1978) contains some errors, the main discrepancies have been here inferred from the legs illustration and are detailed in Table 3. Moreover, in the Laubier (1978) description, legs 1-4 have all setae naked and all spines smooth, while the setae are plumose and the spines are spinulose in *O. kurambia* sp. nov.

Family Nereicolidae Claus, 1875

Genus *Anomopsyllus* Sars G.O., 1921

Diagnosis: Body with small cephalosome, inflated trunk and 2-segmented urosome, 4-segmented maxilliped with stout terminal segment; unmodified, simple setae on 2- or 3-segmented antenna, and swimming legs reduced or absent. Usually living in association with polychaetes (Kim et al., 2013).

*Anomopsyllus* sp.

Material examined: 1 damaged female, KuramBio expedition, station 2-5A, box corer GKG, sediment fraction 300  $\mu$ m, 4869 m depth. August 2 2012, ID. 356 associated with a non identified Ampharetidae.

Diagnosis: Specimen lacking legs. Oral area seriously damaged. Other body regions damaged too, with non-distinguishable characters.

Distribution: Northwest Pacific, abyssal Kuril-Kamchatka Trench area.

Remarks: *Anomopsyllus* is composed by 5 species: *A. hamiltonae* Kim et al., 2013 (legs 1-5 absent), *A. bifurcusi* Kim et al., 2013 and *A. geminus* Kim et al., 2013 (only legs 4 and 5 absent, the remaining vestigial), *A. abyssorum* Laubier 1988 and *A. pranizoides* Sars 1921 (Legs 1-5 vestigial, the 3 first legs more development than the two latter) (Kim et al., 2013; Laubier, 1988). Our specimen resembles *A. hamiltonae* in lacking the legs, but the damaged oral area prevented us to define the position of this copepod with respect to the known congeners.

### 3.2. Biodiversity of annelidicolous copepods (excluding Montrilloidae)

Symbiotic copepods are known to live symbiotically with virtually all marine metazoan taxa (Huys and Boxshall, 1991). However, those infesting invertebrates are relatively poorly

known in comparison with their piscicolous relatives, likely due to the economic interest of this particular group of hosts (O'Reilly, 1991). More specifically, eleven families of cyclopoid copepods are recorded exclusively from polychaete hosts, but several other families include one or more polychaete symbionts (Boxshall and Halsey, 2004).

Twenty two copepod families and seven *incertae sedis* (excluding Monstrilloida) are currently known to be involved in 235 parasitic relationships with annelid polychaetes (Table 4). These copepods belong 168 species from to 74 genera. The most representative family are the Herpyllobiidae, with 50 relationships (21.5 %), followed by Clausiidae and Nereicolidae with 37 (15.9 %) and 34 (14.6 %), respectively, Sabelliphilidae with 23 (9.9 %), and Gastrodelphyidae and Xenocoelomatidae with 13 (5.6 %). The remaining families are involved in less than 9 relationships. Most families include only one (41 %) or two (23 %) polychaete parasitic copepod genera, and there is only one, the Clausiidae, which include 17 genera. A similar pattern is shown by the species per genera, as most of them include a single polychaete parasitic species (59 %), only a 18 % include two species and the remaining 21 % include from 3 to a maximum of 17 (in *Herpyllobius* Steenstrup & Lutken, 1861) species. This apparently supports a high degree of specificity in the relationships between the annelidicolous copepods and their polychaete hosts, however the observed pattern may also be caused by the lack of adequate observations.

In turn, these copepods are associated to 156 species of polychaetes belonging to 104 genera from 22 families (plus 14 unknown polychaetes). The family most commonly found to be infested by copepods is the Polynoidae (63 relationships, 27 %), followed by Sabellidae and Terebellidae (with 30, 12.7 %, and 24, 10.2 %, relationships, respectively). The remaining polychaete families include less than 9 % of the known relationships and, even, eight of them include less than 1% (five and three families in two and one relationships, respectively). Most families include only one (35 %) or two (17 %) parasitized polychaete genera, and there is only one, the Polynoidae, which include 22 genera. This trend is even more exaggerated when analysing the number of species per genera, as 73 % include a single parasitized species and only 17 % include two species. The remaining 11 % include three or more species, and the maximum is nine in the case of the polynoid genus *Harmothoe* Kinberg, 1856. Again, the family Polynoidae included the highest number of infested species, 36, which represents about 25% of the total. Curiously enough, the Polynoidae is also the family including more symbiotic polychaete species, about 56% of the known ones, this representing about 60% of the relationships (Martin and Britayev, 1998).

Concerning the type of relationships, when reported copepods are most often parasites and only four species, have been reported as commensals (Table 4). In the case of *Bulbamphiascus imuse*, for instance, the polychaete hosts shared their tubes with 1-4 copepods each (males, females and/or copepodites) and the commensals were placed between worm and tube (Moore and O'Reilly, 1993). The parasitic annelidicolous copepods, in turn, may be ectoparasitic (i.e. living on the exterior of the host), mesoparasitic (i.e. living partly embedded in its host, usually with the anterior end forming an anchor process) or endoparasitic (i.e. living on the interior of the host). Most parasitic annelidicolous copepods are ectoparasites (158, 68 %), followed by mesoparasites (48, 20.5 %) and endoparasites (14, 6 %). Moreover, there are 12 species whose type of relationship has either not been reported or we have been unable to locate them. In some cases, the association with polychaete hosts may be circumstantial or has been just inferred, as the putative symbiotic copepods were obtained from washing of other organisms. This may be the case, for instance, of *Pseudanthessius gracilis*, *Parangium abstrusum* and *Stokella indica*, in which the association with polychaetes is assumed by comparing them with the most closely related species.

*Ophelicola kurambia* sp. nov. is a typical ectoparasitic species, which remain attached to its polychaete host with the help of the maxillae. Nothing more is known on this species

except the depth range and the geographical location of the collection site, and this is a common situation for most annelidicolous copepods, as some species remain unrecorded since their original description (often very old dated). As stated by O'Reilly (1991) a careful examination of appropriate hosts is often all that is required to rediscover these species or to find new species to be described, as in the case of *O. kurambia* sp. nov.

Like our new species, some annelidicolous copepods were also reported from very deep waters, i.e. deeper than 2000 m. Among them, there are *Trophonophila bradii*, *Gottoniella andeepi*, *Anomopsyllus abyssorum*, *Vectoriella ramosae* and the other known species of *Ophelicola*, *O. drachi* (Table 4), but *O. kurambia* sp. nov. and the damaged specimen of *Anomopsyllus* found at the Kuril-Kamchatka Trench occurred at the deepest known bottoms (almost 5000 m depth). It must be pointed out, however, that we have not been able to find references to the collection site depths for almost half of the species included in Table 4. Taking this into account, most other species ( $\approx 30$ ) are reported from shallow waters around 100 m depth or less, while the remaining 18 occur between 200 and 700 m depth). Deep waters are by far poorly studied compared with shallow ones, but the few studies addressed to these ecosystems seem to confirm that they may be very favourable to the establishment of such intimate relationships.

Concerning the biogeographical distribution, among the 233 known relationships, almost 44 % have been reported from European waters. Accordingly, more than 58 % of all them occurred in the North Atlantic Ocean (including European and American coasts). The following region including more reports of annelidicolous copepods is the Indian Ocean (including Red Sea) with about 16 %. The Antarctic Ocean, the Mediterranean Sea and the North Pacific Ocean include percentages of around 8 - 10 %, the South Pacific Ocean and the Caribbean Sea around 3 %, and the remaining locations (i.e. Black Sea, South Atlantic Ocean, and Arctic Ocean) include less than 1 % (i.e. single report each). The high number of European reports (as well as those from the North Atlantic Ocean) may likely obey to a bias caused by the fact that these coasts have been more intensively studied. Although relatively less studied, the Indian Ocean also includes a relevant percentage of reports, which confirms this area as a hot spot of biodiversity.

The precise number of copepod species is difficult to determine, while the parasites of fish have been estimated to be around 1,600 -1,800 species (Mariniello, 2010). Taking into account the wide range of hosts and the ubiquity of the invertebrate symbiotic copepods, there is no doubt that the real number of species may be as high or even higher than those associated to fish hosts, and that the annelidicolous species may substantially contribute to this number in the near future.

#### **4. ACKNOWLEDGEMENTS**

The KuranBio project was undertaken with the financial support of the PTJ (German Ministry for Science and Education), grant 03G0223A. We would like to thank Drs. Angelika Brandt and Marina Maljutina for giving us (IM) the opportunity to participate in the project and for processing the material. Thanks must also be extended to the captain and the crew of the RV "SONNE". This study is partly supported by the grants of the President of the Russian Federation MK-4481.2014.4 and the Russian Foundation of Basic Research 12-04-00540-a and 14-04-10183-k, and is a contribution of DM to the Consolidated Research Group 2014SGR120, funded by the "Generalitat de Catalunya", and to the Spanish State Research Project CTM2013-43287-P. This is the KuramBio publication # 19.



## 5. REFERENCES


- Barnard, K.H., 1948. New records and descriptions of new species of parasitic Copepoda from South Africa. *Ann. Mag. Nat. Hist.* 12, 242-254.
- Björnberg, T.K.S., Radashevsky, V.I., 2009. A new genus and a new species of Clausiidae (Crustacea, Copepoda) parasitic on *Dipolydora armata* (Polychaeta, Spionidae) in Brazil. *Pap. Av. Zool.* 49, 249-256.
- Björnberg, T.K.S., Radashevsky, V.I., 2011. A new species of *Rhodinicola* (Copepoda, Clausiidae), parasitic copepod of the shell-boring polychaete *Polydora brevipalpa* (Annelida, Spionidae) from the Sea of Japan. *Invert. Zool.* 8, 103-114.
- Bocquet, C., Bocquet-Vedrine, J., L'Hardy, J., 1968. Analyse des rapports du copépode parasite *Xenocoeloma alieni* (Brumpt) et de son hôte *Polycirrus caliendrum* Claparède. *Cah. Biol. Mar.* 9, 285-296.
- Bocquet, C., Stock, J.H., 1958. Copépodes parasites d'invertébrés des côtes de France. VI. Description de *Paranthesius myxicolae* nov. sp., copépode semi-parasite du Sabellidae *Myxicola infundibulum* (Rénier). *Proc. Koninklijke Nederl. Akad. Wetensch., Ser. C, Biol. Sci., Amsterdam* 61, 243-253.
- Bocquet, C., Stock, J.H., 1963. Copépodes parasites d'invertébrés des côtes de France. XVI. Description de *Pseudoclausia longiseta* n. sp. (Copépode Cyclopoïde, famille des Clausiidae). *Proc. Koninklijke Nederl. Akad. Wetensch., Ser. C, Biol. Sci., Amsterdam* 66, 139-152.
- Bocquet, C., Stock, J.H., Kleeton, G., 1963. Copepodes parasites d'invertebres des cotes de la Manche. X. Cyclopoïdes poecilostomes associés aux annélides polychètes dans la région de Roscoff. *Archs. Zool. Exp. Gén.* 102, 20-40.
- Boxshall, G.A., 2001. Copepoda (excl. Harpacticoida), in: Costello, M.J., Emblow, C.S., White, R.J. (Eds.), *European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification*. Muséum National d'Histoire Naturelle, Paris, pp. 252-268.
- Boxshall, G.A., Halsey, S.H., 2004. *An introduction to copepod diversity*. Ray Society, London.
- Boxshall, G., Humes, A.G., 1987. A new species of *Hemicyclops* (Copepoda: Poecilostomatoida) associated with echiuran worm in Hong Kong. *Asian Mar. Biol.*, 61-66.
- Bresciani, J., 1964. Redescription of *Rhodinicola elongata* Levinsen and description of *Rhodinicola gibbosa* sp. nov., parasitic copepods of maldanid polychaetes. *Ophelia* 1, 223-234.
- Bresciani, J., 1967. Redescription du mâle de *Selioides bolbroei* Levinsen, avec une note sur la répartition géographique du genre *Selioides* (Copepoda Cyclopoida). *Crustaceana* 13, 220-226.
- Bresciani, J., Lützen, J., 1966. The anatomy of *Aphanodomus terebellae* (Levinsen) with remarks on the sexuality of the family Xenocoelomidae nov. fam. (Parasitic Copepoda). *Bull. Mus. Nat. Hist. Nat.* 37, 787-806.
- Bresciani, J., Lützen, J., 1972. The sexuality of *Aphanodomus* (Parasitic copepod) and the phenomenon of cryptogonochorism. *Vidensk. Meddr. dansk. naturh. Foren.* 135, 7-20.
- Bresciani, J., Lützen, J., 1974a. *Melinnacheres ergasiloides* M. Sars, a parasitic copepod of the polychaete *Melinna cristata*, with notes on multiple infections caused by annelidicolous copepods. *Ophelia* 13, 31-41.
- Bresciani, J., Lützen, J., 1974b. On the biology and development of *Aphanodomus* Wilson (Xenocoelomidae), a parasitic copepod of the polychaete *Thelepus cincinnatus*. *Vidensk. Meddr. dansk. naturh. Foren.* 137, 25-63.

- Brumpt, E., 1897. Sur un copépode nouveau (*Saccopsis alleni* n. sp.) parasite de *Polycirrus aurantiacus* Grube. C. R. Acad. Sci., Paris 124, 1464-1467.
- Carton, Y., 1971. Copépodes parasites de Madagascar I.: Description de *Sabellacheres aenigmatopygus* n. sp., Copépode parasite de *Potamilla reniformis* (Polychètes, Sabellida3e). Crustaceana 21, 145-152.
- Carton, Y., Laubier, L., 1974. Description de *Selioides guineensis* sp. n., copépode cyclopoïde parasite d'aphroditidae. Archs. Zool. Exp. Gén. 115, 129-139.
- Cordell, J.R., 2007. Copepoda, in: Light, S.F. (Ed.), The Light and Smith manual: intertidal invertebrates from central California to Oregon. Univ of California Press, Oregon.
- Costanzo, G., Crescenti, N., Calafiore, N., 1996. Copepodid development of *Pseudanthessius gracilis* Claus, 1889 (Copepoda, Poecilostomatoida, Pseudanthessiidae), a copepod associated with *Hydroïdes elegans* (Haswell, 1883)(Annelida, Polychaeta) of Lake Faro (Messina, Italy) as obtained in the laboratory. Crustaceana 69, 295-305.
- Dantan, J.L., 1929. Recherches sur le *Nereicola ovata* Keferstein. Ann. Inst. Océanogr., Monaco 7, 175-197, figs. 171-120.
- Delamarre-Deboutville, C., Laubier, L., 1960. Les 'Phyllocolidae', une famille nouvelle de Copépodes parasites d'Annélides Polychètes. C. R. hebd. Séanc. Acad. Sci., Paris 251, 2083-2085.
- Delamarre-Deboutville, C., Laubier, L., 1961. Note rectificative. Vie Milieu 11, 690.
- Dogiel, V., 1908. *Entobius loimiae* ng, n. sp., eine endoparasitische Copepode. Zool. Anz. 33, 561-567.
- Dudley, P.L., 1964. Some gastrodelphyid copepods from the Pacific Coast of North America.; no. 2194. Am. Mus. Novit. 2194, 1-51.
- Gooding, R.U., 1960. North and South American copepods of the genus *Hemicyclops* (Cyclopoida: Clausidiidae). Proc. U. S. Natn. Mus. 112, 159-195.
- Gotto, R.V., 1963. Observations on the structure, affinities and biology of a rare copepod, *Eunicicola* (formerly *Eurynotus*) *insolens* (T. & A. Scott). Proc. Zool. Soc. Lond. 140, 47-56.
- Gotto, R.V., 1966. *Entobius hamondi* n. sp., a copepod associated with a terebellid worm., 11(2): 156-162. Crustaceana 11, 156-162.
- Gotto, R.V., Leahy, Y., 1988. A new annelidicolous copepod, *Cyclorhiza megalova* n. sp., with comments on its functional biology and possible phylogenetic relationships. Hydrobiologia 167-168, 533-538.
- Gravier, C., 1912a. Sur un nouveau genre de Crustacé parasite d'un Syllidien de l'Antarctique sud-américaine (*Thylacoides* nov. g. *sarsi* n. sp.). Bull. Mus. Nat. Hist. Nat., Paris 18, 71-74.
- Gravier, C., 1912b. Sur un type nouveau de Crustacé parasite d'un Serpulier de l'Antarctique sudamericaine (*Bactropus* ng *cystopomati* n. sp.). Bull. Mus. Nat. Hist. Nat., Paris 18, 67-71.
- Gravier, C., 1918a. Contribution à l'étude d'un copépode (*Flabellicola neapolitana* Gravier) parasite d'un annélide polychète [*Flabelligera* (*Siphonostoma*) *diplochaitos* (Otto)]. Publ. Staz. Zool. Napoli 2, 209-222.
- Gravier, C., 1918b. Sur un nouveau copépode (*Flabellicola* ng *neapolitana* n. sp.) parasite d'un annélide polychète [*Flabelligera diplochaitos* (Otto)]. C. R. hébd. Séanc. Acad. Sci., Paris 166, 502-505.
- Hesse, E., 1869. Observations sur les Crustacés rares ou nouveaux des côtes de France (dix-septième article). Ann. Sci.s Nat., Sér. Zool. Biol. Anim., Paris 5, 275-308.
- Ho, J., 1984. New family of poecilostomatoid copepods (Spiophanicolidae) parasitic on polychaetes from southern California, with a phylogenetic analysis of nereicoliform families. J. Crust. Biol. 4, 134-146.


- Ho, J.-S., Kim, I.H., 1990. *Hemicyclops ctenidis*, a new Poecilostomatoid copepod (Clausidiidae) associated with a Polychaete in Korea. Korean J. Zool. 33, 231-237.
- Ho, J.-S., Kim, I.H., 2003. New clausiid copepods (Poecilostomatoida) associated with polychaetes of Korea, with cladistic analysis of the family Clausiidae. J. Crust. Biol. 23, 568-581.
- Ho, J.-S., Kim, I.H., 2004. A new genus of the Clausidiidae (Copepoda: Poecilostomatoida) associated with a polychaete from Korea, with discussion of the taxonomic status of *Hersiliodes* Canu 1988. Proceedings of the Biological Society of Washington. Proc. Biol. Soc. Wash. 117, 95-105.
- Humes, A.G., 1975. Cyclopoid copepods associated with marine invertebrates in Mauritius. Zool. J. Linn. Soc. 56, 171-181.
- Humes, A.G., 1985. Poecilostomatoid copepods parasitic in the scleractinian coral genus *Goniastrea* in the Moluccas. Publs. Seto Mar. Biol. Lab. 30, 277-286.
- Humes, A.G., 1994. How many copepods? Hydrobiologia 292-293, 1-7.
- Humes, A.G., Doriji, M., 1980. A siphonostome copepod associated with a vestimentiferan from the Galapagos Rift and the East Pacific Rise. Proc. Biol. Soc. Wash. 93, 697-707.
- Humes, A.G., Grassle, J.F., 1979. *Serpulidicola josephellae* sp. nov. (Copepoda, Cyclopoida) from a deep-water polychaete west of Ireland. Crustaceana 36, 309-315.
- Humes, A.G., Ho, J.-S., 1967. New cyclopoid copepods associated with polychaete annelids in Madagascar. Bulletin of the Museum of comparative Zoology, 135(7):377-413. Bull. Mus. Comp. Zool. Harvard Coll. 135, 377-413.
- Humes, A.G., Stock, J.H., 1973. A revision of the family Lichomolgidae Kossmann, 1877, cyclopoid copepods mainly associated with marine invertebrates. Smithson. Contr. Zool. 127, 1-368.
- Huys, R., Boxshall, G., 1991. Copepod evolution. The Ray Society, London.
- Kim, I.H., 2000. Poecilostomatoid copepods from an intertidal mud flat in the Yellow Sea. J. Nat. Hist. 34, 367-432.
- Kim, I.H., 2001a. A new genus and two new species of Copepoda (Poecilostomatoida, Sabelliphilidae) associated with the tubicolous polychaetes in the Yellow Sea. Korean J. Biol. Sci. 5, 1-9.
- Kim, I.H., 2001b. A new species of *Clausia* (Copepoda, Poecilostomatoida, Clausiidae) associated with the polychaete *Arenicola brasiliensis* Nonato in Korea Hydrobiologia 452, 217-223.
- Kim, I.H., 2001c. Redescription of *Catinia plana* Bocquet and Stock, 1957 and description of two new species of *Myzomolgus* associated with the sipunculans in Korea. Korean J. Syst. Zool. 17, 71-89.
- Kim, I.H., 2005. Two new species of copepods (Crustacea) associated with the sponge *Phyllospongia foliascens* (Pallas) from the Moluccas. Integr. Biosci. 9, 229-238.
- Kim, I.H., Sikorski, A., O'Reilly, M.G., Boxshall, G.A., 2013. Copepods associated with polychaete worms in European seas. Zootaxa 3651, 1-62.
- Laubier, L., 1961. *Phyllodicola petiti* (Delamare et Laubier, 1960) et la famille des Phyllodicolidae, Copépodes parasites d'Annélides Polychètes en Méditerranée occidentale. Crustaceana 2, 228-242.
- Laubier, L., 1965. Présence de *Nereicola ovatus* Keferstein à Banyuls-sur-Mer. Données morphologiques nouvelles. Bull. Mus. Nat. Hist. Nat. 36, 631-640.
- Laubier, L., 1970. *Rhodinicola thomassini* sp. n., un nouveau Copépode parasite d'Annélides Polychètes Maldanidae de l'Océan Indien. Archs. Zool. Exp. Gén. 111, 559-571.
- Laubier, L., 1971. Description du mâle du genre *Rhodinicola* Levinsen (Copépode Clausiidae). Archs. Zool. Exp. Gén. 112, 351-359.

- Laubier, L., 1978. *Ophelicola drachi* gen. sp. n. un nouveau copépode cyclopoïde abyssal ectoparasite d'Annélides polychètes Opheliidae. Archs. Zool. Exp. Gén., 39-50.
- Laubier, L., 1988. Le genre *Anomopsyllus* Sars, 1921 copépode parasite d'annélides polychètes: *A. pranizoides* Sars, 1921 et *A. abyssorum* nov. sp. Crustaceana 55, 180-192.
- Laubier, L., Carton, Y., 1973. *Vectoriella ramosae* sp. n., un copépode parasite d'annélide polychète en Méditerranée profonde. Archs. Zool. Exp. Gén. 114, 149-158.
- López-González, P.J., Bresciani, J., Conradi, M., 2000. Two new species of *Herpyllobius* Steenstrup & Lutken, 1861 and a new record of *Herpyllobius antarcticus* Vanhoffen, 1913 (parasitic Copepoda) from the Weddell Sea, Antarctica. . Polar Biol. 23, 265-271.
- López-González, P.J., Bresciani, J., Conradi, M., 2006. New genus, three new species and new records of Herpyllobiidae Hansen, 1892 (Crustacea, Copepoda) parasites of polychaetes from Antarctica. Sci. Mar. 70, 243-259.
- Lützen, J., 1964. A revision of the family Herpyllobiidae (parasitic copepods) with notes on hosts and distribution. Ophelia 1, 241-274.
- Marchenov, A.V., 2002. Bradophilidae fam. nov. – The new family of mesoparasitic Copepod (Copepoda: Poecilostomatoida) collected from the polychaete *Brada villosa* from the White Sea. Parasitologia 36, 514-517.
- Mariniello, L., 2010. Parasitic and associated copepods. Biol.Mar.Medit. 17, 438-451.
- Martin, D., Britayev, T.A., 1998. Symbiotic polychaetes: Review of known species. Oceanogr. Mar. Biol. Ann. Rev. 36, 217-340.
- Moom, S. ., Kim, I.H., 2010. Three new species of *Hemicyclops* (Copepoda, Cyclopoida, Clausidiidae) from Korea. Korean J. Syst. Zool. 26, 279-293.
- Moore, C.G., O'Reilly, M.G., 1993. Commensalism between the polychaete, *Capitella capitata* (Fabricius), and the copepod, *Bulbamphiascus imus* (Brady)? Mar. Poll. Bull. 26, 653-654.
- O'Reilly, M.G., 1991. A guide to polychaete-infesting copepods from British waters. Porcupine Newsl. 5, 63-70.
- O'Reilly, M.G., 1995. A new genus of copepod (Copepoda: Poecilostomatoida) commensal with the maldanid polychaete *Rhodine gracilior*, with a review of the Family Clausiidae. J. Nat. Hist. 29, 47-64.
- O'Reilly, M.G., 1999. Notes on copepod parasites of polychaete worms in Scottish waters; Including the first UK records of the Californian copepod *Spiophanicola spinosus* Ho, 1984 (Poecilostomatoida: Spiophanicolidae). Glasgow Naturalist 23, 46-47.
- Sars, G.O., 1870. Bidrag til Kundskab om Christianiafjordens Fauna: Efter Forfatterens efterladte Manuskripter samlet og udgivet af hans Son G. O. Sars. Johan Dahl, Christiania.
- Sars, M., 1862. Beskrivelse med Afbildninger af fire nye parasitiske Copepoder. Beretning om et nyt lernaalignende Krebsdyr, *Sabellacheres gracilis* Sars. . Forhandl. Vidensk. Selsk. Christiana Forh. 1862, 134-143.
- Sebastian, M.J., Pillai, N.K., 1974. Two New Genera of Clausiid Copepods, *Indoclausia* and *Stockia*. Crustaceana, 80-88.
- Southward, E.C., 1964. On three new cyclopoid copepods associated with deep-water polychaetes in the north-east Atlantic. Crustaceana 6, 207-219.
- Stock, J.H., 1959. Copepoda associated with Neapolitan invertebrates. Naples, Stat. Zool. Pubbl. 31, 59-75.
- Stock, J.H., 1960. Sur quelques copépodes associés aux invertébrés des côtes du Roussillon. Crustaceana 1, 218-257.
- Stock, J.H., 1979. Serpulidicolidae, a new family of Copepoda associated with tubicolous polychaetes, with descriptions of a new genus and species from the Gulf of Mexico. Mem. Hourglass Cruises 5, 1-11.


- Stock, J.H., 1986. Cases of hyperassociation in the Copepoda (Herpyllobiidae and Nereicolidae). *Syst. Parasitol.* 8, 71-81.
- Stock, J.H., 1988. A bizarre parasitic copepod (nereicoliform Poecilostomatoida) from the Great Barrier Reef. *Tropical Zool.* 1, 217-222.
- Stock, J.H., 1989. Présence de la famille des Serpulicolidae, Copépodes Poecilostomatoida associés aux Polychètes Serpulidés, dans l'Indo-Pacifique: description d'une nouvelle espèce d'Indonésie. *Indo-Malayan Zool.* 6, 165-171.
- Stock, J.H., 1995. Two new poecilostomatoid copepods (Crustacea) associated with the polychaeta *Spirobranchus*) from The Seychelles. *J. African Zool.* 209, 77-88.
- Stock, J.H., 1996. Two new species of copepoda parasitic on polynoid polychaetes. *Crustaceana* 69, 438-445.
- Stock, J.H., Humes, A.G., Gooding, R.U., 1964. Copepoda Associated with West Indian Invertebrates. IV. The Genera *Octopicola*, *Pseudanthessius* and *Meomicola* (Cyclopoida, Lichomolgidae). *Stud. Fauna Curaçao Carib. Isl.* 18, 1-74, figs. 71-24.
- Suárez-Morales, E., Boxshall, G.A., 2012. A new species of *Sabellacheres* M. Sars, 1862 (Copepoda: Gastrodelpyidae) from a deep-water benthic polychaete in Antarctic waters, with a key to the species of the genus. *Syst. Parasitol.* 83, 65-75.
- Suarez-Morales, E., Carrera-Parra, L.F., 2012. A new species of the rare endoparasitic copepod *Entobius* (Copepoda: Entobiidae) from Mexico with a key to the species of the genus. *Folia Parasitol.* 59, 221-228.
- Walter, T.C., Boxshall, G., 2014. World of Copepods database. <http://www.marinespecies.org/copepoda/aphia.php?p=taxdetails&id=347551>.
- Williams, L.W., 1907. List of the Rhode Island Copepoda, Phyllopoda, and Ostracoda, with new species of Copepoda. *Ann. Rept. Comm. Inland Fish. Rhode Island.* 37, 69-79.
- Wilson, C.B., 1923. A new genus and species of parasitic copepod from Lower California. *Am. Mus. Novit.* 81, 1-4.
- Yáñez-Rivera, B., Suárez-Morales, E., 2008. *Pseudanthessius tortuosus* Stock, 1967 (Copepoda: Cyclopoida) from the amphinomid polychaete *Hermodice carunculata* (Pallas) in the western Caribbean. *Syst. Parasitol.* 69, 211-220.




**Figure 1.** *Ophelicola kurambia* sp. nov. (female). A, habitus dorsal; B, habitus ventral; C, urosome ventral; D, leg 5; E, antennule; F, genital area.



**Figure 2.** *Ophelicola kurambia* sp. nov. (female). A, antenna; B, rostrum; C, D, maxillule; E, labrum; F, detail of rounded lobe of labrum; G, mandible; H, maxilla; I, maxilliped.



**Figure 3.** *Ophelicola kurambia* sp. nov. (female). A, leg 1; B, leg 2; C, leg 4.



**Figure 4.** Living specimens of the parasitic copepods attached to their polychaete hosts. a – *Ophelicola kurambia* sp. nov. (female) and its host Opheliidae; b – *Anomopsyllus* sp. and its host Ampharetidae.



Table 1.- Genus *Ophelicola* (redefinition). Armature formula of legs 1 – 4.

	Coxa	Basis	Exopodal segments	Endopodal segments
Leg 1	0-0	1-0	I-0;I-1;III,I,4	0-1;0-1;II,4
Leg 2	0-0	1-0	I-0;I-1;III,I,5	0-1;0-2;II,4 (I,II,3)
Leg 3	0-0	1-0	I-0;I-1;III,I,5	0-1;0-2;II,4 (I,II,3)
Leg 4	0-0	1-0	I-0;I-1;III,I,5 (4)	0-1;0-2;III,2 (I,III,1)

Table 2.- *Ophelicola kurambia* sp. nov. Armature formula of legs 1 – 4.

	Coxa	Basis	Exopodal segments	Endopodal segments
Leg 1	0-0	1-0	I-0; I-1, III, I, 4	0-1; 0-1; II, 4
Leg 2	0-0	1-0	I-0; I-1; III, I, 5	0-1; 0-2; I, II, 3
Leg 3	0-0	1-0	I-0; I-1; III, I, 5	0-1; 0-2; I, II, 3
Leg 4	0-0	1-0	I-0; I-2; III, I, 4	0-1; 0-2; I, III, 1

Table 3.- Comparison of the armature formula for legs 1-4 according to Laubier (1978), Boxshall and Halsey (2004) and the specimen of *Ophelicola kurambia* sp. nov.; end: endopodal segment; ex: exopodal segment.

	Laubier (1978)	Boxshall & Halsey (2004)	<i>Ophelicola kurambia</i> sp. nov.
Leg 2 end.	0-1;0-2;II,4	0-1;0-2;II,4	0-1;0-2;I,II,3
Leg 3 ex.	I-0;I-1;II,II,6	I-0;I-1;III,I,5	I-0;I-1;III,I,5
Leg 3 end.	0-1;0-2;III,3	0-1;0-2;II,4	0-1;0-2;I,II,3
Leg 4 ex.	I-0;I-1; III,3,3	I-0;I-1;III,I,5	I-0;I-2;III,I,4
Leg 4 end.	0-1;0-2;III,2	0-1;0-2;III,2	0-1;0-2;I,III,1

Table 4.- List of known annelidicolous copepods (excluding monstrolloids) including the type of association, the polychaete hosts, known depth and geographical ranges of distribution and main references. Gl: *Genus inquerendum*.

Anelidicolous Copepods		Type	Polychaete hosts	Depth	Distribution	References	
Anomoclausidae	<i>Anomoclausia andrehusae</i> Gotto, 1964	?	Spionidae	<i>Pseudopolydora paucibranchiata</i> (Okuda, 1937)	180 - 300 m	Norway	Kim et al. (2013)
Bradophiliidae	<i>Bradophila pygmaea</i> Levensen, 1878	Mesoparasite	Flabelligeridae	<i>Brada villosa</i> (Rathke, 1843)		White Sea, Groenland	Marchenov (2002)
Bradophiliidae	<i>Trophonophila bradii</i> McIntosh, 1885 (Gl)	?	Flabelligeridae	<i>Ilyphagus wyvillei</i> (McIntosh, 1885)	3,566 m	Antarctic	Boxshall & Hasley (2004)
Catinidae	<i>Cotylemyzon vervoorti</i> Stock, 1882	Ectoparasite	Acetoidae	<i>Eupolydotes amboinensis</i> Malaquijn & Dehorne, 1907	40 m	Indonesia	Stock, 1981; Boxshall & Hasley (2004)
Catinidae	<i>Cotylomolgus lepidonoti</i> Humes & Ho, 1967	Ectoparasite	Unknown	Unknown		Madagascar	Boxshall & Hasley (2004)
Clausidiidae	<i>Foliomolgus cucullus</i> Kim, 2001	Ectoparasite	Eunicidae	<i>Marphysa sanguinea</i> (Montagu, 1815)	intertidal	Korea	Kim (2001b)
Clausidiidae	<i>Hersiliodes latericia</i> (Grube, 1869)	Ectoparasite	Maldanidae	<i>Praxillura longissima</i> Arwidsson, 1906		Channel and Atlantic coasts of Europe and Mediterranean Sea	O'Reilly (1995)
			Maldanidae	<i>Leiochone leiopygos</i> (Grube, 1860)		Channel and Atlantic coasts of Europe and Mediterranean Sea	O'Reilly (1995)
			Maldanidae	<i>Euclymene oerstedii</i> (Claparède, 1863)		Channel and Atlantic coasts of Europe and Mediterranean Sea	O'Reilly (1995)
Clausidiidae	<i>Goodingius adhaerens</i> (Williams, 1907)	?	Unknown	Unknown			Williams (1907)
Clausidiidae	<i>Goodingius arenicolae</i> (Gooding, 1960)	?	Arenicolidae	<i>Arenicola cristata</i> Stimpson, 1856	intertidal	USA (Massachusetts)	Gooding (1960)
Clausidiidae	<i>Hemicyclops ctenidis</i> Ho & Kim, 1990	Ectoparasite	Nereididae	<i>Neanthes japonica</i> (Izuka, 1908)	shallow	Korea, Sea of Japan, brackish lagoon	Ho & Kim (1990)
Clausidiidae	<i>Hemicyclops nausatus</i> Moom & Kim 2010	?	Unknown	Unknown		Korea	Moom & Kim (2010)
Clausidiidae	<i>Hemicyclops membranatus</i> Moom & Kim 2010	?	Unknown	Unknown		Korea	Moom & Kim (2010)
Clausidiidae	<i>Clausia lubbocki</i> Claparède, 1863	Ectoparasite	Spionidae	<i>Dipolydora</i> sp.	16 m	Europe	Kim et al. (2013)
			Maldanidae	Unknown		Europe	Kim et al. (2013)
Clausidae	<i>Indoclausia baecescui</i> Sebastian & Pillai, 1974	Ectoparasite	Maldanidae	Unknown			Sebastian & Pillai (1974)
Clausidae	<i>Likroclausia nanhaensis</i> Ho & Kim, 2003	Ectoparasite	Capitellidae	<i>Dasybranchus caducus</i> (Grube, 1846)		Kyokpo, Korea, Yellow Sea	Ho & Kim (2003), host as <i>Dasybranchus caudatus</i> ?
Clausidae	<i>Megaclausia mirabilis</i> O'Reilly, 1995	Commensal	Maldanidae	<i>Rhodine gracilior</i> Tauber, 1879	40 - 67 m	British waters of the North Sea	O'Reilly (1995)
Clausidae	<i>Mesnilia culthae</i> (T. & A. Scott, 1986)	Ectoparasite	Spionidae	<i>Dipolydora flava</i> (Claparède, 1870)	15 - 105 m	British waters	O'Reilly (1991), host as <i>Polydora flava</i> Claparède, 1870
			Spionidae	<i>Polydora ciliata</i> (Johnston, 1838)	15 - 105 m	British waters	O'Reilly (1991)
			Spionidae	<i>Dipolydora flava</i> (Claparède, 1870)	15 - 105 m	Channel coasts of France	O'Reilly (1991); Kim et al. (2013)
			Spionidae	<i>Polydora</i> sp.		British waters	O'Reilly (1991); Kim et al. (2013)
Clausidae	<i>Pontoclausia antiqua</i> (Kim, 2001)	Ectoparasite	Arenicolidae	<i>Arenicola brasiliensis</i> Nonato, 1958		Korea	Kim (2001b)
Clausidae	<i>Pontoclausia lobata</i> (Kim, 2000)	Ectoparasite	Eunicidae	<i>Marphysa sanguinea</i> (Montagu, 1815)		Yellow Sea	Kim, (2000)
Clausidae	<i>Pontoclausia prima</i> (Rocha, 1986)	Ectoparasite	Unknown	Unknown		Brazil	Kim (2001c)
Clausidae	<i>Pontoclausia wilsoni</i> (Gooding, 1963)	Ectoparasite	Unknown	Unknown			Kim (2001c)
Clausidae	<i>Pseudoclausia giesbrechti</i> Bocquet & Stock, 1963	Ectoparasite	Unknown	Unknown		France	Bocquet & Stock (1963)
Clausidae	<i>Pseudoclausia longiseta</i> Bocquet & Stock, 1963	Ectoparasite	Unknown	Unknown		France	Bocquet & Stock (1963)
Clausidae	<i>Rhodnicola elongata</i> Levensen, 1878	Ectoparasite	Maldanidae	<i>Rhodine gracilior</i> Tauber, 1879		Denmark	O'Reilly (1991, 1999)
			Maldanidae	<i>Rhodine loveni</i> Malmgren, 1865		British waters	Williams (1907)
Clausidae	<i>Rhodnicola gibbosum</i> Bresciani, 1964	Ectoparasite	Maldanidae	<i>Praxillella praetermissa</i> (Montagu, 1865)	74 - 327 m	Denmark	Bresciani (1964)
Clausidae	<i>Rhodnicola laticauda</i> Ho & Kim, 2003	Ectoparasite	Unknown	Unknown		Yellow Sea	Ho & Kim (2003)
Clausidae	<i>Rhodnicola rugosum</i> (Giesbrecht, 1897)	Ectoparasite	Maldanidae	<i>Microclymene tricirrata</i> Arwidsson, 1906	104 m	Central North Sea	Kim et al. (2013), host as <i>Chlymenura tricirrata</i> (Arwidsson, 1906)
			Maldanidae	<i>Euclymene</i> sp.	104 m	Central North Sea	Kim et al. (2013)
			Maldanidae	<i>Praxillella affinis</i> (Sars in G.O. Sars, 1872)	35 m	England, Suffolk and East Sussex	Kim et al. (2013)
			Maldanidae	<i>Leiochone johnsoni</i> McIntosh, 1915	64 m	North Sea	Kim et al. (2013), host as <i>Chlymenura johnsoni</i> (McIntosh, 1915)
Clausidae	<i>Rhodnicola tenuis</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Unknown	Unknown	69 m	Europe	Kim et al. (2013)
Clausidae	<i>Rhodnicola thomassini</i> Laubier, 1970	Ectoparasite	Maldanidae	<i>Leiochone tenuis</i> Day, 1957		Tulear Reef, Madagascar	Laubier, (1970, 1971); Ho & Kim (2003)
Clausidae	<i>Rhodnicola similis</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Maldanidae	<i>Rhodine gracilior</i> Tauber, 1879	15 - 38 m	Scotland	Kim et al. (2013)
Clausidae	<i>Rhodnicola sp.</i>	Ectoparasite	Maldanidae	<i>Chlymenura clypeata</i> (Saint-Joseph, 1894)			O'Reilly (1991)
Clausidae	<i>Rhodnicola polydora</i> Björnberg & Radashevsky, 2011	Ectoparasite	Spionidae	<i>Polydora brevipalpa</i> Zask, 1993		Peter the Great Bay, Sea of Japan	Björnberg & Radashevsky (2011)
Clausidae	<i>Boreoclausia recta</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Owenidae	<i>Galathowenia fragilis</i> (Nilson & Holthe, 1985)	350 m	Europe	Kim et al. (2013)
Clausidae	<i>Boreoclausia holmesi</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Eunicidae	<i>Myriochele danielsen</i> Hansen, 1879	11 - 178 m	Europe	Kim et al. (2013)
Clausidae	<i>Sheaderia bifida</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Maldanidae	<i>Euclymene oerstedii</i> (Claparède, 1863)	146 m	Europe	Kim et al. (2013)
Clausidae	<i>Vivgotia garwoodi</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Terebellidae	<i>Phasida aurea</i> Southward, 1956	32 m	Europe	Kim et al. (2013)
Clausidae	<i>Hemadona claviculara</i> Ho & Kim, 2004	Ectoparasite	Capitellidae	<i>Dasybranchus caducus</i> (Grube, 1846)		Namhae-do Island, Korea	Ho & Kim (2004), host as <i>Dasybranchus caudatus</i> ?
Clausidae	<i>Donusa clymenicola</i> von Nordmann, 1864	Ectoparasite	Maldanidae	<i>Nicomache lumbricalis</i> (Fabricius, 1780)		Aegean Sea	Chad & Boxshall (2014), host as <i>Clymene lumbricalis</i> Savigny in Lamarck, 1818
Clausidae	<i>Flabelligerulus inersus</i> Bresciani & Lützen, 1962	Ectoparasite	Flabelligeridae	<i>Flabelligera affinis</i> Sars, 1829		Coasts of Sweden	Chad & Boxshall (2014)
Clausidae	<i>Jeannelia</i> sp.	Ectoparasite	Maldanidae	<i>Praxillella abyssorum</i> (McIntosh, 1885)		British Sea	Chad & Boxshall (2014)
Clausidae	<i>Praxillincola kroyeri</i> McIntosh, 1885 (Gl)	?	Maldanidae	Unknown			O'Reilly (1995)
Dirivulidae	<i>Ceuthoecetes aliger</i> Humes, 1980	Ectoparasite	Siboglinidae	<i>Riftia pachyptila</i> Jones, 1981		Galapagos Rift & East Pacific Rise	Humes & Dorji (1980)
Ectinosomatidae (Harpacticoida)	??	?	Serpulidae	<i>Hydroides norvegica</i> Gunnerus, 1768		British waters	O'Reilly (1995)
Entobidae	<i>Entobius eulpsii</i> Barnard, 1948	Endoparasite	Terebellidae	Unknown		British waters; off the South African coast to Red Sea	Dogiel (1908); Barnard (1948); Gotto (1966)
Entobidae	<i>Entobius hamondi</i> Gotto, 1966	Endoparasite	Terebellidae	<i>Polycirrus calidrum</i> Claparède, 1869	12 m	Asis shoal, Plymouth	Gotto (1966)
Entobidae	<i>Entobius loimiae</i> Dogiel, 1908	?	Terebellidae	<i>Loimia medusa</i> (Savigny in Lamarck, 1818)			Dogiel (1908)
Entobidae	<i>Entobius scondides</i> Suárez-Morales & Carrera-Parra, 2012	Endoparasite	Terebellidae	<i>Scionides reticulata</i> (Elhers, 1887)		Gulf of Mexico, Caribbean Sea	Suárez-Morales & Carrera-Parra (2012)
Entobidae	<i>Entobius</i> sp.	Endoparasite	Terebellidae	<i>Polycirrus plumosus</i> (Wolfebaek, 1912)			O'Reilly (1991)
Entobidae	<i>Entobius</i> sp.	Endoparasite	Terebellidae	<i>Polycirrus medusa</i> Grube, 1850			O'Reilly (1991)
Eunicicolidae	<i>Eunicicola clausi</i> Kurz, 1877	Ectoparasite	Eunicidae	<i>Eunice torquata</i> Quatrefages, 1866		Adriatic, Norway, Faroes Island, and British waters	Kim (2005), host as <i>Eunice claparedi</i> Quatrefages, 1866
Eunicicolidae	<i>Eunicicola insolens</i> (T. & A. Scott, 1913)	Ectoparasite	Eunicidae	<i>Eunice harassii</i> Andouin & Milne Edwards, 1834		north-east coast of Ireland	Gotto (1963); Kim (2005)
Gastrodelpyidae	<i>Gastrodelpys clausii</i> Graeffe, 1883	Ectoparasite	Sabellidae	<i>Bispira volutacornis</i> (Montagu, 1804)		English Channel and Mediterranean	Boxshall & Hasley (2004)
			Sabellidae	<i>Bispira</i> sp.		Adriatic Sea	Boxshall & Hasley (2004), host as <i>Distyla josephinae</i> ?
Gastrodelpyidae	<i>Gastrodelpys dalesi</i> (Greeb, 1861)	Ectoparasite	Sabellidae	<i>Eudystilia polymorpha</i> Johnson, 1901		California coast	Boxshall & Hasley (2004)
Gastrodelpyidae	<i>Gastrodelpys fernaldi</i> Dudley, 1964	Ectoparasite	Sabellidae	<i>Bispira crassicornis</i> (Sars, 1851)		Pacific coast, USA	Boxshall & Hasley (2004)
			Sabellidae	<i>Bispira</i> sp.		Pacific coast, USA	Boxshall & Hasley (2004), host as <i>Sabella crassicornis</i> Sars, 1851
Gastrodelpyidae	<i>Gastrodelpys myxicalae</i> List, 1889	Ectoparasite	Sabellidae	<i>Myxicola infundibulum</i> (Montagu, 1808)		Adriatic Sea	Boxshall & Hasley (2004)
Gastrodelpyidae	<i>Sabellacheres aenigmatopygus</i> Carton, 1971	Ectoparasite	Sabellidae	<i>Potamilla reniformis</i> (Bruguère, 1789)		Madagascar	Carton (1971)
Gastrodelpyidae	<i>Sabellacheres antarcticus</i> Suárez-Morales & Boxshal 2012	Ectoparasite	Sabellidae	<i>Perkinsiana brigittae</i> Tovar-Hernández et al. 2012		Antarctic	Suárez-Morales & Boxshal (2012)
Gastrodelpyidae	<i>Sabellacheres drachi</i> Laubier, 1868	Ectoparasite	Sabellidae	<i>Potamilla thorelli</i> (Malmgren, 1866)		Spain	Boxshall & Hasley (2004)
Gastrodelpyidae	<i>Sabellacheres gracilis</i> Sars, 1862	Ectoparasite	Sabellidae	<i>Myxicola infundibulum</i> (Montagu, 1808)		Pacific coast, USA	Boxshall & Hasley (2004)

Gastrodelpyidae	<i>Sabellacheres illigi</i> Dudley, 1964	Ectoparasite	Sabellidae	<i>Megalomma splendida</i> (Moore, 1905)		Pacific coast, USA	Boxshall & Hasley (2004), host as <i>Branchiomma burrardum</i> Berkeley, 1930
Gastrodelpyidae	<i>Chonephilus dispar</i> Sars, 1861 (GI)	Ectoparasite	Sabellidae	<i>Parasabella rugosa</i> (Moore, 1904)		Pacific coast, USA	Boxshall & Hasley (2004), host as <i>Disilydida rugosa</i> Moore, 1904
Herpyllobiidae	<i>Eurysilenum fungosum</i> Stock, 1996	Mesoparasite	Sabellidae	<i>Pseudopotamilla ocellata</i> Moore, 1919		Pacific coast, USA	Boxshall & Hasley (2004), host as <i>Potamilla ocellata</i> ζ G. O. Sars (1870); Sars (1862); Dudley (1964)
Herpyllobiidae	<i>Eurysilenum intermedium</i> Stock, 1996	Mesoparasite	Sabellidae	<i>Euchine papillosa</i> (Sars, 1851)		North Sea, Norway	Stock (1996)
Herpyllobiidae	<i>Eurysilenum oblongum</i> Hansen, 1886	Mesoparasite	Polynoidae	<i>Hemilipidia versuysi</i> (Horst, 1915)		Iles Kai, Indonesia	Stock (1986), host as <i>Harmothoe corralophila</i> Day, 1960
Herpyllobiidae	<i>Eurysilenum truncatum</i> Sars, 1870	Mesoparasite	Polynoidae	<i>Harmothoe corralophila</i> (Day, 1960)		New Caledonia	Cordell (2007)
Herpyllobiidae	<i>Eurysilenum australis</i> López-González, Bresciani & Conradi, 2006	Mesoparasite	Polynoidae	<i>Bylgides promamme</i> (Malmgren, 1867)		North Atlantic Ocean	Cordell (2007)
Herpyllobiidae	<i>Herpyllobius antarcticus</i> Vanhöffen, 1913	Mesoparasite	Polynoidae	<i>Harmothoe imbricata</i> (Linnaeus, 1767)		North Atlantic Ocean	Cordell (2007)
			Polynoidae	<i>Gatryana cirrhosa</i> (Pallas, 1766)	190 - 286 m	North Atlantic Ocean	Cordell (2007)
			Polynoidae	<i>Harmothoe imbricata</i> (Linnaeus, 1776)		Antarctic	López-González et al. (2006)
			Polynoidae	<i>Polyeunoa sp</i>			López-González et al. (2006), host as <i>Enipo rhombigera</i> Elhers, 1908
			Polynoidae	<i>Polyeunoa laevis</i> McIntosh, 1885			López-González et al. (2006), host as <i>Harmothoe gourdoni</i> Gravier 1911
			Polynoidae	<i>Harmothoe fullo</i> (Grube, 1878)			López-González et al. (2006)
			Polynoidae	<i>Harmothoe gourdoni</i> Gravier 1911		Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Herpyllobius antepositus</i> Stock, 1986	Mesoparasite	Polynoidae	<i>Harmothoe spinosa</i> Kinberg, 1856	380 m	Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Herpyllobius arcticus</i> Steenstrup & Lütken, 1861	Mesoparasite	Polynoidae	<i>Lagisca irritans</i> Marenzeller, 1904	365 - 485 m	Crozet Island	Stock (1986); López-González et al. (2006)
			Polynoidae	<i>Austrolaenilla mollis</i> (Sars, 1872)		south-west England; Skagerrak; Faroes; Kattegat; south Norway; east and west Greenland; Kara Sea	Chad & Boxshall (2014)
			Polynoidae	<i>Gatryana cirrhosa</i> (Pallas, 1766)			O'Reilly (1991)
			Polynoidae	<i>Harmothoe extenuata</i> (Grube, 1840)			Lützen, 1964
			Polynoidae	<i>Harmothoe imbricata</i> (Linnaeus, 1776)			O'Reilly (1991); Cordell (2007)
			Phyllococidae	<i>Eumida sanguinea</i> (Oersted, 1843)			O'Reilly, 2000
			Phyllococidae	<i>Pterocirrus macroceros</i> (Grube, 1860)			O'Reilly, 2000
			Polynoidae	<i>Harmothoe impar</i> (Johnston, 1839)			O'Reilly (1991)
Herpyllobiidae	<i>Herpyllobius australis</i> Lützen, 1964	Mesoparasite	Polynoidae	<i>Polyeunoa laevis</i> McIntosh, 1885	666 - 673 m	Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Herpyllobius cordiformis</i> Lützen, 1964	Mesoparasite	Polynoidae	<i>Harmothoe spinosa</i> Kinberg, 1856		East America, Inglefield Bay, Greenland	Lützen, 1964
Herpyllobiidae	<i>Herpyllobius elongata</i> Lützen, 1967	Mesoparasite	Polynoidae	<i>Eunoe nodosa</i> (Sars, 1861)		Southern British, Columbia & Northern Washington	Lützen, 1964
			Polynoidae	<i>Grubeopolyeone tuta</i> (Grube, 1855)			Lützen, 1967, host as <i>Hololepidella tuta</i> (Grube, 1855)
Herpyllobiidae	<i>Herpyllobius gravieri</i> Lützen, 1964	Mesoparasite	Polynoidae	<i>Harmothoe spinosa</i> Kinberg, 1856	380 m	Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Herpyllobius haddoni</i> Lützen, 1964	Mesoparasite	Polynoidae	<i>Harmothoe imbricata</i> (Linnaeus, 1776)			Lützen, 1964; Cordell (2007)
Herpyllobiidae	<i>Herpyllobius hartmanae</i> Lützen & Jones, 1976	Mesoparasite	Polynoidae	<i>Laetmonice producta</i> (Grube, 1876)	476 - 496 m	Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Herpyllobius lobosaccus</i> Stock, 1986	Mesoparasite	Polynoidae	<i>Lagisca irritans</i> Marenzeller, 1904	365 - 485 m	Crozet Island	Stock (1986); López-González et al. (2006)
Herpyllobiidae	<i>Herpyllobius luetzeni</i> López-González, Bresciani & Conradi, 2000	Mesoparasite	Polynoidae	<i>Harmothoe cf spinosa</i> Kinberg, 1856	93 - 94 m	Antarctic	López-González et al. (2000)
Herpyllobiidae	<i>Herpyllobius nipponicus</i> Lützen, 1964	Mesoparasite	Polynoidae	<i>Paralosalysdina pleioplepis</i> (Marenzeller, 1879)		East America, Kara sea, Greenland, Faroes Isles, Noway, England	Lützen, 1964
Herpyllobiidae	<i>Herpyllobius polastermi</i> López-González, Bresciani & Conradi, 2000	Mesoparasite	Polynoidae	<i>Eulagisca gigantea</i> Monro, 1939	391 - 673 m	Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Herpyllobius polyneoes</i> (Kroyer, 1863)	Mesoparasite	Polynoidae	<i>Austrolaenilla mollis</i> (Sars, 1878)		British Isles, North Sea, Skagerrak, North Noway, Kattegat, Iceland, Spitzbergen, east and west Greenland, Baffin Island, Labrador, Gulf of St Lawrence, northeast USA, Kara sea, Alaska, northeast Pacific	O'Reilly (1991), Chad & Boxshall (2014)
			Polynoidae	<i>Bylgides promamme</i> (Malmgren, 1867)		European waters	Lützen (1964), host as <i>Anthoche badia</i> (Théel, 1879)
			Polynoidae	<i>Bylgides sarsi</i> (Kinberg in Malmgren, 1866)		European waters	Lützen (1964)
			Polynoidae	<i>Eunoe nodosa</i> (Sars, 1861)		European waters	Lützen (1964)
			Polynoidae	<i>Gatryana ammodseni</i> (Malmgren, 1867)		European waters	Lützen (1964)
			Polynoidae	<i>Gatryana cirrhosa</i> (Pallas, 1766)		Scotland	Lützen (1964); O'Reilly (O'Reilly, 1999)
			Polynoidae	<i>Gaudichaudius iphionellioides</i> (Johnson, 1901)		European waters	Lützen (1964)
			Polynoidae	<i>Harmothoe aspera</i> (Hansen, 1878)		European waters	Lützen (1964)
			Polynoidae	<i>Harmothoe extenuata</i> (Grube, 1840)		European waters	Lützen (1964)
			Polynoidae	<i>Harmothoe imbricata</i> (Linnaeus, 1767)		European waters	Lützen (1964); Cordell (2007)
			Polynoidae	<i>Harmothoe impar</i> (Johnston, 1839)		European waters	Lützen (1964)
			Polynoidae	<i>Malmgreniella lunulata</i> (Delle Chiaje, 1830)		British waters	O'Reilly (1991)
			Polynoidae	<i>Malmgrenia andreapolis</i> McIntosh, 1874		British waters	O'Reilly (1991)
Herpyllobiidae	<i>Herpyllobius rotundus</i> Lützen & Jones, 1976	Mesoparasite	Polynoidae	<i>Harmothoe sp.</i>	640 - 658 m	Cook Strait	López-González et al. (2006)
Herpyllobiidae	<i>Herpyllobius stocki</i> López-González, Bresciani & Conradi, 2000	Mesoparasite	Polynoidae	<i>Austrolaenilla antarctica</i> Bergström, 1916	395 - 417 m	Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Herpyllobius vanhoffeni</i> López-González, Bresciani & Conradi, 2000	Mesoparasite	Polynoidae	<i>Eulagisca corrientis</i> McIntosh, 1885	666 - 673 m	Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Phallusiella psalidota</i> Leigh-Sharpe, 1926	Mesoparasite	Polynoidae	<i>Harmothoe extenuata</i> (Grube, 1840)		Southwest England	Chad & Boxshall (2014)
Herpyllobiidae	<i>Phallusiella vera</i> Leigh-Sharpe, 1926	Mesoparasite	Polynoidae	<i>Malmgreniella castanea</i> (McIntosh, 1876)			O'Reilly (1991), host as <i>Malmgrenia castanea</i> McIntosh, 1876 and <i>Harmothoe castanea</i> (McIntosh, 1876)
Herpyllobiidae	<i>Gotoniella antarctica</i> López-González, Bresciani & Conradi, 2006	?	Polynoidae	Unknown	374 - 597 m	Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Gotoniella andeepi</i> López-González, Bresciani & Conradi, 2006	?	Polynoidae	Unknown	2,895 m	Antarctic	López-González et al. (2006)
Herpyllobiidae	<i>Thylacoides sarsi</i> Gravier, 1912	?	Syllidae	Unknown		Antarctic	López-González et al. (2006)
							Gravier (1912a)
Miracidae (Harpacticoida)	<i>Bulbamphiascus imus</i> (Brady, 1872)	Commensal	Capitellidae	<i>Capitella sp.</i>		Irvine Bay, Scotland	Moore & O'Reilly (1993), host as <i>Capitella capitata</i> (Fabricius, 1780)
Nereicolidae	<i>Anomopsyllus abyssorum</i> Laubier, 1968	Ectoparasite	Ampharetidae	Unknown	3,992 m	Gulf of Gasconne	Laubier (1988)
Nereicolidae	<i>Anomopsyllus pranztoi</i> Sars, 1921	Ectoparasite	Ampharetidae	<i>Amphictetes gunneri</i> (Sars, 1835)	300 m	Norway	Laubier (1988)
Nereicolidae	<i>Anomopsyllus bifurcus</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Capitellidae	<i>Notomastus latericius</i> Sars, 1851	105 m	Norway	Kim et al. (2013)
Nereicolidae	<i>Anomopsyllus geminus</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Ampharetidae	<i>Ampharete lindstroemi</i> Malmgren, 1867	45 - 112 m	Norway	Kim et al. (2013)
Nereicolidae	<i>Anomopsyllus hamiltonae</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Ampharetidae	<i>Sasane wahrbergi</i> (Eliasson, 1955)	290 m	North Sea, Norway	Kim et al. (2013)
Nereicolidae	<i>Anomopsyllus sp.</i>	Ectoparasite	Ampharetidae	Unknown	4,869 m	Kurile Kamchatka Trench and abyssal plain	This paper
Nereicolidae	<i>Nereicola ovata</i> Keferstein, 1863	Ectoparasite	Nereididae	<i>Nereis eltoralis</i> Eliason, 1962		English Channel, Mediterranean and Black Seas	Dantan (1929); Laubier (1965); O'Reilly (1995)
			Nereididae	<i>Nereis rava</i> Ehlers, 1864		Northsouth Atlantic, North Pacific, Antarctic	Boxshall & Hasley (2004)
			Nereididae	<i>Nereis zonata</i> Malmgren, 1867			Boxshall & Hasley (2004)
			Nereididae	<i>Perinereis cultrifera</i> (Grube, 1840)		North Sea, Norway	Boxshall & Hasley (2004)
			Nereididae	<i>Platynereis dumerilii</i> (Audouin & Milne Edwards, 1834)			Boxshall & Hasley (2004)
Nereicolidae	<i>Pherma curticaudatum</i> Wilson, C. B. 1923	Ectoparasite	Unknown	Unknown		Tortugas Islands, Gulf of Mexico	Wilson (1923)
Nereicolidae	<i>Selioides boqueti</i> Carton, 1963	Ectoparasite	Polynoidae	<i>Alyte assimilis</i> (McIntosh, 1874)			Boxshall & Hasley (2004), host as <i>Subalyte assimilis</i> ?
			Polynoidae	<i>Gatryana cirrhosa</i> (Pallas, 1766)		Swedish and British coasts	O'Reilly (1995)
			Polynoidae	<i>Harmothoe sp.</i>		British waters	O'Reilly (1995)
			Polynoidae	<i>Malmgreniella castanea</i> (McIntosh, 1876)		British waters	O'Reilly (1995)

Nereicoidae	<i>Selioides bolbroei</i> Linvsen, 1878	Ectoparasite	Polynoïdae Polynoïdae Polynoïdae	<i>Bylgides sarsi</i> (Kinberg in Malmgren, 1866) <i>Eunoe nodosa</i> (Sars, 1861) <i>Gatryana cirrhosa</i> (Pallas, 1766)		British waters, Arctic, Denmark, Sweden and eastern North Sea	Bresciani (1967), host as <i>Anthinoe sarsi</i> (Th�el, 1879) Bresciani (1967) O'Reilly (1995)
Nereicoidae	<i>Selioides bulbifer</i> Stock, 1986	Ectoparasite	Polynoïdae	<i>Harmothoe imbricata</i> (Linnaeus, 1767)		Denmark	Bresciani (1967)
Nereicoidae	<i>Selioides capensis</i> Stock, 1986	Ectoparasite	Polynoïdae	<i>Lagisca rarisipina</i> (Sars, 1861)	30 m	Indian Ocean	Bresciani (1967)
Nereicoidae	<i>Selioides guineensis</i> Carton & Laubier, 1974	Ectoparasite	Polynoïdae	<i>Gorgoniapolyone corralophila</i> (Day, 1960)		Sudafrica	Stock (1986), host as <i>Harmothoe corralophila</i> Day, 1960
Nereicoidae	<i>Selioides tardus</i> Gravis, 1912	Ectoparasite	Polynoïdae	<i>Subadyte pellicuda</i> (Ehlers, 1864)		Golfe Guin�e (Atlantic)	Stock (1996)
Nereicoidae	<i>Selioides bilobus</i> Kroyer, 1837	Ectoparasite	Polynoïdae	<i>Subadyte sp.</i>		Antarctic	Carton & Laubier (1974)
Nereicoidae	<i>Sigecheres brittae</i> Bresciani, 1964	Ectoparasite	Phyllocodiidae	<i>Lepidonotus squamatus</i> (Linnaeus, 1758)		Kattegat	Carton & Laubier (1974), host as <i>Hermadion rouchi</i> Gravier, 1911 and <i>Hermadion ferox</i> Baird, 1865
Nereicoidae	<i>Sigecheres conica</i> (T. Scott, 1902)	Ectoparasite	Phyllocodiidae	<i>Eulalia viridis</i> Malmgren, 1865	102 m	Denmark and Sweden	Bresciani (1967)
Nereicoidae	<i>Vectoriella marinovi</i> Stock, 1968	Ectoparasite	Paraonidae	<i>Sige fusigera</i> (Linnaeus, 1767)	12-15 m	Norway	O'Reilly (1991); Kim et al. (2013)
Nereicoidae	<i>Vectoriella ramosae</i> Laubier & Carton, 1973	Ectoparasite	Paraonidae	<i>Aricidea (Acmira) cerrutti</i> Laubier, 1966		Black Sea	O'Reilly (1991); Kim et al. (2013)
Nereicoidae	<i>Vectoriella gabesensis</i> Kim, Sikosky, O'Reilly & Boxhall, 2013	Ectoparasite	Paraonidae	<i>Aedicira mediterranea</i> Laubier & Ramos, 1974	2,000 – 3,000 m	Mediterranean	Laubier & Carton (1973), host as <i>Aricidea jeffreysi</i> [Auctt. (Non McIntosh, 1879)]
Nereicoidae	<i>Chelonidiformis typicus</i> Hesse, 1869 (GI)	Ectoparasite	Unknown	<i>Aricidea catherinae</i> Laubier, 1967	13 m	Mediterranean	Laubier & Carton (1973)
Nereicoidae	<i>Leaniricula roundata</i> McIntosh, 1885	Ectoparasite	Sigalionidae	Unknown			Kim et al. (2013)
Nereicoidae	<i>Octophora lacertae</i> Stock, 1988	Ectoparasite	Serpulidae	<i>Sthenolepis areolata</i> (McIntosh, 1885)		low tidal	Hesse (1869)
Phyllocodiidae	<i>Phyllocicola petiti</i> Delamare Deboutteville & Laubier, 1961	Ectoparasite	Phyllocodiidae	<i>Eulalia expusilla</i> Pleijel, 1987		Great Barrier Reef	Boxshall & Hasley (2004), host as <i>Leanira areolata</i> McIntosh, 1885
		Ectoparasite	Phyllocodiidae	<i>Eumida bahusiensis</i> Bergstrom, 1914		Mediterranean	Stock (1988)
		Ectoparasite	Phyllocodiidae	<i>Eumida sanguinea</i> (Oersted, 1843)		Mediterranean	Delamare-Deboutteville & Laubier (1961), host as <i>Eulalia pusilla</i> Oerstedt, 1843
		Ectoparasite	Phyllocodiidae	<i>Phyllococe sp.</i>		Mediterranean	Delamare-Deboutteville & Laubier (1960, 1961); Laubier (1961)
		Ectoparasite	Phyllocodiidae	<i>Phyllococe rosea</i> (McIntosh, 1877)		Mediterranean	Delamare-Deboutteville & Laubier (1960, 1961); Laubier (1961)
		Ectoparasite	Phyllocodiidae	<i>Pirakia punctifera</i> (Grube, 1860)		Mediterranean	Delamare-Deboutteville & Laubier (1960, 1961); Laubier (1961)
Pseudanthessidae	<i>Spiranthessius pleurocephalus</i> Stock, 1995	Ectoparasite	Serpulidae	<i>Spirobranchus corniculatus</i> (Grube, 1862)		Seychelles	Stock (1995)
Pseudanthessidae	<i>Pseudanthessius tortuosus</i> Stock, Humes & Gooding, 1964	Ectoparasite	Amphinomidae	<i>Hermodice carunculata</i> (Pallas, 1766)		US Virgin Islands; western Caribbean and in Mexican waters	Stock et al. (1964); Y�nez-Rivera et al. (2008)
Pseudanthessidae	<i>Pseudanthessius gracilis</i> Claus, 1889	Ectoparasite	Serpulidae	<i>Hydroides elegans</i> (Haswell, 1883)		Scotland, Italy, Sri Lanka, England, sweden, Norway	Costanzo et al. (1996)
Pseudanthessidae	<i>Pseudanthessius aestheticus</i> Stock, Humes & Gooding, 1964	Ectoparasite	Serpulidae	<i>Filograna sp.</i>		British waters	O'Reilly (1991)
Pseudanthessidae	<i>Pseudanthessius ferox</i> Humes & Ho, 1967	Ectoparasite	Amphinomidae	<i>Spirobranchus triqueter</i> (Linnaeus, 1758)		British waters	O'Reilly (1991), host as <i>Pomatoceros triqueter</i> (Linnaeus, 1758)
Sabelliphilidae	<i>Acaenomolgus gotoi</i> Stock, 1995	Ectoparasite	Sabellidae	<i>Hermodice carunculata</i> (Pallas, 1766)		Jamaica	Stock et al. (1964)
Sabelliphilidae	<i>Acaenomolgus protulae</i> (Stock, 1959)	Ectoparasite	Serpulidae	<i>Sabella fusca</i> Grube, 1870		Indian Ocean, Red Sea	Humes & Ho (1967)
Sabelliphilidae	<i>Acaenomolgus serpulae</i> (Stock, 1960)	Ectoparasite	Serpulidae	<i>Spirobranchus sp.</i>		Desroches Atoll, Seychelles	Stock (1995)
Sabelliphilidae	<i>Doridicola hirsutipes</i> (T. Scot, 1893)	Ectoparasite	Serpulidae	<i>Protella intestinum</i> (Lamarck, 1818)		Naples, Italy; Banyuls, France; Strangford Lough, Northern Ireland	Stock (1959)
Sabelliphilidae	<i>Doridicola agilis</i> Leydig, 1853	Ectoparasite	Serpulidae	<i>Serpula vermicularis</i> Linnaeus, 1767		Banuyls, France	Stock (1960)
Sabelliphilidae	<i>Eupolympniphilus finnarchicus</i> (Scott, 1903)	Ectoparasite	Sabellidae	<i>Sabella sp.</i>		British waters	O'Reilly (1991)
Sabelliphilidae	<i>Myxomolgoides mauritanus</i> Humes, 1975	Ectoparasite	Polynoïdae	<i>Polynoe sp.</i>		English Channel	Bocquet et al. (1963)
Sabelliphilidae	<i>Myxomolgus invulgas</i> Kim, 2001	Ectoparasite	Terebellidae	<i>Eupolymnia nebulosa</i> (Montagu, 1818)		Roscoff	O'Reilly (1991)
Sabelliphilidae	<i>Myxomolgus myxocolae</i> (Bocquet & Stock, 1958)	Ectoparasite	Sabellidae	<i>Sabellastarte magnifica</i> (Shaw, 1800)		Yellow sea	Humes (1975)
Sabelliphilidae	<i>Myxomolgus proximus</i> Humes & Stock, 1973	Commensal	Sabellidae	<i>Myxicola sp.</i>		Plymouth; Roscoff	Kim (2001a)
Sabelliphilidae	<i>Nasomolgus firmus</i> Humes & Ho, 1967	Commensal	Sabellidae	<i>Myxicola infundibulum</i> (Montagu, 1808)		Finisterre, France	Bocquet & Stock (1958)
Sabelliphilidae	<i>Nasomolgus leptus</i> Humes & Ho, 1967	Ectoparasite	Sabellidae	<i>Myxicola aesthetica</i> (Clapar�de, 1870)		Nosy Be, Madagascar	Humes & Stock (1973)
Sabelliphilidae	<i>Nasomolgus parvulus</i> Humes & Ho, 1967	Ectoparasite	Sabellidae	<i>Sabellastarte spectabilis</i> (Grube, 1878)		Nosy Be, Madagascar	Humes & Ho (1967)
Sabelliphilidae	<i>Nasomolgus rudis</i> Humes & Ho, 1967	Ectoparasite	Sabellidae	<i>Sabellastarte magnifica</i> (Shaw, 1800)		Nosy Be, Madagascar	Humes & Ho (1967)
Sabelliphilidae	<i>Sabelliphilus elongatus</i> Sars, 1962	Ectoparasite	Sabellidae	<i>Sabellastarte magnifica</i> (Shaw, 1800)		Nosy Be, Madagascar	Humes & Ho (1967)
Sabelliphilidae	<i>Sabelliphilus sarsi</i> Clapar�de, 1870	Ectoparasite	Sabellidae	<i>Sabella spallanzanii</i> (Gmelin, 1791)		Norway, Sweden, Northern Ireland, Ireland, England, France, Italy, Mediterranean coast of France, Northwestern Spain	O'Reilly (1995), host as <i>Sabella sarsi</i> Kroyer, 1856
Sabelliphilidae	<i>Sabelliphilus bispirae</i> McIntosh, 1904 (GI)	Ectoparasite	Sabellidae	<i>Sabella pavonina</i> Savigny, 1822		Saint George Chanel; British and European waters	O'Reilly (1995)
Sabelliphilidae	<i>Sabelliphilus leuckarti</i> Kossmann, 1877 (GI)	Ectoparasite	Sabellidae	<i>Sabella spallanzanii</i> (Gmelin, 1791)		British waters to Mediterranean Sea	O'Reilly (1995), host as <i>Spirographis spallanzanii</i> (Viviani, 1805)
Sabelliphilidae	<i>Serpuliphilus duplus</i> Humes & Stock, 1973	Ectoparasite	Serpulidae	<i>Bispira volutacornis</i> (Montagu, 1804)			Humes (1975)
Sabelliphilidae	<i>Serpuliphilus tenax</i> Humes & Stock, 1973	Ectoparasite	Serpulidae	<i>Sabella sp.</i>			Humes (1975)
Sabelliphilidae	<i>Terebelliphilus simplex</i> Kim, 2001	Ectoparasite	Serpulidae	<i>Pomatostegus stellatus</i> (Abildgaard, 1789)		Curacao	Humes & Stock (1973)
Saccopsidae	<i>Melinnacheres ergasiloides</i> Sars, 1870	Ectoparasite	Terebellidae	<i>Spirobranchus giganteus</i> (Pallas, 1766)		Curacao, Bonaire, Puerto Rico, Jamaica, Barbados, Bahamas	Humes & Stock (1973)
Saccopsidae	<i>Melinnacheres leviseni</i> (McIntosh, 1885)	Ectoparasite	Terebellidae	<i>Terebella ehrenbergi</i> Gravier, 1906		Yellow Sea	Kim (2001a)
Saccopsidae	<i>Melinnacheres terebellidis</i> (Levinson, 1878)	Ectoparasite	Ampharetidae	<i>Melinna cristata</i> (Sars, 1851)		Scandinavian waters, Massachusetts	Bresciani & L�tzen (1974a)
Saccopsidae	<i>Melinnacheres steenstrupi</i> (Bresciani & L�tzen, 1961)	Ectoparasite	Terebellidae	<i>Ehlersiella atlantica</i> McIntosh, 1885		Atlantic Ocean, between Bermuda and Azores	Bresciani & L�tzen (1974a)
Saccopsidae	<i>Melinnacheres terebellidis</i> (Levinson, 1878)	Ectoparasite	Trichobranchidae	<i>Terebellides stroemi</i> Sars, 1835		off Greenland and northern North America	Bresciani & L�tzen (1974a)
Saccopsidae	<i>Melinnacheres steenstrupi</i> (Bresciani & L�tzen, 1961)	Ectoparasite	Trichobranchidae	<i>Terebellides stroemi</i> Sars, 1835		off Greenland and northern North America; Mediterranean	Bresciani & L�tzen (1974a)
Saccopsidae	<i>Flabellicola neapolitana</i> Gravier, 1918 (GI)	Ectoparasite	Flabelligeridae	<i>Flabelligera diplochaitos</i> (Otto, 1821)		Mediterranean	Gravier (1918a, b); Mariniello (2010)
Saprinidae	<i>Terebellicola reptans</i> Sars, 1862	Ectoparasite	Terebellidae	<i>Eupolymnia nebulosa</i> (Montagu, 1818)		North Sea, Norway	Laubier (1970)
Serpulidicolidae	<i>Parangium abstrusum</i> Humes, 1985	Ectoparasite	Serpulidae	Unknown		Moluccas; Deep waters in northeastern Atlantic; shallow waters in the Gulf of Mexico, South Atlantic off South America; Indo-Pacific; Antarctic waters.	Humes (1985)
Serpulidicolidae	<i>Rhabdopus salmacinae</i> Southward, 1964	Ectoparasite	Serpulidae	<i>Salmacina setosa</i> Langerhans, 1884		North Atlantic Ocean	Southward (1964)
Serpulidicolidae	<i>Rhynchopus catinus</i> Stock, 1979	Ectoparasite	Serpulidae	Undescribed		Eastern Gulf of M�xico	Stock (1979)
Serpulidicolidae	<i>Serpulidicola josephellae</i> Humes & Grassle, 1979	Ectoparasite	Serpulidae	<i>Josepella sp.</i>		North Atlantic Ocean	Humes & Grassle (1979)
Serpulidicolidae	<i>Serpulidicola omphalopomae</i> Southward, 1964	Ectoparasite	Serpulidae	<i>Filogranula stellata</i> (Southward, 1963)		North Atlantic Ocean	Southward (1964), host as <i>Omphalopoma stellata</i> Southward, 1963
Serpulidicolidae	<i>Serpulidicola placostegi</i> Southward, 1964	Ectoparasite	Serpulidae	<i>Placostegus tridentatus</i> (Fabricius, 1779)		North Atlantic Ocean	Southward (1964)
Serpulidicolidae	<i>Serpulidicola segmentatus</i> Stock, 1989	Ectoparasite	Serpulidae	<i>Apomatus sp.</i>		Indonesia	Stock (1989)

Serpulidicolidae	<i>Serpulidicoloides cystopomati</i> (Gravier, 1912)	Ectoparasite	Serpulidae	<i>Hyalopomatus macintoshi</i> (Gravier, 1911)		Antarctic Ocean	Gravier (1912b), host as <i>Cystopomatus macintoshi</i> Gravier, 1911
Spiophanicolidae	<i>Spiophanicola spinosus</i> Ho, 1984	Ectoparasite	Spionidae	<i>Spiophanes berkeleyorum</i> Pettibone, 1962		west coast of southern of California	Ho (1984)
			Spionidae	<i>Spiophanes duplex</i> (Chamberlin, 1919)		west coast of southern of California	Ho (1984), host as <i>Spiophanes missionensis</i> Hartman, 1941
Spiophanicolidae	<i>Spiophanicola atlanticus</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Ectoparasite	Spionidae	<i>Spiophanes kroyeri</i> Grube, 1860	50 - 433 m	west coast of southern of California	O'Reilly (1999)
Xenocoelomatidae	<i>Aphanodomus terebellae</i> (Levinsen, 1878)	Endoparasite	Spionidae	<i>Spiophanes kroyeri</i> Grube, 1860		northern North Sea, Scotland	Kim et al. (2013), host as <i>Spiophanes spinosus</i> $\dot{\iota}$
			Terebellidae	<i>Amphitrite cirrata</i> (O. F. Müller, 1771 in 1776)		Irish waters; wide distribution at high latitudes, from Canada, Greenland and Iceland, to the Kara Sea in the Arctic coasts of Russia	Bresciani & Lützen (1966, 1972, 1974b); O'Reilly (1995)
			Terebellidae	<i>Artacama proboscidea</i> Malmgren, 1866		South Irish Sea, Northeast England, widespread in Arctic waters	Bresciani & Lützen (1966, 1972, 1974b)
			Terebellidae	<i>Lanasa venusta</i> (Malm, 1874)		Iceland	Bresciani & Lützen (1966, 1972, 1974b)
			Terebellidae	<i>Nicolea zostericola</i> Örsted, 1844		Scandinavia	Bresciani & Lützen (1966, 1972, 1974b)
			Terebellidae	<i>Thelepus cincinnatus</i> (Fabricius, 1780)		East Greenland	Bresciani & Lützen (1966, 1972, 1974b)
			Terebellidae	<i>Polycirrus medusa</i> Grube, 1850		British waters	O'Reilly (1995)
			Terebellidae	<i>Polycirrus plumosus</i> (Wollebaek, 1912)			Bresciani & Lützen (1966, 1972, 1974b);
Xenocoelomatidae	<i>Xenocoeloma alleni</i> (Brumpt, 1897)	Ectoparasite	Terebellidae	<i>Polycirrus caliadrum</i> Claparède, 1869		English Channel	Brumpt (1897); Bresciani & Lützen (1966, 1972, 1974b);
			Terebellidae	<i>Polycirrus arenivorus</i> (Caulley, 1915)		French coasts	Brumpt (1897); Bocquet et al. (1968)
			Terebellidae	<i>Polycirrus plumosus</i> (Wollebaek, 1912)			Brumpt (1897)
Xenocoelomatidae	<i>Xenocoeloma brumpti</i> Caullery & Mesnil, 1915	Ectoparasite	Terebellidae	<i>Polycirrus arenivorus</i> (Caulley, 1915)		English Channel	Bresciani & Lützen (1966)
Xenocoelomatidae	<i>Xenocoeloma sp.</i>	Ectoparasite	Terebellidae	<i>Polycirrus arcticus</i> Sars, 1865		Greenland	Bresciani & Lützen (1974b)
			Terebellidae	<i>Polycirrus sp.</i>		Hong Kong	Boxshall (2001)
			Unknown	Unknown		Indian Ocean	Sebastian & Pillai (1974)
Incertae sedis (Poecilostomatoida)	<i>Stockella indica</i> (Sebastian & Pillai, 1974)	?	Unknown	Unknown			
Incertae sedis (Poecilostomatoida)	<i>Cyclorhiza eteonicola</i> Heegaard, 1942	Ectoparasite	Phyllocodidae	<i>Eteone longa</i> (Fabricius, 1780)		North Atlantic and Mediterranean (Lower St Lawrence estuary, Isle of Man, Northeast England, West Norway, eastern North America)	Gotto & Leahy (1988)
Incertae sedis (Poecilostomatoida)	<i>Cyclorhiza megalova</i> Gotto & Leahy, 1988	Ectoparasite	Phyllocodidae	<i>Eteone longa</i> (Fabricius, 1780)		British waters	Gotto & Leahy (1988); O'Reilly (1991)
Incertae sedis (Copepoda)	<i>Notomasticola frondosus</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Endoparasite	Spionidae	<i>Pseudopolydora paucibranchiata</i> (Okuda, 1937)	133 m	Europe	Kim et al. (2013)
Incertae sedis (Copepoda)	<i>Notomasticola frondosus</i> Kim, Sikosky, O'Reilly & Boxhall 2013	Endoparasite	Capitellidae	<i>Notomastus latericius</i> Sars, 1851	125 m	Europe	Kim et al. (2013)
Incertae sedis (Cyclopoidea)	<i>Ophelicola drachi</i> Humes, 1978	Ectoparasite	Opheliidae	Unknown	4.500 m	Bay of Biscay (abyssal plain)	Laubier (Laubier, 1978)
Incertae sedis (Cyclopoidea)	<i>Ophelicola kurambia sp. nov.</i>	Ectoparasite	Opheliidae	Unknown	4,987 – 4,991 m	Kurile Kamchatka Trench and abyssal plain	This paper