

NULL PSEUDO-ISOTROPIC LAGRANGIAN SURFACES

ALFONSO CARRIAZO, VERÓNICA MARTÍN-MOLINA, AND LUC VRANCKEN

ABSTRACT. In this paper we will show that a Lagrangian, Lorentzian surface M_1^2 in a complex pseudo space form $\widetilde{M}_1^2(4c)$ is pseudo-isotropic if and only if M is minimal. Next we will obtain a complete classification of all Lagrangian, Lorentzian surfaces which are lightlike pseudo-isotropic but not pseudo-isotropic.

Key words: *Lagrangian submanifold, complex projective space, isotropic submanifold, Lorentzian submanifold.*

Subject class: 53B25, 53B20.

1. INTRODUCTION

The notion of isotropic submanifold was first introduced in [7] by O'Neill for immersions of Riemannian manifolds and recently extended by Cabrerizo, Fernández and Gómez in [2] to the pseudo-Riemannian case. A submanifold is called pseudo-isotropic if, for any point p and any tangent vector v at a point p , we have that

$$(1.1) \quad \langle h(v, v), h(v, v) \rangle = \widetilde{\lambda}(p) \langle v, v \rangle^2,$$

where h denotes the second fundamental form of the immersion and $\widetilde{\lambda}$ is a smooth function on the submanifold.

Note that since the induced metric is pseudo-Riemannian it is natural to distinguish between timelike, spacelike and lightlike (or null) vectors. This leads in a natural way to the notions of

- (i) Timelike pseudo-isotropic if, for any point p and any timelike tangent vector v at a point p , equation (1.1) is satisfied,
- (ii) Spacelike pseudo-isotropic if, for any point p and any spacelike tangent vector v at a point p , equation (1.1) is satisfied,
- (iii) Lightlike isotropic if, for every lightlike vector v at the point p , we have that $h(v, v)$ is again a lightlike vector.

It was shown in [2] that the notions of pseudo-isotropic, timelike pseudo-isotropic and spacelike pseudo-isotropic are equivalent. In the same paper they also included an example of an immersion which is lightlike pseudo-isotropic but not pseudo-isotropic.

Here we are particularly interested in Lagrangian immersions of complex space forms. In the positive definite case, isotropic Lagrangian immersions have been studied in [3], [4], [5], [6] and [9]. In this paper we will consider pseudo-isotropic and lightlike pseudo-isotropic Lagrangian, Lorentzian surfaces M_1^2 in a complex pseudo space form $\widetilde{M}_1^2(4c)$.

The first two authors are partially supported by the MINECO-FEDER grant MTM2014-52197-P. They also belong to the PAIDI groups FQM-327 and FQM-226 (Junta de Andalucía, Spain), respectively.

We will assume that the space form is not definite and hence has real signature 2. By changing the sign of the metric if necessary, it is sufficient to deal with the cases $c = 0$ or $c = 1$.

We will first show in Section 3 that

Theorem 1.1. *Let M be a Lagrangian, Lorentzian surface in a complex space form. Then M is pseudo-isotropic if and only if M is minimal.*

Next we will obtain a complete classification of all Lagrangian, Lorentzian surfaces which are lightlike pseudo-isotropic but not pseudo-isotropic. To do so, we will first show in Section 4 that such a surface can be seen as the union of two surfaces that we will call of Type I and of Type II. These will be classified, case by case, in Sections 5 and 6, respectively.

2. PRELIMINARIES

Throughout this paper we will assume that M is a Lagrangian, Lorentzian submanifold M of a complex space form \widetilde{M} . We use the standard formulas of Gauss and Weingarten for a submanifold, introducing the second fundamental form h and the shape operators A by

$$\begin{aligned}\widetilde{\nabla}_X Y &= \nabla_X Y + h(X, Y), \\ \widetilde{\nabla}_X \xi &= -A_\xi X + \nabla_X^\perp \xi,\end{aligned}$$

where X and Y are tangent vector fields and ξ is normal. Here, as usual, $\widetilde{\nabla}$ denotes the Levi-Civita connection on the ambient space and, if no confusion is possible, we will always identify M with its image in \widetilde{M} .

Since M is Lagrangian, we have that the complex structure J interchanges the tangent and the normal spaces. Using the formulas of Gauss and Codazzi this implies that

$$\begin{aligned}\nabla_X^\perp JY &= J\nabla_X Y, \\ A_{JX} Y &= -Jh(X, Y) = A_{JY} X.\end{aligned}$$

The latter formula implies that the cubic form $\langle h(X, Y), JZ \rangle$ is totally symmetric in all components.

We denote the curvature tensors of ∇ and ∇^\perp by R and R^\perp , respectively. The first covariant derivative of h is defined by

$$(2.1) \quad (\nabla h)(X, Y, Z) = \nabla_X^\perp h(Y, Z) - h(\nabla_X Y, Z) - h(\nabla_X Z, Y),$$

where X, Y, Z and W are tangent vector fields.

The equations of Gauss, Codazzi and Ricci for a Lagrangian submanifold of $\widetilde{M}^n(4c)$ are given by

$$(2.2) \quad \begin{aligned} \langle R(X, Y)Z, W \rangle &= \langle h(Y, Z), h(X, W) \rangle - \langle h(X, Z), h(Y, W) \rangle \\ &\quad + c(\langle X, W \rangle \langle Y, Z \rangle - \langle X, Z \rangle \langle Y, W \rangle), \end{aligned}$$

$$(2.3) \quad (\nabla h)(X, Y, Z) = (\nabla h)(Y, X, Z),$$

$$(2.4) \quad \begin{aligned} \langle R^\perp(X, Y)JZ, JW \rangle &= \langle [A_{JZ}, A_{JW}]X, Y \rangle \\ &\quad + c(\langle X, W \rangle \langle Y, Z \rangle - \langle X, Z \rangle \langle Y, W \rangle), \end{aligned}$$

where X, Y, Z and W are tangent vector fields. Note that for a Lagrangian submanifold the equations of Gauss and Ricci are mutually equivalent.

We refer to [1] for the construction of the standard models of indefinite complex space forms $\mathbb{C}P_s^n(4c)$ when $c > 0$, $\mathbb{C}H_s^n(4c)$ when $c < 0$ and \mathbb{C}_s^n . For our purposes, it is sufficient to know that there exist pseudo-Riemannian submersions, called Hopf fibrations, given by

$$\Pi : S_{2s}^{2n+1}(c) \rightarrow \mathbb{C}P_s^n(4c) : z \mapsto z \cdot \mathbb{C}^*$$

if $c > 0$, and by

$$\Pi : H_{2s+1}^{2n+1}(c) \rightarrow \mathbb{C}H_s^n(4c) : z \mapsto z \cdot \mathbb{C}^*,$$

if $c < 0$, where

$$\begin{aligned} S_{2s}^{2n+1}(c) &= \{z \in \mathbb{C}^{n+1} | b_{s,n+1}(z, z) = \frac{1}{c}\}, \\ H_{s+1}^{2n+1}(c) &= \{z \in \mathbb{C}^{n+1} | b_{s+1,n+1}(z, z) = \frac{1}{c}\} \end{aligned}$$

and $b_{s,q}$ is the standard Hermitian form with index s on \mathbb{C}^q . For our convenience, we will assume that we have chosen an orthonormal basis such that the first s odd terms appear with a minus sign.

In [1] it is shown that locally any indefinite complex space form is holomorphically isometric to either \mathbb{C}_s^n , $\mathbb{C}P_s^n(4c)$, or $\mathbb{C}H_s^n(4c)$. Remark that, by replacing the metric $\langle \cdot, \cdot \rangle$ by $-\langle \cdot, \cdot \rangle$, we have that $\mathbb{C}H_s^n(4c)$ is holomorphically anti-isometric to $\mathbb{C}P_{n-s}^n(-4c)$. For that purpose, as in our case $n = 2$ and $s = 1$, we only need to consider \mathbb{C}_1^2 and $\mathbb{C}P_1^2(4)$.

In order to study or explicitly obtain examples of Lagrangian submanifolds, it is usually more convenient to work with horizontal submanifolds. In that aspect, we first recall some basic facts from [8] which relate Lagrangian submanifolds of $\mathbb{C}P_s^n(4c)$ to horizontal immersions in $S_{2s}^{2n+1}(c)$. Here, a horizontal immersion $f : M_s^n \rightarrow S_{2s}^{2n+1}(c)$ is an immersion which satisfies $if(p) \perp f_*(T_p M_s^n)$ for all $p \in M_s^n$, where $i = \sqrt{-1}$.

Theorem 2.1 ([8]). *Let $\Pi : S_{2s}^{2n+1}(1) \rightarrow \mathbb{C}P_s^n(4)$ be the Hopf fibration. If $f : M_s^n \rightarrow S_{2s}^{2n+1}(c)$ is a horizontal immersion, then $F = \Pi \circ f : M_s^n \rightarrow \mathbb{C}P_s^n(4c)$ is a Lagrangian immersion.*

Conversely, let M_s^n be a simply connected manifold and let $F : M_s^n \rightarrow \mathbb{C}P_s^n(4)$ be a Lagrangian immersion. Then there exists a 1-parameter family of horizontal lifts $f : M_s^n \rightarrow S_{2s}^{2n+1}(1)$ such that $F = \Pi \circ f$. Any two such lifts f_1 and f_2 are related by $f_1 = e^{i\theta} f_2$, where θ is a constant.

Remark that both immersions have the same induced metric and that the second fundamental forms of both immersions are also closely related. For more details, see [8].

3. MINIMALITY AND PSEUDO-ISOTROPY

In this section we will prove Theorem 1.1. Let us suppose that M_1^2 is a Lagrangian, Lorentzian surface of a complex space form \widetilde{M} . We will assume that either $\widetilde{M} = \mathbb{C}_1^2$ or $\mathbb{C}P_1^2(4)$.

Let $p \in M$. We say that $\{e_1, e_2\}$ is a null frame at a point p if it satisfies

$$\langle e_i, e_j \rangle = (1 - \delta_{ij}), \quad i, j \in \{1, 2\}.$$

In terms of a null frame, it is then clear that a Lagrangian immersion is minimal if and only if

$$h(e_1, e_2) = 0.$$

In view of the symmetries of the second fundamental form, this implies that there exist numbers λ and μ such that

$$\begin{aligned} h(e_1, e_1) &= \lambda J e_2, \\ h(e_2, e_2) &= \mu J e_1. \end{aligned}$$

If we now write $v = v_1 e_1 + v_2 e_2$, it follows that

$$\begin{aligned} h(v, v) &= \mu v_2^2 J e_1 + \lambda v_1^2 J e_2, \\ \langle h(v, v), h(v, v) \rangle &= 2\lambda\mu v_1^2 v_2^2 = \frac{1}{2}\lambda\mu \langle v, v \rangle^2, \end{aligned}$$

which shows that a minimal surface is indeed pseudo-isotropic.

In order to show the converse, we will use the following lemma of [2]:

Lemma 3.1. *Let $F : M \rightarrow \widetilde{M}$ be an isometric pseudo-Riemannian immersion. Then the immersion is (pseudo)-isotropic if and only if for any tangent vectors $x, y, z, w \in T_p M$, we have that*

$$\begin{aligned} \langle h(x, y), h(z, w) \rangle + \langle h(y, z), h(x, w) \rangle + \langle h(z, x), h(y, w) \rangle = \\ = \widetilde{\lambda}(p) \{ \langle x, y \rangle \langle z, w \rangle + \langle y, z \rangle \langle x, w \rangle + \langle z, x \rangle \langle y, w \rangle \}. \end{aligned}$$

Note that in [2] the above lemma was formulated only for immersions into pseudo-Euclidean spaces. However, it is clear that it remains valid for arbitrary immersions in pseudo-Riemannian spaces.

Let us assume now that M is a pseudo-isotropic surface. Then it follows from the previous lemma that

- (i) $h(e_1, e_1)$ is a lightlike vector, by taking $x = y = z = w = e_1$,
- (ii) $h(e_2, e_2)$ is a lightlike vector, by taking $x = y = z = w = e_2$,
- (iii) $h(e_1, e_2)$ is orthogonal to $h(e_1, e_1)$, by taking $x = y = z = e_1$ and $w = e_2$,
- (iv) $h(e_1, e_2)$ is orthogonal to $h(e_2, e_2)$, by taking $x = y = z = e_2$ and $w = e_1$.

We now write

$$h(e_1, e_2) = v_1 J e_1 + v_2 J e_2.$$

The fact that the immersion is Lagrangian then implies that

$$h(e_1, e_1) = v_2 J e_1 + v_3 J e_2,$$

$$h(e_2, e_2) = v_4 J e_1 + v_1 J e_2.$$

Let us now assume that M is not minimal. Then, by interchanging e_1 and e_2 if necessary, we may assume that $v_2 \neq 0$. As $h(e_1, e_1)$ is lightlike by (i), this implies that $v_3 = 0$. It then follows from (iii) that $v_2 = 0$ which is a contradiction. This completes the proof of Theorem 1.

4. LIGHTLIKE ISOTROPIC LAGRANGIAN, LORENTZIAN SURFACES

In this section we will assume that M_1^2 is a Lagrangian, Lorentzian lightlike pseudo-isotropic surface of a complex space form \widetilde{M} . We will assume that either $\widetilde{M} = \mathbb{C}_1^2$ or $\mathbb{C}P_1^2(4)$. We will also assume that M is not pseudo-isotropic, i.e. in view of the previous section we will assume that the immersion is not minimal. We call a surface which is lightlike pseudo-isotropic without minimal points a proper lightlike pseudo-isotropic surface.

We again take a null frame at a point p , i.e. a frame $\{e_1, e_2\}$ such that

$$\langle e_i, e_j \rangle = (1 - \delta_{ij}), \quad i, j \in \{1, 2\}.$$

Note that if both $h(e_1, e_1) = h(e_2, e_2) = 0$, then it follows from the fact that M is Lagrangian that also $h(e_1, e_2) = 0$ and therefore that M is pseudo-isotropic.

We say that M is proper lightlike pseudo-isotropic of Type 1 at the point p if there exists a lightlike vector v such that $h(v, v)$ and Jv are independent. This means, after changing e_1 and e_2 if necessary, that we may assume that

$$h(e_1, e_1) = J e_2,$$

$$h(e_2, e_2) = \mu J e_1 + \lambda J e_2.$$

Since M is Lagrangian, we deduce from this that

$$h(e_1, e_2) = \lambda J e_1.$$

Since M is proper, we have that $\lambda \neq 0$. Hence, since $h(e_2, e_2)$ is lightlike, we deduce that $\mu = 0$ and so $h(e_2, e_2) = \lambda J e_2$.

We say that M is proper lightlike pseudo-isotropic of Type 2 if, for every lightlike vector v , $h(v, v)$ and Jv are dependent. This means, after changing e_1 and e_2 if necessary, that we may assume that

$$h(e_1, e_1) = J e_1,$$

$$h(e_2, e_2) = \lambda J e_2.$$

As M is Lagrangian, we deduce from this that

$$h(e_1, e_2) = \lambda J e_1 + J e_2.$$

We see that M is indeed not minimal and the immersion is therefore proper lightlike pseudo-isotropic.

Note that a point which belongs to the closure of Type 2 points needs to be a minimal point automatically. Therefore, it follows from the fact that M is proper that M can be seen as the union of a Type 1 lightlike proper pseudo-isotropic surface and a Type 2 lightlike proper pseudo-isotropic surface. In both cases, it is immediately clear that the null frame can be extended to a neighborhood of the point p .

5. PROPER LIGHTLIKE ISOTROPIC LAGRANGIAN, LORENTZIAN SURFACES OF TYPE 1

We take a null frame in a neighborhood of the point p as constructed in the previous section. So we have a frame $\{E_1, E_2\}$ such that

$$\langle E_i, E_j \rangle = (1 - \delta_{ij}), \quad i, j \in \{1, 2\},$$

and

$$\begin{aligned} h(E_1, E_1) &= JE_2, \\ h(E_2, E_2) &= \lambda JE_2, \\ h(E_1, E_2) &= \lambda JE_1, \end{aligned}$$

where λ is a nowhere vanishing function. We write

$$\begin{aligned} \nabla_{E_1} E_1 &= \alpha E_1, & \nabla_{E_1} E_2 &= -\alpha E_2, \\ \nabla_{E_2} E_1 &= -\beta E_1, & \nabla_{E_2} E_2 &= \beta E_2, \end{aligned}$$

where α and β are functions.

Lemma 5.1. *We have that $\beta = 0$ and λ satisfies the following system of differential equations:*

$$\begin{aligned} E_1(\lambda) &= -\alpha\lambda, \\ E_2(\lambda) &= 0. \end{aligned}$$

Proof. We have that

$$\begin{aligned} (\nabla h)(E_2, E_1, E_1) &= \nabla_{E_2}^\perp JE_2 - 2h(\nabla_{E_2} E_1, E_1) \\ &= \beta JE_2 + 2\beta JE_2 = 3\beta JE_2. \end{aligned}$$

On the other hand, we have that

$$\begin{aligned} (\nabla h)(E_1, E_2, E_1) &= \nabla_{E_1}^\perp \lambda JE_1 - h(\nabla_{E_1} E_2, E_1) - h(E_2, \nabla_{E_1} E_1) \\ &= (E_1(\lambda) + \alpha\lambda)JE_1 + \alpha h(E_2, E_1) - \alpha h(E_2, E_1) \\ &= (E_1(\lambda) + \alpha\lambda)JE_1. \end{aligned}$$

From the Codazzi equation, we therefore obtain that $\beta = 0$ and $E_1(\lambda) = -\alpha\lambda$. Similarly from the Codazzi equation $(\nabla h)(E_1, E_2, E_2) = (\nabla h)(E_2, E_1, E_2)$, we now deduce that $E_2(\lambda) = 0$. \square

Lemma 5.2. *We have that $c = 0$ and α satisfies*

$$E_2(\alpha) = 0.$$

Proof. We compute $[E_1, E_2](\lambda)$ in two different ways. We have that

$$\begin{aligned} [E_1, E_2](\lambda) &= E_1(E_2(\lambda)) - E_2(E_1(\lambda)) \\ &= E_2(\alpha\lambda) \\ &= E_2(\alpha)\lambda \end{aligned}$$

and

$$\begin{aligned} [E_1, E_2](\lambda) &= (\nabla_{E_1} E_2 - \nabla_{E_2} E_1)(\lambda) \\ &= -\alpha E_2(\lambda) = 0. \end{aligned}$$

Since $\lambda \neq 0$, we deduce that $E_2(\alpha) = 0$.

A direct computation then yields that

$$\begin{aligned} R(E_1, E_2)E_1 &= -\nabla_{E_2} \nabla_{E_1} E_1 - \nabla_{\nabla_{E_1} E_2} E_1 \\ &= -\nabla_{E_2}(\alpha E_1) + \alpha \nabla_{E_2} E_1 \\ &= -E_2(\alpha)E_1 = 0. \end{aligned}$$

So from the Gauss equation we obtain that

$$\begin{aligned} 0 &= cE_1 + A_{h(E_1, E_2)}e_1 - A_{h(E_1, E_1)}E_2 \\ &= cE_1 + \lambda A_{JE_1}E_1 - A_{JE_2}E_2 \\ &= cE_1. \end{aligned}$$

Hence the ambient space must be flat. \square

The previous lemma immediately implies:

Theorem 5.3. *There does not exist a proper lightlike isotropic Lagrangian, Lorentzian surface of Type 1 in $\mathbb{C}P_1^2(4)$.*

Moreover, we can also show that

Theorem 5.4. *Let M be a proper lightlike isotropic Lagrangian, Lorentzian surface of Type 1 in \mathbb{C}_1^2 . Then M is locally congruent with*

$$(\alpha(x)\frac{1}{\sqrt{2}}(-i, -i) + \beta(x)\frac{1}{\sqrt{2}}(1, -1))e^{iv},$$

where $\alpha'(x)\beta(x) - \alpha(x)\beta'(x) \neq 0$.

Proof. We introduce vector fields λE_1 and $\frac{1}{\lambda}E_2$. We have that

$$\begin{aligned} [\lambda E_1, \frac{1}{\lambda}E_2] &= -\frac{E_1(\lambda)}{\lambda}E_2 + [E_1, E_2] \\ &= \alpha E_2 - \alpha E_2 = 0. \end{aligned}$$

Therefore, there exist coordinates u and v such that $\partial u = \lambda E_1$ and $\partial v = \frac{1}{\lambda} E_2$. If we denote the immersion by f , it follows that

$$\begin{aligned} f_{vv} &= i f_v, \\ f_{uv} &= i f_u, \\ f_{uu} &= \lambda E_1(\lambda) E_1 + \lambda^2 \nabla_{E_1} E_1 + \lambda^2 h(E_1, E_1) \\ &= -\lambda^2 \alpha + \lambda^2 \alpha + \lambda^2 i E_2 \\ &= \lambda^3 i f_v, \end{aligned}$$

where λ is a function depending only on u . Integrating the first two equations it follows that

$$f(u, v) = A_1(u) e^{iv} + A_2,$$

where A_1 is a vector valued function and A_2 is a constant. Of course, we may assume that A_2 vanishes by applying a translation of \mathbb{C}_1^2 . The third equation then tells us that

$$A_1'' = -\lambda^3 A_1.$$

Note that this is precisely the expression of a curve lying in the plane spanned by $A_1(0)$ and $A_1'(0)$ parametrised in such a way that $|A_1 A_1'|$ is constant. Given that M is a Lagrangian surface we must have that A_1 and A_1' are linearly independent (over \mathbb{C}) and that the plane spanned by A_1 and $i A_1'$ is real. Therefore, the constant is non-vanishing. Since $f_u = \lambda E_1$ and $f_v = \frac{1}{\lambda} E_2$, by choosing the initial conditions we may assume that $A_1(0) = \frac{1}{\sqrt{2}}(-i, -i)$ and $A_1'(0) = \frac{1}{\sqrt{2}}(1, -1)$.

Conversely, if we define a surface by

$$f(x, v) = (\alpha(x) \frac{1}{\sqrt{2}}(-i, -i) + \beta(x) \frac{1}{\sqrt{2}}(1, -1)) e^{iv},$$

where $\alpha'(x)\beta(x) - \alpha(x)\beta'(x) \neq 0$, we see that just as for the Euclidean arc length of a planar curve, it is possible to construct a parameter u for the curve (α, β) such that $\alpha'(u)\beta(u) - \alpha(u)\beta'(u) = 1$. A straightforward computation then shows that the surface $f(u, v)$ has the desired properties. \square

6. PROPER LIGHTLIKE ISOTROPIC LAGRANGIAN, LORENTZIAN SURFACES OF TYPE 2

We take a null frame in a neighborhood of the point p as constructed in the previous section. Then we have a frame $\{E_1, E_2\}$ such that

$$\langle E_i, E_j \rangle = (1 - \delta_{ij}), \quad i, j \in \{1, 2\},$$

and

$$\begin{aligned} h(E_1, E_1) &= J E_1, \\ h(E_1, E_2) &= \lambda J E_1 + J E_2, \\ h(E_2, E_2) &= \lambda J E_2, \end{aligned}$$

where λ is a function on M . We write

$$\begin{aligned}\nabla_{E_1}E_1 &= \alpha E_1, & \nabla_{E_1}E_2 &= -\alpha E_2, \\ \nabla_{E_2}E_1 &= -\beta E_1, & \nabla_{E_2}E_2 &= \beta E_2,\end{aligned}$$

where α and β are functions.

Lemma 6.1. *We have that $\alpha = 0$ and λ satisfies the following system of differential equations:*

$$\begin{aligned}E_1(\lambda) &= \beta, \\ E_2(\lambda) &= \lambda\beta.\end{aligned}$$

Proof. We have that

$$\begin{aligned}(\nabla h)(E_2, E_1, E_1) &= \nabla_{E_2}^\perp J E_1 - 2h(\nabla_{E_2}E_1, E_1) \\ &= -\beta J E_1 + 2\beta J E_1 = \beta J E_1.\end{aligned}$$

On the other hand, we have that

$$\begin{aligned}(\nabla h)(E_1, E_2, E_1) &= \nabla_{E_1}^\perp (\lambda J E_1 + J E_2) - h(\nabla_{E_1}E_2, E_1) - h(E_2, \nabla_{E_1}E_1) \\ &= (E_1(\lambda) + \alpha\lambda) J E_1 - \alpha J E_2 + \alpha h(E_2, E_1) - \alpha h(E_2, E_1) \\ &= (E_1(\lambda) + \alpha\lambda) J E_1 - \alpha J E_2.\end{aligned}$$

From the Codazzi equation, we therefore obtain that $\alpha = 0$ and $E_1(\lambda) = \beta$. Similarly from the Codazzi equation $(\nabla h)(E_1, E_2, E_2) = (\nabla h)(E_2, E_1, E_2)$, we now deduce that $E_2(\lambda) = \lambda\beta$. \square

Lemma 6.2. *We have that β satisfies*

$$\begin{aligned}E_1(\beta) &= -c - \lambda, \\ E_2(\beta) &= \lambda(-c - \lambda).\end{aligned}$$

Proof. A direct computation yields that

$$\begin{aligned}R(E_1, E_2)E_1 &= \nabla_{E_1}\nabla_{E_2}E_1 + \nabla_{\nabla_{E_2}E_1}E_1 \\ &= -\nabla_{E_1}(\beta E_1) - \beta\nabla_{E_1}E_1 \\ &= -E_1(\beta)E_1.\end{aligned}$$

So from the Gauss equation we obtain that

$$\begin{aligned}-E_1(\beta)E_1 &= cE_1 + A_{h(E_1, E_2)}E_1 - A_{h(E_1, E_1)}E_2 \\ &= cE_1 + \lambda A_{J E_1}E_1 + A_{J E_2}E_1 - A_{J E_1}E_2 \\ &= cE_1 + \lambda E_1,\end{aligned}$$

which reduces to $E_1(\beta) = -(c + \lambda)$. In order to obtain the E_2 derivative of β , we compute $[E_1, E_2](\lambda)$ in two different ways. We have that

$$\begin{aligned}[E_1, E_2](\lambda) &= E_1(E_2(\lambda)) - E_2(E_1(\lambda)) \\ &= E_1(\lambda\beta) - E_2(\beta) \\ &= \beta^2 + \lambda(-c - \lambda) - E_2(\beta)\end{aligned}$$

and

$$\begin{aligned} [E_1, E_2](\lambda) &= (\nabla_{E_1} E_2 - \nabla_{E_2} E_1)(\lambda) \\ &= \beta E_1(\lambda) = \beta^2, \end{aligned}$$

which clearly concludes the proof. \square

It follows by a direct computation that

Corollary 6.3. *There exists a constant r such that*

$$(\lambda + c)^2 + \beta^2 = r^2.$$

Lemma 6.4. *There exist local coordinates u and v such that*

$$\begin{aligned} \frac{\partial}{\partial u} &= E_1, \\ \frac{\partial}{\partial v} &= E_2 - \lambda E_1. \end{aligned}$$

Proof. We define vector fields

$$\begin{aligned} U &= E_1, \\ V &= E_2 - \lambda E_1, \end{aligned}$$

and compute

$$\begin{aligned} [U, V] &= [E_1, E_2 - \lambda E_1] \\ &= [E_1, E_2] - E_1(\lambda) E_1 \\ &= \beta E_1 - E_1(\lambda) E_1 = 0, \end{aligned}$$

which proves the result. \square

It then follows immediately from the previous systems of differential equations that β and λ do not depend on the variable v and are determined by

$$\begin{aligned} \frac{\partial \lambda}{\partial u} &= \beta, \\ \frac{\partial \beta}{\partial u} &= -(c + \lambda). \end{aligned}$$

Therefore, note that, after a translation of the u coordinate if necessary, we may suppose that

$$\begin{aligned} \lambda &= -c + r \sin u, \\ \beta &= r \cos u. \end{aligned}$$

In the above equations, the constant r is allowed to be zero. In that case, we get the special solution $\beta = 0$ and $\lambda = -c$.

In view of the dimension, if necessary changing the sign of the metric on the ambient space, we only have to consider the cases $c = 0$ or $c = 1$.

6.1. **Case $c = 0$: Lightlike isotropic Lagrangian, Lorentzian surfaces in \mathbb{C}_1^2 .** We denote the immersion by f . It follows from the previous equations that f is determined by the system of differential equations:

$$\begin{aligned} f_{uu} &= if_u, \\ f_{uv} &= -re^{-iu}f_u + if_v, \\ f_{vv} &= re^{-iu}f_v + ir^2(e^{-2iu} - 1)f_u. \end{aligned}$$

It follows from the first equation that there exist vector valued functions g_1 and g_2 such that

$$f(u, v) = g_1(v)e^{iu} + g_2(v).$$

Substituting this into the second equation gives

$$g_2'(v) = rg_1(v),$$

and the final equation now reduces to

$$g_1''(v)e^{iu} + g_2''(v) = re^{-iu}(g_1'(v)e^{iu} + g_2'(v)) - r^2(e^{-2iu} - 1)e^{iu}g_1(v).$$

Looking at the different powers of e^{iu} , we deduce that

$$\begin{aligned} g_1''(v) &= r^2g_1(v), \\ g_2''(v) &= rg_1'(v), \\ 0 &= rg_2'(v) - r^2g_1(v). \end{aligned}$$

So the remaining equations are

$$\begin{aligned} g_2'(v) &= rg_1(v), \\ g_1''(v) &= r^2g_1(v). \end{aligned}$$

The solution of the above system depends on the value of r .

6.1.1. *Case $r = 0$.* If $r = 0$ we have that $g_2(v)$ is a constant vector. Hence by applying a translation we may assume that this vector vanishes. Therefore, we have that

$$f(u, v) = (vA_1 + A_2)e^{iu},$$

for some constant vectors A_1 and A_2 . We take an initial point $p = (0, 0)$. Since $\lambda(0, 0) = 0$, it follows that

$$\begin{aligned} E_1(0, 0) &= \frac{\partial f}{\partial u}(0, 0) = iA_2, \\ E_2(0, 0) &= \frac{\partial f}{\partial v}(0, 0) = A_1. \end{aligned}$$

It then follows from the choice of E_1 and E_2 , together with the Lagrangian condition, that $\frac{1}{\sqrt{2}}(A_1 - iA_2)$, $\frac{i}{\sqrt{2}}(A_1 - iA_2)$, $\frac{1}{\sqrt{2}}(A_1 + iA_2)$, $\frac{i}{\sqrt{2}}(A_1 + iA_2)$ can be identified with $(1, 0)$, $(i, 0)$, $(0, 1)$, $(0, i)$. This implies that

$$\begin{aligned} A_1 &= \left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right), \\ A_2 &= \left(\frac{\sqrt{2}}{2}i, -\frac{\sqrt{2}}{2}i\right). \end{aligned}$$

6.1.2. *Case $r \neq 0$.* In this case we have that

$$g_1(v) = A_1 e^{rv} + A_2 e^{-rv}.$$

Therefore

$$g_2'(v) = rA_1 e^{rv} + rA_2 e^{-rv},$$

which implies that, after applying a suitable translation, we have that

$$g_2(v) = A_1 e^{rv} - A_2 e^{-rv}.$$

So we find that

$$f(u, v) = (A_1 e^{rv} + A_2 e^{-rv})e^{iu} + (A_1 e^{rv} - A_2 e^{-rv}),$$

for some constant vectors A_1 and A_2 . We take again as initial point $p = (0, 0)$. Since $\lambda(0, 0) = 0$, it follows that

$$\begin{aligned} E_1(0, 0) &= \frac{\partial f}{\partial u}(0, 0) = i(A_1 + A_2), \\ E_2(0, 0) &= \frac{\partial f}{\partial v}(0, 0) = 2rA_1. \end{aligned}$$

Or equivalently

$$\begin{aligned} A_1 &= \frac{1}{2r}E_2, \\ A_2 &= -iE_1 - \frac{1}{2r}E_2. \end{aligned}$$

It then follows from the choice of E_1 and E_2 , together with the Lagrangian condition, that we may assume that $E_1 = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ and $E_2 = (-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$, which implies that

$$\begin{aligned} A_1 &= \left(-\frac{\sqrt{2}}{4r}, \frac{\sqrt{2}}{4r}\right), \\ A_2 &= \left(\frac{(1-2ir)}{2\sqrt{2}r}, \frac{(-1-2ir)}{\sqrt{2}r}\right). \end{aligned}$$

6.1.3. *Summary.* Combining the previous results, we get

Theorem 6.5. *Let M be a proper lightlike isotropic Lagrangian, Lorentzian surface of Type 2 in \mathbb{C}_1^2 . Then M is congruent with one of the following surfaces:*

(1) *the surface*

$$f(u, v) = (vA_1 + A_2)e^{iu},$$

where $A_1 = (\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})$ and $A_2 = (-\frac{\sqrt{2}}{2}i, \frac{\sqrt{2}}{2}i)$,

(2) *the surface*

$$f(u, v) = (A_1 e^{rv} + A_2 e^{-rv})e^{iu} + (A_1 e^{rv} - A_2 e^{-rv}),$$

where r is a positive constant, and $A_1 = (-\frac{\sqrt{2}}{4r}, \frac{\sqrt{2}}{4r})$ and $A_2 = (\frac{(1-2ir)}{2\sqrt{2}r}, \frac{(-1-2ir)}{\sqrt{2}r})$.

6.2. Case $c = 1$: Lightlike isotropic Lagrangian, Lorentzian surfaces in $\mathbb{C}P_1^2(4)$. We denote the horizontal lift of the immersion into $S_2^5(1)$ by f . It follows from the previous equations that f is determined by the system of differential equations:

$$\begin{aligned} f_{uu} &= if_u, \\ f_{uv} &= -(i + re^{-iu})f_u + if_v - f, \\ f_{vv} &= (i + re^{-iu})(-2f_u + f_v + 2f_u r \sin(u)) - 2(1 - r \sin(u))f. \end{aligned}$$

It follows from the first equation that there exists vector valued functions a_1 and a_2 such that

$$f(u, v) = a_1(v)e^{iu} + a_2(v).$$

Substituting this into the second equation gives

$$a_2'(v) = -ia_2(v) + ra_1(v).$$

The final equation now reduces to

$$a_1''(v) = a_1(v)r^2 + i(a_1'(v) - a_2(v)r).$$

The solution of this differential equation depends on the value of r .

6.2.1. Case $0 \leq r < 1$. In this case we can write $r = \cos(t)$, where $t \in]0, \frac{\pi}{2}]$. We find that

$$\begin{aligned} a_1(v) &= -\csc^2(t) (-c_3 \sin(t) \sin(v \sin(t)) \\ &\quad + (c_1 \cos^2(t) + ic_3) \cos(v \sin(t)) \\ &\quad + 2ic_2 \cos(t) \sin^2(\frac{1}{2}v \sin(t)) - c_1 - ic_3) \end{aligned} \tag{6.1}$$

$$\begin{aligned} a_2(v) &= -\csc^2(t) (\cos(t) (-c_1 \sin(t) \sin(v \sin(t)) \\ &\quad + (c_3 - ic_1) (\cos(v \sin(t)) - 1)) \\ &\quad - c_2 (\cos(v \sin(t)) - i \sin(t) \sin(v \sin(t))) + c_2 \cos^2(t)) \end{aligned} \tag{6.2}$$

We take again as initial point $p = (0, 0)$. We have that $\lambda(0, 0) = -1$. It follows that

$$\begin{aligned} f(0, 0) &= c_1 + c_2, \\ E_1(0, 0) &= \frac{\partial f}{\partial u}(0, 0) = ic_1, \\ E_2(0, 0) &= \frac{\partial f}{\partial v}(0, 0) + \lambda \frac{\partial f}{\partial u}(0, 0) = -i(c_1 + c_2) + c_3 + c_1 \cos t. \end{aligned}$$

So if we pick the initial conditions $f(0, 0) = (0, 0, 1)$, $E_1(0, 0) = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0)$ and $E_2(0, 0) = (-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0)$, we find that

$$c_1 = \left(-\frac{i}{\sqrt{2}}, -\frac{i}{\sqrt{2}}, 0 \right), \tag{6.3}$$

$$c_2 = \left(\frac{i}{\sqrt{2}}, \frac{i}{\sqrt{2}}, 1 \right), \tag{6.4}$$

$$c_3 = \left(i \left(\frac{\cos(t)}{\sqrt{2}} + \frac{i}{\sqrt{2}} \right), i \left(\frac{\cos(t)}{\sqrt{2}} - \frac{i}{\sqrt{2}} \right), i \right). \tag{6.5}$$

6.2.2. *Case $r = 1$.* We obtain as solution of the differential equation that

$$(6.6) \quad a_1(v) = \frac{1}{2} \left(c_1 (v^2 + 2) + v (2c_3 - i(c_2 - c_3)v) \right),$$

$$(6.7) \quad a_2(v) = \frac{1}{2} (2c_2 + v(c_3v + (c_1 - ic_2)(2 - iv))).$$

We take again as initial point $p = (0, 0)$. We have that $\lambda(0, 0) = -1$. It follows that

$$\begin{aligned} f(0, 0) &= c_1 + c_2, \\ E_1(0, 0) &= \frac{\partial f}{\partial u}(0, 0) = ic_1, \\ E_2(0, 0) &= (1 - i)c_1 - ic_2 + c_3. \end{aligned}$$

So, if we pick the initial conditions $f(0, 0) = (0, 0, 1)$, $E_1(0, 0) = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0)$ and $E_2 = (-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0)$, we find that

$$(6.8) \quad c_1 = \left(-\frac{i}{\sqrt{2}}, -\frac{i}{\sqrt{2}}, 0 \right),$$

$$(6.9) \quad c_2 = \left(\frac{i}{\sqrt{2}}, \frac{i}{\sqrt{2}}, 1 \right),$$

$$(6.10) \quad c_3 = \left(\frac{i-1}{\sqrt{2}}, \frac{1+i}{\sqrt{2}}, i \right).$$

6.2.3. *Case $r > 1$.* We obtain as solution of the differential equation that

$$(6.11) \quad a_1(v) = \frac{c_3\sqrt{r^2-1} \sinh(\sqrt{r^2-1}v) + (c_1r^2 - ic_2r + ic_3) \cosh(\sqrt{r^2-1}v) + ic_2r - c_1 - ic_3}{r^2 - 1},$$

$$(6.12) \quad a_2(v) = \frac{\sqrt{r^2-1}(c_1r - ic_2) \sinh(\sqrt{r^2-1}v) + (-c_2 + (c_3 - ic_1)r) \cosh(\sqrt{r^2-1}v) + r(c_2r + ic_1 - c_3)}{r^2 - 1}.$$

We take again as initial point $p = (0, 0)$. We have that $\lambda(0, 0) = -1$. It follows that

$$\begin{aligned} f(0, 0) &= c_1 + c_2, \\ E_1(0, 0) &= \frac{\partial f}{\partial u}(0, 0) = ic_1, \\ E_2(0, 0) &= -i(c_1 + c_2) + c_3 + c_1r. \end{aligned}$$

So, if we pick the initial conditions $f(0, 0) = (0, 0, 1)$, $E_1(0, 0) = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0)$ and $E_2 = (-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0)$, we find that

$$(6.13) \quad c_1 = \left(-\frac{i}{\sqrt{2}}, -\frac{i}{\sqrt{2}}, 0 \right),$$

$$(6.14) \quad c_2 = \left(\frac{i}{\sqrt{2}}, \frac{i}{\sqrt{2}}, 1 \right),$$

$$(6.15) \quad c_3 = \left(\frac{i(i+r)}{\sqrt{2}}, \frac{(1+ir)}{\sqrt{2}}, i \right).$$

6.2.4. *Summary.* Combining the previous results we get the following theorem which finishes our classification.

Theorem 6.6. *Let M be a proper lightlike isotropic Lagrangian, Lorentzian surface in $\mathbb{C}P_1^2(4)$. Then the Hopf lift of M is congruent with one of the following immersions into $S_1^5(1)$ given by $f(u, v) = a_1(v)e^{iu} + a_2(v)$, where either*

- (1) a_1, a_2, c_1, c_2, c_3 are as described in (6.1)–(6.5), or
- (2) a_1, a_2, c_1, c_2, c_3 are as described in (6.6)–(6.10), or
- (3) a_1, a_2, c_1, c_2, c_3 are as described in (6.11)–(6.15).

REFERENCES

- [1] M. Barros, A. Romero, Indefinite Kähler manifolds, *Math. Ann.*, 261: 55–62, 1982.
- [2] J. L. Cabrerizo, M. Fernández, J. S. Gómez, Rigidity of pseudo-isotropic immersions, *J. Geom. Phys.*, 59: 834–842, 2009.
- [3] B.-Y. Chen, Complex extensors and Lagrangian submanifolds in complex Euclidean spaces, *Tôhoku Mathematical Journal. Second Series*, 4: 277–297, 1997.
- [4] B.-Y. Chen, Complex extensors and Lagrangian submanifolds in indefinite complex Euclidean spaces, *Bull. Inst. Math. Acad. Sinica*, 31: 151–179, 2003.
- [5] H. Li, X. Wang, Isotropic Lagrangian submanifolds in complex Euclidean space and complex hyperbolic space *Results Math.* 56: 387–403, 2009.
- [6] S. Montiel, F. Urbano, Isotropic totally real submanifolds, *Math.Z.*, 199: 55–60, 1988.
- [7] B. O’Neill, Isotropic and Kaehler immersions, *Canad. J. Math.*, 17: 907–915, 1965.
- [8] H. Reckziegel, Horizontal lifts of isometric immersions into the bundle space of a pseudo-Riemannian submersion, In D. Ferus, R. B. Gardner, S. Helgason, and U. Simon, editors, *Global Differential Geometry and Global Analysis 1984. Proceedings of a Conference held in Berlin, June 10–14, 1984*, volume 1156 of *Lecture notes in Mathematics*, pages 264–279. Springer-Verlag, 1984.
- [9] L. Vrancken, Some remarks on isotropic submanifolds, *Publ. Inst. Math. (Beograd), N.S.*, 51: 94–200, 1992.

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE MATEMÁTICAS, C/ TARFIA S/N,
UNIVERSIDAD DE SEVILLA, 41012, SEVILLA, SPAIN
E-mail address: `carriazo@us.es`

DEPARTAMENTO DE DIDÁCTICA DE LAS MATEMÁTICAS, FACULTAD DE CIENCIAS DE LA EDUCACIÓN,
C/ PIROTECNIA S/N, UNIVERSIDAD DE SEVILLA, 41013, SEVILLA, SPAIN
E-mail address: `veronicamartin@us.es`

LAMAV, UNIVERSITÉ DE VALENCIENNES, 59313, VALENCIENNES CEDEX 9, FRANCE AND DE-
PARTEMENT WISKUNDE, KU LEUVEN, CELESTIJNENLAAN 200 B, 3001, LEUVEN, BELGIUM
E-mail address: `luc.vrancken@univ-valenciennes.fr`