



Depósito de Investigación
Universidad de Sevilla

Depósito de investigación de la Universidad de Sevilla

<https://idus.us.es/>

“This is an Accepted Manuscript of an article published by Journal of Pure and Applied Algebra on 7 January 2020, available at: <https://doi.org/10.1016/j.jpaa.2020.106312>.”

Normally ordered forms of powers of differential operators and their combinatorics

Emmanuel Briand^{*1}, Samuel A. Lopes^{†2}, and Mercedes Rosas^{*3}

¹*Departamento Matemática Aplicada I, Universidad de Sevilla, España.*

²*CMUP, Faculdade de Ciências da Universidade do Porto, Rua do Campo Alegre 687, 4169-007 Porto, Portugal.*

³*Departamento de Álgebra, Universidad de Sevilla, España.*

Abstract

We investigate the combinatorics of the general formulas for the powers of the operator $h\partial^d$, where ∂ is a differential operator on an arbitrary noncommutative ring in which h is central. New formulas for the generalized Stirling numbers are obtained, as well as results on the divisibility by primes of the coefficients which occur in the normally ordered form of $h\partial^d$. All of the above applies to the theory of formal differential operator rings.

MSC Numbers (2010): 16W25, 16S32, 05A10, 11B73.

Keywords: Derivations, rings of differential operators, Weyl algebra, normal ordering, generalized Stirling numbers.

1 Introduction

1.1 Overview

There exist *universal* polynomials U_n for computing the normally ordered form of $(h\partial)^n$, where ∂ is a derivation of a (noncommutative) ring A and h is a central element of A . For instance, since, as operators on A , we have:

$$\begin{aligned}(h\partial)^2 &= h\partial(h)\partial + h^2\partial^2, \\(h\partial)^3 &= h(\partial(h))^2\partial + h^2\partial^2(h)\partial + 3h^2\partial(h)\partial^2 + h^3\partial^3,\end{aligned}$$

the polynomials U_2 and U_3 are

$$\begin{aligned}U_2 &= y_0y_1t + y_0^2t^2, \\U_3 &= y_0y_1^2t + y_0^2y_2t + 3y_0^2y_1t^2 + y_0^3t^3,\end{aligned}$$

^{*}Partially supported by MTM2016-75024-P, funded by MICIU (España) and European structural funds (FEDER), and the Junta de Andalucía under grant FQM-333.

[†]Partially supported by CMUP (UID/MAT/00144/2019), which is funded by FCT (Portugal) with national (MEC) and European structural funds (FEDER), under the partnership agreement PT2020.

so that $(h\partial)^n$ is obtained from U_n by replacing t with ∂ and y_k with $\partial^k(h)$.

The coefficients occurring in U_n form a family c_λ^n of nonnegative integers indexed by partitions λ and displaying very interesting combinatorial properties. To whet the reader's interest, Table 1 below lists the values c_λ^n for $n \leq 5$ and hints at the connections with Stirling numbers of both kinds and Eulerian numbers.

λ	\emptyset	\square	$\square\square$	$\square\square\square$	$\square\square\square\square$	$\square\square\square\square\square$	$\square\square\square\square\square\square$	$\square\square\square\square\square\square\square$	$\square\square\square\square\square\square\square\square$	$\square\square\square\square\square\square\square\square\square$	$\square\square\square\square\square\square\square\square\square\square$	
c_λ^1	1											
c_λ^2	1	1										
c_λ^3	1	3	1	1								
c_λ^4	1	6	4	7	1	4	1					
c_λ^5	1	10	10	25	5	30	15	1	11	4	7	1

Table 1: The coefficients c_λ^n of the polynomials U_n . Each partition λ is represented by its Young diagram. This table conceals the signless Stirling numbers of the first kind (as the sums $\sum_{\lambda+n-k} c_\lambda^n$), the Stirling numbers of the second kind (as the coefficients $c_{(1^{n-k})}^n$ indexed by one-column shapes) and the Eulerian polynomials (whose coefficients are the sums $\sum_{\ell(\lambda)=k} c_\lambda^n$).

After reviewing in detail formulas and properties of the polynomials U_n and their coefficients, we generalize these results to a broader family of polynomials $U_{n,d}$, providing the normally ordered form of the operator $(h\partial^d)^n$ (Theorems 3.2 and 3.5). In addition, we present a new formula and combinatorial interpretation for the generalized Stirling numbers. We apply our study to the theory of formal differential operator rings. The latter are ubiquitous in representation theory: e.g., over the complex field, the enveloping algebra of a finite-dimensional solvable Lie algebra can be presented as an iterated differential operator ring (see [10]).

The motivation for this work comes from [1], where the powers of the derivation $h \frac{d}{dx}$ appear in the formulas for the action of a certain subalgebra A_h of the Weyl algebra A_1 on its irreducible modules, in case the base field has prime characteristic (see [1, Sec. 8] for more details). This also motivated our study of the modular behavior of the coefficients of U_n and, more generally, of the coefficients of $U_{n,d}$ in case n or d are prime powers, generalizing known results for Stirling numbers of both kinds.

1.2 Organization of the paper.

Section 2 formally defines the polynomials U_n and their coefficients, and presents their main combinatorial and arithmetical properties. It also prepares and motivates the generalizations of Section 3. Several simple descriptions of the polynomials U_n are provided, each leading naturally to the next one. Among these we give an “umbral formula” for U_n , i.e., a formula from which U_n is obtained by applying a linear map $x^\alpha \mapsto y_\alpha$ that “lowers” exponents to turn them into indices. This allows for an explicit presentation of the formula in terms of simple combinatorial objects. While some of these descriptions appear in previous works, we think we have been able to give here a rather clear presentation for them and explain well the natural connections between them. As an example, it is remarked in [6, appx. A] that “the symbolic problem [of calculating U_n] is indeed difficult”. They provide a recipe for computer calculation but it involves several steps, among them taking the compositional inverse of a power series by Lagrange inversion and extracting coefficients from powers of power series.

The various descriptions of the U_n given in 2.2 show that this calculation can be done in simpler ways.

Section 3 generalizes all the results and combinatorial interpretations in Section 2 to a family of polynomials $U_{n,d}$ related to the operator $(h\partial^d)^n$. There, generalizations of Comtet's formula for the coefficients of U_n are obtained (Corollary 3.11). Also, as a byproduct, a new formula and a new combinatorial interpretation of the generalized Stirling numbers are derived (Subsection 3.3). The section ends with the study in Theorem 3.23 of the congruence classes modulo a prime of the coefficients of U_n and $U_{n,d}$.

Lastly, in Section 4 it is shown how the polynomials $U_{n,d}$ provide the normally ordered form of elements in formal differential operator rings $A[z; \partial]$, including as special cases the Weyl algebra A_1 and its family of subalgebras A_h studied in [1, 2, 3].

1.3 Notations.

Let n be a nonnegative integer. Define $[n] = \{1, \dots, n\}$, so in particular $[0] = \emptyset$. The notation $(q)_n$ stands for the *falling factorial* $q(q-1)\cdots(q-n+1)$. An integer partition λ is a weakly decreasing sequence $\lambda_1 \geq \dots \geq \lambda_\ell > 0$ whose terms are called the parts of λ . If the sum of the parts of λ is n , then we write $|\lambda| = n$ or $\lambda \vdash n$. The number ℓ of parts of λ is the *length* of λ , denoted $\ell(\lambda)$.

Our rings are assumed to be associative and unital, but not necessarily commutative. A derivation of a ring A is an additive endomorphism ∂ of A satisfying the Leibniz rule

$$\partial(ab) = \partial(a)b + a\partial(b), \quad \forall a, b \in A.$$

In case A is an algebra over a field \mathbb{F} , it is further assumed that ∂ is linear. If $A = D[x]$ is a polynomial ring over a ring D , we denote its derivation $\frac{d}{dx}$ by ∂_x , so that ∂_x is zero on D and $\partial_x(x) = 1$. For $h \in A$, we denote $\partial^i(h)$ by $h^{[i]}$, reserving the classical notations h', h'' and $h^{(i)}$ for the special case $\partial = \partial_x$. In particular, $h^{[0]} = h$. Given an integer partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$, we define $h^{[\lambda]} = \prod_{i=1}^{\ell} h^{[\lambda_i]}$. In the same spirit, given commuting variables y_1, y_2, y_3, \dots , we define $y_\lambda = \prod_{i=1}^{\ell} y_{\lambda_i}$. In case $\lambda = \emptyset$ (the empty partition), then $y_\emptyset = 1 = h^{[0]}$.

2 The universal polynomials U_n and their coefficients c_λ^n

Let A be an arbitrary (noncommutative) ring and ∂ a derivation of A . For $h \in A$, the map $h\partial$ is not in general a derivation, except if h is central in A . **Thus, we assume throughout that h is central in A .**

Remark 2.1. Note that the derivation ∂ stabilizes the center of A . In particular, h central implies that $\partial^i(h)$ and $\partial^j(h)$ commute, for all $i, j \geq 0$. In fact, all the results in this paper hold if we replace the hypothesis that h is central with the weaker hypothesis that $\partial^i(h)$ and $\partial^j(h)$ commute, for all $i, j \geq 0$.

We will show that there is a *universal* polynomial U_n in the variables y_0, \dots, y_{n-1} and t such that, upon substituting $y_i = h^{[i]} \in A$ and $t = \partial$, we get the normally ordered form of the operator $(h\partial)^n$. By this we mean that we write $(h\partial)^n$ as a sum of terms of the form $a_k \partial^k$ with a_k a monomial in the $h^{[i]}$, for $i \geq 0$, but otherwise independent of A and ∂ (hence the term “universal”). In case A is a polynomial ring and $\partial = \partial_x$, this problem was studied by Scherk [17, para. 8], where formulas for U_n with $n \leq 5$ were obtained (see the account of

Scherk's dissertation in [6, appx. A]). Scherk's results have been revisited or rediscovered in recent times in [8, 4, 15], as pointed out in the detailed survey [13].

Then we proceed to explore the combinatorial properties of the U_n and of its coefficients, including, in 2.2.1, an umbral formula for computing U_n . We conclude the section with a closed formula for the coefficients of U_n and with a study of its modular behavior in case n is the power of a prime.

2.1 The normally ordered form of $(h\partial)^n$

Let $R = \mathbb{Z}[y_i \mid i \geq 0]$ be the integral commutative polynomial ring in the variables $\{y_i\}_{i \geq 0}$. As a free abelian group, R has a basis given by the monomials in the variables y_i . Care is needed when doing the substitution $t = \partial$, as ∂ does not commute with multiplication (e.g., $h\partial_x$ and $\partial_x h$ are different operators on $\mathbb{Z}[x]$). Thus, let $R\langle t \rangle$ be the ring obtained from R by adjoining a new variable t which does not commute with the variables y_i ; in other words, $R\langle t \rangle$ is the unital ring generated over \mathbb{Z} by $\{y_i\}_{i \geq 0}$ and t , subject only to the relations $y_i y_j = y_j y_i$ for all $i, j \geq 0$. Let Δ be the derivation of R defined on the generators by $\Delta(y_i) = y_{i+1}$ for all $i \geq 0$. Specifically, $\Delta = \sum_{i \geq 0} y_{i+1} \partial_{y_i}$. This map can be extended uniquely to a derivation of $R\langle t \rangle$, still denoted Δ , satisfying $\Delta(t) = 0$.

The polynomials U_n are defined recursively, as elements of $R\langle t \rangle$, as follows:

$$U_0 = 1 \quad \text{and} \quad \forall n \geq 0, \quad U_{n+1} = y_0(\Delta + \rho_t)U_n = y_0 U_n t + y_0 \Delta(U_n), \quad (2.2)$$

where ρ_t denotes the right multiplication by t operator on $R\langle t \rangle$. It is clear from the definition that $U_n = (y_0 \Delta + y_0 \rho_t)^n(1)$, and that this polynomial depends only on the variables y_0, \dots, y_{n-1} and t . For $n \geq 1$, we can write $U_n = \sum_{i=1}^n P_{n,i} t^i$, for some $P_{n,i} \in R$. Moreover, it can be seen that U_n is homogeneous of degree n relative to two different gradings on $R\langle t \rangle$:

- (i) the grading in which y_i has degree 1 and t has degree 0;
- (ii) the grading in which y_i has degree i and t has degree 1.

With this we can show that the polynomials U_n determine the normally ordered form of $(h\partial)^n$, for any ring A and derivation ∂ , and as such are universal, in an appropriate sense.

Theorem 2.3. *For any $n \geq 0$, there is a unique polynomial $U_n \in \bigoplus_{i \geq 0} R t^i$ such that, for any ring A , derivation ∂ of A and central element h in A ,*

$$(h\partial)^n = U_n|_{y_i=h^{[i]}, t=\partial} = U_n(h, h^{[1]}, h^{[2]}, \dots; \partial), \quad (2.4)$$

as endomorphisms of A .

Proof. Formally, the evaluation on (2.4) is the result of applying to U_n the ring homomorphism $\text{ev}_h : R\langle t \rangle \rightarrow \text{End}(A)$ sending y_i to $h^{[i]}$, seen as left multiplication by $h^{[i]}$, and t to ∂ . This is a well-defined map since the elements $h^{[i]}$ commute. Moreover,

$$\partial \circ \text{ev}_h(w) = \text{ev}_h((\Delta + \rho_t)(w)) \quad (2.5)$$

holds in $\text{End}(A)$, for all $w \in R\langle t \rangle$. (This could be seen as a version of the chain rule, adapted for our context.) Indeed, using the additive and multiplicative properties of ev_h and Δ we see that it suffices to check this identity on a generating set of $R\langle t \rangle$. In case $w = t$, the identity

is trivial because $\Delta(t) = 0$; in case $w = y_i$ we obtain the equality $\partial \circ h^{[i]} = h^{[i+1]} + h^{[i]} \circ \partial$. Thus, the latter equality follows from the fact that ∂ is a derivation of A .

It is now an easy matter to prove (2.4) by induction on n , using (2.5). The base case is clear, as $U_0 = 1$, and for the inductive step we have

$$(h\partial)^{n+1} = h \circ \partial \circ \text{ev}_h(U_n) = h \circ \text{ev}_h((\Delta + \rho_t)(U_n)) = \text{ev}_h(y_0\Delta(U_n) + y_0U_nt) = \text{ev}_h(U_{n+1}).$$

To show uniqueness, suppose $V_n \in \bigoplus_{i \geq 0} Rt^i$ satisfies (2.4). Take $A = R$, $h = y_0$ and $\partial = \Delta$, so that $\partial(y_i) = y_{i+1}$ for all $i \geq 0$. Let $T = U_n - V_n$. Then we can write $T = \sum_{i=0}^m P_i t^i$, for some $m \geq 0$ and $P_i \in R$. Thus, as in R we have $h^{[i]} = \Delta^i(y_0) = y_i$, we obtain $0 = T|_{y_i=h^{[i]}, t=\Delta} = \sum_{i=0}^m P_i \Delta^i$. Applying this operator to y_k , where k is large enough so that no y_j with $j \geq k$ occurs in any of the P_i , we conclude that $\sum_{i=0}^m P_i y_{k+i} = 0$, from which follows that $P_i = 0$ for all i and $V_n = U_n$. \square

Our next goal is to set a recurrence relation for the coefficients of U_n .

Proposition 2.6 (compare [17, par. 8.I.1], [15, Thm. 1], [1, Sec. 8]). *Assume $n \geq 1$. There exist positive integers c_λ^n , where λ runs through the set of integer partitions of size $0 \leq |\lambda| < n$, such that*

$$U_n = \sum_{k=1}^n \sum_{\lambda \vdash n-k} c_\lambda^n y_0^{n-\ell(\lambda)} y_\lambda t^k. \quad (2.7)$$

Additionally, the coefficients c_λ^n satisfy the recurrence relation

$$c_\emptyset^1 = 1, \quad c_\lambda^{n+1} = c_\lambda^n + \sum_{i=1}^n (\beta_{i-1} + 1) c_{\lambda[i]}^n, \quad (2.8)$$

where:

- $\beta_0 = n - \ell(\lambda)$ and, for $j \geq 1$, β_j is the multiplicity of j in λ ;
- $\lambda[i]$ is obtained from λ by subtracting 1 from a part of λ of size i , provided that $\beta_i > 0$;
- $c_{\lambda[i]}^n = 0$ if $\beta_i = 0$;
- $c_\lambda^n = 0$ if $\lambda \vdash m$ with $m \geq n$.

Proof. As we have observed before, U_n is a sum of integer multiples of monomials of the form $y_0^{a_0} \cdots y_{n-1}^{a_{n-1}} t^k$, with $1 \leq k \leq n$. Given such a monomial occurring in U_n with nonzero coefficient, we need to verify that:

- (a) $0 \leq 1a_1 + 2a_2 + \cdots + (n-1)a_{n-1} \leq n-1$, so that $y_1^{a_1} \cdots y_{n-1}^{a_{n-1}} = y_\lambda$, where λ is the partition with a_i parts of size i ;
- (b) $a_0 = n - \ell(\lambda) = n - (a_1 + \cdots + a_{n-1})$;
- (c) $k = n - |\lambda| = n - (1a_1 + \cdots + (n-1)a_{n-1})$.

These can be easily established by induction on n using (2.2) (actually, they are equivalent to the two homogeneous properties of U_n already mentioned) and so can the recurrence relation (2.8). Finally, the positivity of the coefficients c_λ^n can also be established by induction on n , using (2.8). \square

Example 2.9. For $\nu = (2, 1)$, we have

$$c_\nu^5 = c_\nu^4 + 3c_{\nu[1]}^4 + 2c_{\nu[2]}^4 = c_{2,1}^4 + 3c_2^4 + 2c_{1,1}^4 = 4 + 3 \times 4 + 2 \times 7 = 30,$$

and for $\nu = (2, 1, 1)$, we have

$$c_\nu^5 = c_\nu^4 + 2c_{\nu[1]}^4 + 3c_{\nu[2]}^4 = 2c_{2,1}^4 + 3c_{1,1,1}^4 = 2 \times 4 + 3 \times 1 = 11.$$

Remark 2.10. By Proposition 2.6, the number of monomials in U_n is the number of integer partitions of size at most $n - 1$ (up to an offset, this is sequence A000070 in [16]).

2.2 Combinatorial interpretations of the polynomials U_n

We introduce several descriptions of the polynomials U_n , each leading naturally to the next one. The first description is an “umbral formula” for U_n , i.e., a formula from which U_n is obtained by applying a linear map $x^\alpha \mapsto y_\alpha$ that “lowers” exponents to turn them into indices. The summands happen to be naturally indexed by simple combinatorial objects: strictly subdiagonal maps. This gives an explicit presentation of the formula for U_n . Another presentation and interpretation of the formula is given by increasing trees.

Although our presentation is new, most results in this subsection are known in some other form: they appear in Mohammad–Noori [15] or the (seemingly unrelated) work of Hivert, Novelli and Thibon [11] on the solutions of certain differential equations. We have tried to present them as clearly and explicitly as possible. This serves for setting notations and preparing the generalizations of Section 3.

2.2.1 Umbral formula

We continue to assume that A is an arbitrary ring and ∂ is a derivation of A . Consider the tensor product (over \mathbb{Z}):

$$A^{\otimes n} = A \otimes A \otimes \cdots \otimes A \quad (n \text{ copies of } A),$$

where we index the factors in the tensor product from right to left and from 0 to $n - 1$:

$$n - 1, n - 2, \dots, 1, 0.$$

We denote by $m_n : A^{\otimes n} \rightarrow A$ the n -ary multiplication map and by $\partial_{n,i}$ the map $A^{\otimes n} \rightarrow A^{\otimes n}$ that applies ∂ on factor i and the identity on the others. If $i \leq n - 1$, then $\partial_{n+1,i} = 1_A \otimes \partial_{n,i}$ so, to lighten the notation, we henceforth use ∂_i to denote any of the maps $\partial_{n,i}$, as long as $n \geq i + 1$.

For $n = 2$, the Leibniz identity can be stated as $\partial \circ m_2 = m_2 \circ (\partial_1 + \partial_0)$. By an easy induction, we get that, for any $n \geq 1$,

$$\partial \circ m_n = m_n \circ \sum_{i=0}^{n-1} \partial_i. \tag{2.11}$$

Using this identity, it is derived, again by induction, that for any $n \geq 0$ and any $h_0, h_1, \dots, h_n \in A$,

$$h_n \partial h_{n-1} \partial \cdots \partial h_1 \partial (h_0) = m_{n+1} \circ \left(\prod_{i=0}^{n-1} (\partial_i + \cdots + \partial_1 + \partial_0) \right) (h_n \otimes \cdots \otimes h_1 \otimes h_0).$$

In particular, for any $a, h \in A$ and any $n \geq 0$,

$$(h\partial)^n(a) = m_{n+1} \circ \left(\prod_{i=0}^{n-1} (\partial_i + \cdots + \partial_1 + \partial_0) \right) (h \otimes \cdots \otimes h \otimes h \otimes a).$$

This yields the following recipe for calculating U_n .

Theorem 2.12. *For any $n \geq 0$, the polynomial U_n is obtained by applying to the product*

$$\prod_{i=0}^{n-1} (x_i + \cdots + x_2 + x_1 + x_0) \tag{2.13}$$

the \mathbb{Z} -linear map from $\mathbb{Z}[x_0, x_1, \dots, x_n]$ to $R\langle t \rangle$ defined by

$$x_n^{\alpha_n} x_{n-1}^{\alpha_{n-1}} \cdots x_1^{\alpha_1} x_0^k \longmapsto y_{\alpha_n} y_{\alpha_{n-1}} \cdots y_{\alpha_1} t^k.$$

Remark 2.14. The product in (2.13) contains no occurrence of x_n . This is deliberate so that y_0 is always a factor of U_n , for $n > 0$. Notice in particular that the term x_0^n from (2.13) is mapped to $y_0^n t^n \in R\langle t \rangle$, which is always a term of U_n .

2.2.2 Enumeration of subdiagonal maps

The expansion of the product (2.13) is obtained by choosing a summand $x_{f(i)}$ for the i -th factor $(x_{i-1} + \cdots + x_2 + x_1 + x_0)$ and summing over all possibilities. This expansion is thus

$$\sum_{f \in \text{SD}_n} x_{f(n)} x_{f(n-1)} \cdots x_{f(2)} x_{f(1)}$$

where SD_n is the set of all maps $f : [n] \rightarrow [n] \cup \{0\}$ that are *subdiagonal*, i.e., that satisfy $f(i) < i$ for all $1 \leq i \leq n$. (Although n is not in the image of f , it will be convenient to include n in the codomain of the elements in SD_n .) Gathering the variables, we get that this expansion is:

$$\sum_{f \in \text{SD}_n} \prod_{i=0}^{n-1} x_i^{\#f^{-1}(\{i\})}.$$

In conjunction with Theorem 2.12, the above gives another description of U_n .

Theorem 2.15. *For any $n \geq 0$,*

$$U_n = \sum_{f \in \text{SD}_n} \left(\prod_{i=1}^n y_{\#f^{-1}(\{i\})} \right) t^{\#f^{-1}(\{0\})}.$$

It will be interesting, when studying the coefficients c_λ^n , to consider rather than the functions $f \in \text{SD}_n$, the corresponding partial maps from $[n]$ to $[n]$.

A *partial map* g from a set A to a set B is a map from some subset, $\text{Dom}(g)$, of A to B . Let PD_n be the set of all subdiagonal partial maps from $[n]$ to $[n]$, i.e., all maps g from some subset $\text{Dom}(g)$ of $[n]$ to $[n]$, satisfying $g(i) < i$, for all $i \in \text{Dom}(g)$.

Theorem 2.16. *For any $n \geq 0$,*

$$U_n = \sum_{g \in \text{PD}_n} \left(\prod_{i=1}^n y_{\#g^{-1}(\{i\})} \right) t^{n - \#\text{Dom}(g)}.$$

2.2.3 Expansion in increasing trees

Following Cayley [7], Hivert, Novelli and Thibon [11], Mohammad–Noori [15], and Blasiak and Flajolet [6], we interpret the calculations of iterated derivatives in terms of increasing trees. This is done by interpreting the map $f \in \text{SD}_n$ as the map that associates to each node $j \in [n]$ its father in a tree rooted at 0. This gives a bijection from SD_n to the set \mathbb{T}_n of all increasing rooted trees with vertex set $\{0, 1, \dots, n\}$. Then, for each i , $\#f^{-1}(\{i\})$ is the number of children of node i . From this and Theorem 2.15 follows the following formula.

Theorem 2.17 ([15, 4]). *For any $n \geq 0$,*

$$U_n = \sum_{T \in \mathbb{T}_n} \left(\prod_{i=1}^n y_{\text{ch}(i;T)} \right) t^{\text{ch}(0;T)}, \quad (2.18)$$

where $\text{ch}(i;T)$ stands for the number of children (outdegree) of node i in the rooted tree T .

2.2.4 Solution of the differential equation $x'(u) = y(x(u))$

Given a formal series $y(u) = \sum_{i=0}^{\infty} y_i \frac{u^i}{i!}$, there is a unique formal series $x(u) = \sum_{n=1}^{\infty} x_n \frac{u^n}{n!}$ such that $\frac{dx}{du} = y(x(u))$. Hivert, Novelli and Thibon show in [11, Formula (43)] that its coefficients are given by

$$x_n = \sum_{T \in \mathbb{T}_{n-1}} \prod_{i=0}^{n-1} y_{\text{ch}(i;T)},$$

which is obtained from U_n by replacing t^k with y_k , for all $k \geq 0$.

2.3 Combinatorial interpretations of the coefficients c_{λ}^n

We derive some interesting specializations and properties of the polynomials U_n and their coefficients c_{λ}^n . After recalling from [8, 15, 4] the relation to Stirling numbers of both kinds and Eulerian numbers, we show that the generalized Stirling numbers $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,1}$ appear in specializations of the polynomials U_n .

2.3.1 Stirling numbers of the first kind

If we set $y_i = 1$ for all $i \geq 0$ in U_n then, by Theorem 2.12, this corresponds to taking $x_j = 1$ for all $j \geq 1$ and $x_0 = t$ in (2.13), so we get

$$U_n(1, 1, 1, \dots; t) = \prod_{i=0}^{n-1} (t+i) = t(t+1) \cdots (t+n-1). \quad (2.19)$$

Since the latter expression coincides with the generating function $\sum_{k=1}^n c(n, k) t^k$ for the (sign-less) Stirling numbers of the first kind, where $c(n, k)$ counts the number of permutations in S_n with exactly k cycles, we deduce from (2.7) that

$$\sum_{\lambda \vdash n-k} c_{\lambda}^n = c(n, k). \quad (2.20)$$

This result appears in [8, §5], [15, Prop. 9 (iii)] and [4, p. 274].

2.3.2 Generalized Stirling numbers, Stirling numbers of the second kind and Bell numbers

The *partition Stirling numbers*, or *Stirling numbers of the second kind*, $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$, count the set partitions of $[n]$ having k blocks. It is well known that the partition Stirling numbers are precisely the coefficients of the normally ordered form of $(x\partial_x)^n$, for $A = \mathbb{Z}[x]$:

$$(x\partial_x)^n = \sum_k \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k \partial_x^k.$$

This identity leads to a natural generalization of the partition Stirling numbers, simply called *generalized Stirling numbers* and denoted $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d}$: these are the coefficients in the normally ordered form of $(x^q \partial_x^d)^n$ in $\mathbb{Z}[x]$. More precisely:

$$\begin{aligned} (x^q \partial_x^d)^n &= x^{(q-d)n} \sum_k \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d} x^k \partial_x^k && \text{if } q \geq d, \\ (x^q \partial_x^d)^n &= \left(\sum_k \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d} x^k \partial_x^k \right) \partial_x^{(d-q)n} && \text{if } q \leq d. \end{aligned} \tag{2.21}$$

The generalized Stirling numbers have been studied extensively (see specially [5, 4.7.1], [6, Note 18] and the references in the survey [13]), and in particular the question of finding a simple combinatorial interpretation for them has been raised. This problem was solved (even for more general monomials in x and ∂_x) in [19], in terms of rook placements on chessboards that are Young diagrams and, in [9], in terms of constrained partitions of vertex sets of graphs.

Observe that, for $d = 1$, the right-hand side of (2.21) is equal to the specialization $U_n(h, h', h'', \dots; \partial_x)$, with $h = x^q$. Evaluate at $x = 1$, and use that, for all k , $h^{(k)}(1) = (q)_k$, the falling factorial, to get

$$U_n((q)_0, (q)_1, (q)_2, \dots; \partial_x) = \sum_k \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,1} \partial_x^k. \tag{2.22}$$

Therefore, the specialization of U_n at $y_k = (q)_k$, for q a nonnegative integer, is the ordinary generating series for the generalized Stirling numbers $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,1}$ and it follows that

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,1} = \sum_{\lambda \vdash n-k} c_\lambda^n \prod_i (q)_{\lambda_i}. \tag{2.23}$$

The generalized Stirling numbers $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d}$, for arbitrary d , will be obtained in the same fashion from the polynomials $U_{n,d}$ in Subsection 3.3.

Further specialization of (2.22) at $q = 1$ gives

$$U_n|_{y_0=y_1=1, y_i=0 \forall i>1} = \sum_{k=1}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} t^k,$$

which is the generating function for the partition Stirling numbers (*Touchard polynomials*). In particular, $c_{1^{n-k}}^n = \left\{ \begin{matrix} n \\ k \end{matrix} \right\}$ (compare [8, §5], [15, Prop. 9 (ii)] and [4, p. 275]). Taking $t = 1$ we obtain the Bell number

$$B_n = \sum_{k=1}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} = U_n|_{y_0=y_1=t=1, y_i=0 \ \forall i>1},$$

which counts the total number of set partitions of $[n]$.

2.3.3 Eulerian polynomials

The coefficient of q^k in $U_n(q, 1, 1, \dots; 1)$ is the number of increasing trees $T \in \mathbb{T}_n$ with exactly k leaves. By [18, Prop. 1.5.5], this is precisely the Eulerian number $A(n, k)$, equal to the number of permutations in S_n with exactly $k - 1$ descents. Thence, this specialization of U_n is the Eulerian polynomial $A_n(q)$:

$$U_n(q, 1, 1, \dots; 1) = \sum_{\pi \in S_n} q^{1+\text{no. of descents of } \pi} = A_n(q).$$

The same argument also shows that the coefficient of q^k in $U_n(1, q, q, \dots; 1)$ is the Eulerian number $A(n, n - k)$. Since the Eulerian polynomials are palindromic (i.e., $q^{n+1}A_n(1/q) = A_n(q)$), these properties, in conjunction, show that we have

$$A_n(q) = U_n(q, 1, 1, \dots; 1) = qU_n(1, q, q, \dots; 1).$$

In particular, we obtain

$$\sum_{\lambda: \ell(\lambda)=k} c_\lambda^n = A(n, k+1) = \text{no. of } \pi \in S_n \text{ with exactly } k \text{ descents}, \quad (2.24)$$

a result which is implicit in [15, Prop. 9 (iv)] and [4, p. 275].

2.4 A closed formula for the c_λ^n

In this subsection we investigate combinatorial interpretations of the coefficients c_λ^n . A quite useful one follows directly from (2.18): c_λ^n is the number of trees $T \in \mathbb{T}_n$ whose internal vertices, other than the root, have as numbers of children the parts of λ .

In what follows, we will investigate another kind of combinatorial interpretation, based on the description of U_n in 2.2.2. Our starting point is Theorem 2.16. Define the *type* of a subdiagonal partial map $g \in \text{PD}_n$ as the partition whose parts are the cardinalities of its fibers.

Corollary 2.25. *For any n and any partition λ , the coefficient c_λ^n is the number of subdiagonal partial maps from $[n]$ to $[n]$ of type λ .*

This interpretation provides an explicit formula for the coefficients c_λ^n (see [15, Prop. 2] for a “balls in urns” description). The map

$$g \in \text{PD}_n \mapsto (g^{-1}(\{n-1\}), g^{-1}(\{n-2\}), \dots, g^{-1}(\{1\}))$$

establishes a bijection from PD_n to the set of all sequences $(R_1, R_2, \dots, R_{n-1})$ of pairwise disjoint subsets of $[n]$ satisfying $R_j \subseteq [n] \setminus [n-j]$, for all j . The type of g is the multiset of the nonzero cardinalities of the corresponding sets R_i . The number of sequences $(R_1, R_2, \dots, R_{n-1})$ with given cardinalities i_1, i_2, \dots, i_{n-1} is

$$\prod_{j=1}^{n-1} \binom{j - i_1 - i_2 - \dots - i_{j-1}}{i_j}, \quad (2.26)$$

since one can build any such sequence by choosing first R_1 in $\{n\}$ with i_1 elements, then $R_2 \subseteq \{n-1, n\}$ disjoint from R_1 and with i_2 elements. Once R_1, R_2, \dots, R_{j-1} have been chosen, the i_j elements of R_j have to be selected from the set

$$\{n-j+1, n-j+2, \dots, n\} \setminus (R_1 \sqcup R_2 \sqcup \dots \sqcup R_{j-1}),$$

whose cardinality is $j - i_1 - i_2 - \dots - i_{j-1}$. For this, there are $\binom{j - i_1 - i_2 - \dots - i_{j-1}}{i_j}$ possibilities.

This yields the following formula.

Corollary 2.27. *Let $n \geq k \geq 1$ and λ be a partition of $n - k$. Then*

$$c_\lambda^n = \sum_{i_1, \dots, i_{n-1}} \prod_{j=1}^{n-1} \binom{j - i_1 - \dots - i_{j-1}}{i_j}. \quad (2.28)$$

where the sum is carried over all sequences of nonnegative integers whose nonzero terms are the parts of λ .

Of course, the sum in the corollary above can be restricted to the sequences $(i_1, i_2, \dots, i_{n-1})$ such that $i_1 + i_2 + \dots + i_j \leq j$, for all j , since the other sequences have a zero contribution. Under this hypothesis, the binomial coefficients $\binom{j - i_1 - \dots - i_{j-1}}{i_j}$ in (2.28) can be expanded as usual. Cancellations in the product of binomial coefficients leads to the following formula, due to Comtet [8, Formula (8)]:

Corollary 2.29 ([8, 4, 15]). *Let $n \geq k \geq 1$ and λ be a partition of $n - k$. Then*

$$c_\lambda^n = \frac{1}{(k-1)!} \frac{1}{\prod_i (\lambda_i)!} \sum_{i_1, \dots, i_{n-1}} \prod_{j=1}^{n-1} \binom{j - i_1 - i_2 - \dots - i_{j-1}}{i_j}, \quad (2.30)$$

where the sum is carried over all sequences (i_1, \dots, i_{n-1}) of nonnegative integers whose nonzero terms are the parts of λ , up to reordering, and such that $i_1 + i_2 + \dots + i_j \leq j$, for all j .

Corollaries 2.27 and 2.29 will be generalized in Corollary 3.11.

Example 2.31. Consider the case when λ has $n - k$ parts, all equal to 1. We have established in 2.3.2 that $c_{1^{n-k}}^n$ is the partition Stirling number $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$. But $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = c_{1^{n-k}}^n$ also counts the subdiagonal partial maps from $[n]$ to $[n]$ of type 1^{n-k} , which correspond to the *rook placements* of $n - k$ rooks, all below the diagonal, in a $n \times n$ chessboard. This interpretation of the partition Stirling numbers appears in [19], and Corollary 2.25 is thus a generalization of it.

2.5 The coefficients c_λ^n modulo a prime

Let p be a prime. It is well known that most Stirling numbers of both kinds $c(p, k)$ and $\left\{ \begin{matrix} p \\ k \end{matrix} \right\}$ vanish modulo p . We will see that this property is shared by the coefficients c_λ^p and, more generally, by c_λ^n , in case n is a power of p .

Theorem 2.32. *For any prime p , prime power $n = p^m$ and partition λ with $|\lambda| \neq n - 1$ and $|\lambda|$ not a multiple of p , we have $c_\lambda^n \equiv 0 \pmod{p}$. In particular, if $|\lambda| \neq 0, p - 1$, then $c_\lambda^p \equiv 0 \pmod{p}$.*

Recall from 2.3.1 and 2.3.2 that

$$c(n, k) = \sum_{\lambda \vdash n-k} c_\lambda^n, \quad \left\{ \begin{matrix} n \\ k \end{matrix} \right\} = c_{1^{n-k}}^n \quad \text{and} \quad \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,1} = \sum_{\lambda \vdash n-k} c_\lambda^n \prod_i (q)_{\lambda_i}.$$

Thus, Theorem 2.32 says in particular that for any prime p , $n = p^m$ and $1 < k < n$ not a multiple of p , the Stirling numbers of both kinds $c(n, k)$ and $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$ as well as the generalized Stirling numbers $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,1}$ are multiples of p .

Proof. For each $n > 0$, we will build an action of the cyclic group \mathbb{Z}_n on T_n . This action will preserve the outdegree (number of children) of vertex 0, and will preserve as well the multiset of outdegrees of all other vertices. Therefore, it will be possible to write Formula (2.18) as

$$U_n = \sum_{\omega \in \mathsf{T}_n / \mathbb{Z}_n} \#\omega \cdot \left(\prod_{i=1}^n y_{\text{ch}(i; T_\omega)} \right) t^{\text{ch}(0; T_\omega)}.$$

The sum is carried over all orbits ω for this action and T_ω is an arbitrarily chosen tree in ω .

Fix a prime p and let $n = p^m$. We can assume that $m > 0$. The cardinality of any orbit ω will also be a power of p , hence either 1 or a multiple of p . We will show that if $T \in \mathsf{T}_n$ is fixed under the action, then either $\text{ch}(0; T) = 1$ or $\text{ch}(0; T)$ is a multiple of p , from which the theorem will follow.

The action of \mathbb{Z}_n that we will define is based on the bijection between T_n and the set Σ_n of all sets

$$\{(t_1, A_1), (t_2, A_2), \dots, (t_k, A_k)\}$$

where the A_i are the blocks of a set partition of $[n]$ and, for each i , t_i is an increasing tree whose number of vertices is $\#A_i$. This bijection is defined as follows: given $T \in \mathsf{T}_n$, delete its root 0. An increasing forest with vertex set $[n]$ is obtained. Let t'_1, t'_2, \dots, t'_k be its components (each is an increasing tree). Define A_i as the set of vertices of t'_i . Enumerate the elements of A_i in increasing order: $A_i = \{v_1 < v_2 < \dots < v_r\}$; and define t_i as the tree obtained from t'_i by replacing, for each j , the number v_j with its index j .

The natural action of the cyclic group \mathbb{Z}_n on $[n]$ induces naturally an action on the set of all set partitions of $[n]$, and this in turn induces an action on Σ_n . The bijection $\Sigma_n \cong \mathsf{T}_n$ is then used to transport this action to an action on T_n .

Assume $T \in \mathsf{T}_n$ is fixed under this action of \mathbb{Z}_n and let $\{A_1, \dots, A_k\}$ be the corresponding set partition of $[n]$. In particular, $\text{ch}(0; T) = k$ and it remains to show that either $k = 1$ or

p divides k . Since T is fixed, \mathbb{Z}_n permutes the blocks of the partition and we consider this induced action next. If a certain block A_i is fixed, then necessarily $A_i = [n]$ and thus $k = 1$. Otherwise, the size of the orbit of each block is divisible by p , thence so is k . \square

In Subsection 3.4 below we generalize Theorem 2.32 to the coefficients $c_\lambda^{n,d}$.

3 Generalization: normal ordering for $(h\partial^d)^n$

Our approach generalizes naturally to the study of the normal ordering of the operator $(h\partial^d)^n$, for any positive integer d .

3.1 Formulas for the normal ordering of $(h\partial^d)^n$

Recall the derivation $\Delta = \sum_{i \geq 0} y_{i+1} \partial_{y_i}$ of $R\langle t \rangle$. Given any $d \geq 1$, we recursively define a family $U_{n,d}$ of elements of $R\langle t \rangle$, extending the polynomials U_n , as follows:

$$U_{0,d} = 1 \quad \text{and} \quad \forall n \geq 0, \quad U_{n+1,d} = y_0(\Delta + \rho_t)^d U_{n,d}, \quad (3.1)$$

where, as in (2.2), ρ_t stands for right multiplication by t .

When $d = 1$ we retrieve the polynomials U_n , and it is not surprising that many of the properties which we have established for the U_n generalize to the $U_{n,d}$. Among these we have that, for $n \geq 1$, $U_{n,d}$ is a sum of monomials of the form $P(y_0, \dots, y_{(n-1)d})t^k$ with $P \in R$ and $d \leq k \leq nd$ and that $U_{n,d}$ is homogeneous:

- (i) of degree n relative to the grading in which y_i has degree 1 and t has degree 0;
- (ii) of degree nd relative to the grading in which y_i has degree i and t has degree 1.

The main property of these polynomials is the analogue of Theorem 2.3, which we state below.

Theorem 3.2. *For any $n \geq 0$ and $d \geq 1$, there is a unique polynomial $U_{n,d} \in \bigoplus_{i \geq 0} R t^i$ such that, for any ring A , derivation ∂ of A and central element h in A ,*

$$(h\partial^d)^n = U_{n,d}|_{y_i=h^{[i]}, t=\partial} = U_{n,d}(h, h^{[1]}, h^{[2]}, \dots; \partial), \quad (3.3)$$

as endomorphisms of A .

Proof. The proof is identical to the proof of Theorem 2.3 and is based on

$$\partial^d \circ \text{ev}_h = \text{ev}_h \circ (\Delta + \rho_t)^d \quad (3.4)$$

which follows from (2.5). \square

To avoid repetition, we sketch briefly how the properties of U_n covered in Subection 2.2 generalize to $U_{n,d}$.

Theorem 3.5. *Let $n \geq 0$ and $d \geq 1$.*

(a) The polynomial $U_{n,d}$ is obtained by applying to the product

$$\prod_{i=0}^{n-1} (x_i + \cdots + x_2 + x_1 + x_0)^d$$

the \mathbb{Z} -linear map from $\mathbb{Z}[x_0, x_1, \dots, x_n]$ to $R\langle t \rangle$ defined by

$$x_n^{\alpha_n} x_{n-1}^{\alpha_{n-1}} \cdots x_1^{\alpha_1} x_0^k \mapsto y_{\alpha_n} y_{\alpha_{n-1}} \cdots y_{\alpha_1} t^k.$$

(b) Let $\text{PD}_{n,d}$ be the set of all partial maps $[n] \times [d] \rightarrow [n]$ such that $\forall i, j, f(i, j) < i$. Then

$$U_{n,d} = \sum_{f \in \text{PD}_{n,d}} \left(\prod_{i=1}^n y_{\#f^{-1}(\{i\})} \right) t^{nd - \#\text{Dom}(f)}. \quad (3.6)$$

(c) For $F = (T_1, \dots, T_d) \in \mathbb{T}_n^d$, set $\text{ch}(i; F) = \sum_{j=1}^d \text{ch}(i; T_j)$. Then

$$U_{n,d} = \sum_{F \in \mathbb{T}_n^d} \left(\prod_{i=1}^n y_{\text{ch}(i; F)} \right) t^{\text{ch}(0; F)}. \quad (3.7)$$

Proof. Firstly, as before, by (2.11) and induction, we obtain

$$\begin{aligned} h_n \partial^{d_{n-1}} h_{n-1} \partial^{d_{n-2}} \cdots h_1 \partial^{d_0} (h_0) \\ = m_{n+1} \circ \left(\prod_{i=0}^{n-1} (\partial_i + \cdots + \partial_1 + \partial_0)^{d_i} \right) (h_n \otimes \cdots \otimes h_1 \otimes h_0). \end{aligned}$$

In particular, for any $a, h \in A$, and any $n, d \geq 0$,

$$(h \partial^d)^n (a) = m_{n+1} \circ \left(\prod_{i=0}^{n-1} (\partial_i + \cdots + \partial_1 + \partial_0)^d \right) (h \otimes \cdots \otimes h \otimes a).$$

Description (a) of $U_{n,d}$ follows from this. Now,

$$\begin{aligned} \prod_{i=0}^{n-1} (\partial_i + \cdots + \partial_1 + \partial_0)^d &= \left(\sum_{g \in \text{PD}_n} \partial_0^{n - \#\text{Dom}(g)} \prod_{i=1}^{n-1} \partial_i^{\#g^{-1}(\{i\})} \right)^d \\ &= \sum_{(g_1, \dots, g_d) \in (\text{PD}_n)^d} \partial_0^{nd - \sum_i \#\text{Dom}(g_i)} \prod_{i=1}^{n-1} \partial_i^{\sum_j \#g_j^{-1}(\{i\})}. \end{aligned}$$

This formula becomes (3.6) after making use of the bijection $(\text{PD}_n)^d \cong \text{PD}_{n,d}$ that sends (g_1, \dots, g_d) to the partial map $(i, j) \mapsto g_j(i)$ defined on the pairs (i, j) such that $i \in \text{Dom}(g_j)$. Finally, we have also

$$\prod_{i=0}^{n-1} (\partial_i + \cdots + \partial_1 + \partial_0)^d = \left(\sum_{T \in \mathbb{T}_n} \prod_{i=0}^{n-1} \partial_i^{\text{ch}(i; T)} \right)^d = \sum_{(T_1, \dots, T_d) \in \mathbb{T}_n^d} \prod_{i=0}^{n-1} \partial_i^{\sum_j \text{ch}(i; T_j)}.$$

Formula (3.7) follows from this. □

3.2 The coefficients $c_\lambda^{n,d}$

Let us single out the coefficients of $U_{n,d}$, as in the first part of Proposition 2.6.

Proposition 3.8. *Assume $n, d \geq 1$. There exist positive integers $c_\lambda^{n,d}$, where λ runs through the set of partitions of size $0 \leq |\lambda| \leq (n-1)d$ with at most $n-1$ parts, such that*

$$U_{n,d} = \sum_{k=d}^{nd} \sum_{\substack{\lambda \vdash nd-k \\ \ell(\lambda) \leq n-1}} c_\lambda^{n,d} y_0^{n-\ell(\lambda)} y_\lambda t^k. \quad (3.9)$$

It is obvious from the definition that $c_\lambda^{n,1} = c_\lambda^n$. By Theorem 3.5, we can give several combinatorial descriptions of the coefficients $c_\lambda^{n,d}$.

Here is a description of $c_\lambda^{n,d}$ that generalizes the description of the c_λ^n as counting subdiagonal partial maps (Theorem 2.16).

Proposition 3.10. *The coefficient $c_\lambda^{n,d}$ counts the elements of $\text{PD}_{n,d}$ of type λ , i.e., the partial maps g from $[n] \times [d]$ to $[n]$ fulfilling the condition*

$$\forall i, j, \quad g(i, j) < i,$$

and whose cardinalities of the fibers form the parts of λ .

Below we generalize Corollaries 2.27 and 2.29.

Corollary 3.11. *Let $n, d \geq 1$, and k such that $nd \geq k \geq d$. Let λ be a partition of $nd - k$ with length at most $n-1$.*

(a) *We have*

$$c_\lambda^{n,d} = \sum_{i_1, \dots, i_{n-1}} \prod_{j=1}^{n-1} \binom{jd - i_1 - \dots - i_{j-1}}{i_j}, \quad (3.12)$$

where the sum is carried over all sequences i_1, \dots, i_{n-1} of nonnegative integers whose nonzero terms are the parts of λ (up to reordering).

(b) *We have the following generalization of Comtet's formula (Corollary 2.29):*

$$c_\lambda^{n,d} = \frac{1}{(k-1)!} \frac{1}{\prod_i (\lambda_i!)} \sum_{i_1, \dots, i_{n-1}} \prod_{j=1}^{n-1} \binom{jd - i_1 - \dots - i_{j-1}}{i_j}, \quad (3.13)$$

where the sum is carried over all sequences i_1, i_2, \dots, i_{n-1} of nonnegative integers whose nonzero terms are the parts of λ (up to reordering) such that $i_1 + i_2 + \dots + i_j \leq jd$, for all j .

(c) *Let $\mathbf{M}(n)$ be the set of all lower triangular arrays $a = (a_{i,j})_{1 \leq j < i \leq n}$ of nonnegative integers. To any such array associate the sequence $(r_2(a), r_3(a), \dots, r_n(a))$ of its row sums ($r_i(a) = \sum_j a_{i,j}$), and the sequence $(c_1(a), c_2(a), \dots, c_{n-1}(a))$ of its column sums ($c_j(a) = \sum_i a_{i,j}$). Then:*

$$c_\lambda^{n,d} = \sum_a \frac{1}{\prod_{i,j} (a_{i,j}!)} \prod_{i=2}^n (d)_{r_i(a)}, \quad (3.14)$$

where the sum is carried over all $a \in \mathbf{M}(n)$ whose nonzero column sums equal the parts of λ , up to reordering.

Proof. The proofs of (a) and (b) are straightforward adaptations of those of Corollaries 2.27 and 2.29, and as such are omitted.

Let us prove (c). Associate to any $g \in \text{PD}_{n,d}$ the array $a = (a_{i,j}) \in \mathbf{M}(n)$ such that $a_{i,j} = \#\{k \in [d] \mid g(i,k) = j\}$. Given $a \in \mathbf{M}(n)$, the elements $g \in \text{PD}_{n,d}$ corresponding to a are obtained by:

- Firstly, choosing for each i , a subset $S_i \subseteq [d]$ with $r_i(a)$ elements (to be the set of all k such that $(i,k) \in \text{Dom}(g)$). There are $\binom{d}{r_i(a)}$ possibilities for this choice.
- Next, choosing for each i , and each $k \in S_i$, the value of $g(i,k)$ in $[i-1]$, so that each $j \in [i-1]$ is chosen $a_{i,j}$ times. There are $\binom{r_i(a)}{a_{i,1}, a_{i,2}, \dots, a_{i,i-1}}$ possibilities for this choice.

This yields the formula

$$c_\lambda^{n,d} = \sum_a \prod_i \binom{d}{r_i(a)} \binom{r_i(a)}{a_{i,1}, a_{i,2}, \dots, a_{i,i-1}},$$

where the sum is carried over all $a \in \mathbf{M}(n)$ with nonzero column sums equal to the parts of λ . This formula simplifies into that of the corollary. \square

Example 3.15. Let us compute $c_\lambda^{3,3}$, i.e., the coefficients occurring in the normally ordered form of $(h\partial^3)^3$, using (3.12). In general, we need to determine all different $(n-1)$ -tuples (i_1, \dots, i_{n-1}) which can be obtained by permuting the parts of λ and adding zero parts, if necessary. However, these terms will give a zero contribution unless

$$i_1 + \dots + i_j \leq jd, \quad \text{for all } 1 \leq j \leq n-1, \quad (3.16)$$

which narrows down the number of $(n-1)$ -tuples to consider, as will now be illustrated.

We need to consider partitions of size at most 6 with no more than 2 parts. For example, if $\lambda = 4$, then the possibilities are $(4,0)$ and $(0,4)$, but the former violates (3.16), so we get

$$c_4^{3,3} = \binom{3}{0} \binom{6}{4} = 15.$$

Similarly, for $\lambda = (2,1)$ the possibilities are $(2,1)$ and $(1,2)$, giving

$$c_{2,1}^{3,3} = \binom{3}{2} \binom{4}{1} + \binom{3}{1} \binom{5}{2} = 42.$$

Proceeding as above, we obtain all of the coefficients $c_\lambda^{3,3}$ shown below.

λ	\emptyset	\square	$\begin{array}{ c } \hline \square \\ \hline \end{array}$	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c } \hline \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c } \hline \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c c } \hline \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c } \hline \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c c } \hline \square & \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c c c } \hline \square & \square & \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c c c c } \hline \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c c c c c } \hline \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c c c c c c } \hline \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c c c c c c c } \hline \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$\begin{array}{ c c c c c c c c c c c c c c c } \hline \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$
$c_\lambda^{3,3}$	1	9	15	18	42	21	33	18	15	15	15	6	1	3	3	1	

Note that $\sum_\lambda c_\lambda^{3,3} = 216 = (3!)^3$.

Corollary 3.17. *Let $n, d \geq 1$. Then*

$$\sum_{k=d}^{nd} \sum_{\lambda \vdash nd-k} c_\lambda^{n,d} = n!^d \quad \text{and} \quad \sum_{\lambda \vdash (n-1)d} c_\lambda^{n,d} = (n-1)!^d.$$

Proof. We will use the description of $c_\lambda^{n,d}$ provided in Proposition 3.10. The first sum is just $\#\text{PD}_{n,d} = n!^d$. For the second sum, observe that if $g \in \text{PD}_{n,d}$ has type λ then $\#\text{Dom}(g) = |\lambda|$ and $(n-1)d$ is the maxim possible cardinality of the domain of an element in $\text{PD}_{n,d}$. So, the second sum counts the number of functions g from $([n] \setminus \{1\}) \times [d]$ to $[n]$ satisfying $g(i, k) < i$ for all i, k , which equals $(n-1)!^d$. \square

3.3 Application to generalized Stirling numbers

Now we extend the result of Subsection 2.3 giving the generalized Stirling numbers $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,1}$ as the coefficients of $U_n((q)_0, (q)_1, (q)_2, \dots; t)$.

Proposition 3.18. *Let q, d, n, k be nonnegative integers, with $q \geq d$. The generalized Stirling number $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d}$, defined by (2.21), is given by*

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d} = \sum_{\substack{\lambda \vdash nd-k \\ \ell(\lambda) \leq n-1}} c_\lambda^{n,d} \prod_i (q)_{\lambda_i}, \quad (3.19)$$

where $(q)_j$ stands for the falling factorial $q(q-1)\cdots(q-j+1)$.

Proof. Take $A = \mathbb{Z}[x]$ and $h = x^q$ in (3.3), and compare with (2.21). This yields

$$U_{n,d}((x^q), (x^q)', (x^q)'', \dots; \partial_x) = x^{(q-d)n} \sum_k \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d} x^k \partial_x^k.$$

Setting $x = 1$, this gives

$$U_{n,d}((q)_0, (q)_1, (q)_2, \dots; \partial_x) = \sum_k \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d} \partial_x^k,$$

from which the result follows. \square

The Stirling numbers $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d}$ present a symmetry in q and d , a fact that is not reflected by Formula (3.19). To make this symmetry explicit, it is convenient to set

$$S_{n,k}(q, d) = \left\{ \begin{matrix} n \\ nd-k \end{matrix} \right\}_{q,d} \text{ when } q \geq d, \quad \text{and } S_{n,k}(q, d) = \left\{ \begin{matrix} n \\ nq-k \end{matrix} \right\}_{q,d} \text{ when } q \leq d,$$

so that we have always

$$(x^q \partial_x^d)^n = \sum_k S_{n,k}(q, d) x^{qn-k} \partial_x^{dn-k}.$$

Then we obtain from Formula (3.19) the following nice formula, that seems to be new.

Corollary 3.20. *Let q, d, n, k be nonnegative integers. The generalized Stirling number $S_{n,k}(q, d)$ is given by*

$$S_{n,k}(q, d) = \sum_{a \in \mathcal{M}(n): |a|=k} \frac{\prod_i (d)_{r_i(a)} \cdot \prod_j (q)_{c_j(a)}}{\prod_{i,j} (a_{i,j}!)}, \quad (3.21)$$

where $\mathbf{M}(n)$ is the set of all lower triangular arrays $a = (a_{i,j})_{1 \leq j < i \leq n}$ of nonnegative integers, $|a|$ is the sum of the entries of a , and $c_j(a) = \sum_i a_{i,j}$ (column sum) and $r_i(a) = \sum_j a_{i,j}$ (row sum).

Proof. This is the immediate consequence of Proposition 3.18 and (3.14). \square

It is natural to ask for a combinatorial interpretation of the right-hand side of (3.19). We will see that we recover Varvak's description in [19, Cor. 3.2] of the generalized Stirling

numbers $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d}$.

We recall that a *partial bijection* from a set A to a set B is a bijection g from some subset $\text{Dom}(g)$ of A to some subset of B .

Corollary 3.22. *Let n, k, d, q be nonnegative integers, with $q \geq d$. The generalized Stirling number $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d}$ counts the partial bijections g from $[n] \times [d]$ to $[n] \times [q]$ with the property*

$$\forall (i, a, j, b) \in [n] \times [d] \times [n] \times [q], \quad g(i, a) = (j, b) \Rightarrow j < i,$$

and such that $\#\text{Dom}(g) = nd - k$.

Proof. The number $c_\lambda^{n,d}$ counts the partial maps $g : [n] \times [d] \rightarrow [n]$ that are in $\text{PD}_{n,d}$, and whose fibers have cardinalities equal to the parts of λ . For each such g , the product $\prod_i (q)_{\lambda_i}$ counts the maps ϕ from $\text{Dom}(g)$ to $[q]$ that are injective on each fiber of g . They are exactly the maps $\phi : \text{Dom}(g) \rightarrow [q]$ such that the map $(i, j) \in \text{Dom}(g) \mapsto (g(i, j), \phi(i, j)) \in [n] \times [q]$ is injective. The corollary follows from this. \square

The bijections

$$\begin{array}{ccc} [n] \times [d] & \longrightarrow & [nd] \\ (i, a) & \longmapsto & (i-1)d + a \end{array} \quad \text{and} \quad \begin{array}{ccc} [n] \times [q] & \longrightarrow & [nq] \\ (j, b) & \longmapsto & (j-1)q + b \end{array}$$

turn the graphs of the partial bijections from Corollary 3.22 into the rook placements of $nd - k$ rooks in a chessboard whose shape is the Young diagram of the partition whose parts are $q, 2q, \dots, (n-1)q, nq$, each with multiplicity d . Therefore, the Stirling number $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d}$ counts

these rook placements. This description of $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{q,d}$ agrees with [19, Cor. 3.2].

3.4 The coefficients $c_\lambda^{n,d}$ modulo a prime

Theorem 2.32 admits the following generalization.

Theorem 3.23. *Let p be a prime and assume that d is a power of p . Given a partition λ , let $d \cdot \lambda$ be the partition obtained by multiplying every part of λ by d . Then:*

- (a) $c_{d \cdot \lambda}^{n,d} \equiv c_\lambda^n \pmod{p}$;
- (b) $c_\mu^{n,d} \equiv 0 \pmod{p}$ if $\mu \neq d \cdot \lambda$, for all λ .

In particular, if both n and d are powers of p , then $c_\mu^{n,d} \equiv 0 \pmod{p}$, as long as $|\mu| \neq d(n-1)$ and $|\mu|$ is not a multiple of dp .

Proof. The cyclic group \mathbb{Z}_d acts on the set \mathbb{T}_n^d of d -tuples of increasing trees (T_1, \dots, T_d) by cyclic place permutation. The singular orbits are those of the form (T, T, \dots, T) , for $T \in \mathbb{T}_n$. If d is a power of a prime p , then all other orbits have size divisible by p . Thus,

$$U_{n,d} \equiv \sum_{T \in \mathbb{T}_n} \left(\prod_{i=1}^n y_{d \operatorname{ch}(i;T)} \right) t^{d \operatorname{ch}(0;T)} \pmod{p}. \quad (3.24)$$

We deduce (a) and (b) by equating coefficients in (3.24). The last statement of the theorem follows from the above and Theorem 2.32. \square

Remark 3.25. Observe that Theorem 3.23 above correctly computes the congruence classes modulo 3 of all the coefficients $c_\lambda^{3,3}$, knowing only that $c_\emptyset^3 = c_{(2)}^3 = c_{(1^2)}^3 = 1$ (see Example 3.15).

4 Normal ordering in formal differential operator rings

It is well known that, over the polynomial ring $\mathbb{F}[x]$ and assuming that the base field \mathbb{F} has characteristic 0, then the operators ∂_x and x (multiplication by x) generate the Weyl algebra $\mathbf{A}_1 = \mathbb{F}\langle x, y \mid [y, x] = 1 \rangle$. Hence, the normally ordered form of the operator $(h\partial^d)^n$, which we have been discussing, yields in particular known expressions for the normally ordered form of elements of this type in the Weyl algebra, with $\partial = \partial_x$ and $h \in \mathbb{F}[x]$ (see [5], [6] and [13]). In this section we will apply our results to the more general setting of formal differential operator rings, which include in particular the subalgebras \mathbf{A}_h of the Weyl algebra studied in [1, 2, 3] and defined below in (4.1).

Recall that a *formal differential operator ring* (or *skew polynomial ring*) is a ring, denoted $A[z; \partial]$, constructed from a ring A and a derivation ∂ of A , whose elements can be expressed as polynomials $\sum_{i=0}^n a_i z^i$ (with $n \geq 0$) in a new variable z and uniquely determined coefficients $a_i \in A$. Thus, as a left A -module, $A[z; \partial]$ can be identified with the polynomial ring $A[z]$. However, the variable z does not necessarily commute with the coefficients in A and multiplication in $A[z; \partial]$ is determined by the multiplication in A , associativity, the distributive law and the commutation relation $za = az + \partial(a)$, for all $a \in A$. In particular, the adjoint map $[z, -] : A \rightarrow A$ given by the commutator $[z, a] = za - az$ for $a \in A$ coincides with the derivation ∂ . See [10] or [14] for more details.

An important example of a formal differential operator ring is the first Weyl algebra $\mathbf{A}_1 = \mathbb{F}[x][y; \partial_x]$ over the field \mathbb{F} where, in case $\operatorname{char}(\mathbb{F}) = 0$, x can be identified with left multiplication by x on $\mathbb{F}[x]$ and y with ∂_x . In case $\operatorname{char}(\mathbb{F}) = p > 0$, then this correspondence does not give a faithful representation of \mathbf{A}_1 on $\mathbb{F}[x]$ because, as an operator, $\partial_x^p = 0$, whereas in \mathbf{A}_1 we have $y^p \neq 0$. Another example is given by the family of algebras $\mathbf{A}_h = \mathbb{F}[x][\hat{y}; \delta]$, studied in [1, 2, 3] and determined by the derivation $\delta = h\partial_x$, where $h \in \mathbb{F}[x]$. Then \mathbf{A}_h admits the presentation

$$\mathbf{A}_h = \mathbb{F}\langle x, \hat{y} \mid \hat{y}x - x\hat{y} = h \rangle. \quad (4.1)$$

By taking $h = 1$ we retrieve the Weyl algebra \mathbf{A}_1 and, for an arbitrary nonzero $h \in \mathbb{F}[x]$, we can view \mathbf{A}_h as the (unital) \mathbb{F} -subalgebra of \mathbf{A}_1 generated by x and yh , under the identification $\hat{y} = yh$. As h ranges over $\mathbb{F}[x]$, the family of algebras \mathbf{A}_h runs through all formal differential operator rings over $\mathbb{F}[x]$.

As before, let $R = \mathbb{Z}[y_i \mid i \geq 0]$ and $\Delta = \sum_{i \geq 0} y_{i+1} \partial_{y_i}$ be the derivation of R satisfying $\Delta(y_i) = y_{i+1}$ for all $i \geq 0$. The differential operator ring $R[z; \Delta]$ can also be seen as the

ring generated by the commuting variables y_i , for $i \in \mathbb{Z}$, and a new (non-central) variable z satisfying the commutation relations $zy_i = y_i z + y_{i+1}$, for all $i \geq 0$. In fact, it is natural to view $R[z; \Delta]$ as a homomorphic image of $R\langle t \rangle$ under the epimorphism $R\langle t \rangle \rightarrow R[z; \Delta]$ sending y_i to y_i and t to z . Under this map we can, and will, think of the polynomials $U_{n,d}$ as elements of $R[z; \Delta]$, substituting z for t . Fixing a base field \mathbb{F} (of arbitrary characteristic), we can also consider the corresponding \mathbb{F} -algebras $R_{\mathbb{F}} := \mathbb{F} \otimes_{\mathbb{Z}} R = \mathbb{F}[y_i \mid i \geq 0]$ and $\mathbb{F} \otimes_{\mathbb{Z}} R[z; \Delta] = R_{\mathbb{F}}[z; \Delta]$.

The main result of this section, presented below, shows that the polynomials $U_{n,d}$ give the normally ordered form of all powers of monomials over a formal differential operator ring.

Theorem 4.2. *Let A be a ring, h a central element of A and ∂ a derivation of A . Then, for all $n, d \geq 1$ we have the following normal ordering identity in the formal differential operator ring $A[z; \partial]$:*

$$(hz^d)^n = U_{n,d}|_{y_i=h^{[i]}, t=z} = \sum_{k=d}^{nd} \sum_{\lambda \vdash nd-k} c_{\lambda}^{n,d} h^{n-\ell(\lambda)} h^{[\lambda]} z^k, \quad (4.3)$$

where $h^{[\lambda]} = h^{[\lambda_1]} \dots h^{[\lambda_{\ell}]}$, for $\lambda = (\lambda_1, \dots, \lambda_{\ell})$.

Proof. We will first prove the special case of the Theorem for the ring R , the derivation Δ and the central element $h = y_0 \in R$ and then deduce the general case.

There is a representation $\rho : R[z; \Delta] \rightarrow \text{End}_{\mathbb{Z}}(R)$ sending z to Δ and $r \in R$ to the corresponding multiplication by r endomorphism. This is a ring endomorphism precisely because Δ is a derivation of R .

Then, by Theorem 3.2, we have the identity

$$\rho\left(\left(y_0 z^d\right)^n\right) = \left(y_0 \Delta^d\right)^n = U_{n,d}|_{t=\Delta} = \rho\left(U_{n,d}|_{t=z}\right)$$

in $\text{End}_{\mathbb{Z}}(R)$, as $y_0^{[i]} = y_i$. Hence, it remains to show that ρ is injective. Suppose $\rho\left(\sum_{i=0}^m P_i z^i\right) = 0$. Then $\sum_{i=0}^m P_i \Delta^i = 0$ as an endomorphism of R . By the uniqueness argument from the proof of Theorem 2.3, we conclude that $P_i = 0$ for all i and so $\sum_{i=0}^m P_i z^i = 0$, thus establishing the injectivity of ρ .

Now consider the general case of the Theorem, with A , h and ∂ as in the statement. There is a ring homomorphism $\phi : R[z; \Delta] \rightarrow A[z; \partial]$ defined on the generators by $y_i \mapsto h^{[i]}$ and $z \mapsto z$. This is well defined precisely because $h^{[i]}$ and $h^{[j]}$ commute, for all $i, j \geq 0$, and by the multiplication rule in a formal differential operator ring. Applying ϕ to the identity $(y_0 z^d)^n = U_{n,d}|_{t=z}$ yields precisely $(hz^d)^n = U_{n,d}|_{y_i=h^{[i]}, t=z}$. The right-hand side of (4.3) follows directly from (3.9). \square

Fix a field \mathbb{F} . For $0 \neq h \in \mathbb{F}[x]$, view the algebra \mathbf{A}_h defined in (4.1) as a subalgebra of the Weyl algebra \mathbf{A}_1 by identifying $\hat{y} \in \mathbf{A}_h$ with $yh \in \mathbf{A}_1$, as explained above.

Then \mathbf{A}_h is a free (left) $\mathbb{F}[x]$ -module and each of the following is a free basis (compare [3]):

$$\left(\hat{y}^n\right)_{n \geq 0}, \quad \left((hy)^n\right)_{n \geq 0} \quad \text{and} \quad \left(h^n y^n\right)_{n \geq 0}. \quad (4.4)$$

In fact, fixing $n \geq 0$, each of $(\hat{y}^k)_{0 \leq k \leq n}$, $((hy)^k)_{0 \leq k \leq n}$, and $(h^k y^k)_{0 \leq k \leq n}$ spans the same (left) $\mathbb{F}[x]$ -submodule of \mathbf{A}_h .

The Weyl algebra identity

$$(hy)^n = \sum_{k=1}^n \sum_{\lambda \vdash n-k} c_\lambda^n h^{n-k-\ell(\lambda)} h^{(\lambda)} h^k y^k, \quad (4.5)$$

which follows by direct application of Theorem 4.2, relates the second and third bases in (4.4). Furthermore, the relation $(hy)^{n+1} = h(yh)^n y = h(\hat{y}^n)y$ and the fact that A_1 is a domain yield

$$\hat{y}^n = \sum_{k=0}^n \sum_{\lambda \vdash n-k} c_\lambda^{n+1} h^{n-k-\ell(\lambda)} h^{(\lambda)} h^k y^k, \quad (4.6)$$

relating the first and the third bases in (4.4).

Example 4.7. Take $h = x$. The algebra A_x is the universal enveloping algebra of the non-abelian 2-dimensional Lie algebra. We have, by the binomial theorem:

$$\hat{y}^n = (yx)^n = (xy + 1)^n = \sum_{k=0}^n \binom{n}{k} (xy)^k.$$

On the other hand, we can also apply the formula (4.6), taking into account that $h^{(\lambda)} = 0$ unless $\lambda = 1^\ell$ is a partition with no parts greater than 1. Thus,

$$\hat{y}^n = \sum_{k=0}^n c_{1^{n-k}}^{n+1} x^k y^k = \sum_{k=0}^n \left\{ \begin{matrix} n+1 \\ k+1 \end{matrix} \right\} x^k y^k, \quad (4.8)$$

yielding the identity

$$\sum_{k=0}^n \binom{n}{k} (xy)^k = \sum_{k=0}^n \left\{ \begin{matrix} n+1 \\ k+1 \end{matrix} \right\} x^k y^k.$$

Now, using (4.5) we get, for $k \geq 1$,

$$(xy)^k = \sum_{j=1}^k c_{1^{k-j}}^k x^j y^j = \sum_{j=1}^k \left\{ \begin{matrix} k \\ j \end{matrix} \right\} x^j y^j.$$

Therefore, we deduce the identity

$$\begin{aligned} \hat{y}^n &= 1 + \sum_{k=1}^n \binom{n}{k} (xy)^k = 1 + \sum_{k=1}^n \sum_{j=1}^k \binom{n}{k} \left\{ \begin{matrix} k \\ j \end{matrix} \right\} x^j y^j \\ &= 1 + \sum_{j=1}^n \left(\sum_{k=j}^n \binom{n}{k} \left\{ \begin{matrix} k \\ j \end{matrix} \right\} \right) x^j y^j. \end{aligned}$$

Comparing with (4.8) we obtain the identity

$$\left\{ \begin{matrix} n+1 \\ k+1 \end{matrix} \right\} = \sum_{j=k}^n \binom{n}{j} \left\{ \begin{matrix} j \\ k \end{matrix} \right\}, \quad (4.9)$$

which appears in [12, Sec. 1.2.6 (52)].

In terms of the algebra A_x , equation (4.9) is a consequence of the change of basis relations between the $\mathbb{F}[x]$ -bases of A_x mentioned in (4.4). Indeed, the matrices

$$A_n = (a_{i,j})_{0 \leq i,j \leq n}, B_n = (b_{i,j})_{0 \leq i,j \leq n}, \text{ and } C_n = (c_{i,j})_{0 \leq i,j \leq n},$$

with

$$a_{i,j} = \binom{j}{i}, b_{i,j} = \left\{ \begin{matrix} j \\ i \end{matrix} \right\} \text{ and } c_{i,j} = \left\{ \begin{matrix} j+1 \\ i+1 \end{matrix} \right\},$$

are the transition matrices from $(\hat{y}^k)_{0 \leq k \leq n}$ to $((xy)^k)_{0 \leq k \leq n}$, from $((xy)^k)_{0 \leq k \leq n}$ to $(x^k y^k)_{0 \leq k \leq n}$ and from $(\hat{y}^k)_{0 \leq k \leq n}$ to $(x^k y^k)_{0 \leq k \leq n}$, respectively. Thus, (4.9) is just the statement that

$$\left(\left\{ \begin{matrix} j \\ i \end{matrix} \right\} \right)_{0 \leq i,j \leq n} \cdot \left(\binom{j}{i} \right)_{0 \leq i,j \leq n} = \left(\left\{ \begin{matrix} j+1 \\ i+1 \end{matrix} \right\} \right)_{0 \leq i,j \leq n}, \quad \text{for all } n \geq 0.$$

5 Final remarks

Similarities with the Faà Di Bruno Formulas. The polynomials U_n are defined by the simple formula: $U_n = (y_0(\Delta + \rho_t))^n(1)$, with $\Delta = \sum_i y_{i+1} \frac{d}{dy_i}$. The *Faà Di Bruno formulas* giving the derivatives $z^{(n)}$ of the composition $z = x \circ y$ of two functions, in terms of x and the derivatives of y , can be described similarly: for each n , there is a polynomial $F_n(y_1, y_2, y_3, \dots; t)$ such that

$$z^{(n)} = F_n(y', y'', y''', \dots; \partial_u)(x) \circ y.$$

The polynomials F_n are defined by $F_n = (\Delta + ty_1)^n(1)$.

Noncommutative setting. It is possible to drop the hypothesis “ h central”. We obtain that there is a unique polynomial $V_n(y_0, y_1, \dots; t)$ in non-commuting variables y_i , such that for any ring A , derivation ∂ of A and element $h \in A$, $(h\partial)^n = V_n(h, h^{[1]}, \dots; \partial)$. For instance:

$$\begin{aligned} V_0 &= 1, & V_1 &= y_0 t, & V_2 &= y_0 y_1 t + y_0^2 t^2, \\ V_3 &= y_0 y_1^2 t + y_0^2 y_2 t + 2y_0^2 y_1 t^2 + y_0 y_1 y_0 t^2 + y_0^3 t^3. \end{aligned}$$

Then many of the descriptions of the U_n are still valid for the V_n with simple changes:

- Inductive formula: $V_0 = 1, V_{n+1} = y_0(\Delta + \rho_t)V_n$, where now $\Delta = \sum_{i=1}^{\infty} D_i$, and D_i is the operator on words in y_0, y_1, y_2, \dots that replaces the letter y_j in position i by the next letter y_{j+1} .
- Umbral formula: V_n is obtained from $\prod_{i=0}^{n-1} (x_i + \dots + x_1 + x_0)$ by applying the linear map sending $x_n^{\alpha_n} x_{n-1}^{\alpha_{n-1}} \dots x_1^{\alpha_1} x_0^k$ to $y_{\alpha_n} y_{\alpha_{n-1}} \dots y_{\alpha_1} t^k$.
- Formula from increasing trees:

$$V_n = \sum_{T \in \mathbb{T}_n} y_{\text{ch}(n,T)} y_{\text{ch}(n-1,T)} \dots y_{\text{ch}(1,T)} t^{\text{ch}(0,T)}.$$

The polynomial V_n has an expansion of the form

$$V_n = \sum c_\alpha^n y_0 y_{\alpha_{n-1}} \cdots y_{\alpha_1} t^{n - \sum_i \alpha_i}$$

where the sum is carried over all sequences $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{n-1})$ of nonnegative integers with sum at most n . The coefficient c_α^n interprets as the number of subdiagonal partial maps $[n] \rightarrow [n]$ with α_1 elements mapped to 1, α_2 elements mapped to 2, \dots , α_{n-1} mapped to $n - 1$. The same straightforward extensions hold for the polynomials $U_{n,d}$ as well.

Acknowledgments. S. Lopes would like to thank G. Benkart and M. Ondrus for the discussions motivating the topics of this paper. He would also like to express his gratitude for the hospitality received at the Universidad de Sevilla during his visit at the onset of this collaboration.

References

- [1] G. Benkart, S. A. Lopes, and M. Ondrus. A parametric family of subalgebras of the Weyl algebra II. Irreducible modules. In *Recent developments in algebraic and combinatorial aspects of representation theory*, volume 602 of *Contemp. Math.*, pages 73–98. Amer. Math. Soc., Providence, RI, 2013.
- [2] G. Benkart, S. A. Lopes, and M. Ondrus. Derivations of a parametric family of subalgebras of the Weyl algebra. *J. Algebra*, 424:46–97, 2015.
- [3] G. Benkart, S. A. Lopes, and M. Ondrus. A parametric family of subalgebras of the Weyl algebra I. Structure and automorphisms. *Trans. Amer. Math. Soc.*, 367(3):1993–2021, 2015.
- [4] F. Bergeron and C. Reutenauer. Une interprétation combinatoire des puissances d’un opérateur différentiel linéaire. *Ann. Sci. Math. Québec*, 11(2):269–278, 1987.
- [5] P. Błasiak. *Combinatorics of boson normal ordering and some applications*. PhD thesis, Institute of Nuclear Physics of Polish Academy of Science, Krakow, and Université Pierre et Marie Curie, Paris, 2005.
- [6] P. Błasiak and P. Flajolet. Combinatorial models of creation-annihilation. *Sém. Lothar. Combin.*, 65:Art. B65c, 78, 2010/12.
- [7] A. Cayley. On the theory of the analytical forms called trees. *Philosophical Magazine*, 4(13):172–176, 1857.
- [8] L. Comtet. Une formule explicite pour les puissances successives de l’opérateur de dérivation de Lie. *C. R. Acad. Sci. Paris Sér. A-B*, 276:A165–A168, janvier 1973.
- [9] J. Engbers, D. Galvin, and J. Hilyard. Combinatorially interpreting generalized Stirling numbers. *European Journal of Combinatorics*, 43:32–54, 2015.
- [10] K. R. Goodearl and R. B. Warfield, Jr. *An introduction to noncommutative Noetherian rings*, volume 16 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1989.

- [11] F. Hivert, J.-C. Novelli, and J.-Y. Thibon. Multivariate generalizations of the Foata-Schützenberger equidistribution. In *Fourth Colloquium on Mathematics and Computer Science Algorithms, Trees, Combinatorics and Probabilities*, volume AG of *DMTCS Proceedings*, pages 289–299, Nancy, 2006. Assoc. Discrete Math. Theor. Comput. Sci.
- [12] D. E. Knuth. *The art of computer programming. Vol. 1*. Addison-Wesley, Reading, MA, 1997. Fundamental algorithms, Third edition.
- [13] T. Mansour and M. Schork. *Commutation relations, normal ordering, and Stirling numbers*. Discrete Mathematics and its Applications (Boca Raton). CRC Press, Boca Raton, FL, 2016.
- [14] J. C. McConnell and J. C. Robson. *Noncommutative Noetherian rings*, volume 30 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, revised edition, 2001. With the cooperation of L. W. Small.
- [15] M. Mohammad-Noori. Some remarks about the derivation operator and generalized Stirling numbers. *Ars Combin.*, 100:177–192, 2011.
- [16] OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, 2018.
- [17] H. F. Scherk. *De evolvenda functione ($yd.yd.yd\dots ydX/dx^n$) disquisitiones nonnullae analyticae*. PhD thesis, Berlin, 1823. Available online from Göttinger Digitalisierungszentrum (GDZ).
- [18] R. P. Stanley. *Enumerative combinatorics. Volume 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2012.
- [19] A. Varvak. Rook numbers and the normal ordering problem. *J. Combin. Theory Ser. A*, 112(2):292–307, 2005.